
2009

Using Calculus to Model the Growth of *L. Plantarum* Bacteria

Erin Carey
University of South Florida

Advisors:

Arcadii Grinshpan, Mathematics and Statistics
Scott Campbell, Chemical & Biomedical Engineering

Problem Suggested By: Scott Campbell

Follow this and additional works at: <https://digitalcommons.usf.edu/ujmm>



Part of the [Mathematics Commons](#)

UJMM is an open access journal, free to authors and readers, and relies on your support:

[Donate Now](#)

Recommended Citation

Carey, Erin (2009) "Using Calculus to Model the Growth of *L. Plantarum* Bacteria," *Undergraduate Journal of Mathematical Modeling: One + Two*: Vol. 1: Iss. 2, Article 2.

DOI: <http://dx.doi.org/10.5038/2326-3652.1.2.2>

Available at: <https://digitalcommons.usf.edu/ujmm/vol1/iss2/2>

Using Calculus to Model the Growth of *L. Plantarum* Bacteria

Abstract

Experimental data for the growth of *Lactobacillus plantarum* bacteria have been obtained over time, creating the need for mathematical means to model this data. We use the Gompertz model because it is a sigmoid function for a time series, where growth is slowest at the start and end of a time period. The Gompertz model is especially useful because it defines specific parameters that characterize the S-shaped curve. In addition, the Gompertz model uses relative growth, which is the logarithm of the given population compared to the initial population. This reflects the fact that bacteria grow exponentially. The important parameters that were found were the lag time and the asymptote.

Keywords


Lactobacillus plantarum, Gompertz mode, Asymptote

Creative Commons License



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/).

Motivation



Lactobacillus plantarum (Wessels)


Lactobacillus plantarum bacteria are aerotolerant bacteria that are used in silage, fermented food products and therapeutics. The high levels of this organism in human foods also make it an ideal candidate for the development of probiotics. Furthermore, this bacteria has been shown to reduce Irritable Bowel Syndrome in humans because of its ability to produce antimicrobial substances that help it survive in the gastrointestinal tract of humans (Niedzielin 1144). Thus, the bacteria have the potential to strongly impact the health of both humans and farm animals through the use of food biotechnology.

“Since bacteria reproduce exponentially, it is often useful to plot the logarithm of the relative population size ($y = \ln N/N_0$) against time. The three phases of the growth curve can be described by three parameters: The maximum specific growth rate is defined as the slope of the tangent at the inflection point, the lag time is defined as the x-axis intercept of this tangent, and the asymptote ($A = \lim_{t \rightarrow \infty} \ln(N/N_0)$) is the maximal value reached” (Zwietering 1876). This theory exemplifies the close relationship between math and science. The ability to model the growth of *Lactobacillus plantarum* using calculus provides a significant foundation for the development of future products that improve health through the use of this bacteria.

Problem Statement

Growth curve of *L plantarum* bacteria

The typical growth curve for bacteria is S-shaped as shown in the figure below. N/N_0 is the ratio of number of bacteria, N , to the initial population, N_0 , and t is the time. The curve shows an initial lag, followed by exponential growth and then a leveling off.



There are several parameters that are useful in characterizing the growth curve. One is the asymptote A , which represents the ultimate value of $\ln N/N_0$. Another is the lag time t_L . The lag time can be found as follows:

- (1) Obtain the time of the inflection point t_i .
- (2) Find the slope μ_m of the tangent line at the inflection point. This represents the maximum growth rate.
- (3) The lag time t_L is obtained by finding the x-axis intercept of this tangent line.

A number of models have been used to represent the growth curve. One is the Gompertz model:

$$\ln \frac{N}{N_0} = ae^{-e^{(b-ct)}},$$

where a , b and c are constants, which are obtained by fitting experimental data. We will use this model here. The table below provides experimental data for the growth rate of an aerotolerant bacteria, *L plantarum*, at 41.5°C . Our goal is to obtain the asymptote A and the lag time t_L . A reasonable approach is as follows:

- (a) Fit the Gompertz equation to the data to obtain a , b and c .
- (b) Show how the parameters A and t_L relate to a , b and c .
- (c) Obtain numerical values of A and t_L from step (b).

Growth data for *L. plantarum* at 41.5° C

t (hr)	$\ln N/N_0$
2	0.2
3	0.5
4	1.1
5	1.8
6	2.5
8	4.0
12	6.4
13	6.6

Mathematical Description and Solution Approach

We begin by finding the constants in the Gompertz model that best fit the given data. The Gompertz model is given by

$$f(t) = ae^{-e^{(b-ct)}}.$$

To determine the asymptote, we take the limit as t approaches infinity:

$$\text{Asymptote} = \lim_{t \rightarrow \infty} f(t) = \lim_{t \rightarrow \infty} ae^{-e^{(b-ct)}}.$$

Lag time is obtained by taking the first and second derivatives:

$$f'(t) = ae^{-e^{(b-ct)}} \cdot ce^{(b-ct)} = ace^{(b-ct)-e^{(b-ct)}};$$

$$f''(t) = ac^2 e^{(b-ct)-e^{(b-ct)}} (e^{(b-ct)} - 1).$$

To find the point of inflection, set the second derivative equal to zero. The x-axis intercept of the tangent line gives the lag time.


Discussion

The first step to finding the growth curve was to find the constants in the Gompertz model that best fit the curve given by the above data. Microsoft Excel was used by inputting the given data and a guess was made as to the three constants. The square in the error of the guesses was calculated. The solver application was then used by manipulating the constants until the sum of the squared errors (sos) was as small as possible. The table below tabulates the results of this procedure. The first two columns give the experimental data; the third column gives the value of $\ln(N/N_0)$ as determined by the Gompertz model

with constants a , b , and c given at the bottom of the table. The final column gives the squared difference. The row with “sos” gives the sum of the square errors. A graph of the curve with the constants given in the table is also shown, along with the given experimental data.

t	$\ln(N/N_0)$	Calc	sqerror
2	0.2	0.24169	0.001738
3	0.5	0.562623	0.003922
4	1.1	1.066686	0.00111
5	1.8	1.731284	0.004722
6	2.5	2.498084	3.67E-06
8	4	4.068512	0.004694
12	6.4	6.316004	0.007055
13	6.6	6.654936	0.003018
		sos	0.026262
a	7.832422		
b	1.803128		
c	0.278282		

$$f(x) = 7.8324 \cdot \exp\{-e^{(1.8031-0.27828t)}\}$$



The next step in analyzing the growth curve was to find the parameters indicated above, the asymptote A and the lag time t_L .

The horizontal asymptote can be found by determining the limit of the given function as t approaches infinity. (See Figure 1 in the appendix.)

$$A = \lim_{t \rightarrow \infty} 7.8324 \exp(-e^{(1.8031-0.27828t)}) = 7.8324.$$

It was also necessary to take the first and second derivatives of the equation in order to find the required parameters. The second derivative was necessary to find the point of

inflection. An inflection point is a point on a curve at which the curvature changes from convex to concave or vice versa. Thus, setting the second derivative equal to zero and solving for t gives the point of inflection. (See Figure 2)

$$0 = ac^2 e^{(b-ct)-e^{(b-ct)}} (e^{(b-ct)} - 1)$$

$$\Rightarrow t_i = \frac{b}{c} = 6.479.$$

The slope of the tangent line at the point of inflection, which represents the maximum growth rate, was found using the first derivative. (See Figure 3)

$$\text{Max growth rate: } f'(t_i) = 0.8018.$$

The lag time, t_L , is obtained by finding the x-axis intercept of the tangent line. The slope-intercept form of a line, $y = mx + b$, was used to solve for b at the point (6.4286, 2.8404). (See Figure 4)

$$2.8404 = (0.80184)6.4286 + b$$

$$b = -2.3146$$

$$0 = 0.80184 t_L - 2.3146$$

$$t_L = 2.8867$$

Conclusions and Recommendations

At 41.5 degrees Celsius, the growth of *Lactobacillus plantarum* bacteria was successfully modeled using the Gompertz curve. The asymptote, which represents the maximum relative population of the bacteria, was found to be 7.8324. The lag time was found to be 2.8867 hours. The maximum rate of growth of *Lactobacillus plantarum* bacteria was found to be 0.80184. This means that approximately eight-tenths of the relative population increased per hour during the fastest period of growth.

While the Gompertz model was a very realistic model of the growth of *Lactobacillus plantarum*, it was not perfect. The constants a , b and c that were found did give some error when compared to the experimental data. The sum of the squares of this error was approximately 0.26. A model that is able to reduce the sum of the squared error even further would notably improve further studies of this problem.

The objective of the paper was met and the Gompertz curve was able to model the experimental data. The results have direct implications for biology and medicine because they provide the fastest rate of growth of bacteria. In addition, the results showed the asymptote and the lag time before the bacteria reaches its maximum growth rate. These results are vital to the development of products using *Lactobacillus plantarum* bacteria and have improved the understanding of the relationship between bacterial growth and Calculus.

References

- Banks, Robert. Growth and Diffusion Phenomena: Mathematical Frameworks and Applications. Texts in applied mathematics, 14. Berlin: Springer-Verlag, 1994.
- Bielecki, Stanisław, J. Tramper, and Jacek Polak. Food Biotechnology. Progress in Biotechnology, 17. Amsterdam: Elsevier, 2000.
- Niedzielin K, Kordecki H, and Birkenfeld B. “A controlled, double-blind, randomized study on the efficacy of Lactobacillus Plantarum 299V in patients with irritable bowel syndrome.” Eur J Gastroenterol Hepatol. 2001 Oct;13(10):1143-7
- Wessels, Stephen. “The Microorganisms We Eat – Who decides Which Ones?” Human Health and Safety. www.teknologiportalen.dk/.../Foodstuffs/QPS.htm. 27 April 2009.
- Zwietering, M. H., Jongenburger, I., Rombouts, F. M., VAN 'T Riet, K. Modeling of the Bacterial Growth Curve. Appl. Environ. Microbiol. 1990 56: 1875-1881.

Appendix

Figure 1




Figure 2




Figure 3




Figure 4

