

8-2018

## Does the Mexican sugar-sweetened beverage tax have a signaling effect? ENSANUT 2016

Cristina Álvarez-Sánchez  
*Teachers College Columbia University*

Isobel Contento  
*Teachers College Columbia University*

Alejandra Jiménez-Aguilar  
*Mexican National Institute of Public Health*

Pamela Koch  
*Teachers College Columbia University*

Heewon L. Gray  
*University of South Florida, hlgray@usf.edu*

*See next page for additional authors*

Follow this and additional works at: [https://scholarcommons.usf.edu/cfh\\_facpub](https://scholarcommons.usf.edu/cfh_facpub)

---

### Scholar Commons Citation

Álvarez-Sánchez, Cristina; Contento, Isobel; Jiménez-Aguilar, Alejandra; Koch, Pamela; Gray, Heewon L.; Guerra, Laura A.; Rivera-Dommarco, Juan; Uribe-Carvajal, Rebeca; and Shamah-Levy, Teresa, "Does the Mexican sugar-sweetened beverage tax have a signaling effect? ENSANUT 2016" (2018). *Community and Family Health Faculty Publications*. 27.

[https://scholarcommons.usf.edu/cfh\\_facpub/27](https://scholarcommons.usf.edu/cfh_facpub/27)

This Article is brought to you for free and open access by the Community and Family Health at Scholar Commons. It has been accepted for inclusion in Community and Family Health Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact [scholarcommons@usf.edu](mailto:scholarcommons@usf.edu).

---

**Authors**

Cristina Álvarez-Sánchez, Isobel Contento, Alejandra Jiménez-Aguilar, Pamela Koch, Heewon L. Gray, Laura A. Guerra, Juan Rivera-Dommarco, Rebeca Uribe-Carvajal, and Teresa Shamah-Levy

RESEARCH ARTICLE

# Does the Mexican sugar-sweetened beverage tax have a signaling effect? ENSANUT 2016

Cristina Álvarez-Sánchez<sup>1\*</sup>, Isobel Contento<sup>1</sup>, Alejandra Jiménez-Aguilar<sup>2</sup>, Pamela Koch<sup>1</sup>, Heewon Lee Gray<sup>1a</sup>, Laura A. Guerra<sup>1</sup>, Juan Rivera-Dommarco<sup>2</sup>, Rebeca Uribe-Carvajal<sup>2</sup>, Teresa Shamah-Levy<sup>2</sup>

**1** Program in Nutrition, Department of Health and Behavior Studies, Teachers College Columbia University, New York, NY, United States of America, **2** Mexican National Institute of Public Health (INSP), Cuernavaca, Morelos, Mexico

<sup>a</sup> Current address: Department of Community and Family Health, College of Public Health, University of South Florida, Tampa, Florida, United States of America.

\* [cristina.alvarez@tc.columbia.edu](mailto:cristina.alvarez@tc.columbia.edu)



**OPEN ACCESS**

**Citation:** Álvarez-Sánchez C, Contento I, Jiménez-Aguilar A, Koch P, Gray HL, Guerra LA, et al. (2018) Does the Mexican sugar-sweetened beverage tax have a signaling effect? ENSANUT 2016. PLoS ONE 13(8): e0199337. <https://doi.org/10.1371/journal.pone.0199337>

**Editor:** Rodrigo Huerta-Quintanilla, Cinvestav-Merida, MEXICO

**Received:** February 20, 2018

**Accepted:** June 5, 2018

**Published:** August 22, 2018

**Copyright:** © 2018 Álvarez-Sánchez et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Data are freely available from the Instituto Nacional de Salud Publica (National Institute of Public Health) database: [http://ensanut.insp.mx/ensanut2016/descarga\\_bases.php](http://ensanut.insp.mx/ensanut2016/descarga_bases.php).

**Funding:** This article was possible due to an unrestricted research grant from Bloomberg Philanthropies (project CINY5/1305) and the Tisch Doctoral Scholar Fund, Program in Nutrition, Teachers College, Columbia University. The founding sponsors had no role in the design of the

## Abstract

### Objective

To evaluate the potential signaling effect of the Mexican tax on sugar-sweetened beverages (SSBs) by analyzing the association between awareness of and opinions about its effectiveness with current consumption of taxed SSBs and with a self-reported change in consumption of SSBs since the implementation of the tax. We also examined the association between psychosocial and environmental determinants of SSB consumption with current consumption of taxed SSBs and with a reported change in consumption of SSBs.

### Methods

Cross-sectional analyses of survey and food-frequency questionnaire data from the Mexican National Health and Nutrition Survey 2016. Participants were Mexican adults (20–59 years, N = 6,650). Logistic regression was used to evaluate the probability of a person reporting a decrease in SSB consumption, given their awareness of the tax, opinion about its effectiveness, psychosocial (SSB health-related beliefs, self-efficacy, and liking of SSBs) and environmental (availability of potable water) determinants. Multiple linear regression analysis was utilized to examine the association between the aforementioned factors and current consumption of taxed SSBs.

### Results

Compared with adults not aware, adults who were aware of the SSB tax were more likely (*OR* = 1.30) to report a decrease in SSB consumption (*p* = .012). In urban areas, adults aware of the tax drank a significantly lower amount of taxed SSBs (-15.7%; *p* = .023) than those not aware. Self-efficacy and liking of SSBs were significantly associated with a reported decrease in consumption and with current consumption (*p* < .001), while health beliefs and availability of potable water were not significantly associated with either reported change in SSB consumption or current consumption of taxed SSBs.

study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; and in the decision to publish the results.

**Competing interests:** The authors have declared that no competing interests exist.

## Conclusions

Implementation of an SSB tax accompanied by highly visible campaigns may further influence the impact of taxes on SSBs consumption. Future public health and nutrition education campaigns designed to increase knowledge and enhance motivation should be complemented by programs to assist individuals develop self-efficacy and self-regulation skills.

## Introduction

In Mexico, 73% of adults and 36% of children and adolescents (aged 2–19 years) have overweight or obesity [1]. Nearly 15% of adults are estimated to have type 2 diabetes—being the principal cause of mortality [2]. Frequent consumption of SSBs has been linked to an increased risk of a number of adverse health outcomes, including obesity [3–5], type 2 diabetes [6–9], coronary heart disease [10], dental caries [11], and tooth loss [12].

Consumption of SSBs, soda in particular, in Mexico is widespread [13] and has a strong cultural component [14]. Among many groups, soda drinking is a sign of conviviality, hospitality, and even of social status [14]. Currently, SSBs contribute about 69% of added sugars, 45% of total sugar intake, and 10% of total energy intake to the Mexican diet [13], more than three times the level recommended by the American Heart Association and approximately 3% of total energy intake [15, 16].

Due to this context, public health professionals advocated for the passage of an excise SSB tax and carried out strong and focused public awareness campaigns about the sugar content in SSBs, the health consequences of a high SSB consumption, and the rationale of a SSB tax; they also proposed that the SSB tax revenue be used to pay for purified water fountains in schools [17]. The debate around the Mexican SSB tax attracted a considerable amount of media attention and raised the profile of these issues among the public [17]. This culminated in the passing of a nationwide one-peso per liter (equivalent to a 10% increase) excise tax on SSBs [18], levied on manufacturers and effective from January 1, 2014—along with the implementation of other public health actions—as a public health measure to counteract obesity.

Studies conducted since the implementation of the tax indicate that SSB purchases by Mexican households declined by 7.6% on average in 2014 and 2015, even more than trends predicted [19, 20]. The decrease in purchases suggests a corresponding reduction in SSB consumption and therefore of caloric intake. The decrease in purchases and consumption may not be fully explained by the (economic) elastic nature of SSBs [19], but may be the result of an increased awareness of the detrimental health effects of SSBs. One study conducted prior to the implementation of the tax had already found declines in sales of SSBs in Mexico which, the authors hypothesize may have been due to “[a very] visible and well-funded media campaign linking [SSBs] with diabetes” [21].

Behavioral economics research suggests that the way in which taxes are presented or framed matters and could influence their impact [22]. SSB taxes are believed to provide consumers a behavioral rationale for changes (like nudges), in addition to traditional economic justification [23]. According to Adbukadirov [23], SSB taxes can increase the prominence of beverage choice to consumers through two mechanisms, first, “[SSB taxes] and the publicity that surround[s] [them may] trigger consumers to think about their health goals and to choose healthier drink[s],” and second, “attaching higher costs to unhealthy choices at the time of purchase may help undercut consumers’ myopia by countering the immediate benefits of enjoying a [SSB] with the immediate costs of the [SSB] tax.” There is emerging evidence supporting the hypothesis that “junk food” and SSB taxes imposed with public health goals in mind may

contribute to enhancing people's awareness about the negative health consequences of highly processed, less healthy foods and beverages [24, 25]. In economic theory, this is known as the "signaling effect" of tax policy, which poses that in addition to the tasks of raising public funds and correcting external effects, tax policies signal missing information to individuals about the effect of their consumption of the taxed product [26].

Understanding the impact of the Mexican SSB tax is further complicated by the fact that there were other initiatives undertaken during the same period, including the regulation of unhealthy food and beverages in schools [27, 28], the partial voluntary self-regulation of foods and beverages advertising directed at children [29], and the regulation of advertisement of foods and non-alcoholic beverages during children's television viewing time [30] that may have had an impact on SSB purchases over the same time period.

While it would be very difficult to evaluate the separate effects of the SSB tax and other simultaneous public health initiatives aimed at curbing SSB consumption, it is important to explore whether awareness of the SSB tax and opinion about its potential to reduce SSB intake, as well as psychosocial and environmental determinants of SSB consumption, are associated with current consumption of taxed SSBs, and with self-reported changes in consumption of SSBs since the SSB tax was passed. To our knowledge, no study has examined these associations after the implementation of a nation-wide SSB tax. Therefore, the current study addressed the following research questions:

1. Are Mexican adults aware of the SSB tax? What is their opinion about the effectiveness of the SSB tax in decreasing purchases of taxed SSB? Do awareness of and opinion about the SSB tax differ by socio-demographic characteristics?
2. Are awareness of and opinion about the effectiveness of the SSB tax, and psychosocial and environmental factors of SSB consumption, associated with a reported decrease in SSB consumption?
3. Are awareness and opinion about the effectiveness of the SSB tax and psychosocial and environmental factors of SSB consumption associated with current consumption of taxed SSBs?

Overall, we hypothesized that a higher percentage of adults living in Mexico City and of higher socio-economic status (SES) would be aware of the tax, and that those who were aware and expressed a positive opinion about the effectiveness of the SSB tax in reducing purchases of SSBs would, in effect, drink less SSBs and/or report a decrease in SSB consumption, compared to those who were not aware and/or expressed a negative opinion about the effectiveness of the SSB tax. These findings would be useful for health advocates and policy makers when considering passing a SSB tax.

## Materials and methods

### Population and study design

The current study is an analysis of data collected with the 2016 Mexican National Health and Nutrition Survey (ENSANUT by its Spanish acronym). The ENSANUT is a nationally representative probabilistic multistage stratified cluster survey constructed with sufficient sampling power to make distinctions between urban ( $\geq 2,500$  inhabitants) and rural ( $< 2,500$  inhabitants) areas, and among four geographic regions (categorized as North, Central, Mexico City, and South). Sampling weights are used to estimate nationally representative values. (A detailed description of the sampling procedures and survey methodology has been described elsewhere [31].) The ENSANUT 2016 was approved by the Research, Ethics and Biosafety Committees at

the National Institute of Public Health. Written informed consent was obtained from all study participants. Trained personnel administered all questionnaires and measures face-to-face.

We primarily used data from the *Perception of Obesity, Physical Activity and Diet Questionnaire* (POCAA-Q, by its Spanish acronym) [32], which had been applied to a random subsample of 6,550 adults aged 20–59 years. A description about the development and validation of the POCAA-Q can be found elsewhere [33]. The aim of the POCAA-Q was to explore adult Mexicans' perceptions of their dietary and physical activity habits as well and knowledge about causes and consequences of obesity. The POCAA-Q also explores the population's awareness of and opinion about the effectiveness of governmental legislation (including the SSB tax) to prevent and control obesity. The specific questions from the POCAA-Q that were used in this study are described below in measures. Additional data were obtained from other ENSANUT 2016 files: the semi-quantitative food frequency questionnaire (SFFQ), and the demographic file (i.e. demographic, socio-economic characteristics and sample weights).

## Measures

**Awareness of the tax and opinion about the effectiveness of the tax.** The variables awareness of the SSB and opinion about the effectiveness of the SSB tax come from the POCAA-Q. Their operational definitions can be found in [S1 Table](#).

**Self-perception of change in consumption of SSBs.** The variable self-perception of change in consumption of SSBs in the two years prior to the survey is a proxy for the time when the SSB tax was implemented. It also comes from the POCAA-Q. Its description can be found in [S1 Table](#).

**Consumption of taxed SSBs.** Beverage consumption was assessed using a SFFQ [34] which was validated for use with Mexican adolescents and adults [35]. The questionnaire includes 140 food items including a variety of sugar-sweetened, unsweetened, or artificially sweetened beverages. To assess consumption of each food item, reported frequency of consumption was converted into grams. To calculate consumption of taxed industrialized SSBs, we summed quantities (g/person/day) of all SSBs subject to the excise tax included in the SFFQ: regular carbonated SSBs, industrialized flavored waters with added sugar, and industrialized fruit nectars with added sugar. Sweetened energy and sports beverages are subject to the SSB tax, but they are not captured by the FFQ; thus they were not contemplated in this study. The data from the SFFQ had already been cleaned and processed following the steps detailed by Ramirez Silva and colleagues [36]; we excluded an additional three individuals with extreme observations (more than 3 SDs the log taxed SSB consumption).

**Psychosocial and environmental determinants of SSB consumption.** The selection of psychosocial and environmental variables from the POCAA-Q was informed by the health literature and includes SSB health-related beliefs [37] (measured with four questions about beliefs on whether SSB consumption contributes to high blood pressure, obesity, diabetes, and dental caries), self-efficacy to decreased consumption of SSBs [38, 39] (measured with one question), degree of liking of SSBs [40] (measured with one question; with a higher score indicating lower preference of SSBs), and availability of free/low-cost water [41] (measured with one question). [S1 Table](#) presents the definitions and rationale for choice of each variable.

For SSB health-related beliefs, a composite scale/measure was constructed based on the four health beliefs questions, with one additional point for a “yes” response regarding the belief about each condition. The scale ranged from 0 (reporting “no” to all four health beliefs questions) to 4 (reported “yes” to all four questions), with a higher score indicating an incremental agreement with the statements regarding the health damage of SSBs. (Cronbach's alpha for the scale was 0.844.)

**Covariates.** Socio-demographic variables included were sex (men and women), age (continuous variable), and a validated socio-economic status index [33] (with terciles derived from principal components analysis of eight variables: household building materials; number of bedrooms; basic services infrastructure; ownership of a car, television, radio, and refrigerator). Body mass index (BMI) was calculated as the weight in kilograms divided by the square height in meters ( $\text{kg}/\text{m}^2$ ) [42]. Height and weight were measured using standardized procedures [43, 44]. Values between 10 and  $58 \text{ kg}/\text{m}^2$  were considered as valid data [1]. We used the WHO BMI classification: underweight:  $< 18.5$ , normal weight:  $18.5\text{--}24.9$ , overweight:  $25.0\text{--}29.9$ , and obesity:  $\geq 30.0$  [42].

We also included self-reporting of diabetes diagnosis, in response to the question: “¿Algún médico le ha dicho que tiene diabetes o alta el azúcar en la sangre?” (Has a doctor told you that you have diabetes or high blood sugar?).

## Statistical analyses

**Relationship between categorical variables.** Chi-square tests were run to examine the relationship between categorical variables, such as awareness of the SSB tax and socio-demographic variables; with a  $p$ -value  $< .05$  as the cutoff point for statistical significance. Differences between subcategories of socio-demographic variables (e.g. sex: male/female, location: urban/rural) were considered to be statistically significant if their 95% confidence intervals (CIs) did not overlap; we used this approach recognizing its limitation, namely, that when the CIs of two statistics do not overlap, they are necessarily significantly different, but they could be significantly different even if their CIs overlap [45].

**Binary logistic regression.** A binary logistic regression was conducted to evaluate the probability that a given person would report a decrease in their SSB consumption in the two years prior, given their: (1) awareness of the SSB tax, (2) opinion about the effectiveness of the SSB tax in reducing purchases of SSBs, (3) health beliefs scale (psychosocial determinant), (4) self-efficacy (psychosocial determinant), (5) liking of SSBs, and (6) availability of potable water for free or at a low cost. We constructed the binary outcome variable (consumption of SSBs decreased and consumption did not decrease) from the three-category perception of change in the SSB consumption variable by keeping the “consumption decreased” category and combining the “consumption stayed the same” and “consumption increased” categories. Covariates sex, BMI, SES, geographic region, urban-rural location, and diabetes diagnosis were entered as categorical variables; age was entered as a continuous variable. As indicated by Bursac and colleagues, selection of variables for the logistic regression was conducted purposefully using (manual) backward elimination [46]. We started with a full model with the six variables and seven covariates, tested for interactions, and subsequently eliminated insignificant predictors at four different steps to arrive at a parsimonious final model [46]. The final logistic regression model includes three variables: awareness of the tax, self-efficacy, and liking of SSBs; and two covariates: age and diabetes diagnosis.

Results are expressed as adjusted *Odds Ratio* (OR) and their corresponding 95% CIs. Results were considered to be statistically significant if the 95% CI excluded the value of 1 [47].

**Multiple linear regression.** Multiple linear regression analysis was utilized to examine the association between six variables: (1) awareness of the tax, (2) opinion about the effectiveness of the SSB tax, (3) health beliefs scale (psychosocial determinant), (4) liking of SSBs (psychosocial determinant), (5) self-efficacy (psychosocial determinant) and (6) availability of potable water for free or at a low cost; with current consumption of taxed SSBs ( $\log \text{ g}/\text{d}$ ), after controlling for seven covariates (sex, age, BMI, diabetes diagnosis, SES, urban-rural location, and region). The outcome variable (consumption of taxed SSBs ( $\log \text{ g}/\text{person}/\text{day}$ )) was strongly,

positively skewed, and thus log-transformed. For the purpose of improving interpretability of the beta estimates, we calculated the percentage change for each estimate in the outcome variable per one unit change in the independent variable while all other variables in the model were held constant; we used the equation: % change in consumption of taxed SSBs =  $(e^{\beta} - 1) * 100$  [48]. We started with a full model with the six variables and seven covariates, tested for interactions, and subsequently performed a manual backward elimination of insignificant predictors, at three different steps, to arrive at a parsimonious final model where all variables included were significant [47]. The final multiple regression model includes three variables: awareness of the tax, self-efficacy, and liking of SSBs; and the seven covariates: urban-rural location, sex, age, region, diabetes diagnosis, and BMI. Multiple regression results are expressed as: regression coefficients, percent changes in consumption of taxed SSBs in relation to changes in independent variables, and standard errors. The (adjusted) R square is presented to indicate the estimated amount of explained variance. Results were considered significant at  $p < .05$  [47].

Lastly, we estimated mean taxed SSB consumption for the total sample and by the theoretical variables of interest (which include awareness and opinion about the tax, as well as psychosocial and environmental factors). All statistical analyses were performed with IBM SPSS, version 24.0. Calculations were weighted by expansion factors and adjusted for the complex sampling survey design using the SPSS command for complex surveys. Data for the  $\chi^2$  tests met the assumptions of sample size and independence of observations. The binary logistic regression model was checked for linearity and multicollinearity. The multiple regression model was checked for multicollinearity, linearity and for normality, homoscedasticity, and independence of residuals.

## Results

Study population characteristics are presented in [Table 1](#).

### Awareness and opinion about the SSB tax

At national level, 65.2% of the respondents reported being aware of the existence of the SSB tax, however, only 20.3% indicated that they thought the SSB tax was helping to decrease the purchase of the SSBs ([Fig 1](#)); the majority of those who reported being aware of the tax (53.1%) indicated that they thought it was not reducing purchases of SSBs. The percentage of respondents who thought that the SSB tax was reducing purchases of SSBs was significantly greater among individuals aware of the SSB tax (12.1%) than among those not aware of the SSB tax (8.2%).

In the analyses stratified by socio-demographic characteristics ([Table 2](#)), chi-squared tests of independence revealed statistical significant differences between awareness and opinion about the SSB tax and sex ( $\chi^2 = 30.366$ ,  $p = 0.019$ ), SES ( $\chi^2 = 306.593$ ,  $p < .001$ ), area ( $\chi^2 = 87.617$ ,  $p < .001$ ), region ( $\chi^2 = 113.116$ ,  $p = 0.002$ ), and age ( $\chi^2 = 178.097$ ,  $p < .001$ ). The percentages of respondents who were aware of the SSB tax were significantly higher among people of high SES (74.4%), and living in Mexico City (76.6%) and in urban areas (67.5%).

### Factors associated with a self-reported decrease in SSB consumption

The final (parsimonious) logistic regression model only included statistically significant (or nearly significant) predictors and covariates. The final model was significant,  $F(10, 278) = 15.110$ ,  $p < .001$ , and explained 9.4% (Nagelkerke pseudo- $R^2$ ) of the variance in change ([Table 3](#)).



Table 1. Socio-demographic characteristics.

		Un-weighted <i>n</i>	Weighted <i>n</i> (in millions)	Weighted percentages (%)
Sex				
	Male	2,152	25.7	47.8
	Female	4,498	28.5	52.2
SES				
	Low	2,276	12.4	20.8
	Medium	2,266	17.3	29.1
	High	2,108	29.8	50.1
Location				
	Urban	3,323	46.0	77.3
	Rural	3,327	13.5	22.7
Region				
	North	1,434	12.5	21.0
	Center	2,171	19.6	33.0
	Mexico City	763	10.4	17.6
	South	2,282	17.0	28.4
Age ( <i>mean ± SEM</i> )		38.6 ± 0.1	36.6 ± 0.3	NA
	20–29	1,647	19.1	32.1
	30–39	1,936	17.2	28.9
	40–49	1,694	13.2	22.2
	50–59	1,373	10.0	16.8
BMI ( <i>mean ± SEM</i> )		28.7 ± 0.1	28.5 ± 0.1	NA
	Normal weight	1,582	14.6	26.3
	Overweight	2,423	21.9	39.5
	Obesity	2,316	19.0	34.2
Total		6,650	59.5	100

Notes.

SES, socio-economic status; SEM, standard error of the mean; BMI, body mass index; NA; non-applicable.

Data are from the ENSANUT 2016: Mexican adults (20–59 years old), *n* = 6,650.

<https://doi.org/10.1371/journal.pone.0199337.t001>

Among the six independent variables, three were statistically significant: awareness of the SSB tax, self-efficacy, and liking of SSBs. Respondents who were aware of the SSB tax were 30% more likely to report a decrease in consumption of SSBs in the two years prior. High self-efficacy and low liking of SSBs were also individually associated with a reported decrease in SSBs (*OR* = 1.68) and (*OR* = 4.29), respectively.

**Factors associated with current consumption of taxed SSBs**

The final (parsimonious) model significantly predicted consumption of taxed SSBs, *F*(18, 262) = 32.51, *p* < .001, with *R*<sup>2</sup> = 21.1% (Table 4).

Self-efficacy and liking of SSBs added significantly to the prediction (*p* < .001). Respondents who were very confident or confident in limiting their consumption of SSBs to <1 glass/week consumed less taxed SSBs (53.2% and 36.9%, respectively) than those who did not feel confident. Individuals who dislike SSBs consumed less (42.3%) than those who like them. A significant interaction between urban-rural location and awareness of the tax was found (*p* = .017), indicating that location of residence moderated the relationship between awareness of the SSB tax on consumption of taxed SSB beverages. In particular, only individuals living in urban areas a significant difference between those aware and not aware of the SSB tax was




Fig 1. Awareness of the SSB tax and opinion about whether it was reducing the purchases of SSBs (n = 6,321).

<https://doi.org/10.1371/journal.pone.0199337.g001>

observed (a 15.7% decrease in taxed SSB consumption among those aware, compared to those who were not aware;  $p = .023$ ).

## Discussion

### Awareness of the SSB tax and opinion about the impact of the SSB tax

Results show slightly more than sixty-five percent (65.2%) of adults reported being aware of the SSB tax. And while there are no equivalent country-level data, a study conducted about a year after the passing of an excise tax on SSBs in Berkley, California found a similar figure—68% of people interviewed knew that the tax had been on their city’s ballot [25]. It is possible that some respondents may have indicated they recalled the tax because it was the socially desirable response, but it’s worth noting that the tax was passed in the midst of very visible and controversial campaigns from proponents and opponents of the fiscal measure [17, 49]. According to Donaldson [17], the media campaign put forth by health advocates “generated over 1,000 media articles in the five-month period leading up to the vote on the tax. . . reaching the public as well as key decision-makers”. According to a Pan American Health Organization report [49] “the entire industry involved presented a united front against the tax, with very significant activism in the media—television, radio, press and advertising campaigns”.

In the current study, the largest percentage of respondents aware of the SSB tax was again found among people living in Mexico City and in urban areas, and of high SES. This finding is congruent with our hypothesis and can be explained by the fact that Mexico City was the stage of most of the advocacy and opposition campaigns, and that people of high SES living in urban

Table 2. Awareness of the SSB tax and opinion about whether it was reducing purchases of SSBs, stratified by socio-demographic characteristics\*.

	Total		Aware of the SSB tax				NOT aware of the SSB tax			
			Believe the SSBs tax is reducing purchases of SSBs		DO NOT believe the SSB tax is reducing purchases of SSBs		Believe the SSBs tax is reducing purchases of SSBs		DO NOT believe the SSB tax is reducing purchases of SSBs	
			Unweighted n	Weighted (in MM) n	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)	% (95% CI)
National	6,321	6.9	12.1 (10.7–13.7)	53.1 (50.3–55.9)	8.2 (7.0–9.5)	26.6 (24.5–28.8)				
Sex †										
Male	2,054	27.6	12.0 (9.8–14.6)	55.9 (51.9–59.9)	6.6 (5.2–8.3)	25.5 (22.2–29.0)				
Female	4,267	29.8	12.2 (10.7–13.8)	50.5 (47.5–53.5)	9.7 (8.1–11.5)	27.7 (25.3–30.1)				
SES †										
Low <sup>a</sup>	2,088	11.7	14.1 (12.2–16.2)	39.0 (35.2–42.9)	12.6 (10.1–15.6)	34.3 (30.8–37.9)				
Medium <sup>b</sup>	2,175	16.8	12.7 (10.4–15.3)	45.3 (41.4–49.2)	9.0 (7.2–11.2)	33.0 (29.1–37.2)				
High <sup>c</sup>	2,058	28.9	11.0 (8.8–13.5)	63.4 <sup>a,b</sup> (59.0–67.5)	5.9 <sup>a</sup> (4.3–8.0)	19.8 <sup>a,b</sup> (16.9–23.0)				
Area †										
Urban <sup>a</sup>	3,213	44.6	11.4 (9.7–13.4)	56.1 (52.7–59.4)	7.1 (5.8–8.7)	25.3 (22.8–28.0)				
Rural	3,108	12.8	14.3 (12.7–16.1)	42.8 <sup>a</sup> (39.4–46.2)	11.9 <sup>a</sup> (9.9–14.2)	31.0 (27.8–34.4)				
Region <sup>*</sup>										
North <sup>a</sup>	1,372	12.2	11.6 (8.1–16.4)	50.7 (43.4–58.0)	9.2 (6.1–13.7)	28.5 (23.6–34.0)				
Center <sup>b</sup>	2,083	18.7	12.9 (10.6–15.8)	53.4 (47.7–59.1)	6.9 (5.3–8.8)	26.8 (22.8–31.1)				
Mexico City <sup>c</sup>	751	10.4	12.8 (9.6–16.8)	63.8 <sup>b,c</sup> (58.9–68.4)	5.8 (3.8–8.9)	17.6 <sup>a,b,c</sup> (13.5–22.6)				
South <sup>d</sup>	2,115	16.1	11.0 (9.2–13.2)	47.7 (44.4–51.1)	10.5 (8.3–13.1)	30.8 (28.2–33.6)				
Age †										
20–29 <sup>a</sup>	1,575	18.6	10.6 (8.6–13.0)	44.1 <sup>b,c,d</sup> (39.8–48.4)	10.4 (7.7–13.9)	34.9 <sup>b,c,d</sup> (31.0–39.1)				
30–39 <sup>b</sup>	1,837	16.7	10.5 (8.2–13.3)	56.4 (50.6–62.0)	7.6 (5.9–9.7)	25.5 (21.5–30.0)				
40–49 <sup>c</sup>	1,621	12.8	14.9 (11.7–18.9)	58.2 (54.0–62.3)	7.3 (5.5–9.7)	19.5 (15.9–23.6)				
50–59 <sup>d</sup>	1,288	9.3	14.1 (11.5–17.1)	58.2 (53.5–62.7)	6.0 (4.6–7.8)	21.8 (18.2–25.8)				
BMI										
Normal weight	1,489	14.2	11.6 (9.0–14.7)	52.1 (47.3–56.9)	8.6 (6.6–11.1)	27.8 (23.5–32.6)				
Overweight	2,304	20.7	12.8 (10.7–15.1)	53.8 (49.6–57.8)	7.7 (6.9–9.7)	25.8 (22.7–29.2)				
Obesity	2,220	18.6	11.5 (9.4–13.9)	54.0 (48.7–59.2)	8.5 (6.1–11.7)	26.1 (22.5–30.0)				

Notes.

SSB, sugar-sweetened beverages; MM, millions; SES, socio-economic status; BMI, body mass index.

Data are from the ENSANUT 2016: Mexican adults (20–59 years old).

\* Values are percentages and 95% CIs. Percentages across a row sum up to 100.

† p < .05 based on  $\chi^2$  test across categories. For each socio-demographic variable, different subscripts down a column (a, b, c, d) indicate statistically significant differences based on the 95% CIs (i.e. the CIs do not overlap).

<https://doi.org/10.1371/journal.pone.0199337.t002>

areas might have had increased exposure and attentiveness to all the health messaging (print media, television, radio debates, etc.) about SSBs that went with the tax.

At national level, only 20.3% of respondents (combining those aware and unaware of the tax) thought that the fiscal measure was helping to decrease the purchase of SSBs. This finding may be explained by several potential factors. First, respondents reporting that the SSB tax was not reducing their purchases of SSBs could have made their judgment based on negative reports, articles and/or debates about the impact of the SSB tax. Second, they could have based their response on judgments of their own behavior and/or that of their peers. Third, in the past few decades, consumption of SSBs became deeply rooted in Mexican dietary habits [50], thus, in spite of an average 7.6% decrease in purchases of SSBs over the first two years [20], the perception might be that SSBs are still ubiquitous. Further qualitative/mixed methods studies

**Table 3. Final, parsimonious, model for self-reporting a decrease in consumption of SSBs since the year the SSB tax was implemented, obtained by binary logistic regression\* †.**

		Consumption decreased vs consumption did not decrease since the year the SSB tax was implemented		
		Odds Ratio	95% CI	P-value
Awareness of the SSB tax				
	Aware	1.30	1.06, 1.59	.012
	Not aware	Reference		
Self-efficacy				< .001
	Very confident	1.68‡	1.15, 2.46	
	Confident	1.12	0.77, 1.64	
	Slightly confident	0.88	0.58, 1.35	
	Not confident	Reference		
Liking of SSBs				< .001
	Completely disagree	4.29‡	1.90, 9.70	
	Disagree	3.33‡	2.19, 5.01	
	Agree	1.68‡	1.23, 2.30	
	Completely agree	Reference		
Age		1.00	1.00, 1.02	.056
Diabetes				< .001
	Yes	1.77‡	1.33, 2.35	
	Yes—gestational	1.24	0.25, 6.10	
	No	Reference		
$R^2$ Cox and Snell = 0.063				
$R^2$ Nagelkerke = 0.094				

Notes.

SSB, sugar-sweetened beverages.

Data are from the ENSANUT 2016: Mexican adults (20–59 years old),  $n = 6,349$ .

\* The full binary logistic regression model included seven variables (awareness of the tax, opinion about the effectiveness of the tax, self-efficacy, liking of SSBs, health-beliefs, and availability of free/low-cost potable water) and covariates (age, diabetes diagnosis, sex, socio-economic status, geographic region, and area). The reference category included adults who reported that their SSB consumption in the two years prior had decreased. The statistics presented are from the parsimonious model, which includes only statistically significant (or nearly significant) predictors and covariates.

† Values are Odd Ratios, 95% CIs, and  $P$ -values of variable effect in overall model based on Wald F test.

‡ Significant findings of subcategories based on the 95% CI (i.e. the CI does not include 1).

<https://doi.org/10.1371/journal.pone.0199337.t003>

are warranted to explore the reasons why most Mexican adults think the SSB tax is not working.

### Factors associated with a self-reported decrease in SSB consumption and with current consumption of taxed SSBs

Results of the binary logistic regression analysis showed that factors associated with a self-reported decrease in SSB consumption in the 2 years prior are: awareness of the SSB tax, high self-efficacy, and not liking of SSBs. Results of the multiple regression analysis showed that factors significantly associated with current consumption of taxed SSBs (log g/person/day) are: self-efficacy, liking of SSBs; and the interaction between awareness of the SSB tax and urban-rural area. In none of the models were opinion about the impact of the tax, health beliefs, and drinkable water availability significant.

**Table 4. Final, parsimonious, model of factors associated with current consumption of taxed SSBs, obtained by multiple regression \* † ‡.**

	$\beta$	% change in consumption of taxed SSBs §	SE of the $\beta$	P-value (F test)	P-value (t tests)
Awareness of the SSB tax AND Area					
Urban (aware vs not aware)	-0.17	-15.72	0.08		.023
Rural (aware vs not aware)	0.08	8.76	0.08		.250
Awareness of the SSB tax	Reference			.394	
Self-efficacy					
Very confident	-0.76	-53.23	0.212		< .001
Confident	-0.46	-36.87	0.11		< .001
Slightly confident	-0.23	-20.55	0.13		.076
Not confident	Reference				
Liking of SSBs					
Completely disagree	-0.55	-42.31	0.28		.046
Disagree	-0.73	-51.81	0.13		< .001
Agree	-0.34	-28.82	0.10		.001
Completely agree	Reference				
Urban-rural location	0.37	44.20	.090	< .001	
Sex (female vs male)	-0.60	-45.12	0.05	< .001	
Age	-0.02	-1.98	<0.01	< .001	
Region					
North	0.42	50.20	0.11		< .001
Center	0.20	22.14	0.08		.008
Mexico City	-0.01	-1.00	0.09		.891
South	Reference				
Diabetes					
Yes	-0.28	-24.42	0.09		.003
Yes—gestational	1.04	182.92	0.96		.141
No	Reference				
BMI					
Obesity	0.31	36.34	0.09		< .000
Overweight	0.10	10.52	0.09		.285
Normal weight	Reference				
$R^2 = 0.211$					

Notes.  
 SSBs, sugar-sweetened beverages;  $\beta$ , regression coefficient; SE, standard error;  $R^2$  proportion variance explained.  
 Data are from the ENSANUT 2016: Mexican adults (20–59 years old),  $n = 4,624$   
 \* The outcome variable, taxed SSB consumption (log g/person/day), was created combining the following variables: regular soda, industrialized nectars, industrialized fruit waters.  
 † The full multiple regression included seven variables (awareness of the tax, opinion about the effectiveness of the tax, self-efficacy, liking of SSBs, health-beliefs, and availability of free/low-cost potable water) and covariates (age, BMI, diabetes diagnosis, sex, socio-economic status, geographic region, and area). The statistics presented are from the final (parsimonious) model which includes only statistically significant variables.  
 ‡ Values are  $\beta$  coefficients, % change in consumption of taxed SSBs (log g/person SSBs\* (log g/person/day), SEs of the  $\beta$ s,  $p$ -values of each coefficient estimate in the Wald F test, and  $p$ -values of the t tests for each of the coefficients of the sub-categories within each factor. Lower  $\beta$ s indicate expectation of less SSB consumption/lower score on unfavorable behavior.  
 § % change in consumption of taxed SSBs (log g/person/day) was calculated as: % change in consumption of taxed SSBs =  $(e^\beta - 1) * 100$ .

<https://doi.org/10.1371/journal.pone.0199337.t004>

Individuals aware of the SSB tax were 23% more likely to report a decrease in SSB consumption that those who were not aware. In addition, those aware of the tax, and living in urban

areas, consumed 16.6% less taxed SSBs than people not aware. These findings suggest that the SSB tax and the publicity that surrounded it may have had a “signaling effect” thereby making people more conscious about their beverage choices. Our findings agree with the results of two prior studies that examined the impact of taxes on unhealthy food [24, 25]. An impact assessment of a tax on unhealthy non-staple food products passed in Hungary, found that 22–38% of consumers (depending on food categories) had reduced their intake of taxed products due to an increased health consciousness [24]. In the US city of Berkeley, a stronger than expected reduction in consumption of SSBs after the passing of SSB tax was partly attributed to the pro-tax media campaign, which, according to the study authors, may have shifted social norms and increased overall health consciousness [25]. Nevertheless, causality between awareness of a SSB tax and consumption of SSBs cannot be established, as people with a priori favorable attitudes and behaviors might have been more likely to pay attention to campaigns and debates.

*Opinion about whether the SSB tax was reducing SSB purchases* was not a significant predictor of reported change in SSB consumption since the year the tax was implemented. Two plausible explanations for this finding are that even if there has been a considerable decrease in purchases of SSBs—7.6% on average over the first 2 years since the introduction of the tax [20]—the change in participants’ purchases (in number of units or volume) of taxed beverages may have been small and not clearly noticeable to them, or perhaps there has not been a large enough critical mass who have changed their behaviors so as to have precipitated a change in a social norm that is so deeply entrenched [51, 52]. In this regard, it should be noted that when the ENSANUT 2016 was conducted the tax had already lost a small percent of its value because of inflation—the tax was adjusted in January 2018 after it rose 10 percent inflation from the time of implementation.

*Liking of SSBs* was a strong significant predictor of a self-reported decrease in SSB consumption in the 2 years prior to the survey, and also of current consumption of taxed SSBs. Studies have found that taste is one of the primary drivers of SSBs consumption [40, 53]. This is not surprising given that humans are genetically predisposed to prefer sweet taste [54]. However, preference is also learned [54]. In Mexico, there is a high exposure to sweetened beverages, starting from infancy [55, 56]. Therefore, interventions and programs should focus on reducing children’s repeated exposure to SSBs to prevent heightened SSB preferences early in life from developing. In addition, it is not certain whether a liking for sweet taste can be reduced, thus efforts should be aimed at improving individuals’ self-efficacy and self-regulation skills.

*SSB-health related beliefs* were not associated with either a self-reported change in consumption of SSBs or current consumption of taxed SSBs. There are two plausible reasons for the absence of significance. One is that while beliefs about health outcomes or risks of behavior are a precondition for change, they are not enough on their own, and self-efficacy is needed to overcome impediments or barriers to adopting and maintaining healthy behaviors [57]. A second reason could be that there was little variation in the health beliefs data: 83% of all survey respondents believed that drinking SSBs is associated with the four diseases/conditions they were asked about (See S2 Table).

*Self-efficacy* was also a strong, significant predictor in both regression models. This suggests that people may have felt that they had the confidence to limit their SSB consumption if they wanted to, and, that if they have decreased SSB consumption, it might have been because they had a high sense of self-control. To explore whether people who reported being self-efficacious were those who did not drink SSBs, we conducted further analyses eliminating individuals with low consumption of SSBs ( $\leq 50$  g/person/day), and the results remained significant (data not shown). Self-efficacy has been shown to be significantly associated with SSB consumption in other studies conducted in adults. For example, a study with US parents ( $n = 66$ ) of

adolescents found that among parents perceived behavior control was a significant predictor of SSB consumption and was significantly correlated with intention to decrease SSB consumption [38]. In another study with adults ( $n = 199$ ), Zoellner et al. [39] found that perceived behavioral control was significantly associated with SSB consumption. While drinking (or stopping drinking) SSBs is not a complex behavior in itself, the innate preference for sweet taste and the important sociocultural aspects of SSB drinking in Mexico makes drinking less SSBs a challenging behavior change. Given that the awareness about the detrimental health consequences of drinking SSBs in this population is high (83% of respondents believed that SSBs contribute to obesity, HPB, diabetes, and dental caries), future public health efforts to increase knowledge and enhance motivation should be complemented by programs to assist individuals develop self-efficacy and self-regulation skills [57]. Nutrition education is therefore called for to help individuals develop self-efficacy and self-regulation skills, as well as to help them recognize their susceptibility to disease based on their current SSB consumption.

Overall, the regression models explained a modest amount of variation in the data (9% in self-reported change of SSB consumption, and 21% in current consumption of taxed SSBs). Nevertheless, the existing quantitative psychosocial models of dietary behavior change report a predictive validity less than 30% [58] suggesting that the results of this study are in line with the literature and indicates that the processes underlying food choice are complex and influenced by many factors.

### Limitations and strengths

There are several limitations to this study that should be considered when interpreting its results. First, the data from the POCAA-Q survey are self-reported, and thus could be subject to recall and social desirability response biases. Second, the associations are cross-sectional and do not permit assessment of causality or ascertaining the direction of the association. Third, the study did not use a pre-post design; thus, it was unable to assess change in measures before and after the SSB tax. Fourth, a post-only comparison of outcomes between those aware and not aware of the SSB tax does not fully take into account individuals with a priori favorable attitudes and behaviors who might have been more likely to pay more attention to the campaign. Fifth, there were other public health interventions aimed at decreasing consumption of SSBs that were implemented around the same time as the SSB tax. Lastly, the preference and self-efficacy constructs were assessed with only one item each; according to some researchers this may not adequately define a construct that is stable enough to use in future studies [59, 60].

Despite these limitations, the study has several strengths. Foremost, it provides the first analysis of awareness of the Mexican SSB tax and opinion about its effectiveness in reducing purchases of SSBs in addition to its relationships with a self-reported change in SSB consumption and with current consumption among Mexican adults. It is also the first to assess the association of self-efficacy, taste preference, and health beliefs with SSB consumption in Mexico on a national scale. Findings are generalizable nationally because the ENSANUT 2016 survey uses a probabilistic representative sample.

### Conclusions

Our findings suggest that accompanying SSB taxes with highly visible educational/informational campaigns may contribute to amplifying their effect by further reducing consumption of SSBs. Similarly, studies of tobacco control initiatives have suggested that while tobacco taxation and smoke-free places were two of the key elements of tobacco control strategies [61], part of the success could also be attributed to a shift in social norms and attitudes that

emanated from policy initiatives and health education campaigns [62]. Further research is needed to understand the signaling effect of taxes and the influence of the publicity of taxes on SSB consumption but the aforementioned research lends support to suggested educational campaigns. Researchers in countries that are about to pass SSB taxes could more thoroughly examine this phenomenon by employing pre/post designs. The use of mixed-method approaches for the study of this complex phenomenon—beverage choice in the context of SSB taxes—is advised.

In addition, we found that SSB health-related beliefs were not significantly associated with either a self-reported decrease in SSB consumption after the implementation of the SSB tax, or to current consumption of taxed SSBs. Self-efficacy, on the other hand, and liking of SSBs, were significantly associated. In this context, where a majority of the Mexican adult population likes SSBs, drinks them frequently, and possesses knowledge about the detrimental consequences of SSBs consumption, public health and nutrition education campaigns designed to increase knowledge and enhance motivation should be complemented by programs to assist individuals develop self-efficacy and self-regulation skills.

## Supporting information

**S1 Table. Operational definitions and rationale of variables of choice from the perception of obesity, physical activity and questionnaire (POCAA-Q). ENSANUT 2016.**

(DOCX)

**S2 Table. Percentages and (unadjusted) mean consumption of taxed SSBs in people self-reporting a decrease (or no decrease) in consumption of SSBs in the 2 years prior, by various characteristics.**

(DOCX)

## Author Contributions

**Conceptualization:** Cristina Álvarez-Sánchez, Isobel Contento, Alejandra Jiménez-Aguilar, Pamela Koch, Heewon Lee Gray, Juan Rivera-Dommarco, Rebeca Uribe-Carvajal, Teresa Shamah-Levy.

**Data curation:** Cristina Álvarez-Sánchez.

**Formal analysis:** Cristina Álvarez-Sánchez, Heewon Lee Gray, Laura A. Guerra.

**Funding acquisition:** Pamela Koch, Juan Rivera-Dommarco.

**Methodology:** Cristina Álvarez-Sánchez, Heewon Lee Gray, Laura A. Guerra.

**Project administration:** Cristina Álvarez-Sánchez, Teresa Shamah-Levy.

**Resources:** Pamela Koch, Juan Rivera-Dommarco, Teresa Shamah-Levy.

**Supervision:** Isobel Contento, Pamela Koch, Teresa Shamah-Levy.

**Validation:** Laura A. Guerra.

**Visualization:** Cristina Álvarez-Sánchez.

**Writing – original draft:** Cristina Álvarez-Sánchez.

**Writing – review & editing:** Isobel Contento, Alejandra Jiménez-Aguilar, Pamela Koch, Heewon Lee Gray, Laura A. Guerra, Juan Rivera-Dommarco, Rebeca Uribe-Carvajal, Teresa Shamah-Levy.



## References

1. Shamah-Levy T, Ruiz-Matus C, Rivera-Dommarco J, Kuri-Morales P, Cuevas-Nasu L, Jiménez-Corona M, et al. Encuesta Nacional de Salud y Nutrición de Medio Camino 2016. Resultados Nacionales. Cuernavaca, Mexico: Instituto Nacional de Salud Pública (México), 2017.
2. Barquera S, Campos-Nonato I, Hernández-Barrera L, Pedroza-Tobías A, Rivera-Dommarco JA. Prevalencia de obesidad en adultos mexicanos, ENSANUT 2012. *Salud Pública México*. 2013; 55(sup 2):151–60.
3. Hu FB. Resolved: there is sufficient scientific evidence that decreasing sugar-sweetened beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obes Rev*. 2013; 14(8):606–19. <https://doi.org/10.1111/obr.12040> PMID: 23763695; PubMed Central PMCID: PMC5325726.
4. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ (Clinical research ed)*. 2013; 346:e7492. <https://doi.org/10.1136/bmj.e7492> PMID: 23321486.
5. Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr*. 2006; 84(2):274–88. Epub 2006/08/10. <https://doi.org/10.1093/ajcn/84.1.274> PMID: 16895873; PubMed Central PMCID: PMC53210834.
6. Greenwood DC, Threapleton DE, Evans CE, Cleghorn CL, Nykjaer C, Woodhead C, et al. Association between sugar-sweetened and artificially sweetened soft drinks and type 2 diabetes: systematic review and dose-response meta-analysis of prospective studies. *Br J Nutr*. 2014; 112(5):725–34. <https://doi.org/10.1017/S0007114514001329> PMID: 24932880.
7. Imamura F, O'Connor L, Ye Z, Mursu J, Hayashino Y, Bhupathiraju SN, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. *BMJ (Clinical research ed)*. 2015; 351:h3576. <https://doi.org/10.1136/bmj.h3576> PMID: 26199070; PubMed Central PMCID: PMC53210779.
8. Malik VS, Popkin BM, Bray GA, Despres JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care*. 2010; 33(11):2477–83. <https://doi.org/10.2337/dc10-1079> PMID: 20693348; PubMed Central PMCID: PMC53210779.
9. Wang M, Yu M, Fang L, Hu RY. Association between sugar-sweetened beverages and type 2 diabetes: A meta-analysis. *J Diabetes Investig*. 2015; 6(3):360–6. <https://doi.org/10.1111/jdi.12309> PMID: 25969723; PubMed Central PMCID: PMC53210770.
10. Huang C, Huang J, Tian Y, Yang X, Gu D. Sugar sweetened beverages consumption and risk of coronary heart disease: a meta-analysis of prospective studies. *Atherosclerosis*. 2014; 234(1):11–6. <https://doi.org/10.1016/j.atherosclerosis.2014.01.037> PMID: 24583500.
11. Moynihan PJ, Kelly SA. Effect on caries of restricting sugars intake: systematic review to inform WHO guidelines. *J Dent Res*. 2014; 93(1):8–18. <https://doi.org/10.1177/0022034513508954> PMID: 24323509; PubMed Central PMCID: PMC53210779.
12. Kim S, Park S, Lin M. Permanent tooth loss and sugar-sweetened beverage intake in U.S. young adults. *J Public Health Dent*. 2017; 77(2):148–54. Epub 2016/11/26. <https://doi.org/10.1111/jphd.12192> PMID: 27886383.
13. Sanchez-Pimienta TG, Batis C, Lutter CK, Rivera JA. Sugar-Sweetened Beverages Are the Main Sources of Added Sugar Intake in the Mexican Population. *J Nutr*. 2016; 146(9):1888S–96S. <https://doi.org/10.3945/jn.115.220301> PMID: 27511931.
14. Álvarez-Sánchez C. Mexicans' Consumption of Taxed Sugar-Sweetened Beverages and the Psychosocial Determinants of Consumption in the Context of the 2014 Sugar-Sweetened Beverage Tax—A Mixed Methods Study [Dissertation]. New York City: Columbia University; 2018.
15. Lloyd-Jones DM, Hong Y, Labarthe D, Mozaffarian D, Appel LJ, Van Horn L, et al. Defining and setting national goals for cardiovascular health promotion and disease reduction: the American Heart Association's strategic Impact Goal through 2020 and beyond. *Circulation*. 2010; 121(4):586–613. Epub 2010/01/22. <https://doi.org/10.1161/CIRCULATIONAHA.109.192703> PMID: 20089546.
16. Batis C, Aburto TC, Sanchez-Pimienta TG, Pedraza LS, Rivera JA. Adherence to Dietary Recommendations for Food Group Intakes Is Low in the Mexican Population. *J Nutr*. 2016; 146(9):1897S–906S. Epub 2016/08/12. <https://doi.org/10.3945/jn.115.219626> PMID: 27511940.
17. Donaldson E. Advocating for sugar-sweetened beverage taxation—A case study of Mexico. Johns Hopkins Bloomberg School of Public Health, 2015.
18. Ley del impuesto especial sobre producción y servicios [Excise Tax Law for Production and Services], Última Reforma DOF 11-12-2013 (2013).

19. Colchero MA, Popkin BM, Rivera JA, Ng SW. Beverage purchases from stores in Mexico under the excise tax on sugar sweetened beverages: observational study. *BMJ (Clinical research ed)*. 2016; 352: h6704. Epub 2016/01/08. <https://doi.org/10.1136/bmj.h6704> PMID: 26738745.
20. Colchero MA, Rivera-Dommarco J, Popkin BM, Ng SW. In Mexico, Evidence Of Sustained Consumer Response Two Years After Implementing A Sugar-Sweetened Beverage Tax. *Health affairs (Project Hope)*. 2017; 36(3):564–71. Epub 2017/02/24. <https://doi.org/10.1377/hlthaff.2016.1231> PMID: 28228484; PubMed Central PMCID: PMC5442881.
21. Popkin BM, Hawkes C. Sweetening of the global diet, particularly beverages: patterns, trends, and policy responses. *Lancet Diabetes Endocrinol*. 2016; 4(2):174–86. [https://doi.org/10.1016/S2213-8587\(15\)00419-2](https://doi.org/10.1016/S2213-8587(15)00419-2) PMID: 26654575; PubMed Central PMCID: PMC4733620.
22. Leicester A, Levell P, Rasul I. *Tax and Benefit Policy: Insights from Behavioural Economics*. London: Institute for Fiscal Studies, 2012.
23. Abdulkadrov S. *Who Should Nudge? Nudge Theory in Action: Behavioral Design in Policy and Markets*. Palgrave Advances in Behavioral Economics: Palgrave Macmillan; 2016.
24. World Health Organization. *Fiscal policies for diet and the prevention of noncommunicable diseases*. Geneva: Switzerland: 2016.
25. Falbe J, Thompson HR, Becker CM, Rojas N, McCulloch CE, Madsen KA. Impact of the Berkeley Excise Tax on Sugar-Sweetened Beverage Consumption. *American journal of public health*. 2016; 106(10):1865–71. Epub 2016/08/24. <https://doi.org/10.2105/AJPH.2016.303362> PMID: 27552267; PubMed Central PMCID: PMC45024386.
26. Barigozzi F, Villeneuve B. The signalling effect of tax policy. *Journal of Public Economic Theory*. 2006; 8(4):611–30.
27. Secretaría de Educación Pública, Secretaría de Salud. AGREEMENT which establishes the general guidelines for the sale or distribution of foods and beverages in primary schools [ACUERDO mediante el cual se establecen los lineamientos generales para el expendio o distribución de alimentos y bebidas en los establecimientos de consumo escolar de los planteles de educación básica]. In: Secretaría de Educación Pública y Secretaría de Salud, editor. Mexico: Diario Oficial de la Federación; 2010.
28. Secretaría de Educación Pública, Secretaría de Salud. AGREEMENT which establishes the general guidelines for the sale and distribution of foods and beverages prepared and processed in schools of the National Educational System [ACUERDO mediante el cual se establecen los lineamientos generales para el expendio y distribución de alimentos y bebidas preparados y procesados en las escuelas del Sistema Educativo Nacional.]. In: Salud SadEPySd, editor. Mexico: Diario Oficial de la Federación; 2014.
29. CONAR. Code of Self-Regulation of Food and Non-Alcoholic Beverages Advertising Directed at Children [Código de Autoregulación de Publicidad de Alimentos y Bebidas No Alcohólicas dirigida al Público Infantil]. Mexico: Consejo de Autorregulación y Ética Publicitaria (CONAR), 2009.
30. Secretaría de Salud. Guidelines presenting the nutritional and advertising criteria to be observed by food and non-alcoholic beverage advertisers to advertise their products on open and restricted television, and movie theaters, in accordance with the provisions in articles 22 Bis, 79, section X and 86, section VI, of the Regulation of the General Health Law in relation to Advertising. [LINEAMIENTOS por los que se dan a conocer los criterios nutrimentales y de publicidad que deberán observar los anunciantes de alimentos y bebidas no alcohólicas para publicitar sus productos en televisión abierta y restringida, así como en salas de exhibición cinematográfica, conforme a lo dispuesto en los artículos 22 Bis, 79, fracción X y 86, fracción VI, del Reglamento de la Ley General de Salud en Materia de Publicidad.]. In: Secretaría de Salud, editor. Mexico: Diario Oficial de la Federación; 2014.
31. Romero-Martinez M, Shamah-Levy T, Cuevas-Nasu L, Gomez-Humaran IM, Gaona-Pineda EB, Gomez-Acosta LM, et al. [Methodological design of the National Health and Nutrition Survey 2016]. *Salud Publica Mex*. 2017; 59(3):299–305. Epub 2017/09/14. <https://doi.org/10.21149/8593> PMID: 28902317.
32. INSP. Perception of Obesity, Physical Activity and Diet Questionnaire—National Health and Nutrition Survey 2016 [Cuestionario de percepciones de la población respecto al daño a la salud por la obesidad para adultos de 20 a 59 años de edad]2016. Available from: [https://ensanut.insp.mx/ensanut2016/descarga\\_bases.php](https://ensanut.insp.mx/ensanut2016/descarga_bases.php).
33. Gutierrez JP. Household socioeconomic classification in the National Health and Nutrition Survey 2012. *Salud pública Méx*. 2013; 55(suppl.2).
34. INSP. Food Frequency Questionnaire for Adolescents and Adults (>12 years old)—National Health and Nutrition Survey 2016 [Cuestionario de Frecuencia de Consumo de Adolescentes-Adultos (>12 años de edad)—Encuesta Nacional de Salud Pública y Nutrición de Medio Camino 2016]2016. Available from: [https://ensanut.insp.mx/ensanut2016/descarga\\_bases.php](https://ensanut.insp.mx/ensanut2016/descarga_bases.php).

35. Denova-Gutierrez E, Ramirez-Silva I, Rodríguez-Ramirez S, Jimenez-Aguilar A, Shamah-Levy T, Rivera-Dommarco JA. Validity of a food frequency questionnaire to assess food intake in Mexican adolescent and adult population. *Salud Publica Mex.* 2016; 58(6):617–28. Epub 2017/02/23. <https://doi.org/10.21149/spm.v58i6.7862> PMID: 28225938.
36. Ramirez-Silva I, Jimenez-Aguilar A, Valenzuela-Bravo D, Martinez-Tapia B, Rodriguez-Ramirez S, Gaona-Pineda EB, et al. Methodology for estimating dietary data from the semi-quantitative food frequency questionnaire of the Mexican National Health and Nutrition Survey 2012. *Salud Publica Mex.* 2016; 58(6):629–38. Epub 2017/02/23. <https://doi.org/10.21149/spm.v58i6.7974> PMID: 28225939.
37. Park S, Onufrak S, Sherry B, Blanck HM. The relationship between health-related knowledge and sugar-sweetened beverage intake among US adults. *J Acad Nutr Diet.* 2014; 114(7):1059–66. <https://doi.org/10.1016/j.jand.2013.11.003> PMID: 24360502; PubMed Central PMCID: PMC4470487.
38. Riebl SK, MacDougal C, Hill C, Estabrooks PA, Dunsmore JC, Savla J, et al. Beverage Choices of Adolescents and Their Parents Using the Theory of Planned Behavior: A Mixed Methods Analysis. *J Acad Nutr Diet.* 2016; 116(2):226–39 e1. Epub 2015/12/22. <https://doi.org/10.1016/j.jand.2015.10.019> PMID: 26686818; PubMed Central PMCID: PMC4746018.
39. Zoellner J, Estabrooks PA, Davy BM, Chen YC, You W. Exploring the theory of planned behavior to explain sugar-sweetened beverage consumption. *J Nutr Educ Behav.* 2012; 44(2):172–7. Epub 2011/12/14. <https://doi.org/10.1016/j.jneb.2011.06.010> PMID: 22154130; PubMed Central PMCID: PMC3290682.
40. Zoellner J, Krzeski E, Harden S, Cook E, Allen K, Estabrooks PA. Qualitative application of the theory of planned behavior to understand beverage consumption behaviors among adults. *J Acad Nutr Diet.* 2012; 112(11):1774–84. <https://doi.org/10.1016/j.jand.2012.06.368> PMID: 23102176; PubMed Central PMCID: PMC3500704.
41. Onufrak SJ, Park S, Sharkey JR, Sherry B. The relationship of perceptions of tap water safety with intake of sugar-sweetened beverages and plain water among US adults. *Public Health Nutr.* 2014; 17(1):179–85. Epub 2012/10/27. <https://doi.org/10.1017/S1368980012004600> PMID: 23098620; PubMed Central PMCID: PMC4521760.
42. WHO. Body Mass Index—BMI 2017 [cited 2017 September 24th, 2017]. Available from: <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>.
43. Habicht JP. [Standardization of quantitative epidemiological methods in the field]. *Bol Oficina Sanit Panam.* 1974; 76(5):375–84. Epub 1974/05/01. PMID: 4277063.
44. Lohman TG, Roche AF, Martorell R. Anthropometric standardization reference manual. Human Kinetics Books 1991.
45. Knezevic A. StatNews # 73: Overlapping Confidence Intervals and Statistical Significance 2008. Available from: <https://www.cscu.cornell.edu/news/statnews/stnews73.pdf>.
46. Bursac Z, Gauss CH, Williams DK, Hosmer DW. Purposeful selection of variables in logistic regression. *Source Code Biol Med.* 2008; 3:17. Epub 2008/12/18. <https://doi.org/10.1186/1751-0473-3-17> PMID: 19087314; PubMed Central PMCID: PMC2633005.
47. Field A. *Discovering statistics using SPSS.* 4th Ed. London: SAGE Publications; 2014.
48. University of California Los Angeles (UCLA) Institute for Digital Research and Education (IDRE). FAQ How do I interpret a regression model when some variables are log transformed? California, USA: UCLA Institute for Digital Research and Education (IDRE); 2017 [cited 2017 November 20, 2017]. Available from: <https://stats.idre.ucla.edu/other/mult-pkg/faq/general/faqhow-do-i-interpret-a-regression-model-when-some-variables-are-log-transformed/>.
49. PAHO (Pan American Health Organization). *Taxes on Sugar-sweetened Beverages as a Public Health Strategy: The Experience of Mexico.* Mexico: PAHO, 2015.
50. Barquera S, Hernandez-Barrera L, Tolentino ML, Espinosa J, Ng SW, Rivera JA, et al. Energy intake from beverages is increasing among Mexican adolescents and adults. *J Nutr.* 2008; 138(12):2454–61. <https://doi.org/10.3945/jn.108.092163> PMID: 19022972.
51. Rogers EM. *Diffusion of Innovations*, ed. 5. New York 2003.
52. Xie J, Sreenivasan S, Korniss G, Zhang W, Lim C, Szymanski BK. Social consensus through the influence of committed minorities. *Phys Rev E Stat Nonlin Soft Matter Phys.* 2011; 84(1 Pt 1):011130. Epub 2011/08/27. <https://doi.org/10.1103/PhysRevE.84.011130> PMID: 21867136.
53. Block JP, Gillman MW, Linakis SK, Goldman RE. "If it tastes good, I'm drinking it": qualitative study of beverage consumption among college students. *J Adolesc Health.* 2013; 52(6):702–6. <https://doi.org/10.1016/j.jadohealth.2012.11.017> PMID: 23415754; PubMed Central PMCID: PMC3657589.
54. Ventura AK, Mennella JA. Innate and learned preferences for sweet taste during childhood. *Curr Opin Clin Nutr Metab Care.* 2011; 14(4):379–84. Epub 2011/04/22. <https://doi.org/10.1097/MCO.0b013e328346df65> PMID: 21508837.

55. Barquera S, Campirano F, Bonvecchio A, Hernandez-Barrera L, Rivera JA, Popkin BM. Caloric beverage consumption patterns in Mexican children. *Nutr J*. 2010; 9:47. Epub 2010/10/23. <https://doi.org/10.1186/1475-2891-9-47> PMID: 20964842; PubMed Central PMCID: PMCPMC2987771.
56. Rodriguez-Ramirez S, Munoz-Espinosa A, Rivera JA, Gonzalez-Castell D, Gonzalez de Cosio T. Mexican Children under 2 Years of Age Consume Food Groups High in Energy and Low in Micronutrients. *J Nutr*. 2016; 146(9):1916S–23S. Epub 2016/08/12. <https://doi.org/10.3945/jn.115.220145> PMID: 27511938.
57. Bandura A. Health promotion by social cognitive means. *Health Educ Behav*. 2004; 31(2):143–64. Epub 2004/04/20. <https://doi.org/10.1177/1090198104263660> PMID: 15090118.
58. Baranowski T, Cullen KW, Baranowski J. Psychosocial correlates of dietary intake: advancing dietary intervention. *Annu Rev Nutr*. 1999; 19:17–40. Epub 1999/08/17. <https://doi.org/10.1146/annurev.nutr.19.1.17> PMID: 10448515.
59. Tabachnick BG, Fidell LS. *Using Multivariate Statistics* 4th ed. Boston, MA: Allyn & Bacon; 2001.
60. Velicer WF, Fava JS. The effects of variable and subject sampling on factor pattern recovery. *Psychol Methods*. 1998; 3:231–51.
61. Bader P, Boisclair D, Ferrence R. Effects of tobacco taxation and pricing on smoking behavior in high risk populations: a knowledge synthesis. *Int J Environ Res Public Health*. 2011; 8(11):4118–39. <https://doi.org/10.3390/ijerph8114118> PMID: 22163198; PubMed Central PMCID: PMCPMC3228562.
62. Centers for Disease C, Prevention. Decline in smoking prevalence—New York City, 2002–2006. *MMWR Morb Mortal Wkly Rep*. 2007; 56(24):604–8. PMID: 17585290.