

2019

Diversity of Interaction Solutions of a Shallow Water Wave Equation

Jian-Ping Yu

University of Science and Technology Beijing

Wen-Xiu Ma

University of South Florida

Bo Ren

Shaoxing University

Yong-Li Sun

Beijing University of Chemical Technology

Chaudry Masood Khalique

North-West University

Follow this and additional works at: https://digitalcommons.usf.edu/mth_facpub

Scholar Commons Citation

Yu, Jian-Ping; Ma, Wen-Xiu; Ren, Bo; Sun, Yong-Li; and Khalique, Chaudry Masood, "Diversity of Interaction Solutions of a Shallow Water Wave Equation" (2019). *Mathematics and Statistics Faculty Publications*. 51. https://digitalcommons.usf.edu/mth_facpub/51

This Article is brought to you for free and open access by the Mathematics and Statistics at Digital Commons @ University of South Florida. It has been accepted for inclusion in Mathematics and Statistics Faculty Publications by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact scholarcommons@usf.edu.

Research Article

Diversity of Interaction Solutions of a Shallow Water Wave Equation

Jian-Ping Yu,¹ Wen-Xiu Ma ,^{2,3,4,5,6} Bo Ren ,⁷ Yong-Li Sun ,^{8,9}
and Chaudry Masood Khalique ⁶

¹Department of Applied Mathematics, University of Science and Technology Beijing, Beijing 100083, China

²Department of Mathematics, Zhejiang Normal University, Jinhua 321004, Zhejiang, China

³Department of Mathematics and Statistics, University of South Florida, Tampa, FL 33620, USA

⁴School of Mathematics, South China University of Technology, Guangzhou 510640, China

⁵College of Mathematics and Systems Science, Shandong University of Science and Technology, Qingdao 266590, Shandong, China

⁶International Institute for Symmetry Analysis and Mathematical Modelling, Department of Mathematical Sciences, North-West University, Mafikeng Campus, Private Bag X2046, Mmabatho 2735, South Africa

⁷Institute of Nonlinear Science, Shaoxing University, Shaoxing 312000, China

⁸Department of Mathematics, Beijing University of Chemical Technology, Beijing 100029, China

⁹Department of Mathematics and Statistics, Brock University, Saint Catharines, ON, Canada

Correspondence should be addressed to Wen-Xiu Ma; wma3@usf.edu and Yong-Li Sun; sunyl@mail.buct.edu.cn

Received 28 May 2019; Accepted 16 October 2019; Published 22 November 2019

Academic Editor: Rosa M. Lopez Gutierrez

Copyright © 2019 Jian-Ping Yu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this paper, we study the diversity of interaction solutions of a shallow water wave equation, the generalized Hirota–Satsuma–Ito (gHSI) equation. Using the Hirota direct method, we establish a general theory for the diversity of interaction solutions, which can be applied to generate many important solutions, such as lumps and lump-soliton solutions. This is an interesting feature of this research. In addition, we prove this new model is integrable in Painlevé sense. Finally, the diversity of interactive wave solutions of the gHSI is graphically displayed by selecting specific parameters. All the obtained results can be applied to the research of fluid dynamics.

1. Introduction

The Hirota method played an important role in solving partial differential equations [1]. And, we can solve the corresponding Hirota bilinear equations using many efficient techniques, for example, applying the Wronskian technique [2, 3], we can get positons and complexitons [4]. And, if we take a long wave limit, the lumps, which are locally rationalized along all spacial directions, can be obtained [5–8]. Since the interaction solutions among different classes of solutions can describe more diverse nonlinear phenomena [3], studying interaction solutions is a hot topic for the researchers of mathematical physics [9–16]. Particularly, the interactions between the lumps and kinks [17, 18].

A lot of useful references can be found in [19–27]. Reference [1] presented a shallow water wave equation as follows:

$$\begin{aligned}u_t &= u_{xxt} + 3uu_t - 3u_x v_t - u_x, \\v_x &= -u,\end{aligned}\tag{1}$$

of which the Hirota bilinear form is

$$(D_t D_x^3 - D_t D_x - D_x^2) f \cdot f = 0,\tag{2}$$

via the transformation $u = 2(\ln f)_{xx}$. This kind of transformations is an important part of Bell polynomial theory of partial differential equations [21].

In this study, we will investigate the diversity of a (2 + 1)-dimensional generalized HSI equation that reads

$$\begin{aligned} \text{gHSI} := & u_{tt} + u_{xxxxt} + 6u_x u_t + 3uu_{xt} - 3u_{xx} v_t \\ & + \beta u_{xt} + u_{yt} + \alpha u_{xx} = 0, \end{aligned} \quad (3)$$

$$v_x = -u,$$

which has the following Hirota bilinear form:

$$(D_x^m D_y^n D_t^k f \cdot g)(x, y, t) = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'} \right)^m \left(\frac{\partial}{\partial y} - \frac{\partial}{\partial y'} \right)^n \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'} \right)^k \Big|_{x=x', y=y', t=t'}, \quad (5)$$

with integers $m, n, k \geq 0$.

We will establish the general theory of interaction solutions of equation (3) so that we can build a general method to find the interaction solutions between lumps and other types of solutions of the (2+1)-dimensional gHSI equation by using the Hirota direct approach. Lump solutions and interaction solutions are presented to show diverse nonlinear phenomena. In Section 2, we derive the general approach for finding lumps and interaction solutions. Some applications are presented in Section 3 to illustrate obtained method in Section 2. In the meantime, the diversity of the interaction solutions of the gHSI equation is illustrated vividly by some graphs. In Section 4, the gHSI equation (3) showed that it is integrable in Painlevé sense. Finally, some remarks will be given in the conclusion part.

2. Diversity of Interaction Solutions

There are many ways to find solutions, for example, the symmetry method, the Hirota direct method, and the generalized bilinear method [21–26]. In this section, we will apply the Hirota direct method to establish the theory for the diversity of interaction solutions of the (2+1)-dimensional gHSI equation (3). Hence, the combined solutions of the HSI equation can be found efficiently.

Assume that the (2+1)-dimensional general bilinear equation be as follows:

$$P(D_x, D_y, D_t)(f \cdot f) = 0, \quad (6)$$

where $P(x, y, t)$ is a polynomial of even degree and satisfies $P(0, 0, 0) = 0$. Let

$$\begin{aligned} f &= G + \sum_{i=1}^n (d_i e^{\eta_i}), \\ \eta_i &= a_i x + b_i y + c_i t, \end{aligned} \quad (7)$$

where $G = G(x, y, t)$ is a function of x, y , and t and d_i 's are all real constant to be determined. Moreover, we assume

- (1) $\eta_i, \eta_i + \eta_j \neq 0$ and η_i, η_{j+k} are all distinct for all $i, j, k = 1, \dots, n$.
- (2) G is a positive polynomial and $d_i \geq 0$ and $H = \sum_{i=1}^n (d_i e^{\eta_i})$ with $\eta_i = a_i x + b_i y + c_i t$. According to the Hirota derivatives, we obtain

$$(D_t^2 + D_x D_x^3 + \beta D_t D_x + D_t D_y + \alpha D_x^2) f \cdot f = 0, \quad (4)$$

through the transformation $u = 2(\ln f)_{xx}$. The parameters $\alpha, \beta \neq 0$ are all real constant and D_x, D_t , and D_y are Hirota derivatives [1] which are

$$\begin{aligned} P(D_x, D_y, D_t)(f \cdot f) &= P(D_x, D_y, D_t)(G \cdot G) \\ &\quad + P(D_x, D_y, D_t)(G \cdot H) \\ &\quad + P(D_x, D_y, D_t)(H \cdot H), \\ P(D_x, D_y, D_t)(e^{\eta_i} \cdot e^{\eta_j}) &= 0, \\ P(D_x, D_y, D_t)(e^{\eta_j} \cdot e^{\eta_k}) &= P(a_j - a_k, b_j - b_k, c_j - c_k) e^{\eta_j + \eta_k}, \end{aligned} \quad (8)$$

which implies that (8) can be rewritten as follows:

$$\begin{aligned} P(D_x, D_y, D_t)(f \cdot f) &= P(D_x, D_y, D_t)(G \cdot G) \\ &\quad + 2 \sum_{i=1}^n (d_i P(D_x, D_y, D_t)(G \cdot e^{\eta_i})) \\ &\quad + 2 \sum_{1 \leq j < k < n} (d_j d_k P(D_x, D_y, D_t)(e^{\eta_j + \eta_k})). \end{aligned} \quad (9)$$

Hence, if

$$\begin{aligned} P(D_x, D_y, D_t)(G \cdot e^{\eta_i}) &= 0, \quad i = 1, \dots, n, \\ P(D_x, D_y, D_t)(e^{\eta_j} \cdot e^{\eta_k}) &= 0, \end{aligned} \quad (10)$$

where $i, j, k = 1, \dots, n$ and $j \neq k$, then f is a solution of equation (6) if and only if G is also a solution of equation (6). Therefore, using the transformations $u = 2(\ln f)_x$ or $u = 2(\ln f)_{xx}$, we can get the interact solutions: lump-solution solutions of the gHSI equation (3).

Remark. (1) If we further let

$$f = g^2 + h^2 + d + ke^l, \quad (11)$$

where

$$\begin{aligned} g &= a_1 x + a_2 y + a_3 t + a_4, \\ h &= b_1 x + b_2 y + b_3 t + b_4, \\ l &= c_1 x + c_2 y + c_3 t, \end{aligned} \quad (12)$$

and $d, k \geq 0$, then f is a solution of equation (6) if and only if $g^2 + h^2 + d$ is also a solution of equation (6) under the condition

$$P(D_x, D_y, D_t)((g^2 + h^2 + d) \cdot e^l) = 0. \quad (13)$$

(2) If $G = g^2 + h^2 + d$ is a solution of equation (6), then we have

- (i) $u = 2(\ln G)_x$ or $u = 2(\ln G)_{xx}$ is a lump solution
- (ii) Moreover, if $k > 0$, then $u = 2(\ln f)_x$ or $u = 2(\ln f)_{xx}$ is a lump-soliton solution if and only if

$$P(D_x, D_y, D_t)(G \cdot e^l) = 0. \quad (14)$$

3. Application to Shallow Water Wave Equation

3.1. Lump Solution of the gHSI Equation. Firstly, we consider the lump solutions of equation (4). We suppose that

$$G = g^2 + h^2 + d = (a_1x + a_2y + a_3t + a_4)^2 + (b_1x + b_2y + b_3t + b_4)^2 + d, \quad (15)$$

where g and h are linearly independent and $d > 0$. The parameters a_i 's are obtained via the direct computation as follows:

$$\begin{aligned} d &= -\frac{3a_1(a_1^2 + b_1^2)a_3}{b_1^2(a_1^2 - b_1^2)\alpha}, \\ a_2 &= \frac{-\alpha a_1^2 + \alpha b_1^2 - \beta a_1 a_3 - a_3^2}{a_3}, \\ b_2 &= \frac{-b_1(\beta a_1^2 - \beta b_1^2 + 2a_1 a_3)}{a_1^2 - b_1^2}, \\ b_3 &= \frac{2a_3 b_1 a_1}{a_1^2 - b_1^2}, \end{aligned} \quad (16)$$

where $\alpha, \beta \neq 0$. Then, we can get the lump solution of equation (3) as

$$u = \frac{4((a_1^2 + b_1^2)f - 2(a_1g + b_1h)^2)}{f^2}, \quad (17)$$

with $\alpha a_1 a_3 < 0$ and $a_1^2 - b_1^2 \neq 0$. It is observed that, at any given time t , the extremum points can be obtained by direct computation, from which the traveling speeds, along x -direction and y -direction, and the changes of waveform can be obtained. The amplitude of u is also attained. We also noted that the lump wave is analytic in the XY -plane if and only if $d > 0$. Moreover, it is easy to find the aforementioned lump solution $u \rightarrow 0$ if and only if the sum of squares $g^2 + h^2 \rightarrow \infty$, or equivalently, $x^2 + y^2 \rightarrow \infty$ at any given time. The evolution profile, density plot, and contour plots of solution (15) with specific parameters are shown in Figure 1, from which we can see that the waveforms of (15) change only a little bit at different time.

3.2. Interaction Solutions of the gHSI Equation. In this part, we will find some lump-soliton solutions of the gHSI equation (3). Assume $f = g + h + d + ke^l$ with g, h, d , and k defined as in equation (11). By the logarithm transformation $u = 2(\ln f)_{xx}$, we get the lump-soliton solution as

$$u = 2 \frac{f_{xx}f - f_x^2}{f^2}. \quad (18)$$

By theories in Section 2, we can find the solution of all the parameters as follows:

$$\begin{aligned} a_1 &= -\frac{3b_3c_1^2}{2\alpha}, \\ a_2 &= -\frac{-9ab_1c_1^4 - 9\beta b_3c_1^4 + 4\alpha^2b_1}{6\alpha c_1^2}, \\ a_3 &= \frac{2\alpha b_1}{3c_1^2}, \\ c_3 &= -\frac{2\alpha}{3c_1}, \\ b_2 &= -\frac{-9b_3c_1^4 + 4\alpha\beta b_1 + 4\alpha b_3}{4\alpha}, \\ d &= 0, \\ b_4 &= \frac{3a_4b_3c_1^2}{2\alpha b_1}, \\ c_2 &= \frac{3c_1^4 - 6\beta c_1^2 + 4\alpha}{6c_1}, \end{aligned} \quad (19)$$

which yields the following functions:

$$\begin{aligned} g &= -\frac{3b_3c_1^2}{2\alpha}x - \frac{-9ab_1c_1^4 - 9\beta b_3c_1^4 + 4\alpha^2b_1}{6\alpha c_1^2}y + \frac{2\alpha b_1}{3c_1^2}t + a_4, \\ h &= b_1x - \frac{-9b_3c_1^4 + 4\alpha\beta b_1 + 4\alpha b_3}{4\alpha}y + b_3t + b_4, \\ ke^l &= c_1x + \frac{3c_1^4 - 6\beta c_1^2 + 4\alpha}{6c_1}y - \frac{2\alpha}{3c_1}t. \end{aligned} \quad (20)$$

Therefore, we can get the function $f = g^2 + h^2 + d + ke^l$ which implies that the lump-soliton solution of the gHSI equation is also obtained by equation (20). We can also get the extremum points by direct computation in Maple, which play an important role in studying the wave equations, for example, the velocities, along x -direction and y -direction, the amplitude of u , and the changes of waveform can be obtained via the extremum points. We also found that the lump wave is analytic in the XY -plane if and only if $c_1 \neq 0$ and $b_1 \neq 0$. The aforementioned lump-soliton solution is an interactive solution; hence, during the collision, they interact like fusion and fission phenomenon in physics. At first, the energy of the lump wave is stronger than the stripe wave




FIGURE 1: 3D plots, density plots, and contour plots of lump wave solution (19) with the specific parameters $a_1 = 2, a_3 = 1, a_4 = 1, b_1 = 1, b_3 = 1, b_4 = 1, \alpha = -1$, and $\beta = 2$. (a), (d), and (g) are for $t = -2$; (b), (e), and (h) are for $t = 0$; (c), (f), and (i) are for $t = 6$.

described by the exponential function, but finally the lump wave are gradually swallowed by the stripe soliton, which implies that its energy is also transferred to the stripe soliton completely. They become one soliton. The evolution profiles and contour plots of solution (20) with specific parameters are shown in Figure 2, from which we observed that the intersect solution (20) of the gHSI equation change greatly at different time.

4. Painlevé Analysis

It is well known that Painlevé analysis is a very powerful tool for finding the integrable model from given nonlinear equations [27]. Using the WTC-Kruskal approach, we firstly analyze the leading order to the negative integer α , then determine the resonant points, and finally obtain the

compatibility conditions, which must be completely satisfied for all the positive resonant points. Baldwin et al. presented two packages in Mathematica based on the WTC approach and Kruskal's simplification.

Applying the aforementioned packages in Mathematica to test the integrability of the $(2+1)$ -dimensional gHSI equation (3), we find five resonant points $j = -1, 1, 4, 5, 6$. In all the cases, equation (3) does pass the Painlevé test. It is noted that the presence of soliton solutions can indicate the integrability of the tested equation. But, this is not enough since it should be supported by the Painlevé test, or the Lax pair of the examined equation or other approaches. In this study, we formally obtained lump solutions and lump-soliton solutions of the gHSI equation (3) and showed that it passed the Painlevé test, which implies that it is an integrable equation in Painlevé sense.




FIGURE 2: 3D plots, density plots, and contour plots of lump wave solution (20) with the specific parameters $k = 1, a_4 = 1, b_1 = 1, b_3 = 1, b_4 = 1, c[1] = 1, \alpha = 1$, and $\beta = 2$. (a), (d), and (g) are for $t = -5$; (b), (e), and (h) are for $t = 0$; (c), (f), and (i) are for $t = 6$.

5. Conclusions

In this research, we introduced a shallow water wave equation, the gHSI equation (3), and established the theory of its diversity of interactions, the lump solution, and lump-soliton solutions. All the computations are performed in Maple using the Hirota bilinear equations. Moreover, we proved that this gHSI equation (3) is Painlevé integrable. During the study, we found that the waveforms of (20) are completely different if we select different values of α and β . For example, if we choose $\alpha = -2$, the waveform has a unique peak at the maximum point.

The research of the diversity of interaction actions is an interesting and hot topic in mathematical physics since we can get a lot of useful solutions for the physical research. Hence, we will continue studying other interaction solutions, such as the interactions between the periodic function

solutions and the hyperbolic function solutions. In addition, we hope that we can find whether equation (3) is integrable in Liouville sense or not.

In the meantime, this introduced shallow water equation has some applications in physics research. For example, it can be used to describe the flow under a pressure surface (sometimes a free surface) in a fluid, which implies that it can be applied to the research on the fluid dynamics.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (nos. 11101029, 11271362, 11375030, and DMS-1664561), Fundamental Research Funds for the Central Universities (no. 610806), Beijing City Board of Education Science and Technology Key Project (no. KZ201511232034), Beijing Nova Program (no. Z131109000413029), and Beijing Finance Funds of Natural Science Program for Excellent Talents (no. 2014000026833ZK19).

References

- [1] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, New York, NY, USA, 2004.
- [2] N. C. Freeman and J. J. C. Nimmo, "Soliton solutions of the Korteweg-de Vries and Kadomtsev-Petviashvili equations: the wronskian technique," *Physics Letters A*, vol. 95, no. 1, pp. 1-3, 1983.
- [3] W. X. Ma and Y. You, "Solving the Korteweg-de Vries equation by its bilinear form: Wronskian solutions," *Transactions of the American Mathematical Society*, vol. 357, no. 5, pp. 1753-1778, 2005.
- [4] A.-M. Wazwaz and S. A. El-Tantawy, "New (3+1)-dimensional equations of burgers type and Sharma-Tasso-Olver type: multiple-soliton solutions," *Nonlinear Dynamics*, vol. 87, no. 4, pp. 2457-2461, 2017.
- [5] S. V. Manakov, V. E. Zakharov, L. A. Bordag, A. R. Its, and V. B. Matveev, "Two-dimensional solitons of the Kadomtsev-Petviashvili equation and their interaction," *Physics Letters A*, vol. 63, no. 3, pp. 205-206, 1977.
- [6] J. Satsuma and M. J. Ablowitz, "Two-dimensional lumps in nonlinear dispersive systems," *Journal of Mathematical Physics*, vol. 20, no. 7, pp. 1496-1503, 1979.
- [7] M. J. Ablowitz and P. A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, UK, 1991.
- [8] C. R. Gilson and J. J. C. Nimmo, "Lump solutions of the BKP equation," *Physics Letters A*, vol. 147, no. 8-9, pp. 472-476, 1990.
- [9] W.-X. Ma, "Lump solutions to the Kadomtsev-Petviashvili equation," *Physics Letters A*, vol. 379, no. 36, pp. 1975-1978, 2015.
- [10] B. Ren, "Interaction solutions for mKP equation with nonlocal symmetry reductions and CTE method," *Physica Scripta*, vol. 90, no. 6, Article ID 065206, 2015.
- [11] B. Ren, X.-P. Cheng, and J. Lin, "The (2+1)-dimensional Konopelchenko-Dubrovsy equation: nonlocal symmetries and interaction solutions," *Nonlinear Dynamics*, vol. 86, no. 3, pp. 1855-1862, 2016.
- [12] J.-Y. Yang and W.-X. Ma, "Lump solutions of the BKP equation by symbolic computation," *International Journal of Modern Physics B*, vol. 30, no. 28n29, Article ID 1640028, 2016.
- [13] W. X. Ma, Z. Y. Qin, and X. Lü, "Lump solutions to dimensionally reduced p-gKP and p-gBKP equations," *Nonlinear Dynamics*, vol. 84, no. 2, pp. 923-931, 2016.
- [14] J.-P. Yu and Y.-L. Sun, "Lump solutions to dimensionally reduced Kadomtsev-Petviashvili-like equations," *Nonlinear Dynamics*, vol. 87, no. 2, pp. 1405-1412, 2017.
- [15] Y. Zhang, H. H. Dong, X. E. Zhang, and H. W. Yang, "Rational solutions and lump solutions to the generalized (3+1)-dimensional shallow water-like equation," *Computers & Mathematics with Applications*, vol. 73, no. 2, pp. 246-252, 2017.
- [16] Z. H. Xu, H. L. Chen, and Z. D. Dai, "Rogue wave for the (2+1)-dimensional Kadomtsev-Petviashvili equation," *Applied Mathematics Letters*, vol. 37, pp. 34-38, 2014.
- [17] Y. N. Tang, S. Q. Tao, and G. Qing, "Lump solitons and the interaction phenomena of them for two classes of nonlinear evolution equations," *Computers & Mathematics with Applications*, vol. 72, no. 9, pp. 2334-2342, 2016.
- [18] J.-Y. Yang and W.-X. Ma, "Abundant interaction solutions to the KP equation," *Nonlinear Dynamics*, vol. 89, no. 2, pp. 1539-1544, 2017.
- [19] C. Gilson, F. Lambert, J. Nimmo, and R. Willox, "On the combinatorics of the Hirota D-operators," *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, vol. 452, no. 1945, pp. 223-234, 1996.
- [20] A. M. Wazwaz, "New solutions of distinct physical structures to high-dimensional nonlinear evolution equations," *Applied Mathematics and Computation*, vol. 196, pp. 363-370, 2008.
- [21] W.-X. Ma, "Bilinear equations, Bell polynomials and linear superposition principle," *Journal of Physics: Conference Series*, vol. 411, Article ID 012021, 2013.
- [22] W.-X. Ma and W. Strampp, "An explicit symmetry constraint for the Lax pairs and the adjoint Lax pairs of AKNS systems," *Physics Letters A*, vol. 185, no. 3, pp. 277-286, 1994.
- [23] H. Dong, Y. Zhang, and X. Zhang, "The new integrable symplectic map and the symmetry of integrable nonlinear lattice equation," *Communications in Nonlinear Science and Numerical Simulation*, vol. 36, pp. 354-365, 2016.
- [24] W. X. Ma, "Generalized bilinear differential equations," *Studies in Nonlinear Sciences*, vol. 2, no. 4, pp. 140-144, 2011.
- [25] W.-X. Ma, "Bilinear equations and resonant solutions characterized by bell polynomials," *Reports on Mathematical Physics*, vol. 72, no. 1, pp. 41-56, 2013.
- [26] W.-X. Ma, "Trilinear equations, bell polynomials, and resonant solutions," *Frontiers of Mathematics in China*, vol. 8, no. 5, pp. 1139-1156, 2015.
- [27] D. Baldwin and W. Hereman, "Symbolic software for the Painlevé test of nonlinear ordinary and partial differential equations," *Journal of Nonlinear Mathematical Physics*, vol. 13, no. 1, pp. 90-110, 2006.




Hindawi

Submit your manuscripts at
www.hindawi.com

