

A PALEOECOLOGICAL RECONSTRUCTION OF THE LATE GLACIAL AND HOLOCENE BASED ON MULTIDISCIPLINARY STUDIES AT STEREGOIU SITE (GUTÂI MTS., NW ROMANIA)

ANGELICA FEURDEAN¹, LEIF BJÖRKMAN², BARBARA WOHLFARTH²

ABSTRACT. High resolution analyses of pollen, mineral magnetic properties, loss of ignition, lithostratigraphy and AMS ¹⁴C measurements of lake sediments and peat deposits accumulated in the former crater lake of Steregoiu (Gutâiului Mts., NW Romania), gave new and important information about vegetation and climate changes from the period GS-2 to the present. During the Lateglacial, three cold events were recorded: before 14,700 cal. years BP (GS-2), 14,050–13,800 cal. years BP (GI-1d), 12,900-11,500 cal. years BP (GS-1), and a warm climatic event between 13,800-12,950 cal. years BP (GI-1c to GL-1a). The Late Glacial/Holocene transition around 11,500 cal. years BP, was determined by an expansion of *Betula*, *Alnus* and *Picea*, followed by a rapid and strong expansion of *Ulmus*. At 10,700 cal. years BP, dense and highly diverse forests with *Ulmus*, *Quercus*, *Tilia*, *Fraxinus* and a few *Acer* and *Corylus* individuals dominated the area. *Corylus* and *Picea* were the dominant species in the forests from 10,150 to 8,500 cal. years BP. The first occurrence of single *Fagus* pollen grains was around 8,000 cal years BP. Only at 4,700 cal year BP *Fagus* and *Carpinus* became widespread and established trees in the local woodlands.

Keywords: Vegetation dynamic, Late Glacial-Holocene, pollen, AMS ¹⁴C, SIRM, magnetic susceptibility, LOI, Steregoiu, Gutâi Mts., Romania.

INTRODUCTION

The aim of this paper is to present a detailed paleoecological reconstruction of the Late Glacial and Holocene vegetation and climate history in the Gutâiului Mountains and to compare this record with other data from Romania. High-resolution pollen analyses were performed in order to study local long-term vegetation dynamics, to detect glacial refugia of deciduous trees, and to establish a chronology of the immigration and expansion of tree species at the Late Glacial/Holocene transition as a response to climatic change. Apart from pollen analysis we used other paleoenvironmental methods in order to improve the interpretation of the past environment. Thus, loss on ignition (hereafter LOI) was performed to determine the organic production at the time when the sediment was deposited. Magnetic measurements (SIRM and susceptibility) were used to aid the visual correlation of individual cores, and to detect the amount of inorganic allochthonous material in the basin, and also to characterize the composition of the magnetic minerals. AMS ¹⁴C measurements on terrestrial plant macrofossils enabled to establish the chronology of the sediments.

¹ Group for Quaternary Research, Department of Geology, "Babeș-Bolyai" University, Kogălniceanu 1, 3400 Cluj-Napoca, Romania.

² Department of Quaternary Geology, University of Lund, Tornavägen 13, 223 63 Lund, Sweden.

Quaternary paleoenvironment reconstruction in Romania is based mostly on pollen analyses (Pop, 1932, 1942; 1960; Diaconeasa; 1995; Lupşa, 1980; Buz, 1999; Fărcaş, 1999, 2001). Only in the last decade have new modern methods been utilized in an attempt to better understand Quaternary events (Onac & Lauritzen, 1996; Björkman et al., 2001; Onac et al., 2001; Wohlfarth et al., 2001). No published information about Quaternary glacial deposits exists for the study area. According to Woldstedt (1958), alpine glaciers did not reach below 1600 m. a.s.l. in the Carpathians during the Last Glacial Maximum. Thus, Romania could probably support a glacial refuge for different tree species. The presence of scattered pollen grains or low percentages of deciduous trees, such as *Quercus*, *Ulmus*, *Tilia*, and *Corylus* during the Late Glacial were found in few places in Romania and have been attributed to contamination during coring, redeposition, or long distance transport (Buz, 1999; Fărcaş, 1999). Only the southern and southeastern part of Romania is accepted as being glacial refuges for some deciduous tree taxa as *Quercus*, *Tilia* and *Fraxinus*. Other deciduous trees, *Salix*, *Betula*, and *Alnus* are constantly being recorded in Late Glacial deposits, and are known as “cold resistant species”. Together with these species, coniferous trees (*Pinus* and *Picea*) were also common. The rapid establishment and expansion of deciduous trees at the beginning of the Holocene implies that glacial refuges were not located far from Romania, or may have existed within Romania.

In this area several sites were investigated by means of pollen analysis. Romania has one of the longest traditions of palynological work in Europe. Earlier investigations have been concentrated mostly on Holocene forest history. Moreover, little has been done with respect to Late Glacial vegetation dynamics and climate. Pollen analyses have been performed with very low resolution (often more than 10 cm between samples) and none have been supported by radiocarbon dates (Pop, 1932, 1942; 1960; Lupşa, 1980). These authors have established the chronologies of the pollen diagrams by comparison with the German chronology proposed by Firbas (1949, 1952) for central and Eastern Europe. Instead this multidisciplinary study enables comparison of the Romanian paleoenvironment with well-dated sequences from western and northwestern Europe (Björkman et al., 2001; Wohlfarth et al., 2001).

STUDY AREA

The study site, Steregoiu crater (47° 48'48" N, 23° 32'41" E) is situated in the northwestern part of Romania, on the western flank of the volcanic Gutâiului Mountains at an altitude of ca. 800 m.a.s.l (Fig. 1.). This massive belongs to the western extremity of the Eastern Carpathian mountain chain. The bedrock consists of volcanic rocks, mainly andesites rich in pyroxene and quartz.

The present climatic conditions in this region are continental-temperate (cold and humid). The massif is highly forested and has narrow and long valleys. Its southern slope receives high quantities of solar radiation, being at the same time protected from strong winds coming from west and north. Sometimes, warm air masses come from the Pannonian steppe, therefore making it possible for Mediterranean species, such as *Castanea sativa* to grow in Romania.




Fig. 1. Location of the Steregoiu crater site

The mean annual temperature is 2-4°C high up in the mountains and 7-8°C in depressionary zone. The mean winter temperature is 3.5°C, and mean summer temperature is 12-14°C. The area has a high humidity, the precipitation's is around 900 mm/yr, but could sometimes rise to 1200-1400 mm/yr. (Istvan et al., 1990; Mac & Budai, 1992). The crater has an elongate surface of 0,5 ha and is drained by a small stream towards the east. The site lies within the beech (*Fagus silvatica*) forest belt in which rarely *Betula verrucosa*, *Acer pseudoplatanus* and *Carpinus betulus* can be found. The vegetation on the surrounding slopes consists of young beech (*Fagus*) forest, In addition spruce (*Picea*) trees can be found around the site, that is now swamp of a eumesotrophic character, where the mire vegetation consists mainly of grasses, sedges, herbs and mosses.

METHODS

Sample collection was performed with a Russian peat sampler (length: 1 m; diameter: 5 cm). Six overlapping cores were collected from the central part of the site. The cores were preliminary described in field. Laboratory work was performed at the Department of Geology in Lund, Sweden.

In the laboratory, the core surfaces were cleaned carefully in order to avoid contamination and described in detail. The cores were sub-sampled continuously at 2-cm interval for pollen analysis, magnetic analysis, LOI, and at 4-cm intervals for plant macrofossil studies.

The samples were dried at 40°C, in order to calculate magnetic susceptibility and Saturation Isothermal Remanent Magnetization (SIRM). Magnetic susceptibility was measured in a low magnetic field of 0.1 mT using a balanced alternating current bridge circuit. Mass specific units were calculated and expressed as $\mu\text{m}^3\text{kg}^{-1}$. SIRM was induced in a strong magnetic field of 1 Tesla by a Redcliff BSM 700 Puls Magnetic Charger. This magnetic field is strong enough to saturate the samples. The resulting remanent magnetization was measured with a Molspin Spinner Magnetometer. Mass specific units were calculate as $\text{mAm}^2\text{kg}^{-1}$. SIRM reflects

the concentration, composition and grain size distribution of magnetic minerals in the sample, however only minerals that are capable of holding a remanence will contribute to this signal (Sandgren & Snowball, 1999; Thomson & Oldfield, 1986).

LOI was estimated following the methods described by Bengtesson & Enell (1986). The samples were placed in crucibles with a known weight, dried overnight at 100°C, cooled and weighed. The samples were then ashed at 550°C for 3 hours, cooled and weighed again. Previous research has shown that carbonates are not present in the sediments. LOI was used for estimating the amount of organic matter and is presented as the percentage of the weight of the dried sample.

For pollen analysis, 258 sub-samples with 1 cm³ of volume were taken. Five *Lycopodium* tablets with a known number of spores were added to each sample in order to determine the concentration of fossil pollen. The chemical preparation for pollen analysis follows the standard methods of Berglund and Ralska-Jasiewiczowa (1986) and Moore et al. (1991). The slides were mounted in glycerine. Pollen counts were normally made at 400x magnification, whereas 1000x were used for some critical determinations. Pollen identification follows the keys and illustrations in Moore et al. (1991), Reille (1992) and Faegri et al. (1989). In addition, the pollen reference collection at the Department of Geology, Lund University, was also used to check several types. Between 500 and 700 pollen grains were counted in most of the samples, except in the ten bottom most samples, where a sum of 300 grains was accepted due to low pollen concentration. The calculation of pollen percentages is based on the total sum of terrestrial pollen taxa. The pollen diagram was constructed with the TILIA computer program (Grimm, 1987, 1991).

AMS ¹⁴C measurements were performed on terrestrial plant macrofossils. Seventeen samples were dated by the AMS facility in Uppsala, Sweden.

RESULTS

1. Lithostratigraphy, content of organic matter, mineral magnetic measurements

Based on the sediment description of the cores, ten lithological units were described (Table 1). The measurements of magnetic susceptibility, SIRM and carbon content are presented in Figure 2.

Unit 1 in the bottom-part of the sequence is composed of reddish brown silty clay with sandy and gravelly layers. SIRM starts with high values, but decreases from 27.192 to 3.3 mAm²kg⁻¹ towards the top. A few isolated peaks occur due to sandy and gravelly layers. Magnetic susceptibility decreases from 1.8 to 0.22 μm³kg⁻¹ and LOI is around 8%.

Unit 2 is composed of gyttja clay with different colors. SIRM has a maximum value of 2.49 mAm²kg⁻¹, and magnetic susceptibility is 0.14 μm³kg⁻¹. LOI shows a slight, but distinct increase to 14.4%.

Unit 3 consists of grayish brown silt, and slightly sandy and gravelly layers with an erosive lower boundary. In comparison with previous zone, SIRM values show an increase to 5.6 mAm²kg⁻¹, and magnetic susceptibility to 0.34 μm³kg⁻¹, while LOI decreases to 12%.

Unit 4 consists of clay and gyttja with visible macrofossils. Susceptibility decreases throughout the unit. SIRM shows decreasing values up to the middle of

A PALEOECOLOGICAL RECONSTRUCTION OF THE LATE GLACIAL AND HOLOCENE ...

the unit, but increases again toward the top to 2.4 mAm²kg⁻¹. LOI increases to 18%, and upwards to 99% in the middle part.

Unit 5 is composed of grayish gyttja. SIRM and susceptibility values show a slight increase and the LOI remains constant.

Unit 6 is composed of brown gyttja. SIRM and susceptibility value decrease and LOI increases slightly to 25%.

Unit 7 is composed of dark brown, coarse detritus gyttja with wood fragments. SIRM and susceptibility values are low and LOI increase to 62%, but decreases again to 40% in the upper part.

Unit 8 consists of dark brown, drift gyttja. SIRM and susceptibility values are low and LOI increases from 66% to 92%.

Unit 9 is composed of dark brown, carr peat with alternations of more and less humidified layers. SIRM and susceptibility values are low and LOI fluctuates from 77 to 99%.

Unit 10 is a dark brown fen peat with low, medium, and highly humidified horizons. The LOI ranges from 72 to 99%.

Unit	Depth below surface (m)	Description
10	0.07 - 0	Medium humidified brown, fen peat
	0.11 - 0.07	Medium to low humidified reddish brown, fen peat
	0.255 - 0.11	Medium humidified reddish brown, fen peat
	125 - 25.5	Dark brown, highly humidified fen peat
9	2.80 - 1.255	Dark brown carr peat with more humidified and less humidified parts
8	2.865 - 2.80	Dark brown, transition "drift gyttja", g LB
7	3.35 - 2.865	Dark brown coarse detritus gyttja
	3.715 - 3.35	Dark brown, very coarse detritus gyttja with large wood fragments g LB
	4.115 - 3.715	Dark brown, very coarse detritus gyttja, g LB
	4.28 - 4.115	Brown, coarse detritus gyttja, g LB
6	4.57 - 4.28	Brown, gyttja, g LB
5	4.663 - 4.57	"Transition" Dark gray, gyttja
4	4.82 - 4.663	Lighter greyish, brown clay gyttja, g LB
	4.3 - 4.82	Grayish brown, clayey gyttja, s LB
	5.108 - 4.93	Lighter greyish brown, clayey gyttja, g LB
	5.225 - 5.108	Grayish brown, clayey gyttja, g LB, visible macrofossils
3	5.26 - 5.225	Grayish-brown, silty, slightly sandy and gravel erosive LB
2	5.35 - 5.26	Grayish brown, gyttja -clay, g LB
	5.38 - 5.35	Reddish brown, gyttja clay, s LB
	5.405 - 5.38	Bluish brown, gyttja clay, g LB5
	5.44 - 5.405	Reddish brown, gyttja clay, g LB
1	5.70 - 5.44	Reddish brown, partly laminated silty clay, g LB
	5.84 - 5.70	Reddish brown, slightly clayey sandy silt with gravel, s LB
	5.88/87 - 5.84	Reddish, silty clay, sandy gravelly layers at 584.5, s LB
	5.925 - 5.88/87	Brown reddish, silty clay with gravel

Table 1. Lithostratigraphic description of the Steregoiu core (LB = layer boundary; s = sharp; g = gradual).




Fig. 2. SIRM, magnetic susceptibility and organic matter (loss of ignition) of the Steregoiu core sediments.

2. Chronology

The AMS ^{14}C measurements were converted into calibrated years BP, using the radiocarbon calibration curve (Stuvier et al., 1998). Based on the calibrated dates, an age depth curve was established. From this curve accumulation rates of the sediments were estimated (Fig. 3). This rate is calculated to 41.3 years/cm of in the inorganic sediments (between 5.43–4.67 m), 15.2 years/cm in the gyttja (between 4.67–2.80 m). In the peat, the accumulation rate is about 9.5 years/cm (in carr peat between 2.80–1.80 m), 66.2 years/cm in the upper part of carr peat (between 1.80–1.255 m), and 31 years/cm in the fen peat (1.255–0 m).

3. Pollen stratigraphy

The pollen data are presented in a percentage diagram with all terrestrial pollen types included in the calculation sum (Fig. 3). The pollen taxa are plotted against depth and an interpolated age. To facilitate the description and interpretation of the pollen diagram, nineteen local pollen assemblages zones (LPAZ S1-S19) have been established (Table 2). These zones have been established visually, and each zone boundary denotes significant changes in pollen deposition and hence, major changes in vegetation cover.

A PALEOECOLOGICAL RECONSTRUCTION OF THE LATE GLACIAL AND HOLOCENE ...

Table 2. Pollen stratigraphy.

Age	Depth	Sedimentary units	Pollen zones	Dominating pollen types
0 – 300	0.14 – 0.005	10	19	<i>Fagus – Poaceae – Quercus</i>
300 – 1050	0.39 – 0.14	10	18	<i>Fagus – Quercus – Carpinus</i>
1050 – 2.200	0.73 – 0.39	10	17	<i>Fagus – Poaceae – Carpinus</i>
2.200 – 3.400	1.11 – 0.73	10	16	<i>Fagus – Picea – Carpinus</i>
3.400 – 4.800	1.41 – 1.11	10	15	<i>Corylus – Picea – Fagus</i>
4.800 – 7.500	1.89 – 1.41	9	14	<i>Corylus – Picea – Ulmus – Quercus</i>
7.500 – 8.200	2.5 – 1.89	9	13	<i>Corylus – Picea – Ulmus – Filipendula</i>
8.200 – 8.600	2.82 – 2.5	9	12	<i>Corylus – Picea – Ulmus</i>
8.600 – 9.300	3.35 – 2.82	7, 8	11	<i>Corylus – Ulmus – Picea</i>
9.300 – 10.200	3.98 – 3.35	7	10	<i>Ulmus – Corylus – Picea</i>
10.200 – 10.750	4.36 – 3.98	6, 7	9	<i>Ulmus – Picea – Fraxinus</i>
10.750 – 11.250	4.59 – 4.36	6	8	<i>Ulmus – Pinus – Betula</i>
11.250 – 11.500	4.69 – 4.59	5	7	<i>Pinus – Betula – Alnus</i>
11.500 – 12.600	4.99 – 4.69	4	6	<i>Pinus – Artemisia – Poaceae</i>
12.600 – 12.950	5.09 – 4.99	4	5	<i>Betula – Pinus – Alnus</i>
12.950 – 13.750	5.25 – 5.09	4	4	<i>Betula – Pinus – Picea</i>
13.750 – 14.150	5.32 – 5.25	2, 3	3	<i>Pinus – Alnus – Betula</i>
14.150 – 14.700	5.44 – 5.32	2	2	<i>Pinus – Poaceae – Betula</i>
> 14.700	5.92 – 5.44	1	1	<i>Pinus – Poaceae – Artemisia</i>

DISCUSSION

1. Reconstruction of the sedimentary history

The sediments from Steregoiu extend more than 14700 cal. years BP back in time. The infilling of the Steregoiu crater was made by allochthonous material (clay and sand), and by autochthonous material (as gyttja and peat). The bottom sediments, below 5.44 m contain inorganic and compact sediments. They were probably they were deposited due to slope erosion, and strong winds in a mainly dry climate. Between 5.44 m to 5.26 m, the sediment becomes more organogenic as gyttja clay was observed in the lake. Increasing amounts of organogenic material suggests higher productivity in the lake, but still made shallow conditions. Between 5.26 –4.57 m an increase of inwashed minerogenic material, is recorded, as clayey gyttja was deposited. From 4.57 m to 2.80 m, gyttja with a more or less coarse content (particularly wood fragments) was deposited. High values of organic material record an increase in lake productivity. From 2.80 m, the lake had been infilled with sediments and carr peat. The most common carr species were *Alnus*, *Betula*, *Salix*, sedges, grasses and herbs, such as *Filipendula*, *Lysimachia*, Ranunculaceae and *Equisetum* were also common. At 1,25m to the top fen peat replaced carr peat.

2. Reconstruction of the vegetation development and climate change

>14,700 cal. years BP

During this period the vegetation in the area had an open structure. The vegetation was a mosaic with areas dominated by low shrubs, such as *Salix*, *Juniperus*, and *Betula*, and areas dominated by grasses, sedges and herbs, such as *Artemisia*, Chenopodiaceae, *Helianthemum*, Asteraceae, Caryophyllaceae and Ranunculaceae. With *Pinus* percentages represented around 50 % it is difficult to interpret these values as indicating a local presence. *Pinus* is known as a species which produces enormous amounts of pollen, which can be, transported very long distances. Usually percentages values above 20 % can be attributed to a local presence (Huntley, 1983). Most likely, *Pinus* had a regional presence in valleys at lower altitudes with favorable microclimate. Scattered areas with little or no-vegetation around the site. Low pollen concentration and minerogenic sediments indicate this fact. Strong winds and slope erosion during the end of the Full Glacial period (GS-2) caused the transfer of large amounts of allochthonous material into the crater (Fig. 2).

14,700–14,050 cal. years BP

During this period open boreal forest dominated by *Pinus* and *Betula* expanded. Open vegetation with shrubs (mainly *Salix* and *Juniperus*) and herbs, such as Poaceae, Cyperaceae, *Artemisia*, Chenopodiaceae, Asteraceae and Ranunculaceae were also common in the area. *Pinus* has its highest values during this period. This certainly must indicate that *Pinus* expanded regionally, as well as locally, simultaneously as the open vegetation diminished. *Alnus* was common around the basin. The development of a denser vegetation cover corresponds with a warm event at the beginning of the Late Glacial (GI-1e in the GRIP ice core stratigraphy).

14,050–13,800 cal. years BP

There is strong evidence for a development of a rather open forest type during this period. *Pinus* percentages had become lower at the same time as herb pollen values have increased. Very high pollen values of *Alnus* and *Betula* can most likely be assigned to local occurrences in the vicinity of the basin. The increase of minerogenic input into the basin may be the effect of a reduction in vegetation cover as a response to a colder climate (GL-1d, in the GRIP ice core stratigraphy).

13,800–12,600 cal. years BP

The vegetation during this period was dominated by coniferous trees, such as *Pinus* and *Picea*, but, deciduous trees, such as *Betula*, *Ulmus* and *Alnus* were also abundant. *Picea* is recorded by pollen values with a maximum of around 15 %, indicating it was an important component in the forests. Knowing that values of 20–30 % are usually suggesting a local presence (Tzedakis, 1993). *Ulmus* has a short and rapid appearance, and reaches maximum values of 5.3 %. *Ulmus* pollen percentage above 2 % is commonly attributed as indicative for a local presence (Huntley, 1983). *Ulmus* and *Picea* showed rapid expansions that could suggest that their glacial refugia was close to the studied area. Intriguing is the presence of scattered or single pollen grains of other deciduous trees, such as *Quercus*, *Fraxinus* and *Tilia*. Most likely their presence is due to long distance transportation. Areas with open vegetation were present in the region, but were probably confined to higher elevations. The expansion of tree vegetation likely corresponds to a warmer climatic phase, which in time corresponds approximately to GI-1c – GI-1a in the GRIP event stratigraphy.

12,600–11,500 cal. years BP

This period is characterized by a re-expansion of open vegetation communities around the basin. The open vegetation was mainly dominated by herb species, such as Poaceae, *Artemisia*, Chenopodiaceae, Asteraceae Tubuliflorae, Asteraceae Liguliflorae and *Helianthemum*, and low shrubs, such as *Juniperus* and *Salix*. Most likely, *Betula* individuals also occurred. Un-vegetated areas and unstable ground were probably widespread in the area. The forest around this area was reduced by the colder climate, some trees probably still occurred in the region, at lower altitude but, only on favourable sites in valleys.

The high percentages for herbaceous taxa, particularly *Artemisia* and Chenopodiaceae, the reduction of trees and increased accumulation of inorganic material, are a clear signal for the cool phase known as GS-1 (Younger Dryas).

11,500–10,700 cal. years BP

At the Late Glacial/Holocene transition *Betula* and *Pinus* expanded and formed open forests. *Ulmus* and *Picea* probably had scattered occurrences at the beginning of this period, but at ca. 11,250 cal. years BP they rapidly expanded becoming the dominant trees in the forest canopy. The sudden and large increase of *Ulmus* (from scattered pollen grains to values around 40 %) may indicate it had survived during the Late Glacial in the close vicinity from where it could rapidly expand during the early Holocene. Previous palynological records from Romania

show similar patterns of vegetation development in early Holocene (Farcas et al., 1999; Buz, 1999). Approximately 100 km to the west from our study area, at Bartoliget Marsh site (Hungary), strong reduction of coniferous trees and expansion of deciduous woodland marked the Late Glacial/Holocene transition. *Tilia* dominated, followed by an expansion of *Quercus*. The evolution in Hungary it is very different from our findings. This situation was probably caused by differences in altitude and regional climate. The high values of herbs and shrubs pollen types indicate areas with open vegetation still occurred in the region, but were confined to higher altitudes. Between 11,250–10,700 cal. years BP the density of *Ulmus* woodland increased while herb pollen values significantly decreased.

10,700-8,600 cal. years BP

From 10,700 cal. years BP onwards, the forest in the area was of a species-rich nemoral type. *Ulmus* was the dominant tree, but *Quercus*, *Fraxinus* and *Tilia* had also become established and started to expand in the region. The low but continuous presence of *Acer* pollen grains suggests it occurred locally. Around 10,750 cal. years BP *Corylus* had a slight presence in region. Later *Corylus* it significantly expanded, simultaneously *Ulmus* was reduced. Over a period of 1,300 years a mixed deciduous forest type developed, where *Ulmus* and *Corylus* were dominant species. Around 9,300 cal. years BP *Corylus* reached its maximum abundance in the forests.

8,600-4,800 cal. years BP

Around 8,600 cal. years BP an important change in the forest composition is indicated by a significant expansion of *Picea* at the expense of *Corylus*. For a short period (around 250 years) *Picea* had high percentages mirrored by lower values for *Corylus*. Most of the broad leaves trees, such as *Tilia*, *Quercus* and *Fraxinus*, show slight declines. Only *Ulmus* has more or less constant frequencies throughout this period.

The regular presence of *Acer* pollen grains, suggests that locally it was probably common. *Fagus* pollen grains occur regularly from around 8,100 cal. years BP. *Fagus* also shows an isolated peak value around 7,900 cal. years BP. At this point, *Fagus* has percentages that are high enough to indicate a local presence. However, *Fagus* seems to have failed to establish a larger population in the area, and appears to have diminished quite soon. The pteridophytes are well represented in the pollen samples, and often clumps of spores, or sporangia with attached spores are found. Most likely, pteridophytes were important in the field layer of the forest. Few areas with open vegetation in the region existed.

4,800-3,300 cal. years BP

During this period, a dense mixed forest type with many tree and shrub taxa, such as *Quercus*, *Ulmus*, *Fraxinus*, *Acer*, *Carpinus*, *Fagus*, *Tilia*, and *Corylus* occurred in the area. *Corylus* and *Picea* were still dominant in the forest, however *Fagus* and *Carpinus* had become established and had started to expand in forest.

Carpinus seems to have established and expanded at the same time as *Fagus*. Around 3,300 cal. years BP *Fagus* became the dominant tree in the canopy. This development is in good agreement with Lupsa's study (1980). Pop (1943) also found an early and abundant presence of *Fagus* in the Oas-Maramures area. He suggested that this area could have supported glacial refugia. Our pollen data does not show a presence of *Fagus* during the Late glacial, but its presence in this area is recorded earlier than in other regions of Romania. The appearance of *Carpinus* happened however, later in study area, when compared to other regions in Romania. In northeastern Romania *Carpinus* expanded around 5,500 cal. years BP. In the southeastern part of the country (close to the Black Sea) it started to expand at the beginning of the Holocene. Single or scattered pollen grains of *Plantago lanceolata*, *Rumex acetosa/R. acetosella* and *Urtica* may be interpreted as signs of a local human influence on the vegetation.

3,300-2,200 cal. years BP.

The composition and structure of the forest changed dramatically during this period. *Fagus* was the dominant tree in the forest, but *Carpinus* and *Quercus* were also common. *Quercus* and *Carpinus* most likely dominated the woodlands at lower altitudes, while *Fagus* occurred at higher elevations together with *Picea*. The presence of *Ulmus*, *Tilia* and *Fraxinus* were strongly reduced and they never attained higher values again. Most likely they were rare in the forest comparing with today situation. *Corylus* occurred with low values in the shrub layer. The regularly presence of *Abies* pollen grains in the upper part of period may suggest that *Abies* occurred in the region, but it was certainly not present near the study site. Previous investigations by Pop (1943) showed that *Abies* had minimum poor presence during the period whereas *Fagus* had maximum values. He postulated that the *Fagus-Abies* community is new, and has originated during the late Holocene. During the Quaternary period, the dynamics of these taxa have followed different pathway. The field layer in the forest may have been dominated by pteridophytes.

2,200-0 cal. years BP

During the last 2,200 years, the forest in the area was dense and dominated by *Fagus*, in addition but some *Quercus* and *Carpinus* individuals did occur. The *Fagus* dominance is extraordinary, its pollen percentages reaching values up to 75%. This dominance reduced the species diversity in the forest, most of the nemoral trees, such as *Ulmus*, *Tilia* and *Fraxinus* were out-competed by *Fagus*.

From 300 cal. years BP to the present small change in the forest occurred, as *Carpinus* became less common. *Corylus* shows a slightly increasing trend during the last few hundred years, and was probably favoured by human influence. This is the first period where the pollen data indicate a significant influence of humans in area. For instance, the first evidence of cultivated cereals, such as *Hordeum* and *Secale*, appear however, with very low pollen values. There are other indicators of human influence. For example, pollen grains indicating pasture and fallow land, i.e. *Juglans-type*, Poaceae undiff. >40µm, *Plantago lanceolata*, Chenopodiaceae, *Rumex acetosa/R. acetosella*, *Artemisia*, *Cannabis-type* and *Urtica*. Some of these pollen types have appeared earlier in our site but, only during this period does their

percentages show increasing trends. Increases of Poaceae <40 µm and *Artemisia* in the region may indicate areas with expanded open vegetation. These open areas were possibly used for grazing. Today, the open vegetation, including the forest, is successfully used for grazing. The surface vegetation of the mires dominated by Poaceae (particularly *Molinia* and *Glyceria*), are used by farmers in order to provide hay to feed domestic animals.

CONCLUSIONS

The multidisciplinary analyses of Steregoiu shows that the sediments in the former crater lake cover a time span of over 14,700 years (Fig. 4).

The bottom sediments, which were deposited > 14,700 cal. years BP, belong to the Last Glacial Maximum. The Last Glacial Maximum is equivalent with GS-2 in the GRIP ice core event stratigraphy (Björck et al., 1998; Walker et al., 1999). These sediments were minerogenic and poor in pollen. Most of the pollen deposited during this time belongs to the herbs, grasses, sedge, and shrubs, such as *Artemisia*, Chenopodiaceae, Asteraceae, Ranunculaceae, *Salix*, *Juniperus* and *Ephedra*. *Pinus* and *Betula* have fairly high percentages of pollen. These high percentages indicate, *Pinus* and *Betula* but were present in the area but most likely occurred in lower altitudes.

The climatic events during the Late Glacial (14.700-11.500 cal. years BP) are well expressed in the vegetation development. Successions of cold (GI-1d, GS-1) and warm episodes (GI-1e, GI-1c, GI-1a) have had an immediate response in vegetation cover and lithostratigraphy. During cold phases, herbs, grasses, sedges and low shrubs dominated areas of widespread open vegetation. During climatically more favorable phases, this vegetation type was replaced by open woodlands containing *Betula*, *Pinus*, *Picea* and *Alnus*.

Open forest with *Betula*, *Picea*, *Pinus* and *Ulmus* developed around the site during the warm phase (GI-1c-1a), indicated in the GRIP ice core event stratigraphy between 13,800-12,950 cal. years BP. The appearance of *Ulmus* during this period is remarkable (it reaches a maximum of c. 5 %). Most likely, *Ulmus* must have been present on glacial refugia not far from the study area, otherwise, it could not have responded quickly to the warmer conditions. Besides *Ulmus*, *Picea*, *Betula* and *Alnus* must have been surviving in the area during the Late Glacial. For the same period high percentages of *Picea* were recorded by Wohlfarth et al. (2001). In addition, Wohlfarth's study include macrofossils that clearly shows the presence of *Picea*, *Pinus* (*P. sylvestris*, *P. mugo*, *P. cembra*), *Populus*, *Larix*, *Betula*, *Juniperus* and *Salix*. Glacial refugia for *Picea* (at 11,140 BP) is also confirmed by Farcas et al. (1999) in their studies of high altitude sites in the northeast and southwest of the Carpathians Mountains. The cooling signal of the GS-1 (Younger Dryas), between 12,950-11,500 cal. years BP, is clearly recorded in our sediments. The cooling is expressed by a reduction in the tree vegetation, which was eventually replaced by herbs, sedges, and shrubs, particularly *Artemisia*, Chenopodiaceae and Poaceae, respectively.

At the Late Glacial/Holocene transition (around 11,500 cal. years BP), *Betula* and *Alnus* expand slightly later, and *Picea* and *Ulmus* became established. At 11,250 cal. years BP, *Ulmus* became the dominant trees in the forest. The rapid

response of *Ulmus* to the climatic amelioration at the transition to the Holocene suggests that it must have survived on a refugia not very far from the study site. Slightly later, deciduous trees, such as *Quercus*, *Fraxinus* and *Tilia* appeared and expanded. Only at 10,500 cal. years BP they had they become abundant in the forest. From 10,500 cal. years BP onward, a dense mixed forest containing *Ulmus*, *Quercus*, *Tilia*, *Fraxinus*, *Picea* a few *Corylus* and *Acer* occurred in the area. *Corylus* expanded further in the area around 9,300 cal. years BP. At 8,600 cal. years BP *Picea* had become co-dominant with *Corylus*, when the broad leaves species were considerably reduced in the forest. This type of vegetation dominated the area until 4,800 cal. years BP when *Fagus* and *Carpinus* became established and rapidly expanded in the local forest.

The regular presence of *Fagus* pollen grains start around 8,000 cal years BP, but it was first at c. 4,800 cal. year BP, *Fagus* suddenly became widespread in the local forest. Around 4,000 cal. years BP, *Fagus* became the dominant tree in the canopy. *Fagus* has held this dominance until the present day. The presence of *Fagus* is recorded earlier in other regions of country. For instance, it started to appear around 3,000 years ago in the north-eastern and south-western Carpathians Mountains (Farcas, 1999). *Carpinus* instead, seems to have occurred later (around 4,700 cal. years BP) in the study site, than in other parts of Romania. Expansion of *Carpinus* occurred around 5,500 cal. years B.P in the north-eastern part of Romania, and in the south-eastern part (close to the Black Sea) at the beginning of the Holocene. *Abies* only occurred between 3,000-1,500 cal. years BP, indicated by scattered pollen grains.

Signs of human influence on the local vegetation are strong only during the last period from about 300 cal. years BP. At this time the grazing pressure seems to have increased and agriculture may have expanded in the Talna Valley below 500 m a.s.l. Several pollen grains of cultivated species are found during this period, but they most likely originated from arable fields on lower altitudes. An introduction of forest grazing in the area may have occurred around 1000 cal. years BP, as *Plantago lanceolata* starts to occur regularly around this time.

A PALEOECOLOGICAL RECONSTRUCTION OF THE LATE GLACIAL AND HOLOCENE ...

Years (cal.BP)	Sediments	Sedimentary units	Vegetation history	Periods
0	Fen peat OM 70-98% SIRM & Susceptibility very low.	10	Introduction of forest grazing Dominance of <i>Fagus</i> forest	Subatlantic
1000			Slightly increase of humans indicators, particularly <i>Juglans</i> -type, Poaceae undiff > 40mm, <i>Plantago lanceolata</i> Chenopodiaceae, <i>Rumex acetosa</i> / <i>R. acetosella</i> , <i>Artemisia Cannabis</i> -type and <i>Urtica</i> . Scattered pollen grains of <i>Cerealia</i> as <i>Hordeum</i> and <i>Secale</i> .	
2000			Expansion of <i>Fagus</i> woodland. <i>Carpinus</i> and <i>Quercus</i> were still common. Strong reduction of the <i>Ulmus</i> , <i>Tilia</i> and <i>Fraxinus</i> . Slightly presence of <i>Abies</i> and <i>Acer</i> .	Subboreal
3000	Dense and highly diverse forest with <i>Corylus</i> , <i>Picea</i> , <i>Ulmus</i> and <i>Quercus</i> . <i>Fagus</i> and <i>Carpinus</i> have established and expanded			
4000	Carr peat OM 80-95% SIRM & Susceptibility very low	9	Expansion of <i>Corylus</i> and <i>Picea</i> in the woodland along with reduction of all broad leaves tree. Appearance of new taxa (e.g., <i>Fagus</i>)	Boreal
5000				
6000	Detritus Gytja OM 20-60% SIRM & Susceptibility low with small peak at the beginning of the unit	7	Rich and dense deciduous forest dominated by <i>Ulmus</i> <i>Quercus</i> , <i>Tilia</i> , <i>Fraxinus</i> were also common. Increasing of <i>Corylus</i> and <i>Picea</i> by the end of this period.	Preboreal
7000				
8000	Gyttja SIRM & Susceptibility high	6; 5	Expansion of <i>Betula</i> and <i>Pinus</i> , followed by <i>Ulmus</i> Further appearance of deciduous trees (e.g., <i>Quercus</i> , <i>Tilia</i> and <i>Fraxinus</i>).	GS-1
9000				
10000	Clayey Gytja OM 16-18% SIRM & Susceptibility high	4	Re-expansion of steppe community with dominance of <i>Artemisia</i> , Chenopodiaceae, low shrubs as <i>Juniperus</i> and <i>Salix</i> . Tree are represented by <i>Pinus</i> , <i>Betula</i> and <i>Alnus</i> .	GL-1a-c
11000				
12000	Gyttja Clay OM low SIRM & susceptibility high	2	Expansion of the boreal forest with <i>Pinus</i> , <i>Picea</i> and <i>Betula</i> and also of <i>Ulmus</i> .	GL-1d
13000				
14000	Silty Clay & Sandy Silt OM low SIRM & Susceptibility very high	1	Increase of open vegetation along with decrease of <i>Pinus</i> .	GL-1e
14700				
?			Open vegetation with <i>Salix</i> , <i>Juniperus</i> sedges, grasses and herbs as <i>Artemisia</i> and Chenopodiaceae. Scattered <i>Pinus</i> individuals.	GS-2

Fig. 4. Summary of the vegetation and environmental history of Steregoiu, based on sediment parameters, pollen and calibrated radiocarbon data (OM: organic matter; SIRM: saturation isothermal remanent magnetization).

REFERENCES

- Bengtsson, L., Enell, M. (1986), *Chemical analysis*. In: Berglund, B.E. (Ed.), *Handbook of Holocene Palaeoecology and Palaeohydrology*, Wiley & Sons, Chichester, pp. 423-454.
- Berglund, B. E., Ralska-Jasiewiczowa, M. (1986) Pollen analysis and pollen diagrams. In: Berglund, B. E. (Ed.), *Handbook of Holocene Palaeoecology and Palaeohydrology*, Wiley & Sons, Chichester, p. 455-484.
- Björck, S., Walker, M. J. C., Cwynar, L. C., Johnsen, S., Knudsen, K.-L., Lowe, J. J., Wohlfarth, B., INTIMATE members (1998), *An event stratigraphy for the Last Termination in the North Atlantic region based on the Greenland ice-core record: a proposal by the INTIMATE group*. *Journal of Quaternary Science*, 13, p. 283-292.
- Björckman, L., Feurdean, A., Cinthio, K., Possnert, G. (2001), *Late Glacial and early Holocene vegetation development in the Gutaiului Mountains, NW Romania*, *Quaternary Science Reviews* (in press).
- Buz, Z. (1999), *Cercetări fitosociologice și palinologice în zona Sovata-Praid-Dealul*. Ed. Casa Cartii de Știință, Cluj Napoca.
- Diaconeasa, B., Farcas, S. (1996), *Stejărișurile amestecate, evoluția și dinamica lor în Tardiglaciul și Holocenul din România*. Universitatea "Babeș-Bolyai", Cluj-Napoca, *Contributii Botanice*, 1995–1996, p. 103-115.
- Faegri, K., Iversen, J. (1989), *Textbook of pollen analyses* (4th ed., revised by Faegri, K., Kaland, P.E., Krzywinski, K.). John Wiley, Chichester.
- Fărcaș, S., de Beaulieu, J.L., Reille, M., Coldea, G., Diaconeasa, B., Goslar, T., Jull, T. (1999), *First ¹⁴C datings of Late Glacial and Holocene pollen sequences from the Romanian Carpathians*. *Comptes Rendues de l'Académie des Sciences de Paris (Sciences de la Vie et de la Terre)*, 322, p. 799-807.
- Fărcaș, S. (2001), *Pollen data on the peat bogs in the Căliman Mountains*. Doctoral thesis abstract.
- Firbas, F. (1949), *Spät- und nacheiszeitliche Waldgeschichte Mitteleuropas nördlich der Alpen*. Band I. Fischer, Jena, p. 480.
- Firbas, F. (1952), *Spät- und nacheiszeitliche Waldgeschichte Mitteleuropas nördlich der Alpen*. Band II. Fischer, Jena, p. 256.
- Grimm, E. (1987), *Coniss: a FORTRAN 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares*. *Computers and Geosciences* 13: 13-37.
- Grimm, E. (1991), *Tilia 1.12, Tilia Graph 1.18*. Springfield, Illinois State Museum, Research and Collection Center.
- Huntley, B., Birks, H. J. B. (1983), *An atlas of past and present pollen maps for Europe: 0–13.000 years ago*. Cambridge University Press, Cambridge, p. 667.
- Isvan, D., Popescu, S., Pop, I. (1990), *Munții Gutii*. Ed. Sport Turism, București, p. 7-30.
- Lupsa, V. (1980), *Evoluția postglaciara a ecosistemelor forestiere din "Tara Oașului" (Jud. Satu Mare)*. *Contribuții botanice* 5, p. 63-68.
- Mac, I., Budai, C. (1992), *Munții Oaș-Gutâi-Țibleș*. Casa editorială pentru turism și cultură "Abedona". București, p. 7-45.

A PALEOECOLOGICAL RECONSTRUCTION OF THE LATE GLACIAL AND HOLOCENE ...

- Moore, P.D., Webb, J.A., Collinson, M.E. (1991), *Pollen analysis* (2nd ed). Blackwell, Oxford, p. 216.
- Onac, B. P., Lauritzen, S. E. (1996), *The climate of the last 150,000 years recorded in speleothems: preliminary results from north-western Romania*. Theoretical and Applied Karstology 9: 9-21.
- Onac, B. P., Constantin, S., Lauritzen, S. E., Lundberg, J. (2001), *Isotope-climate record in a Holocene stalagmite from Ursilor Cave (Romania)*. Journal of Quaternary Science (in review).
- Pop, E. (1932), *Contribuții la istoria vegetației cuaternare din Transilvania*. Bul. Grad. Bot. Cluj, 12, p. 29-102.
- Pop, E. (1942) *Contribuții la istoria pădurilor din nordul Transilvaniei*. Bul. Grad. Bot., Cluj, 9, 3-4, p. 81-210.
- Pop, E., Boscaiu, N., Ratiu, F., Diaconeasa B. (1960), *Mlaștinile de turbă din R.P.R.* Ed. Academiei Romane, București, p. 511.
- Reille, M. (1992), *Pollen et spores d'Europe et d'Afrique du nord*. Laboratoire de botanique historique et palynologie, Marseille, p. 520.
- Stuiver, M., Reimer, P. J., Bard, E., Beck, J. W., Burr, G. S., Hughen, K. A., Kromer, B., McCormac, G., van der Plicht, J., Spurk, M. (1998), *INTCAL98 Radiocarbon age calibration, 24,000-0 cal BP*. Radiocarbon, 40, p. 1041-1083.
- Tzedakis, P.C. (1993), *Long-term tree population in northwest Greece through multiple Quaternary climate cycles*. Nature, 364, p. 437-440.
- Walden, J., Oldfield, F. & Smith, J.P. (editors) (1999), *Environmental Magnetism: a practical guide*. Technical Guide, No. 6, p. 35-88. Quaternary Research Association, London.
- Walker, M. J. C., Björck, S., Lowe, J. J., Cwynar, L. C., Johnsen, S., Knudsen, K.-L., Wohlfarth, B., INTIMATE group (1999), *Isotopic 'events' in the GRIP ice core: a stratotype for the Late Pleistocene*. Quaternary Science Reviews, 18, p. 1143-1150.
- Willis, K. J., Sümegei, P., Braun, M., Tóth, A. (1995), *The late Quaternary environmental history of Bátorliget, N.E. Hungary. Palaeogeography, Palaeoclimatology, Palaeoecology*, 118, p. 25-47.
- Wohlfarth, B., Hannon, G., Feurdean, A., Ghergari, L., Onac, B.P., Posnert, G. (2001), *Reconstruction of climatic and environmental changes in NW Romania during the early part of the last deglaciation (15,000–13,600 cal years BP)*. Quaternary Science Reviews (in press).
- Woldstedt, P. (1958), *Das Eiszeitalter*, 2. Ferdinand Enke Verlag, Stuttgart, pp. 438.