

USF Patents

---

June 2013

## Antibiotic-bound poly(caprolactone) polymer

Edward Turos

Michelle Leslie

Follow this and additional works at: [https://digitalcommons.usf.edu/usf\\_patents](https://digitalcommons.usf.edu/usf_patents)

---

### Recommended Citation

Turos, Edward and Leslie, Michelle, "Antibiotic-bound poly(caprolactone) polymer" (2013). *USF Patents*. 226.

[https://digitalcommons.usf.edu/usf\\_patents/226](https://digitalcommons.usf.edu/usf_patents/226)

This Patent is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Patents by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact [digitalcommons@usf.edu](mailto:digitalcommons@usf.edu).



US008470958B1

(12) **United States Patent**  
**Turos et al.**

(10) **Patent No.:** **US 8,470,958 B1**  
(45) **Date of Patent:** **Jun. 25, 2013**

(54) **ANTIBIOTIC-BOUND  
POLY(CAPROLACTONE) POLYMER**

(75) Inventors: **Edward Turos**, Wesley Chapel, FL  
(US); **Michelle Leslie**, Tampa, FL (US)

(73) Assignee: **University of South Florida**, Tampa, FL  
(US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/367,965**

(22) Filed: **Feb. 7, 2012**

**Related U.S. Application Data**

(62) Division of application No. 11/746,704, filed on May  
10, 2007, now Pat. No. 8,110,678.

(60) Provisional application No. 60/747,061, filed on May  
11, 2006.

(51) **Int. Cl.**  
**C08G 63/08** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **528/318**; 525/386; 525/415; 549/266;  
549/271; 549/342

(58) **Field of Classification Search**  
USPC ..... 525/386, 415; 528/318; 549/266,  
549/271, 342  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

3,477,997 A \* 11/1969 Gumboldt et al. .... 528/319  
5,025,025 A 6/1991 Bhagwat et al.

5,629,381 A 5/1997 Isozaki  
5,801,224 A \* 9/1998 Narayan et al. .... 528/357  
5,917,067 A 6/1999 Cotarca et al.  
6,967,208 B2 11/2005 Pinto et al.  
7,091,219 B2 8/2006 Freskos et al.

**OTHER PUBLICATIONS**

USPTO strcucure search; Feb. 2013.\*

Turos, E., et al. 2005. N-Methylthio B-Lactum Antibacterials: Effects  
of the C3/C4 Ring Substituents on Anti-MRSA Activity. Bioorganic  
& Medicinal Chemistry vol. 13. p. 6289-6308.

Turos, E., et al., 2007. Antibiotic-Conjugated Polyacrylate  
Nanoparticles: New Opportunities for Development of Anti-MRSA  
Agents. Bioorganic and Medicinal Chemistry Letters. vol. 17 pp.  
53-56.

Sinha, V.R., et al., 2004. Poly-e-Caprolactone Microspheres and  
Nanospheres: an Overview. International Journal of Pharmaceutics  
vol. 278, pp. 1-23.

Trollsas, M, et al., "Hydrophilic Aliphatic Polyesters: Design, Syn-  
thesis, and Ring-Opening Polymerization of Functional Cyclic  
Esters" Marcomolecules, 2000, 33,4619-4627.

Woster, Journal of Medicinal Chemistry (1989), 32(6), 1300-1307.  
Adger, et al., J. Chem Soc., Chem. Commun., 1995, 1563-1564.

\* cited by examiner

*Primary Examiner* — Gregory Listvoyb

(74) *Attorney, Agent, or Firm* — Robert J. Varkonyi; Smith  
& Hopen, P.A.

(57) **ABSTRACT**

This invention is the design and synthesis of a caprolactone  
monomer which bears a pendant protected carboxyl group.  
This monomer has been copolymerized with caprolactone in  
varying ratios. After polymerization, the protecting group can  
be removed and an antibiotic can be attached as a new pendant  
group. The bioactivity of the antibiotic-bound poly(caprolac-  
tone) polymer is described.

**5 Claims, 4 Drawing Sheets**






FIG. 4




FIG. 5A




FIG. 5B




FIG. 6







FIG. 7



Reagents and conditions: (a)  $\text{PPh}_3$ ,  $\text{C}_6\text{H}_6$ , 12h, rt; (b) 20%  $\text{NaOH}_{(\text{aq})}$ , 5h, rt.

FIG. 8



Drug mass	ZOI
20 ug	11mm
50 ug	20mm
100 ug	20mm

FIG. 9





FIG. 10



1

# ANTIBIOTIC-BOUND POLY(CAPROLACTONE) POLYMER

## CROSS-REFERENCE TO RELATED APPLICATIONS

This divisional application claims priority to U.S. Non-Provisional application Ser. No. 11/746,704 filed on May 10, 2007 and U.S. Provisional Application No. 60/747,061 filed on May 11, 2006, entitled "Antibiotic-Bound Poly(Caprolactone) Polymer".

## STATEMENT OF GOVERNMENT INTEREST

This invention was made with Government support under Grant No. RO1 A151351 awarded by the National Institutes of Health. The Government has certain rights in the invention.

## FIELD OF INVENTION

This invention relates to attaching one or more bioactive molecules to the same polymer, and also for attachment before or after polymerization

## BACKGROUND OF THE INVENTION

The delivery of water-insoluble drugs to targets within the human body is a challenge that presently places strict limitations on what drugs can be applied clinically. The need for methods which overcome this is of high priority in the development of new therapeutics for treatment of human disease.

The development of antibiotics for control of pathogenic bacteria has been of pressing need in this era of drug resistant infections. N-Methylthiolated  $\beta$ -lactams have been identified as a new family of antibacterial agents active against *Staphylococcus* bacteria, including methicillin-resistant *Staphylococcus aureus* (MRSA). (See Turos, E.; Konaklieva, M. I.; Ren, R. X. F.; Shi, H.; Gonzalez, J.; Dickey, S.; Lim, D. *Tetrahedron* 2000, 56, 5571; Bart Heldreth, Timothy E. Long, Seyoung Jang, Suresh K. R. Guntireddygar, Edward Turos, Sonja Dickey, Daniel V. Lim, "N-Thiolated  $\beta$ -Lactam Antibacterials: Effects of the N-Organothio Substituent on anti-MRSA Activity," *Bioorganic and Medicinal Chemistry* 14, 3775-3784 (2006); and Edward Turos, Jeung-Yeop Shim, Yang Wang, Kerriann Greenhalgh, G. Suresh Kumar Reddy, Sonja Dickey, Daniel V. Lim, "Antibiotic-Conjugated Polyacrylate Nanoparticles: New Opportunities for Development of Anti-MRSA Agents," *Bioorganic and Medicinal Chemistry Letters* 16, in press (2006); which are incorporated herein by reference).

The compounds have also displayed promising anticancer properties. These lactams exert their growth inhibitory effects on bacteria through a mode of action that is distinctively different to that of other  $\beta$ -lactam antibiotics, and possess structure—activity patterns unlike those already mapped for other  $\beta$ -lactam antibacterials such as the penicillins. One of the major limitations in the potential application of these N-thiolated  $\beta$ -lactam compounds, however, is their exceedingly low water solubility.

Drug delivery vehicles such as liposomes and gold nanoparticles have been developed to improve bioavailability, efficacy, and specificity of pharmaceutical compounds, particularly for anticancer agents, but nanoparticles have received surprisingly little attention in the antibiotic and infectious disease area. Some of the few notable examples have included

2

antibiotic-encapsulated polymeric nanoparticles and liposomes, biodegradable nanospheres and surface-coated gold and silver nanoparticles.

## SUMMARY OF INVENTION

In one embodiment, the invention provides an effective drug delivery platform that would enhance the water solubility of the lactams, without sacrificing inherent bioactivity.


In another embodiment, the invention provides for the development of antibacterial polyacrylate nanoparticles based on well-precedented emulsion polymerization procedures.

This invention addresses this need, and demonstrates the use of antibiotic-bound poly (caprolactone) polymers as anti-infective materials for biomedical applications in the prevention of bacterial infections.

In a first embodiment, the invention includes a functionalized compound comprising at least one caprolactone ring with an appended functional group. A plurality of methylene groups act as a spacer between the lactone ring and the functional group. In a preferred embodiment, the functional group is an antibiotic, such as a N-thiolated  $\beta$ -lactam. The functional group is preferably covalently bonded.

In an alternate embodiment, a method is provided for producing the functionalized compound of the previous embodiment. The method includes providing a at least one caprolactone ring with a protecting group spaced apart from the caprolactone ring by a plurality of methylene groups. The protecting group is then cleaved from the caprolactone ring and replaced by bonding a functional group to the caprolactone ring. Illustrative reagents for cleaving the protecting group include 10% pd/C as a catalyst in the presence of  $H_{2(g)}$  and ethyl acetate (EtOAc). The protecting group is selected from the group consisting of alcohol protecting groups, amine protecting groups, carbonyl protecting groups and carboxyl protecting groups. In a preferred embodiment, the protecting group is selected from the group consisting of a benzyl ester and a tert-butyl ester. As with the previous embodiment, the functional group is an antibiotic such as a N-thiolated  $\beta$ -lactam.


In another embodiment, the invention includes the compound represented by the formula:



or a pharmaceutically acceptable salt or ester thereof, wherein is selected from the group consisting of water-insoluble drugs, antibiotics, lactams,  $\beta$ -lactams, N-thiolated  $\beta$ -lactams and protecting groups. The protecting group is selected from the group consisting of alcohol protecting groups, amine protecting groups, carbonyl protecting groups and carboxyl-protecting groups.


In yet another embodiment, the invention includes the compound represented by the formula:

3




or a pharmaceutically acceptable salt or ester thereof.

In another embodiment, the invention includes the compound represented by the formula:



or a pharmaceutically acceptable salt or ester thereof.

In still another embodiment, the invention includes the compound represented by the formula:



or a pharmaceutically acceptable salt or ester thereof.

#### BRIEF DESCRIPTION OF THE DRAWINGS

For a fuller understanding of the nature and objects of the invention, reference should be made to the following detailed description, taken in connection with the accompanying drawings, in which:

FIG. 1 shows the polymerization of a caprolactone monomer to polycaprolactone (PCL).

FIG. 2 shows the functionalized caprolactone polymer can be used for covalent binding of drug molecules.

FIG. 3 shows examples of functionalized caprolactones from the literature.

FIG. 4 shows the functionalized lactone of present invention.

FIG. 5A shows drug appendage before polymerization.

FIG. 5B shows drug appendage after polymerization.

FIG. 6 shows lactone resynthesis.

FIG. 7 shows ylide synthesis.

FIG. 8 is an image showing results of biological testing of the lactam-containing monomer vs. MSSA.

4

FIG. 9 demonstrates the anti-Bacillus activity of the  $\beta$ -lactam containing copolymer.


FIG. 10 shows the invention employing an alternate carboxyl protecting group.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

In the following detailed description of the preferred embodiments, reference is made to the accompanying drawings, which form a part hereof, and within which are shown by way of illustration specific embodiments by which the invention may be practiced. It is to be understood that other embodiments may be utilized and structural changes may be made without departing from the scope of the invention.

##### N-thiolated $\beta$ -lactams

New family of anti-MRSA and anti-Bacillus agents that have recently been reported by our laboratory. Extensive SAR studies have shown that changing the N-thioalkyl substituent has a large effect on the bioactivity and that changes at the other positions of the ring exert a more subtle effect. Recent research in our group has been focused on covalent attachment of these and other antibiotics to polymers for drug, delivery and for new biomaterials.



As used herein, the term "drug" to any therapeutic or prophylactic agent other than food which is used in the prevention, diagnosis, alleviation, treatment, or cure of disease in man or animal.

As used herein, the term "antibiotic" refers to any natural, synthetic, and semi-synthetic compound that has been identified as possessing antibacterial, antifungal, antiviral, or antiparasitic activity.

##### Polycaprolactones

Polycaprolactone (PCL) is a biodegradable polyester which can be prepared by ring opening polymerization of  $\epsilon$ -caprolactone using a catalyst such as stannous octanoate, as shown in FIG. 1. PCL is degraded by hydrolysis of its ester linkages in physiological conditions (such as in the human body) and is therefore useful as a biomaterial. PCL has been approved by the Food and Drug Administration (FDA) for use in the human body as (for example) a suture (sold under the brand name Monocryl<sup>TM</sup> or generically). In particular, PCL has been used in long term implantable devices, owing to its degradation which is relatively slow. (See V. R. Sinha, K. Bansal, R. Kaushik, R. Kumria and A. Trehan; Poly- $\epsilon$ -caprolactone microspheres and nanospheres: an overview, *International Journal of Pharmaceutics*, Volume 278, Issue 1, 18 Jun. 2004, Pages 1-23; which is incorporated herein by reference.)

The characteristics of PCL make it useful as a delivery mechanism for antibiotics. For example PCL is biodegradable (bulk hydrolysis of ester bonds), the byproducts of degradation are non-toxic (biocompatible), it is FDA approved and displays high permeability to many drugs. Therefore, the invention provides a functionalized caprolactone polymer can be used for covalent binding of drug molecules (FIG. 2).

The functionalized caprolactones of the prior art (Detrembleur et al *Macromolecules*, 2000, 33, 14-18 and Trollas et al *Macromolecules*, 2000, 33, 4619-4627) are shown in FIG. 3.


## 5

In contrast, the functionalized lactone **10** of present invention is shown in FIG. 4. Lactone **10** comprises at least one lactone ring **12**, functional group **16** and at least one spacer **14**. As it can be seen, functional group **16** is placed away from site of polymerization. Moreover, an additional methylene spacer **14** between functional group **16** and lactone ring **12** enhances further functionalization.

In another embodiment, the invention provides a method of producing an antibiotic-conjugated functionalized caprolactone (FIG. 5A and FIG. 5B). In Step 1 includes providing a caprolactone comprising lactone ring **12**, methylene spacer **14** and protecting group **18**. In Step 2, protecting group **18** is cleaved, preferably under mild conditions. Finally, in Step 3, drug of interest **20** is covalently bonded to the finished compound. FIG. 5A illustrates the method of appending a drug of interest to the functionalized lactone before polymerization. FIG. 5B illustrates the method of appending a drug of interest to the functionalized lactone after polymerization. Lactone retrosynthesis is shown in FIG. 6 and Ylide synthesis is shown in FIG. 7.

## EXAMPLE 1




The following represents an embodiment of the invention wherein 1,4-dioxaspiro[4.5]decan-8-one is used to synthesize a functionalized lactone bearing a pendent benzyl ester as the protecting group.



## 6

Reagents and conditions: (a) 2 eq. ylide,  $C_6H_6$ , 12 h, reflux; (b) 60 psi  $H_{2(g)}$ , cat. 10% Pd/C, MeOH, 12 h; (c) 3LiOH, MeOH, 12 h, rt; 1M HCl; (d)  $3K_2CO_3$ , 1.1 BnBr, MeCN, 12 h, reflux; (e) 70% AcOH, 12-24 h, rt; (f) 1.5mCPBA,  $CHCl_3$ , 3-5 h, reflux.

Next the functionalized lactone is coupled with an antibiotic, here N-thiolated  $\beta$ -lactam, after deprotection of benzyl ester. FIG. 8 shows the results of biological testing of the lactam-containing monomer vs. MSSA.




Reagents and conditions: (a) cat. 10% Pd/C, 60 psi  $H_{2(g)}$ , EtOAc, 12 h; (b) 1.5 eq. EDCI, cat. DMAP, dry  $CH_2Cl_2$ , 12 h, rt.



## EXAMPLE 2

In another embodiment, the invention provides a PCL derived from copolymerization of a functionalized lactone monomer with caprolactone. Copolymers containing 10%, 15%, 20%, 25% and 30% of the substituted lactone were prepared and characterized by TLC,  $^1H$ , NMR,  $^{13}C$  NMR and MALDI-TOF. The Copolymers displayed low molecular, between about 1000 and 4000.

7




Deprotection of the copolymer and coupling with the antibiotic,  $\beta$ -lactam, is shown below and was achieved using the following reagents and conditions: (a) cat. 10% Pd/C,  $H_{2(g)}$ , EtOAc, 12-24 h. (b) 15 EDCI, cat. DMAP, dry  $CH_2Cl_2$ , 12 h. rt. The antibiotic activity of the completed PCL against *Bacillus* is shown in FIG. 9.



8


## EXAMPLE 3

In yet another embodiment, the functionalized lactone comprises a carboxyl protecting group, as shown in FIG. 10. Ylide synthesis is shown below using the reagents and conditions: (a)  $PPh_3$ ,  $C_6H_6$ , 12 h, rt; (b) 20%  $NaOH_{(aq)}$ , 5 h, rt.




## EXAMPLE 4

The following demonstrates the synthesis of lactone bearing pendant tert-butyl ester using the reagents and conditions: (a) 1.5 eq. ylide  $C_6H_6$ , 12 h, reflux, (b)  $H_{2(g)}$ , cat. 10% Pd/C, EtOAc, 24 h; (c) 0.1 eq.  $I_2$ , dry acetone, 1 hr, rt; (d) 1.5mCPBA  $CHCl_3$ , 3 h, reflux.



Copolymerization with caprolactone was achieved as shown below. Copolymers containing 10%, and 20% of the substituted lactone were prepared and characterized by TLC,  $^1H$  NMR and  $^{13}C$  NMR.

9




Functional caprolactone monomers have been synthesized and characterized. The monomers were further copolymerized with caprolactone. Caprolactone monomers and polymers with covalently bound N-thiolated  $\beta$ -lactams have been prepared and shown to possess bioactivity against MSSA and *Bacillus* respectively.

It will be seen that the advantages set forth above, and those made apparent from the foregoing description, are efficiently attained and since certain changes may be made in the above construction without departing from the scope of the invention, it is intended that all matters contained in the foregoing description or shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

It is also to be understood that the following claims are intended to cover all of the generic and specific features of the invention herein described, and all statements of the scope of the invention which, as a matter of language, might be said to fall therebetween. Now that the invention has been described,

What is claimed is:


1. A homopolymer derived from polymerization of a single compound represented by the formula below:



wherein  $\text{R}_1$  is alkyl,  $\text{R}_2$  is an aryl,  $\text{R}_3$  is alkyl or aryl.


2. A homopolymer derived from polymerization of a single compound represented by the formula below:

10




wherein  $\text{R}_1$  is alkyl,  $\text{R}_2$  is an aryl.


3. A homopolymer derived from polymerization of a single compound represented by the formula below:



4. A homopolymer derived from polymerization of a single compound represented by the formula below:




5. A copolymer derived from polymerization of a plurality of compounds represented by the formulas below:



**11****12**

-continued



5

10

wherein R1 is alkyl, R2 is an aryl, R3 is alkyl or aryl.

\* \* \* \* \*