
USF Patents

July 2013

Effective treatment of esophageal adenocarcinoma using tricitiribine and related compounds

Jin Q. Cheng

Follow this and additional works at: https://digitalcommons.usf.edu/usf_patents

Recommended Citation

Cheng, Jin Q., "Effective treatment of esophageal adenocarcinoma using tricitiribine and related compounds" (2013). *USF Patents*. 230.

https://digitalcommons.usf.edu/usf_patents/230

This Patent is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Patents by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.



US008476241B2

(12) **United States Patent**
Cheng

(10) **Patent No.:** **US 8,476,241 B2**
(45) **Date of Patent:** ***Jul. 2, 2013**

(54) **EFFECTIVE TREATMENT OF ESOPHOGEAL
ADENOCARCINOMA USING TRICIRIBINE
AND RELATED COMPOUNDS**

(75) Inventor: **Jin Q. Cheng**, Tampa, FL (US)

(73) Assignee: **University of South Florida**, Tampa, FL
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **13/163,123**

(22) Filed: **Jun. 17, 2011**

(65) **Prior Publication Data**

US 2012/0202765 A1 Aug. 9, 2012

Related U.S. Application Data

(63) Continuation of application No. 12/206,468, filed on
Sep. 8, 2008, now Pat. No. 8,178,502.

(60) Provisional application No. 60/935,940, filed on Sep.
7, 2007.

(51) **Int. Cl.**
A01N 43/04 (2006.01)
A61K 31/70 (2006.01)

(52) **U.S. Cl.**
USPC **514/43; 514/42; 514/52**

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2006/0247188 A1 11/2006 Cheng et al.

OTHER PUBLICATIONS

Sagatys, Elizabeth, et al.: "Activation of the Serine/Threonine Protein
Kinase Akt During the Progression of Barret Neoplasia," Human
Pathology, vol. 38, pp. 1526-1531, 2007.
Office Action from Patrick T. Lewis in co-pending U.S. Appl. No.
10/113,927 mailed Sep. 29, 2010.

Primary Examiner — Patrick Lewis

(74) *Attorney, Agent, or Firm* — Fanelli Hagg & Kilger
PLLC

(57) **ABSTRACT**

The inventors have determined, contrary to the prior art and
experience, how to successfully use triciribine to treat eso-
phogeal adenocarcinoma by one or a combination of (i)
administering triciribine only to patients which according to
a diagnostic test described below, exhibit enhanced sensitiv-
ity to the drug; (ii) use of a described dosage level that mini-
mizes the toxicity of the drug but yet still exhibits efficacy; or
(iii) use of a described dosage regimen that minimizes the
toxicity of the drug.

20 Claims, 23 Drawing Sheets




FIG. 1A




FIG. 1C







FIG. 1B

FIG. 1D








FIG. 3A




FIG. 3B-1




FIG. 3B-4




FIG. 3B-2




FIG. 3B-5




FIG. 3B-3




FIG. 3B-6




FIG. 3B-7




FIG. 3B-9




FIG. 3B-8




FIG. 3B-10




FIG. 3B-11




FIG. 3B-12




FIG. 3C-1




FIG. 3C-2




FIG. 3C-3




FIG. 3C-4




FIG. 3C-5

FIG. 3C-6




FIG. 4A




FIG. 4C




FIG. 4E




FIG. 4D




FIG. 4B-1




FIG. 4B-2




FIG. 4B-3




FIG. 4B-4




FIG. 4B-5

PIP3:	-	+	+	+	+
PDK1:	-	+	+	+	+
Akt:	+	+	+	+	+
API-2:	-	-	-	25	10 (μ M)



FIG. 5

atgaacgaacgtagccattgtgaaggagggtctggctgcacaaacagaggggaatatattaa
1 -----> 60
tacttgcctgcctggtaaacacttccctcccgacacgctgttttgcctcccttataataatt
a M N D V A I V K E S W L H K R G E Y F K -
acctggcgggcacgcctacttccctccctaaagaaacgtggcacctttatttggctacacggaa
61 -----> 120
tggacacgacggtgcgatgaaggagggagtttttgcctacgctggaaataacgaagtctcctt
a T W R F R Y F L L K N D G T F I S Y K E -
cggctccaggatgttggtacagcagagctcccccctccacaccttctccagtggacacatgc
121 -----> 180
gcggagctccacacactagtcgctctcaggggtgagtgttggaaagctccacgctgttacc
a R F Q D V D Q R E S F L N N F S V A Q C -
PacI
cagctgatgaagacagcagcgggcacagggcccaacacctttatccctccgctgcctggctgg
181 -----> 240
gtcgactacttctgtctccgcgggttcgggttctggaaatagtaggcacgggcgctccac
a Q L H K T E R F S P N T F I I R C L Q W -
accacagtcatttgagcgcaccttccctgttagaaacgcctgaggagcgggaagatgggac
241 -----> 300
tgggttcagtaactcgcgtgggaaggtacatcttttgcggactccctcgccttcttaccgg
a T T V I E R T F K V E T F S S R E E W A -
acggcccttcagactgttggcagatggactccagaggccgggaagagagcagctggacttc
301 -----> 360
tggcggtaagttctgacacgggtacctgagttctccgtccctctctctctgtacctgaag
a T A I Q T V A D G L K R Q E K K T M D P -
cgatcaggctacccagtgacacactcaggggtgaagagatggaggtgtccctggccacg
361 -----> 420
gctagtcagggtgggtacatgttgagtaacggactctctctacctccacagggacgggttc
a R S G S F S D N S G A E E M E V S L A K -

FIG. 6A

ccaaagcacacgtgtgacccatgaacagagtttgggtacactgaaactactgggcaaggggcaccc
421 -----+-----+-----+-----+-----+-----+-----+ 480
gggttcgtgggcacacgtggtacttgcctcaaaactcatggactttgatgacacgttcccggtg
a F K H R V T M N S F E Y L K L L G K G Y ~
cttgggaagtgatctctgtgtaaaagagagaggccacaggcccgctactatgcaatgaagatc
481 -----+-----+-----+-----+-----+-----+ 540
aaacacatttccactaagaacattttctctctccggtgtccggcgatgatacggtaacttctctag
a F G K V I L V K E K A T G R Y Y A M K I ~
ctcaagaaggaggtctatgtgtcccaaggatgaggttgcccaacagcttactgaaaacggt
541 -----+-----+-----+-----+-----+-----+ 600
gagttcttccctccagtagcagcggttctctactccaaacgggtgtggaatgacttttggca
a L E K E V I V A K D E V A N T L Y E N S ~
PstI
|
gtcctgcagaactctaggcatcccttcccttaaggccctcaagtactcattccagacccac
601 -----+-----+-----+-----+-----+ 660
caggacgtctttgagatccgtagggaagggaatgcggggagttccatgagtaagggtctcggtg
a V L Q N S R N P F L T A L K Y S F Q T N ~
KhaI
|
gaacgctctctgcttctgtctctggagtatgccaacggggggagagctcttcttccactgtct
661 -----+-----+-----+-----+-----+ 720
ctgggggagacgaacagtaactcatcaggttgcaccccgctcggagagagaggtggacaga
a D R L C F V M E Y A N G G E L F F N L S ~
cgaagcgcggtgttctctcgaaggacggggcccgcttcttatggtgcggagattgtgtctgca
721 -----+-----+-----+-----+-----+ 780
gctctcgcgcacaaaggaggtctcttggcccgggcgagataccacgctctctacacagacgg
a R E R V F S E D R A R F Y G A E I V S A ~
ctggactacttgcactccgagaagacgtggtgtacccgggaactgaagctagagaacctc
781 -----+-----+-----+-----+-----+ 840
gaactgatgaacgtgaggctcttcttgcacccatggaacctggacttcgatctcttggag
a L D Y L N S E E N V V Y R D L K L E N L ~

FIG. 6B

```

      atgtgtggacaaaggaaggacattcaagataaaggacttcgggtgtgtgaaggagggggtatc
841  +-----+-----+-----+-----+-----+-----+-----+-----+ 900
      taagaaactgttctcgtgacccgtgttagttctctattgacctgaagcccgacacgttctctccctag

a   M L D K D G H I K I T D F G L C R E G I ~

      aaggatggtgtgcactatgaagacattctcggggaaagccgggagtaacctggcccccgtgaggtg
901  +-----+-----+-----+-----+-----+-----+-----+-----+ 960
      ttcttaaccacggtgatacttctgttaagaagcccttgaggccctcatggaacgggggactccac

a   K D G A T M E T F C G T P E Y L A P E V ~

      ctggaagacaaagactaaggccggtgcagtggaactggttgggggtgtgggggtgtgtcatgtat
961  +-----+-----+-----+-----+-----+-----+-----+-----+ 1020
      gaccttctgttgtgtgatgcgggacccgttcactgaacaccccccgaccccgaccagttacata

a   L E D R D Y G R A V D W M G L G V V M Y ~

      gagatgtatgtgtggccgctgcaccttctacaaaccaggacacagagaagctgttccagctg
1021  +-----+-----+-----+-----+-----+-----+-----+-----+ 1080
      ctctactacacacccgggggaacgggaagatgttgggtcctgtgtctctcttggacaagctcgac

a   E M M C G R L F F Y N Q D R E K L F E L ~

      atctctatggaggagatccgcttcccccgcacactcgggcccctgaggccaaagtccctgtctc
1081  +-----+-----+-----+-----+-----+-----+-----+-----+ 1140
      taggggtacctctctctaggccgaaggggccgctgtgtgagccgggactccggttccagggaacag

a   I L M E E I R F P R T L G P E A K S L L ~

      tcggggtgtctcaagaaggacccctacacagaggctcgggtgggggctctcagggaatgccaa
1141  +-----+-----+-----+-----+-----+-----+-----+-----+ 1200
      aggcaccgaagattcttctgggaigtgtctctccggagccaccccggagactcctacggttc

a   S G L L K K D F T Q R L G G G S E D A K ~

      gagatcatgcagcacccggttctttgcacacatcgtgttggccaggtatgttatgagaagaag
1201  +-----+-----+-----+-----+-----+-----+-----+-----+ 1260
      ctctagtacgtcgttggccaaagaacgggttgtagcacacccgtctacacatactctctcttc

a   E I M Q H R F F A N I V W Q D V Y E K E ~

```

FIG. 6C


```
ctgagcccccctttcaggccccaggtccctctgagactgacaccaggtatttogatgag
1361 -----> 1368
gactcgggtggcaggttcggggtccagtgagagctctgactgtggtccctcaggctactc
*   L S P P F K P Q V T S E T D T R Y F D E ~

gattccacagctcagatgatccacctcccgccgctgctcagatgacagcatggagtgt
1381 -----> 1388
ctcagggtctcaggtctactgtggttagtgccggcgagtagttctactgtctactcaca
*   E F T Q Q M I T I T P P D Q D D S M E C ~

gtggacagtgagcggaggccgcacttccccagttctctactgacccagtgccacagcc
1381 -----> 1440
cactgtccactcgcctccggcgtgaggggggtcaggaggatgagtcggtcacggtgtgg
*   Y D S E R R P H P P Q F S Y S A S G T A ~

tga
1441 --- 1443
act
*   *
```

FIG. 6D

ATGAATGAGGTGTCTTCATCAAAGAGGGCTGGCTCCACAGGCTGGTGAATACATCAAG
328 ----- 387
TACTTACTCCACAGACAGTAGTTTCTTCGACCGAGGTGTTCGACCACTTATGTAGTTC

* H H E V S V I K E G W L H K R G E Y I K ~

ACCTGAGGGCCACGGTACTTCCTGCTGAAGAGCGACGGCTCCTTCATTGGGTACAAGGAG
388 ----- 447
TGGACCTCCGGTGGCATGAAGGACGACTTCTCGCTGCCGAGGAAGTAACCCATGTTCTCTC

* T W R P R Y F L L K S D G S F I G Y K E ~

AGGCGCGAGGGCCCTGATCAGACTCTACCCCCCTTAACAACCTTCTCGGTAGCAGAATGC
448 ----- 507
TCCGGGCTCCGGGGACTAGTCTGAGATGGGGGAATTTGTTGAAGAGGCATCGTCTTACG

* R P E A P D Q T L P P L H H F S V A E C ~

PstI
|

CAGCTGATGAAGACCGAGAGGGCCGGACCGAACACCTTTGTCAACGCTGCCCTGCAGTGG
508 ----- 567
GTGCACTACTTCTGGCTCTCCGGCGCTGGGTTGTGGAACAGTATGCGACGGACGTCAAC

* Q L M K T E R P R P H T F V I R C L Q N ~

ACCACAGTCATCGAGAGGACCTTCACGTGGATTCTCCAGACGAGAGGGAGGAGTGGATG
568 ----- 627
TGTTGTCACTAGCTCTCCTGGAGGTGCACCTAAGAGGTCTGCTCTCCCTCCTCACCTAC

* T T V I E R T F H V D S P D E R E E W M ~

ApaI
|

CGGGCCATCCAGATGGTCCCAACAGGCTCAAGCAGCGGGCCCCAGGCGAGGACCCCATG
628 ----- 687
GCCCCGTAGCTCTACCAAGCGGTTGTCGGAGTTCGTGCCCCGGGGTCCGCTCCTTGGGGTAC

* R A I Q M V A H S L K Q R A P G E G F M ~

GACTACAAGTGTGGCTCCCCAGTGACTCCTCCACGACTGAGGAGATGGAGTGGCGGTC
688 ----- 747
CTGATGTTCAACCGAGGGGGTCACTGAGGAGGTGCTGACTCCCTCTACCTTCACCGCCAG

* D Y K C G S P S D S S T T E E M E V A V ~

FIG. 7A




FIG. 7B

CTCATGCTGGACAAAGATGSECCACATCAAGATCACTGACTTTGGCCCTCTSCAAGAGGGC
1168 ----- 1227
GAGTACGACCTGTCTTCTACCGGTGTAGTTCTAGTGACTGAAACCGGAGACGTTTCTCCCG
* L M L D E D G H I K I T D F G L C K E G ~
ATCAGTGACGGGGCCACCATGAAAACCTTCTGTGGGACCCCGGAGTACCTGGCGCCTGAG
1228 ----- 1287
TAGTCACTGCCCCGGTGGTACTTTTGGAGACACCTGGGGCCTCATGGACCGCGGACTC
* I S D G A T M K T F C G T P E Y L A F E ~
GTCTGGAGGACAATGACTATGSCCGGGCCGTGGACTGGTGGGGCTGGGTGTGGTCATG
1288 ----- 1347
CACGACCTCTGTGTACTGATACCGGGCCGGCACCTGACCACCCCCGACCCACACCACTAC
* V L E D N D Y G R A V D W W G L G V V M ~
TACGAGATGATGTGCGGCCGCTGCCCCCTCTACAAACAGGACCCAGAGCGCCTCTTCGAG
1348 ----- 1407
ATGCTCTACTACACGCGGCGGACGGGAAGATGTTGGTCTGTGCTGCGGAGAGGCTC
* Y E M M C G R L P F Y N Q D H E R L F E ~
SacI
|
CTCATCTCTATGGAAGAGATCCGCTTCCCGGSCACGCTCAGCCCCGAGGGCCAAATCCCTG
1408 ----- 1467
GAGTAGGAGTACCTTCTCTAGGGCGAAGGGCGGCTGCCAGTCCGGGCTCCGGTTCCAGGGAC
* L I L M E E I R F P R T L S P E A K S L ~
ApaI
|
CTTGTGGGCTGCTTAAGAAGGACCCCAAGCAGAGGCTTGGTGGGGGGCCCGGATGCC
1468 ----- 1527
GAACGACCCGACGAATTCCTTCTGGGGTTCTGCTCCGAACCACCCCCGGGTCGCTACGG
* L A G L L K K D P R Q R L G G G P S D A ~
AAGGAGGTCAATGAGACACAGGTTCTTCTCAGCATCAACTGGCAGGACGTGGTCCAGAG
1528 ----- 1587
TTCTCCAGTACCTCGTGTCAGAAGGAGTCTGTAGTTGACCGCTCTGCACCAAGTCTTC
* K E V M E H R F F L S I N W Q D V V Q R ~

FIG. 7C

SalI
|

1598 AAGCTCCTSCACCCCTTCABACCTCAGGTCACGTCGAGGTCGACACAGGTACTTCGAT
----- 1647
TTCGAGGACGGTGGGAAGTTTGGASTCCAGTGCAGGCTCCAGCTGTGTTCCATGAAGCTA

* K L L P P P R P Q V T S E V D T R Y F D ~

1648 GATGAATTTACCGCCAGTCCATCACAATCACACCCCTGACCGCTATGACAGCCTGGGC
----- 1707
CTACTTAATGGCGGATCAGGTAGTGTAGTGTGGGGGACTGGCGATACTGTCCGAGCCCG

* D E F T A Q S I T I T P P D R Y D S L G ~

1708 TTACTGGAGCTGGACGAGCGGACCCACTTCCCCAGTTCTCTTACTCGGCGAGCATCCGC
----- 1767
AATGACCTCGACCTGGTCGCTGGGTGAAGGGGGTCAGAGGATGAGCCGGTCGTAGGCG

* L L E L D Q R T H F P Q F S Y S A S I R ~

1768 GASTGAGCAGTCTGCCACGCGAGAGGACGACGCTCGCTGCCATCACCCTGGGTGTTTT
----- 1827
CTCACTGGTCAGACGGGTGGCTCTCTGCGTGCAGGAGCGGTAGTGGGAGCCACCAAA

* E * A V C P R R G R T L A A I T A G W F ~

1828 TTTACCCCTGCCCCG
----- 1842
AAATGGGGAGGGGG

* F T F A R ~

FIG. 7D

[illegible]

FIG. 8A

atgctctctacacccctcctataaaagaaagacacatgaatgatttttgactatttgaaactac
421 -----> 480
tccggagatgtttgggtcgtattttctttctgttacttactaaaactgataaactttgatg
* A S T T N N X R * T K N D P S Y L K L L -
taggttaaggacacttttgggaagttattttgtttcgagagagggcagtggaactatc
481 -----> 540
ctccatttccgtgaaaaacactttcactaaaaacacagctctctctccgttccacttttatga
* C K C T P G K V I L V K E K A S G K T T -
atgctatgagatcttctgagaaagacgtcattttgmaaggatgaagtggacacacac
541 -----> 600
tccgatctctctaaagacttctttcttccagttaataagtttctactctccacgtgtgtgag
* A M K I L K E E V I I A K D E V A H T L -
taactgaaggacagatcttaaaagacactagacatccctttttaaactcacttgaactat
601 -----> 660
attgactttcgtctcctaattttcttctgtgatctgttagggaaaaattgtaggaactttata
* T E S R V L K N T R N F F L T S L K Y S -
ccttccagacaaaaagacgttttgtgttttctgtgatggaaatgtttaattggggcagactgt
661 -----> 720
ggagaggtctgttttctgtgcaaacacaaacactacattatatacaattacccccgtctgaca
* F Q T K C K L C F V K E Y V N G G K L F -
ttttccatttctgtcagagagagcgggtgtttctctgaggacacagcagtttctatgtgtgag
721 -----> 780
aaaaggttaacagctctctctcgcacacacagagactccttggcgtgtgcaagataccacgtc
* F R L S K E R V F S E D E T K F Y G A E -
caattgtctctgacttctacattccggaaagattgtgttaacgtgacttcaagt
781 -----> 840
tttaacagagacgggaactgtatgagtgttaaggcctttctaaacacatggcactagagttac
* I V S A L D Y L N S S E I V Y R D L K L -

FIG. 8B

```

      tggagaatctaatgtgtggacaaagatggccacataaaaattacagatttttggaatttgca
941 -----+-----+-----+-----+-----+-----+-----+-----+ 960
      acctcttagattacgacctgtttctacoggtgtattttttaattgtotaaaaacctgaaacgt

*      E N L M L D K D G H I X I T D F G L C K -

      aagaagggatccacagatgcagccacacatgaagacattctgtggcactccagaatatctgg
961 -----+-----+-----+-----+-----+-----+-----+ 980
      ttcttccctagtggtotacgtcgggtggtactctgttaagacacogtgagggtcttatagacc

*      E G I T D A A T M K T F C G T F E Y L A -

      caacagaggtgtttagaagataatgactatggccagacagtagaactggtggggcctagggg
981 -----+-----+-----+-----+-----+-----+-----+ 1020
      gtggtctccacaactcttctattactgtataccggctcgtctctgaccccccggatcccc

*      P E V L E D N D Y G R A V D W W G L G V -

      ttgtcattgtatgaatgatgtgtgtggaggttacctttctacaaacccaggacacatgagaac
1021 -----+-----+-----+-----+-----+-----+-----+ 1080
      aacagtaactactttactacacacccctccaaatggaaagatgttgggtccttggtactctttg

*      V M Y E N M C G R L P F Y N Q D H E K L -

      tttttgaattaatatttaattggaagacattaaatttccctcgaaacactctctcttcagatgcaa
1081 -----+-----+-----+-----+-----+-----+-----+ 1140
      aaaaacttaattataattaccttctgttaatttaaaggagcttgtgagagaagtctacggt

*      F E L I L M E D I K F F R T L S S D A K -

      BamHI
      |
      aatcattgttttcagggctcttggataaaggtccaaataaacgccttgggtggaggaccag
1141 -----+-----+-----+-----+-----+-----+-----+ 1200
      ttagtaacgaaagtcccgagaactatttccctagggtttatttggggaaacacctcctggtc

*      X L L S G L L I K D P N K R L G G G F D -

      atgatgcaaaagaaatttatgagacacagtttcttctctgtggagttaaactggcaagatgtat
1201 -----+-----+-----+-----+-----+-----+-----+ 1260
      tactacggttttctttaactctgtgtcaagagaagagaacctcaatttgacogttctacata

*      D A K E I M R R S F F S G V N W Q D V Y -

```

FIG. 8C


```

      HindIII
      |
atgataaaaaagcttgtacntctttttaacntcaagtaacntctgagacagctactagat
1261 -----+-----+-----+-----+-----+ 1326
tactatthtttgcacatggagggaaaatttggagttcattgtagactctgtctatgatcta

*      D K X L V P P P K P Q V T S E T D T E Y ~

      attttgatgaagaattttacagctcagactattacaaataacacacntgaaaaatatgatg
1321 -----+-----+-----+-----+-----+ 1386
taaaactacttcttaaxtgctcagagtctgataatgtctattgtggtggactthttactact

*      F D E E F T A G T I T I T P P E E Y D E ~

      NotI
      |
eggatggtatggactgcatggcaaatgagagggggcgcatttccctcaattttactact
1381 -----+-----+-----+-----+-----+ 1446
tactaacatacctgaactacctgttactctccgcggcgttaaggggagttaaaaggatga

*      D G N D C M D N E R R P K F P Q P S Y S ~

      ctgcaagtggagagaaataagtcctctttcatttctgatacttcaactgtcctcttcaattta
1441 -----+-----+-----+-----+-----+ 1506
gcglttcaactgtctcttatttcagagaaagttaagcagatgaagtgaagtgagagttasat

*      A S G E E * V S F I L L L N C N L Q P I ~

      ttactgaaaa
1501 -----+ 1516
aatgaactttt

*      T E ~

```

FIG. 8D

1

EFFECTIVE TREATMENT OF ESOPHOGEAL ADENOCARCINOMA USING TRICIRIBINE AND RELATED COMPOUNDS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. provisional patent application No. 60/935,940, filed Sep. 7, 2007, which is incorporated herein by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

This application may claim subject matter that was developed using grants from the National Institute of Health and the Department of Defense.

TECHNICAL FIELD

This application provides particular therapeutic regimens of triciribine and related compounds and compositions with reduced toxicity for the treatment of esophageal adenocarcinoma and other disorders associated with abnormal cell proliferation.

A computer readable text file, entitled "101889-5003-Sequence Listing.txt," created on or about May 19, 2009 having a size of about 34 kb submitted in U.S. patent application Ser. No. 12/206,468 contains the sequence listing for this application and is hereby incorporated by reference.

BACKGROUND

Cancer is an abnormal growth of cells. Cancer cells rapidly reproduce despite restriction of space, nutrients shared by other cells, or signals sent from the body to stop reproduction. Cancer cells are often shaped differently from healthy cells, do not function properly, and can spread into many areas of the body. Abnormal growths of tissue, called tumors, are clusters of cells that are capable of growing and dividing uncontrollably. Tumors can be benign (noncancerous) or malignant (cancerous). Benign tumors tend to grow slowly and do not spread. Malignant tumors can grow rapidly, invade and destroy nearby normal tissues, and spread throughout the body.

Cancers are classified according to the kind of fluid or tissue from which they originate, or according to the location in the body where they first developed. In addition, some cancers are of mixed types. Cancers can be grouped into five broad categories, carcinomas, sarcomas, lymphomas, leukemias, and myelomas, which indicate the tissue and blood classifications of the cancer. Carcinomas are cancers found in body tissue known as epithelial tissue that covers or lines surfaces of organs, glands, or body structures. For example, a cancer of the lining of the stomach is called a carcinoma. Many carcinomas affect organs or glands that are involved with secretion, such as breasts that produce milk. Carcinomas account for approximately eighty to ninety percent of all cancer cases. Sarcomas are malignant tumors growing from connective tissues, such as cartilage, fat, muscle, tendons, and bones. The most common sarcoma, a tumor on the bone, usually occurs in young adults. Examples of sarcoma include osteosarcoma (bone) and chondrosarcoma (cartilage). Lymphoma refers to a cancer that originates in the nodes or glands of the lymphatic system, whose job it is to produce white blood cells and clean body fluids, or in organs such as the brain and breast. Lymphomas are classified into two categories:

2

Hodgkin's lymphoma and non-Hodgkin's lymphoma. Leukemia, also known as blood cancer, is a cancer of the bone marrow that keeps the marrow from producing normal red and white blood cells and platelets. White blood cells are needed to resist infection. Red blood cells are needed to prevent anemia. Platelets keep the body from easily bruising and bleeding. Examples of leukemia include acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, and chronic lymphocytic leukemia. The terms myelogenous and lymphocytic indicate the type of cells that are involved. Finally, myelomas grow in the plasma cells of bone marrow. In some cases, the myeloma cells collect in one bone and form a single tumor, called a plasmacytoma. However, in other cases, the myeloma cells collect in many bones, forming many bone tumors. This is called multiple myeloma.

Tumor induction and progression are often the result of accumulated changes in the tumor-cell genome. Such changes can include inactivation of cell growth inhibiting genes, or tumor suppressor genes, as well as activation of cell growth promoting genes, or oncogenes. Hundreds of activated cellular oncogenes have been identified to date in animal models, however, only a small minority of these genes have proven to be relevant to human cancers (Weinberg et al 1989 *Oncogenes and the Molecular Origins of Cancer* Cold Spring Harbor, N.Y., Stanbridge and Nowell 1990 *Cell* 63 867-874, Godwin et al 1992 *Oncogenes and antioncogenes in gynecological malignancies*. In W J Hoskins, C A Perez and R C Young (eds), *Gynecological oncology: principles and practice*, pp 87-116, Lippincott, Philadelphia). The activation of oncogenes in human cancers can result from factors such as increased gene copy number or structural changes. These factors can cause numerous cellular effects, for example, they can result in overexpression of a gene product. Several oncogenes involved in human cancer can be activated through gene overexpression.

It has become apparent that the successive genetic aberrations acquired by cancer cells result in defects in regulatory signal transduction circuits that govern normal cell proliferation, differentiation and programmed cell death (Hanahan, D. and R. A. Weinberg, *Cell*, 2000. 100(1): p. 57-700). This in turn results in fundamental defects in cell physiology which dictate malignancy. These defects include: a) self sufficiency in growth signals (i.e. overexpression of growth factor receptor tyrosine kinases such as EGFR and aberrant activation of downstream signal transduction pathways such as Ras/Raf/Mek/Erk 1/2 and Ras/PI3K/Akt), b) resistance to anti-growth signals (i.e. lower expression of TGF β and its receptor), c) evading apoptosis (i.e. loss of proapoptotic p53; overexpression of pro-survival Bcl-2; hyperactivation of survival pathways such as those mediated by PI3K/Akt), d) sustained angiogenesis (i.e. high levels of secretion of VEGF) and f) tissue invasion and metastasis (i.e. extracellular proteases and prometastatic integrins) (Hanahan, D. and R. A. Weinberg, *Cell*, 2000. 100(1): p. 57-700).

Receptor tyrosine kinases such as EGFR, ErbB2, VEGFR and insulin-like growth factor I receptor (IGF-1R) are intimately involved in the development of many to human cancers including colorectal pancreatic, breast and ovarian cancers (Khaleghpour, K., et al. *Carcinogenesis*, 2004. 25(2): p. 241-8.; Sekharam, M., et al., *Cancer Res*, 2003. 63(22): p. 7708-16). Binding of ligands such as EGF, VEGF and IGF-1 to their receptors promotes stimulation of the intrinsic tyrosine kinase activity, autophosphorylation of specific tyrosines in the cytoplasmic domain of the receptors and recruitment of signaling proteins that trigger a variety of complex signal transduction pathways (Olayioye, M. A., et

al., *Embo J*, 2000. 19(13): p. 3159-67, Porter, A. G. and R. R. Vaillancourt, *Oncogene*, 1998. 17(11 Reviews): p. 1343-52). This in turn leads to the activation of many tumor survival and oncogenic pathways such as the Ras/Raf/Mek/Erk $\frac{1}{2}$, JAK/STAT3 and PI3K/Akt pathways. Although all three pathways have been implicated in colon, pancreatic, breast and ovarian oncogenesis, those that are mediated by Akt have been shown to be critical in many steps of malignant transformation including cell proliferation, anti-apoptosis/survival, invasion and metastasis and angiogenesis (Dana, S. R. et al. *Genes Dev*, 1999. 13(22): p. 2905-27).

Akt is a serine/threonine protein kinase (also known as PK δ), which has 3 family members Akt1, Akt2 and Akt3. Stimulation of cells with growth or survival factors results in recruitment to the receptors of the lipid kinase phosphoinositide-3-OH-kinase (PI3K) which phosphorylates phosphoinositol-4,5-bisphosphate (PIP $_2$) to PIP $_3$ which recruits Akt to the plasma membrane where it can be activated by phosphorylation on Thr308 and Ser473 (Akt1), Thr308 and Ser474 (Akt2) and Thr308 and Ser472 (Akt3) (Datta, S. R. et al. *Genes Dev*, 1999. 13(22): p. 2905-27). Thus, PI3K activates Akt by phosphorylating PIP $_2$ and converting to PIP $_3$. The phosphatase PTEN dephosphorylates PIP $_3$ to PIP $_2$ and hence prevents the activation of Akt.

The majority of human cancers contain hyperactivated Akt (Datta, S. R. et al. *Genes Dev*, 1999. 13(22): p. 2905-27, Bellacosa, A., et al., *Int J Cancer*, 1995. 64(4): p. 280-5; Sun, M., et al., *Arm J Pathol*, 2001. 159(2): p. 431-7). In particular, Akt is overexpressed and/or hyperactivated in 57%, 32%, 27% and 36% of human colorectal, pancreatic, breast and ovarian cancers, respectively (Roy, H. K., et al. *Carcinogenesis*, 2002. 23(1): p. 201-5, Altomare, D. A., et al., *J Cell Biochem*, 2003. 88(1): p. 470-6., Sun, M., et al., *Cancer Res*, 2001. 61(16): p. 5985-91., Stal, O., et al. *Breast Cancer Res*, 2003. 5(2): p. R37-44, Cheng, J. Q., et al., *Proc Natl Acad Sci USA*, 1992. 89(19): p. 9267-71, Yuan, Z. Q., et al., *Oncogene*, 2000. 19(19): p. 2324-30). Hyperactivation of Akt is due to amplification and/or overexpression of Akt itself as well as genetic alterations upstream of Akt including overexpression of receptor tyrosine kinases and/or their ligands (Khaleghpour, K., et al. *Carcinogenesis*, 2004. 25(2): p. 241-8.; Sekharam, M., et al., *Cancer Res*, 2003. 63(22): p. 7708-16, Cohen, B. D., et al., *Biochem Soc Symp*, 1998. 63: p. 199-210., Muller, W. J., et al. *Biochem Soc Symp*, 1998. 63: p. 149-57, Miller, W. E., et al. *J Virol*, 1995. 69(7): p. 4390-8, Slamon, D. J., et al., *Science*, 1987. 235(4785): p. 177-82, Andrulis, I. L., et al., *J Clin Oncol*, 1998. 16(4): p. 1340-9) and deletion of the phosphatase PTEN. Proof-of-concept of the involvement of Akt in oncogenesis has been demonstrated preclinically by showing that ectopic expression of Akt induces malignant transformation and promotes cell survival (Sun, M., et al. *Am J Pathol*, 2001. 159(2): p. 431-7, Cheng, J. Q., et al., *Oncogene*, 1997. 14(23): p. 2793-801) and that disruption of Akt pathways inhibits cell growth and induces apoptosis (Jetzi, A., et al. *Cancer Res*, 2003. 63(20): p. 6697-706).

Current treatments of cancer and related diseases have limited effectiveness and numerous serious unintended side effects. Despite demonstrated clinical efficacy of many anticancer drugs, severe systemic toxicity often halts the clinical development of promising chemotherapeutic agents. Further, overexpression of receptor tyrosine kinases such as EGFR and their ligands such as IGF-1, Akt overexpression and/or loss of PTEN (all of which result in hyperactivation of Akt) are associated with poor prognosis, resistance to chemotherapy and shortened survival time of cancer patients. Cur-

rent research strategies emphasize the search for effective therapeutic modes with less risk.

Triciribine

The anticancer action of triciribine (TCN, NSC-154202, 3-amino-1,5-dihydro-5-methyl-1- β -ribofuranosyl-1,4,5,6,8-pentaazaacenaphthylene) and its 5'-phosphate ester, triciribine phosphate (TCN-P, NSC-280594) was initially identified in the 19703 (Townsend & Milne (1975) *Ann NY Acad Sci*, 255: 92-103). TCN-P was the chemical entity advanced into clinical trials because it is more soluble than the parent drug. By the early eighties, TCN-P had shown preclinical activity against leukemias and carcinomas. By the early eighties, TCN-P had been identified as an inhibitor of DNA, RNA and protein synthesis, which demonstrated selectivity towards cells in the S phase of the cell cycle (Roti-Roti et al. 1978 *Proc Am Assoc Cancer Res and ASCO* 19:40). It had also been proposed that unlike other nucleoside antitumor agents at the time, TCN-P is not phosphorylated beyond the level of the monophosphate and is not incorporated into polynucleotides (Bennett et al 1978 *Biochem Pharmacol* 27:233-241, Plagemann *JNCI* 1976 57: 1283-95). It was also established that in vivo, TCN-P is dephosphorylated to TCN by a plasma enzyme and by cellular ecto-5'-nucleotidase. Inside the cells, TCN can be rephosphorylated to TCN-P by adenosine kinase (Wotring et al 1981 *Proc Am Assoc Cancer Res* 22: 257, Basseches et al. *J Chromatogr* 1982 233: 227-234).

In 1982, TCN-P was entered into Phase I clinical trials by Mittelman and colleagues in twenty patients with advanced refractory malignancies (Mittelman et al. 1983 *Cancer Treat Rep* 67: 159-162). TCN-P was administered as an intravenous (i.v.) infusion over fifteen minutes once every three weeks at doses from 25 to 350 mg/m 2 . The patients in the trial were diagnosed with breast, head/neck, lung, pancreas, thyroid, melanoma or undetermined cancer. Only limited therapeutic responses were found and significant toxicity was evident. Mittelman's group concluded that further clinical trials employing their dosing schedule were not warranted, but urged other groups to examine the effects of TCN-P in certain specific cancer types. Also in 1983, Lu et al. (*ASCO Abstracts, Clinical Pharmacology*, p 34 C-133) examined the clinical pharmacology of TCN in patients given 30-40 mg/m 2 intravenously by continuous infusion for five days. Lu et al. reported that TCN contributed to liver toxicity and anemia and suggested that patients should be monitored for these toxicities.

Cobb et al (*Cancer Treat Reports* 1983 67: 173) reported the activity of TCN-P against surgical explants of human tumors in the six day subrenal capsule assay in mice. They examined eighty tumor types that represented breast, lung, colon, ovarian and cervical. Cobb et al reported that TCN produced variable response rates in the different tumors, ranging from 21% (breast) to 88% (cervical).

Another Phase I was also reported by Feun et al. in 1984 (*Cancer Research* 44 (8) 3608-12). Feun et al administered 10, 20, 30, and/or 40 mg/m 2 intravenously by continuous infusion for five days, every three to four or six weeks. The patients in the trial had been diagnosed with colon, sarcoma, melanoma, lung or "other" cancer. Feun et al. reported that significant toxicity was seen, including hyperglycemia, hepatotoxicity and thrombocytopenia. The authors recommended a schedule for Phase II trials of 20 mg/m 2 per day for five days for six weeks and also recommended due to the toxicity that the patients should be closely monitored for liver and pancreatic function, and that patients with diabetes, liver dysfunction or massive hepatic metastasis should be excluded.

In 1986, Schilcher et al. (*Cancer Research* 1986 46: 3147-3151) reported the results of a Phase I evaluation of TCN-P

using a weekly intravenous regimen. The study was conducted in twenty-four patients with advanced solid cancers via a slow intravenous injection over five minutes on days 1, 8, 15 and 22 of a 42 day cycle with a two week rest. Five dose levels ranging from 12 to 96 mg/m² were studied with 3-12 patients treated at each level with a total of 106 doses administered. The patients in the trial had been diagnosed with colon, rectal, bladder, adrenal, ovarian, pancreas, sarcoma, melanoma, lung or "other" cancer. Schilcher et al. concluded

"This weekly schedule produced unexpected clinical toxicity and should not be pursued."

"At this time our group is discouraged to conduct further studies with TCN-P given on weekly or intermittent schedules. A future Phase I-II study using a different regimen (e.g., a single application once a month) might be resumed if TCN-P demonstrates a pronounced in vitro activity against therapeutic resistant primary pancreatic and hepatic tumors."

In 1986, Powis et al (Cancer Treatment Reports 70: 359-362) reported the disposition of TCN-P in blood and plasma of patients during Phase I and II clinical trials. The Phase I trial employed a daily dose of 24-55 mg/m² for 5 days, whereas the Phase II clinical trial employed a single dose of 250 mg/m². Powis et al failed to identify a correlation between TCN-P pharmacokinetic parameters and toxicity of TCN-P.

In the late 1980s, early 1990s, TCN-P advanced to Phase II trials for metastatic colorectal adenocarcinoma, non-small cell lung cancer, advanced squamous cell carcinoma of the cervix and metastatic breast cancer. In 1987, O'Connell et al. (Cancer Treat Reports 71, No. 3, 333-34) published the results of a Phase U trial in patients with metastatic colorectal adenocarcinoma. The patients were administered TCN-P i.v. over 15 minutes 165 or 250 mg/m² once a week in three week intervals. O'Connell et al. concluded that the trials show a lack of clinical usefulness of TCN-P in the treatment of patients with metastatic colorectal adenocarcinoma. Further, in 1991, Lyss, et al., (Proc Annu Meet Am Soc Clin Oncol, (1996) 15 A1151) reported the preliminary results of a trial of the administration of 35 mg/m² per day for five days once every six weeks to patients with advanced non-small cell lung cancer.

Feun et al. (Am J Clin Oncol 1993 16: 506-508) reported the results of a Phase II trial of TCN-P in patients with advanced squamous cell carcinoma of the cervix. A 5 day continuous infusion of at least 35 mg/m² was repeated every six weeks. Among the twenty-one evaluable patients, only two responses were observed. Fuen et al. concluded "using this dose and schedule, TCN-P appears to have limited activity in metastatic or recurrent squamous cell cancer of the cervix."

In 1996, Hoffman et al (Cancer Chemother Pharmacol 37: 254-258) reported the results of a Phase I-II study of TCN-P for metastatic breast cancer. In one study, fourteen patients were treated with 20 mg/m² per day via continuous infusion for five days every six weeks. When the authors failed to see a response at this dose, the dose was escalated to at least 35 mg/m² using the same 5 day continuous infusion schedule. Hoffman et al concluded that "TCN is ineffective at all doses tested and at doses of greater than or equal to 35 mg/m² has unacceptable toxic effects."

Thus, the combination of limited efficacy and unacceptable toxicity prevented the further clinical development of TCN-P and related compounds.

WO 03/079748 to the Regents of the University of California disclosed certain ZNF217 inhibitors, such as tricinibine, in combination with additional chemotherapeutic agents, such as doxorubicin.

It is an object of the present invention to provide for the administration of tricinibine and related compounds and compositions with reduced toxicity for the treatment of tumors, cancer, and others disorders associated with abnormal cell proliferation.

It is another object of the present invention to provide improved methods to treat tumors or cancer in the subject with tricinibine and related compounds.

SUMMARY OF THE INVENTION

The present invention provides novel therapeutic regimens of tricinibine, tricinibine phosphate and related compounds to treat esophageal adenocarcinoma in a subject while limiting systemic toxicity. The invention is based on the discovery that esophageal adenocarcinoma, which overexpress Akt kinase are particularly sensitive to the cytotoxic effects of TCN and related compounds. The inventors have determined, contrary to the prior art and experience, how to successfully use tricinibine to treat esophageal adenocarcinoma by one or a combination of (i) administering tricinibine only to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the drug; (ii) use of a described dosage level that minimizes the toxicity of the drug but yet still exhibits efficacy; or (iii) use of a described dosage regimen that minimizes the toxicity of the drug.

In one aspect of the present invention, methods are provided to identify esophageal adenocarcinoma susceptible to the toxic effects of TCN, TCN-P and/or related compounds. In one embodiment, methods are provided for treating esophageal adenocarcinoma in a mammal, particularly a human that includes (i) obtaining a biological sample from the tumor. (ii) determining whether the tumor overexpresses an Akt kinase, and (iii) treating the tumor that overexpresses Akt kinase with tricinibine, tricinibine phosphate or a related compound as described herein. In one embodiment, the level of Akt kinase expression can be determined by assaying the esophageal adenocarcinoma for the presence of a phosphorylated Akt kinase, for example, by using an antibody that can detect the phosphorylated form. In another embodiment, the level of Akt expression can be determined by assaying a esophageal adenocarcinoma cell obtained from a subject and comparing the levels to a control tissue. In certain embodiments, the Akt can be overexpressed at least 2, 2.5, 3 or 5 fold in the esophageal adenocarcinoma sample compared to the control. In certain embodiments, the overexpressed Akt kinase can be a hyperactivated and phosphorylated Akt kinase.

In another aspect of the present invention, dosing regimens are provided that limit the toxic side effects of TCN and related compounds. In one embodiment, such dosing regimens minimize or eliminate toxic side effects, including, but not limited to, hepatotoxicity, thrombocytopenia, hyperglycemia, vomiting, hypocalcemia, anemia, hypoalbuminemia, myelosuppression, hypertriglyceridemia, hyperamylasemia, diarrhea, stomatitis and/or fever. In another embodiment, the administration of TCN, TCN-P Or related compounds provides at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15, 20, or 25% of the subjects.

In one embodiment, a method is provided to treat a subject which has been diagnosed with esophageal adenocarcinoma by administering to the subject an effective amount of TCN, TCN-P or a related compound, for example compounds described herein, according to a dosing schedule that includes administering the drug approximately one time per week for approximately three weeks followed by a one week period

wherein the drug is not administered. In another embodiment, methods are provided to treat esophageal adenocarcinoma in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P or a related compound one time per week. In one embodiment, the compound can be administered as a single bolus dose over a short period of time, for example, about 5, 10 or 15 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the continuous administration can be repeated at least once a week, once every two weeks and/or once a month. In other embodiments, the compounds can be administered at least once every three weeks. In further embodiments, the compounds can be administered at least once a day for at least 2, 3, 4 or 5 days.

In further embodiments, TCN, TCN-P and related compounds as disclosed herein can be administered to patients in an amount that is effective in causing esophageal adenocarcinoma regression. The administration of TCN, TCN-P or related compounds can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a compound disclosed herein can be administered to a subject. The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen can include administering less than 20 mg/m² of TCN and related compounds. In one embodiment, less than 10 mg/m² of TCN or related compounds can be administered once a week. In further embodiments, dosages of or less than 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN or a related compound can be administered to subject. In another embodiment, less than 10 mg/m² can be administered to a subject via continuous infusion for at least five days. In particular embodiments, TCN or a related compound as disclosed herein can be used for the treatment of esophageal adenocarcinoma.

In one embodiment, the compounds and/or therapeutic regimens of the present invention can be used to prevent and/or treat esophageal adenocarcinoma. In particular embodiments, TCN or a related compound as disclosed herein can be used for the treatment of esophageal adenocarcinoma. In further embodiments of the present invention, the compounds disclosed herein can be used in the treatment of angiogenesis-related diseases. In certain embodiments, methods are provided to treat esophageal adenocarcinoma via continuous infusion of TCN, TCN-P or a related compound via continuous infusion for at least 24, 48, 72 or 96 hours. In other embodiments, the continuous infusion can be repeated, for example, at least once every two, three or four weeks.

In a particular embodiment, there is provided a method for the treatment of esophageal adenocarcinoma, and others disorders associated with an abnormal cell proliferation in a host, the method comprising administering to the host an effective amount of a compound disclosed herein optionally in combination with a pharmaceutically acceptable carrier.

In one aspect, the compounds and compositions can be administered in combination or alternation with at least one additional chemotherapeutic agent. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time. In one embodiment, compositions of the invention can be combined with antiangiogenic agents. In other embodiments of the present invention, the compounds and compositions disclosed herein can be used in combination or alterna-

tion with the following types of drugs, including, but not limited to: antiproliferative drugs, antimetabolic agents, antimitotic agents, antineoplastic drugs, alkylating agents or nitrogen mustard; drugs which target topoisomerases, drugs which target signal transduction in tumor cells, gene therapy and antisense agents, antibody therapeutics, steroids, steroid analogues, antiemetic drugs and/or nonsteroidal agents.

In other embodiments, TCN, TCN-P or a related compound as disclosed herein can be used to treat esophageal adenocarcinoma resistant to one or more drugs, including the embodiments of esophageal adenocarcinoma and drugs disclosed herein. In one embodiment, TCN, TCN-P or a related compound as disclosed herein is administered in an effective amount for the treatment of a patient with a drug resistant esophageal adenocarcinoma, for example, multidrug resistant esophageal adenocarcinoma, including but not limited to those resistant to taxol, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa). In one embodiment, the TCN, TCN-P or related compound as disclosed herein can be administered with an additional chemotherapeutic agent that can be a P-glycoprotein inhibitor, such as verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextropropofol, valspodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In certain embodiments, a method is provided including administering to a host in need thereof an effective amount of a compound disclosed herein, or pharmaceutical composition comprising the compound, in an effective amount for the treatment of the treatment of tumors, cancer, and others disorders associated with an abnormal cell proliferation in a host.

In one embodiment, a method for the treatment of a esophageal adenocarcinoma is provided including an effective amount of a compound disclosed herein, or a salt, isomer, prodrug or ester thereof, to an individual in need thereof. The compound, or salt, isomer, prodrug or ester thereof, is optionally provided in a pharmaceutically acceptable composition including the appropriate carriers, such as water, which is formulated for the desired route of administration to an individual in need thereof. Optionally the compound is administered in combination or alternation with at least one additional therapeutic agent for the treatment of esophageal adenocarcinoma.

Also within the scope of the invention is the use of a compound disclosed herein or a salt, prodrug or ester thereof in the treatment of esophageal adenocarcinoma, optionally in a pharmaceutically acceptable carrier; and the use of a compound disclosed herein or a salt, prodrug or ester thereof in the manufacture of a medicament for the treatment of cancer or tumor, optionally in a pharmaceutically acceptable carrier.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 demonstrates the identification of API-2 (tricitiribine) as a candidate of Akt inhibitor from the NCI Diversity Set. A illustrates the chemical structure of API-2 (tricitiribine). B demonstrates that API-2 inhibits phosphorylation levels of AKT2 in AKT2-transformed NIH3T3 cells. Wild type AKT2-transformed NIH3T3 cells were treated with API-2 (1 μ M) for indicated times and subjected to immunoblotting analysis with anti-phospho-Akt-T308 and -S473 antibodies (top and middle panels). The bottom panel shows expression of total AKT2. In C, it is shown that API-2 inhibits three isoforms of Akt. HEK293 cells were transfected with HA-Akt1, -AKT2 and -AKT3 and treated with API-2 (1 μ M) or wortmannin (15 μ M) prior to EGF stimulation, the cells were lysed and immu-

noprecipitated with anti-HA antibody. The immunoprecipitates were subjected to in vitro kinase assay (top) and immunoblotting analysis with anti-phospho-Akt-T308 (bottom) antibody. Middle panel shows expression of transfected Akt1, Akt2 and Akt3. D illustrates that API-2 did not inhibit Akt in vitro. In vitro kinase assay of constitutively active AKT2 recombinant protein in a kinase buffer containing 1 μ M API-2 (lane 3).

FIG. 2 demonstrates that API-2 does not inhibit PI3K, PDK1 and the closely related members of AGC kinase family. A demonstrates an in vitro PI3K kinase assay. HEK293 cells were serum-starved and treated with API-2 (1 μ M) or Wortmannin (15 μ M) from 30 minutes prior to EGF stimulation. Cells were lysed and immunoprecipitated with anti-p110 antibody. The immunoprecipitates were subjected to in vitro kinase assay using PI-4-P as substrate. B illustrates the effect of API-2 on in vitro PDK1 activation (top panel), closed circles show inhibition by API-2. Open circles show inhibition by the positive control staurosporine, which is a potent PDK1 inhibitor (IC₅₀=5 nM). Bottom panels are immunoblotting analysis of HEK293 cells that were transfected with Myc-PDK1 and treated with wortmannin or API-2 prior to EGF stimulation. The immunoblots were detected with indicated antibodies. C illustrates an immunoblot analysis of phosphorylation levels of PKC α with anti-phospho-PKC α T638 (top) and total PKC α (bottom) antibodies following treatment with API-2 or a nonselective PKC inhibitor Ro31-8220. D shows an in vitro SGK kinase assay. HEK293 cells were transfected with HA-SGK and treated with API-2 or wortmannin prior to EGF stimulation. In vitro kinase was performed with HA-SGK immunoprecipitates using MBP as substrate (top). Bottom panel shows the expression of transfected HA-SGK. E illustrates the results of a PKA kinase assay. Immuno-purified PKA was incubated in ADB buffer (Upstate Biotechnology Inc) containing indicated inhibitors (API-2 or PKAI) and substrate Kemptide. The kinase activity was quantified. In F, a western blot is shown. OVCAR3 cells were treated with API-2 for indicated times. Cell lysates were immunoblotted with indicated anti-phospho-antibodies (panels 1-4) and anti-actin antibody (bottom).

FIG. 3 demonstrates that API-2 inhibits Akt activity and cell growth and induces apoptosis in human cancer cells with elevated Akt. A is a western blot, following treatment with API-2, phosphorylation levels of Akt were detected with anti-phospho-Akt-T308 antibody in indicated human cancer cell lines. The blots were reprobbed with anti-total Akt antibody (bottom panels). In B, a cell proliferation assay is shown. Cell lines as indicated in the figure were treated with different doses of API-2 for 24 h and 48 h and then analyzed with CellTiter 96 Cell Proliferation Assay kit (Promega). C provides an apoptosis analysis. Cells were treated with API-2 and stained with annexin V and PI and analyzed by FACSscan.

FIG. 4 shows that API-2 inhibits downstream targets of Akt and exhibits anti-tumor activity in cancer cell lines with elevated Akt in mouse xenograft. In A, it is demonstrated that API-2 inhibits Akt phosphorylation of tuberlin, Bad, AFX and GSK-3 β . Following treatment with API-2, OVAR3 cells were lysed and immunoblotted with indicated antibodies. B shows that API-2 inhibits tumor growth. Tumor cells were subcutaneously injected into nude mice with low level of Akt cells on left side and elevated level of Akt cells on right side. When the tumors reached an average size of about 100-150 mm³, animals were treated with either vehicle or 1 mg/kg/day API-2. Each measurement represents an average of 10 tumors. C illustrates a representation of the mice with OVCAR3 (right) and OVCAR5 (left) xenograft treated with API-2 or vehicle (control). D shows examples of tumor size

(bottom) and weight (top) at the end of experiment. In E, immunoblot analysis of tumor lysates was performed with anti-phospho-Akt-S473 (top) and anti-AKT2 (bottom) antibodies in OVCAR-3-derived tumors that were treated (T3 and T4) and untreated (T1 and T2) with API-2.

FIG. 5 shows that API-2 (tricitiribine) inhibits Mt kinase activity in vitro. In vitro kinase assay was performed with recombinant of PDK1 and Akt in a kinase buffer containing phosphatidylinositol-3,4,5-P₃ (PIP₃), API-2 and histone H2B as substrate. After incubation of 30 min, the reactions were separated by SDS-PAGE and exposed in a film.

FIG. 6 provides the mRNA (SEQ ID NO: 1) and amino acid sequence (SEQ ID NO: 2) of human Akt1, restriction enzyme sites are also noted.

FIG. 7 provides the mRNA (SEQ ID NO: 3) and amino acid sequence (SEQ ID NO: 4) of human Akt2 restriction enzyme sites are also noted.

FIG. 8 provides the mRNA (SEQ ID NO: 5) and amino acid sequence (SEQ ID NO: 6) of human Akt3 restriction enzyme sites are also noted.

DETAILED DESCRIPTION

The inventors have determined, contrary to the prior art and experience, how to successfully use tricitiribine to treat esophageal adenocarcinoma by one or a combination of (i) administering tricitiribine only to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the drug; (ii) using a described dosage level that minimizes the toxicity of the drug but yet still exhibits efficacy; or (iii) using a described dosage regimen that minimizes the toxicity of the drug.


Esophageal adenocarcinoma has demonstrated a rapid increase in incidence over the last 10 years. This increase mirrors a dramatic rise in that of Barrett esophagus, which is associated with esophageal adenocarcinoma in at least 95% of cases. In an attempt to understand the pathogenesis of esophageal adenocarcinoma, attention has turned to the anti-apoptotic and oncogenic pathways. Here it is demonstrated that Akt was frequently activated in Barrett esophagus-related adenocarcinoma. Surprisingly, the levels of Akt activation were associated with tumor progression. After institutional review board ethics approval, 60 archival tissue specimens of esophageal adenocarcinoma arising on a background of Barrett esophagus were selected for immunohistochemical staining with phosphorylated Akt (p-Akt) antibody. The slides were scored by 2 independent observers. Approximately 80% of high-grade dysplasia and esophageal adenocarcinoma cases demonstrated strong to moderate Akt activity. Sixty-two percent of Barrett mucosa revealed low Akt activity, the remaining cases being p-Akt negative. None of the low-grade dysplasia cases exhibited strong p-Akt staining, whereas only weak p-Akt activity is seen in a portion of metaplastic Barrett mucosa, Akt is highly activated in high-grade dysplasia and esophageal adenocarcinoma arising from Barrett esophagus. These findings suggest a role of p-Akt in the progression of Barrett esophagus to esophageal adenocarcinoma and provide the rationale for using p-Akt inhibitor API-2/tricitiribine, which is currently in clinical trial, in the treatment of esophageal adenocarcinoma.

Compounds

The present invention provides for the use of TCN, TCN-P and related compounds for use in particular therapeutic regimens for the treatment of esophageal adenocarcinoma.

In one embodiment, the compounds provided herein have the following structures:

11



wherein each R2', R3' and R5' are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate pro-drug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein R2', R3' or R5' is independently H or mono-, di- or tri-phosphate;


wherein R^x and R^y are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid.

R₁ and R₃ each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl.

12

In one embodiment, R2' and R3' are hydrogen. In another embodiment, R2' and R5' are hydrogen. In yet another embodiment, R2', R3' and R5' are hydrogen. In yet another embodiment, R2', R3', R5', R1 and R2 are hydrogen.


5 In another embodiment, the compound has the following structure:




wherein R₃ is H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, NH₂, NHR⁴, N(R⁴)₂, aryl, alkoxyalkyl, aryloxyalkyl, or substituted aryl; and

each R⁴ independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl, or aryl. In a subembodiment, R₃ is a straight chained C1-11 alkyl, iso-propyl, i-butyl, or phenyl.

In one embodiment, the compounds provided herein have the following structure:




10 In another embodiment, the compounds provided herein have the following structure:




15 In another embodiment, the compounds provided herein have the following structure:

13



14




15 In another embodiment, the compounds provided herein have the following structure:

wherein R_6 is H, alkyl, (including lower alkyl) alkenyl, alkynyl, alkoxyalkyl, hydroxyalkyl, arylalkyl, cycloalkyl, NH_2 , NHR^4 , NR^4R^4 , CF_3 , CH_2OH , CH_2F , CH_2Cl , CH_2CF_3 , $C(Y^3)_3$, $C(Y^3)_2C(Y^3)_3$, $C(=O)OH$, $C(=O)OR^4$, $C(=O)$ -alkyl, $C(=O)$ -aryl, $C(=O)$ -alkoxyalkyl, $C(=O)NH_2$, $C(=O)NHR^4$, $C(=O)N(R^4)_2$, where each Y^3 is independently H or halo; and

each R^4 independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl, or aryl.

In a subembodiment, R_6 is ethyl, CH_2CH_2OH , or CH_2 -phenyl.

In another embodiment, the compounds provided herein have the following structure:




wherein R_7 is H, halo, alkyl (including lower alkyl), alkenyl, alkynyl, alkoxy, alkoxyalkyl, hydroxyalkyl, cycloalkyl, nitro, cyano, OH , OR^4 , NH_2 , NHR^4 , NR^4R^4 , SH , SR^4 , CF_3 , CH_2OH , CH_2F , CH_2Cl , CH_2CF_3 , $C(Y^3)_3$, $C(Y^3)_2C(Y^3)_3$, $C(=O)OH$, $C(=O)OR^4$, $C(=O)$ -alkyl, $C(=O)$ -aryl, $C(=O)$ -alkoxyalkyl, $C(=O)NH_2$, $C(=O)NHR^4$, $C(=O)N(R^4)_2$, or N_3 , where each Y^3 is independently H or halo; and

each R^4 independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, to alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl.

In a subembodiment, R_7 is methyl, ethyl, phenyl, chloro or NH_2 .

In another embodiment, the compounds provided herein have the following structure:



It is to be understood that the compounds disclosed herein may contain chiral centers. Such chiral centers may be of either the (R) or (S) configuration, or may be a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, or be stereoisomeric or diastereomeric mixtures. It is understood that the disclosure of a compound herein encompasses any racemic, optically active, polymorphic, or stereoisomeric form, or mixtures thereof, which preferably possesses the useful properties described herein, it being well known in the art how to prepare optically active forms and how to determine activity using the standard tests described herein, or using other similar tests which are well known in the art. Examples of methods that can be used to obtain optical isomers of the compounds include the following:

physical separation of crystals—a technique whereby macroscopic crystals of the individual enantiomers are manually separated. This technique can be used if crystals of the separate enantiomers exist, i.e., the material is a conglomerate, and the crystals are visually distinct;

simultaneous crystallization—a technique whereby the individual enantiomers are separately crystallized from a solution of the racemate, possible only if the latter is a conglomerate in the solid state;

enzymatic resolutions—a technique whereby partial or complete separation of a racemate by virtue of differing rates of reaction for the enantiomers with an enzyme

enzymatic asymmetric synthesis—a synthetic technique whereby at least one step of the synthesis uses an enzymatic

15

reaction to obtain an enantiomerically pure or enriched synthetic precursor of the desired enantiomer;

chemical asymmetric synthesis—a synthetic technique whereby the desired enantiomer is synthesized from an achiral precursor under conditions that produce asymmetry (i.e., chirality) in the product, which may be achieved using chiral catalysts or chiral auxiliaries;

diastereomer separations—a technique whereby a racemic compound is reacted with an enantiomerically pure reagent (the chiral auxiliary) that converts the individual enantiomers to diastereomers. The resulting diastereomers are then separated by chromatography or crystallization by virtue of their now more distinct structural differences and the chiral auxiliary later removed to obtain the desired enantiomer,

first- and second-order asymmetric transformations—a technique whereby diastereomers from the racemate equilibrate to yield a preponderance in solution of the diastereomer from the desired enantiomer or where preferential crystallization of the diastereomer from the desired enantiomer perturbs the equilibrium such that eventually in principle all the material is converted to the crystalline diastereomer from the desired enantiomer. The desired enantiomer is then released from the diastereomer;

kinetic resolutions—this technique refers to the achievement of partial or complete resolution of a racemate (or of a further resolution of a partially resolved compound) by virtue of unequal reaction rates of the enantiomers with a chiral, non-racemic reagent or catalyst under kinetic conditions;

enantiospecific synthesis from non-racemic precursors—a synthetic technique whereby the desired enantiomer is obtained from non-chiral starting materials and where the stereochemical integrity is not or is only minimally compromised over the course of the synthesis;

chiral liquid chromatography—a technique whereby the enantiomers of a racemate are separated in a liquid mobile phase by virtue of their differing interactions with a stationary phase. The stationary phase can be made of chiral material or the mobile phase can contain an additional chiral material to provoke the differing interactions;

chiral gas chromatography—a technique whereby the racemate is volatilized and enantiomers are separated by virtue of their differing interactions in the gaseous mobile phase with a column containing a fixed non-racemic chiral adsorbent phase;

extraction with chiral solvents—a technique whereby the enantiomers are separated by virtue of preferential dissolution of one enantiomer into a particular chiral solvent;

transport across chiral membranes—a technique whereby a racemate is placed in contact with a thin membrane barrier. The barrier typically separates two miscible fluids, one containing the racemate, and a driving force such as concentration or pressure differential causes preferential transport across the membrane barrier. Separation occurs as a result of the non-racemic chiral nature of the membrane which allows only one enantiomer of the racemate to pass through.

In some embodiments, tricyribine, tricyribine phosphate (TCN-P), tricyribine 5'-phosphate (TCN-P), or the DMF adduct of tricyribine (TCN-DMF) are provided. TCN can be synthesized by any technique known to one skilled in the art, for example, as described in Tetrahedron Letters, vol. 49; pp. 4757-4760 (1971). TCN-P can be prepared by any technique known to one skilled in the art, for example, as described in U.S. Pat. No. 4,123,524. The synthesis of TCN-DMF is described, for example, in INSERM, vol. 81, pp. 37-82 (1978). Other compounds related to TCN as described herein can be synthesized, for example, according to the methods disclosed in Gudmundsson, K. S., et al., "Synthesis of car-

16

bocyclic analogs of 2',3'-dideoxysangivamycin, 2',3'-dideoxytocamycin, and 2',3'-dideoxytricyribine," *Nucleosides Nucleotides Nucleic Acids*, 20(10-11):1823-1830 (October-November 2001); Porcari, A. R., et al., "6-N-Acyltricyribine analogues: structure-activity relationship between acyl carbon chain length and activity against HIV-1," *J. Med. Chem.*, 43(12):2457-2463 (Jun. 15, 2000); Porcari, A. R., et al., "Acyclic sugar analogs of tricyribine: lack of antiviral and antiproliferative activity correlate with low intracellular phosphorylation," *Nucleosides Nucleotides*, 18(11-12):2475-2497 (November-December 1999), Porcari, A. R., et al., "Deoxy sugar analogues of tricyribine: correlation of antiviral and antiproliferative activity with intracellular phosphorylation," *J. Med. Chem.*, 43(12):2438-2448 (Jun. 15, 2000), Porcari, A. R., et al., "Synthesis and antiviral activity of 2-substituted analogs of tricyribine," *Nucleosides Nucleotides Nucleic Acids*, 22(12):2171-2193 (December 2003), Porcari, A. R., et al., "An improved total synthesis of tricyribine: a tricyclic nucleoside with antineoplastic and antiviral properties," *Nucleosides Nucleotides Nucleic Acids*, 23(1-2):31-39 (2004), Schweinsberg, P. D., et al., "Identification of the metabolites of an antitumor tricyclic nucleoside (NSC-154020)," *Biochem. Pharmacol.*, 30(18):2521-2526 (Sep. 15, 1981), Smith, K. L., et al., "Synthesis of new 2'-beta-C-methyl related tricyribine analogues as anti-HCV agents," *Bioorg. Med. Chem. Lett.*, 14(13):3517-3520 (Jul. 5, 2004), Townsend, L. B., et al., "The synthesis and biological activity of certain pentaazaacenaphthylenes, hexaazaacenaphthylenes and their corresponding nucleosides," *Nucleic Acids Symp. Ser.*, 1986(17):41-44 (1986), and/or Wotring, L. L., et al., "Mechanism of activation of tricyribine phosphate (TCN-P) as a prodrug form of TCN," *Cancer Med. Rep.*, 70(4):491-7 (April 1986).

Pharmaceutically Acceptable Salts and Prodrugs

In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compound as a pharmaceutically acceptable salt may be appropriate. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. In particular, examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids, which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Suitable inorganic salts may also be formed, including, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

Any of the nucleotides described herein can be administered as a nucleotide prodrug to increase the activity, bioavailability, stability or otherwise alter the properties of the nucleoside. A number of nucleotide prodrug ligands are known. In general, alkylation, acylation or other lipophilic modification of the mono, di or triphosphate of the nucleoside will increase the stability of the nucleotide. Examples of substituent groups that can replace one or more hydrogens on the phosphate moiety are alkyl, aryl, steroids, carbohydrates, including sugars, 1,2-diacylglycerol and alcohols. Many are

described in R. Jones and N. Bischofberger, *Antiviral Research*, 27 (1995) 1-17. Any of these can be used in combination with the disclosed nucleosides to achieve a desired effect.

In one embodiment, the tricitabine or a related compound is provided as 5'-hydroxyl lipophilic prodrug. Nonlimiting examples of U.S. patents that disclose suitable lipophilic substituents that can be covalently incorporated into the nucleoside, preferably at the 5'-OH position of the nucleoside or lipophilic preparations, include U.S. Pat. Nos. 5,149,794 (Sep. 22, 1992, Yatvin, et al.); 5,194,654 (mar. 16, 1993, Hostetler, et al.); 5,223,263 (Jun. 29, 1993, Hostetler, et al.); 5,256,641 (Oct. 26, 1993, Yatvin, et al.); 5,411,947 (May 2, 1995, Hostetler, et al.); 5,463,092 (Oct. 31, 1995, Hostetler, et al.); 5,543,389 (Aug. 6, 1996, Yatvin, et al.); 5,543,390 (Aug. 6, 1996, Yatvin, et al.); 5,543,391 (Aug. 6, 1996, Yatvin, et al.); and 5,554,728 (Sep. 10, 1996, Basava, et al.), all of which are incorporated herein by reference.

Foreign patent applications that disclose lipophilic substituents that can be attached to the tricitabine or a related compounds of the present invention, or lipophilic preparations, include WO 89/02733, WO 90/00555, WO 91/16920, WO 91/18914, WO 93/00910, WO 94/26273, WO/15132, EP 0 350 287, EP 93917054.4, and WO 91/19721.

Additional nonlimiting examples of derivatives of tricitabine or a related compounds are those that contain substituents as described in the following publications. These derivatized tricitabine or a related compounds can be used for the indications described in the text or otherwise as antiviral agents, including as anti-HIV or anti-HBV agents. Ho, D. H. W. (1973) Distribution of Kinase and deaminase of 1 β -D-arabinofuranosylcytosine in tissues of man and mouse. *Cancer Res.* 33, 2816-2820; Holy, A. (1993) Isopolar phosphorous-modified nucleotide analogues. In: De Clercq (ed.), *Advances in Antiviral Drug Design*, Vol. 1, JAI Press, pp. 179-231; Hong, C. I., Nechaev, A., and West, C. R. (1979a) Synthesis and antitumor activity of 1 β -3-arabinofuranosylcytosine conjugates of cortisol and cortisone. *Biochem. Biophys. Res. Commun.* 88, 1223-1229; Hong, C. I., Nechaev, A., Kirisits, A. J. Buchheit, D. J. and West, C. R. (1980) Nucleoside conjugates as potential antitumor agents. 3. Synthesis and antitumor activity of 1-(β -D-arabinofuranosyl)cytosine conjugates of corticosteroids and selected lipophilic alcohols. *J. Med. Chem.* 28, 171-177; Hostetler, K. Y., Stuhmiller, L. M., Lenting, H. B. M. van den Bosch, H. and Richman, D. D. (1990) Synthesis and antiretroviral activity of phospholipid analogs of azidothymidine and other antiviral nucleosides. *J. Biol. Chem.* 266, 11714-11717; Hostetler, K. Y., Korba, B. Sridhar, C., Gardener, M. (1994a) Antiviral activity of phosphatidyl-dideoxycytidine in hepatitis B-infected cells and enhanced hepatic uptake in mice. *Antiviral Res.* 24, 59-67; Hostetler, K. Y., Richman, D. D., Sridhar, C. N. Felgner, P. L., Felgner, J., Ricci, J., Gardener, M. F. Selleseth, D. W. and Ellis, M. N. (1994b) Phosphatidylazidothymidine and phosphatidyl-ddC: Assessment of uptake in mouse lymphoid tissues and antiviral activities in human immunodeficiency virus-infected cells and in rausher leukemia virus-infected mice. *Antimicrobial Agents Chemother.* 38, 2792-2797; Hunston, R. N., Jones, A. A. McGuigan, C., Walker, R. T., Balzarini, J., and De Clercq, E. (1984) Synthesis and biological properties of some cyclic phosphotriesters derived from 2'-deoxy-5-fluorouridine. *J. Med. Chem.* 27, 440-444; Ji, Y. H., Moog, C., Schmitt, C. Bischoff, P. and Luu, B. (1990) Monophosphoric acid diesters of 7 β -hydroxycholesterol and of pyrimidine nucleosides as potential antitumor agents; synthesis and preliminary evaluation of antitumor activity. *J. Med. Chem.* 33, 2264-2270; Jones, A. S., McGuigan, C., Walter, R. T., Balzarini, J.

and DeClercq, E. (1984) Synthesis, properties, and biological activity) of some nucleoside cyclic phosphoramidates. *J. Chem. Soc. Perkin Trans. I*, 1471-1474; Juodka, B. A. and Smart, J. (1974) Synthesis of ditribonucleoside a (P \square N) amino acid derivatives. *Coll. Czech. Chem. Comm.* 39, 363-968; Kataolca, S., Imai, J., Yamaji, N., Kato, M., Saito, M., Kawada, T. and Imai, S. (1989) Alkylated cAMP derivatives; selective synthesis and biological activities. *Nucleic Acids Res. Sym. Ser.*, 21, 1-2; Kataoka, S., Uchida, R. and Yamaji, N. (1991) A convenient synthesis of adenosine 3',5' cyclic phosphate (cAMP) benzyl and methyl triesters. *Heterocycles* 32, 1351-1356; Kinchington, D., Harvey, J. J., O'Connor, T. J., Jones, B. C. N. M., Devine, K. G., Taylor-Robinson, D., Jeffries, D. J. and McGuigan, C. (1992) Comparison of antiviral effects of zidovudine phosphoramidate and phosphorodiamidate derivatives against HIV and ULV in vitro. *Antiviral Chem. Chemother.* 3, 107-112; Kodama, K., Morozumi, M., Saitoh, K. L., Kuninaka, H., Yoshino, H. and Saneyoshi, M. (1989) Antitumor activity and pharmacology of 1- β -D-arabinofuranosylcytosine-5'-stearylphosphate; an orally active derivative of 1- β -D-arabinofuranosylcytosine. *Jpn. J. Cancer Res.* 80, 679-685; Korty, M. and Engels, J. (1979) The effects of adenosine- and guanosine 3',5'-phosphoric and acid benzyl esters on guinea-pig ventricular myocardium. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 310, 103-111; Kumar, A., Goe, P. L., Jones, A. S. Walker, R. T. Balzarini, J. and De Clercq, E. (1990) Synthesis and biological evaluation of some cyclic phosphoramidate nucleoside derivatives. *J. Med. Chem.* 33, 2368-2375; LeBec, C., and Huynh-dinh, T. (1991) Synthesis of lipophilic phosphate triester derivatives of 5-fluorouridine and arabinocytidine as anticancer prodrugs. *Tetrahedron Lett.* 32, 6553-6556; Lichtenstein, J., Barner, H. D. and Cohen S. S. (1960) The metabolism of exogenously supplied nucleotides by *Escherichia coli*. *J. Biol. Chem.* 235, 457-465; Lucity, J., Von Daeniken, A., Friederich, J. Manthey, B., Zweifel, J., Schlatter, C. and Bann, M. H. (1981) Synthesis and toxicological properties of three naturally occurring cyanoepithioalkanes. *Mitt. Geg. Lebensmittellunters. Hyg.* 72, 131-133 (*Chem. Abstr.* 95, 127093); McGuigan, C., Tollerfield, S. M. and Riley, P. A. (1989) Synthesis and biological evaluation of some phosphate triester derivatives of the antiviral drug Ara. *Nucleic Acids Res.* 17, 6065-6075; McGuigan, C., Devine, K. G., O'Connor, T. J., Galpin, S. A., Jeffries, D. J. and Kinchington, D. (1990a) Synthesis and evaluation of some novel phosphoramidate derivatives of 3'-azido-3'-deoxythymidine (AZT) as anti-HIV compounds. *Antiviral Chem. Chemother.* 1, 107-113; McGuigan, C., O'Connor, T. J., Nicholls, S. R. Nickson, C. and Kinchington, D. (1990b) Synthesis and anti-HIV activity of some novel substituted dialkyl phosphate derivatives of AZT and ddCyd. *Antiviral Chem. Chemother.* 1, 355-360; McGuigan, C., Nicholls, S. R., O'Connor, T. J., and Kinchington, D. (1990c) Synthesis of some novel dialkyl phosphate derivative of 3'-modified nucleosides as potential anti-AIDS drugs. *Antiviral Chem. Chemother.* 1, 25-33; McGuigan, C., Devine, K. G., O'Connor, T. J., and Kinchington, D. (1991) Synthesis and anti-HIV activity of some haloalkyl phosphoramidate derivatives of 3'-azido-3'-deoxythymidine (AZT); potent activity of the trichloroethyl methoxyalaninyl compound. *Antiviral Res.* 15, 255-263; McGuigan, C., Pathirana, R. N., Mahmood, N., Devine, K. G. and Hay, A. J. (1992) Aryl phosphate derivatives of AZT retain activity against HIV1 in cell lines which are resistant to the action of AZT. *Antiviral Res.* 17, 311-321; McGuigan, C., Pathirana, R. N., Choi, S. M., Kinchington, D. and O'Connor, T. J. (1993a) Phosphoramidate derivatives of AZT as inhibitors of HIV; studies on the carboxyl terminus. *Antiviral Chem. Chemother.* 4, 97-101;

McGuigan, C., Pathirana, R. N., Balzarini, J. and De Clercq, E. (1993b) Intracellular delivery of bioactive AZT nucleotides by aryl phosphate derivatives of AZT. *J. Med. Chem.* 36, 1048-1052.

Alkyl hydrogen phosphonate derivatives of the anti-HIV agent AZT may be less toxic than the parent nucleoside analogue. *Antiviral Chem. Chemother.* 5, 271-277; Meyer, R. B., Jr., Shuman, D. A. and Robins, R. K. (1973) Synthesis of purine nucleoside 3',5'-cyclic Phosphoramidates. *Tetrahedron Lett.* 269-272; Nagyvary, J. Gohil, R. N., Kirchner, C. R. and Stevens, J. D. (1973) Studies on neutral esters of cyclic AMP, *Biochem. Biophys. Res. Commun.* 55, 1072-1077; Namane, A. Goyette, C., Fillion, M. P., Fillion, G and Huynh-Dinh, T. (1992) Improved brain delivery of AZT using a glycosyl phosphotriester prodrug. *J. Med. Chem.* 35, 3939-3044; Nargeot, J. Nerbonne, J. M. Engels, J. and Leser, H. A. (1983) *Natl. Acad. Sci. U.S.A.* 80, 2395-2399; Nelson, K. A., Bentrude, W. G., Stser, W. N. and Hutchinson, J. P. (1987) The question of chair-twist equilibria for the phosphate rings of nucleoside cyclic 3',5'-monophosphates. ^1H NMR and x-ray crystallographic study of the diastereomers of thymidine phenyl cyclic 3',5'-monophosphate. *J. Am. Chem. Soc.* 109, 4058-4064; Nerbonne, J. M., Richard, S., Nargeot, J. and Lester, H. A. (1984) New photoactivatable cyclic nucleotides produce intracellular jumps in cyclic AMP and cyclic GMP concentrations. *Nature* 301, 74-76; Neumann, J. M., Hervé, M., Debouzy, J. C., Guerra, F. I., Gouyette, C., Dupraz, B. and Huynh-Dinh, T. (1989) Synthesis and transmembrane transport studies by NMR of a glucosyl phospholipid of thymidine. *J. Am. Chem. Soc.* 111, 4270-4277; Ohno, R., Tatsumi, N., Hirano, M., Imai, K., Mizoguchi, H., Nakamura, T., Kosaka, M., Takatuski, K., Yamaya, T., Toyama, K., Yoshida, T., Masaoka, T., Hashimoto, S., Ohshima, T., Kimura, I., Yamada, K. and Kimura, J. (1991) Treatment of myelodysplastic syndromes with orally administered 1- β -D-rabinofuranosylcytosine-5'-stearylphosphate. *Oncology* 48, 451-455. Palomino, E., Kessle, D. and Horwitz, J. P. (1989) A dihydropyridine carrier system for sustained delivery of 2',3' dideoxynucleosides to the brain. *J. Med. Chem.* 32, 622-625; Perkins, R. M., Barney, S., Wittrock, R., Clark, P. H., Levin, R. Lambert, D. M., Petteway, S. R., Serafinowska, H. T., Bailey, S. M., Jackson, S., Hamden, M. R., Ashton, R., Sutton, D., Harvey, J. J. and Brown, A. G. (1993) Activity of BRL47923 and its oral prodrug, SB203657A against a rauscher murine leukemia virus infection in mice. *Antiviral Res.* 20 (Suppl. I): 84; Piantadosi, C., Marasco, C. J., Jr., Morris-Natschke, S. L., Meyer, K. L., Gumus, F., Surles, J. R., Ishaq, K. S., Kucera, L. S. Iyer, N., Wallen, C. A., Piantadosi, S. and Modest, E. J. (1991) Synthesis and evaluation of novel ether lipid nucleoside conjugates for anti-HIV-1 activity. *J. Med. Chem.* 34, 1408-1414; Pompon, A., Lefebvre, I., Imbach, J. L., Kahn, S. and Farquhar, D. (1994) Decomposition pathways of the mono- and bis(pivaloyloxymethyl) esters of azidothymidine-5'-monophosphate in cell extract and in tissue culture medium; an application of the on-line ISRP-cleaning HPLC technique. *Antiviral Chem. Chemother.* 5, 91-98; Postemark, T. (1974) Cyclic AMP and cyclic GMP. *Anu. Rev. Pharmacol.* 14, 23-33; Prisbe, E. J., Martin, J. C. M., McGee, D. P. C., Barker, M. F., Smee, D. P. Duke, A. E., Matthews, T. R. and Verheyden, J. P. J. (1986) Synthesis and antihherpes virus activity of phosphate and phosphonate derivatives of 9-[(1,3-dihydroxy-2-propoxy)methyl]guanine. *J. Med. Chem.* 29, 671-675; Pucchi, F., Gosselin, G., Lefebvre, I., Pompon, A., Aubertin, A. M. Dim, A. and Imbach, J. L. (1993) Intracellular delivery of nucleoside monophosphate through a reductase-mediated activation process. *Antiviral Res.* 22, 155-174; Pugaeva, V. P., Kochkeva, S. I., Mashbits, F.

D. and Eizengart, R. S. (1969). Toxicological assessment and health standard ratings for ethylene sulfide in the industrial atmosphere. *Gig. Trf. Prof. Zabol.* 13, 47-48 (*Chem. Abstr.* 72, 212); Robins, R. K. (1984) The potential of nucleotide analogs as inhibitors of retroviruses and tumors. *Pharm. Res.* 11-18; Rosowsky, A., Kim, S. H., Ross and J. Wick, M. M. (1982) Lipophilic 5'-(alkylphosphate) esters of 1- β -D-arabinofuranosylcytosine and its N'-acyl and 2,2'-anhydro-3'-O-acyl derivatives as potential prodrugs. *J. Med. Chem.* 25, 171-178; Ross, W. (1961) Increased sensitivity of the walker turnout towards aromatic nitrogen mustards carrying basic side chains following glucose pretreatment. *Biochem. Pharm.* 8, 235-240; Ryu, E. K., Ross, R. J., Matsushita, T., MacCoss, M., Hong, C. I. and West, C. R. (1982). Phospholipid-nucleoside conjugates 3. Synthesis and preliminary biological evaluation of 1- β -D-arabinofuranosylcytosine 5' diphosphate[-], 2-diacylglycerols. *J. Med. Chem.* 25, 1322-1329; Saffhill, R. and Hume, W. J. (1986) The degradation of 5-iododeoxyuridine and 5-bromodeoxyuridine by serum from different sources and its consequences for the use of these compounds for incorporation into DNA. *Chem. Biol. Interact.* 57, 347-355; Saneyoshi, M., Morozumi, M., Kodama, K., Machida, J., Kuninaka, A. and Yoshino, H. (1980) Synthetic nucleosides and nucleotides XVI. Synthesis and biological evaluations of a series of 1- β -D-arabinofuranosylcytosine 5'-alkyl or arylphosphates. *Chem. Pharm. Bull.* 28, 2915-2923; Sastry, J. K., Nehete, P. N., Khan, S., Nowak, B. J., Plunkett, W., Arlinghaus, R. B. and Farquhar, D. (1992) Membrane-permeable dideoxyuridine 5'-monophosphate analogue inhibits human immunodeficiency virus infection. *Mol. Pharmacol.* 41, 441-445; Shaw, J. P., Jones, R. J. Arimilli, M. N., Louie, M. S., Lee, W. A. and Cundy, K. C. (1994) Oral bioavailability of PMEAs from PMEAs prodrugs in male Sprague-Dawley rats. 9th Annual AAPS Meeting. San Diego, Calif. (Abstract). Shuto, S., Ueda, S., Imamura, S., Fukukuawa, K., Matsuda, A. and Ueda, T. (1987) A facile one-step synthesis of 5'-phosphatidyl nucleosides by an enzymatic two-phase reaction. *Tetrahedron Lett.* 28, 199-202; Shuto, S., Itoh, H., Ueda, S., Imamura, S., Kukukawa, K., Tsujino, M., Matsuda, A. and Ueda, T. (1988) A facile enzymatic synthesis of 5'-(3-sn-phosphatidyl)nucleosides and their antileukemic activities. *Chem. Pharm. Bull.* 36, 209-217. One preferred phosphate prodrug group is the S-acyl-2-thioethyl group, also referred to as "SATE."

Additional examples of prodrugs that can be used are those described in the following patents and patent applications: U.S. Pat. Nos. 5,614,548, 5,512,671, 5,770,584, 5,962,437, 5,223,263, 5,817,638, 6,252,060, 6,448,392, 5,411,947, 5,744,592, 5,484,809, 5,827,831, 5,696,277, 6,022,029, 5,780,617, 5,194,654, 5,463,092, 5,744,461, 4,444,766, 4,562,179, 4,599,205, 4,493,832, 4,221,732, 5,116,992, 6,429,227, 5,149,794, 5,703,063, 5,888,990, 4,810,697, 5,512,671, 6,030,960, 2004/0259845, 6,670,341, 2004/0161398, 2002/082242, 5,512,671, 2002/0082242, and or PCT Publication Nos WO 90/11079, WO 96/39197, and/or WO 93/08807.

DEFINITIONS

As used herein, the terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth, i.e., proliferative disorders. Examples of such proliferative disorders include cancers such as carcinoma, lymphoma, blastoma, sarcoma, and leukemia, as well as other cancers disclosed herein. More particular examples of such cancers include breast cancer, prostate cancer, colon cancer, squamous cell

cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, e.g., hepatic carcinoma, bladder cancer, colorectal cancer, endometrial carcinoma, kidney cancer, and thyroid cancer.

Other non-limiting examples of cancers are basal cell carcinoma, biliary tract cancer; bone cancer, brain and CNS cancer, choriocarcinoma; connective tissue cancer; esophageal cancer, eye cancer; cancer of the head and neck; gastric cancer; intra-epithelial neoplasm; larynx cancer; lymphoma including Hodgkin's and Non-Hodgkin's lymphoma; melanoma; myeloma; neuroblastoma; oral cavity cancer (e.g., lip, tongue, mouth, and pharynx); pancreatic cancer, retinoblastoma; rhabdomyosarcoma; rectal cancer; cancer of the respiratory system; sarcoma; skin cancer, stomach cancer, testicular cancer, uterine cancer; cancer of the urinary system, as well as other carcinomas and sarcomas.

As used herein, the term "tumor" refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pro-cancerous and cancerous cells and tissues. For example, a particular cancer may be characterized by a solid mass tumor. The solid tumor mass, if present, may be a primary tumor mass. A primary tumor mass refers to a growth of cancer cells in a tissue resulting from the transformation of a normal cell of that tissue. In most cases, the primary tumor mass is identified by the presence of a cyst, which can be found through visual or palpation methods, or by irregularity in shape, texture or weight of the tissue. However, some primary tumors are not palpable and can be detected only through medical imaging techniques such as X-rays (e.g., mammography), or by needle aspirations. The use of these latter techniques is more common in early detection. Molecular and phenotypic analysis of cancer cells within a tissue will usually confirm if the cancer is endogenous to the tissue or if the lesion is due to metastasis from another site.

The term alkyl, as used herein, unless otherwise specified, includes a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of for example C₁ to C₂₄, and specifically includes methyl, trifluoromethyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, t-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. The alkyl is optionally substituted, e.g., with one or more substituents such as halo (F, Cl, Br or I), (e.g. CF₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃ or CF₃CF₃), hydroxyl (e.g. CH₂OH), amino (e.g. CH₂NH₂, CH₂NHCH₃ or CH₂N(CH₃)₂), alkylamino, arylamino, alkoxy, aryloxy, nitro, azido (e.g. CH₂N₃), cyano (e.g. CH₂CN), sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene, et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, Second Edition, 1991, hereby incorporated by reference.

The term lower alkyl, as used herein, and unless otherwise specified, refers to a C₁ to C₄ saturated straight, branched, or if appropriate, a cyclic (for example, cyclopropyl) alkyl group, including both substituted and unsubstituted forms.

The term alkylamino or arylamino includes an amino group that has one or two alkyl or aryl substituents, respectively.

The term amino acid includes naturally occurring and synthetic α , β , γ or δ amino acids, and includes but is not limited to, amino acids found in proteins, i.e. glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histi-

dine. In a preferred embodiment, the amino acid is in the L-configuration. Alternatively, the amino acid can be a derivative of alanyl, valinyl, leucinyl, isoleuccinyl, prolinyl, phenylalaninyl, tryptophanyl, methioninyl, glycyl, serinyl, threoninyl, cysteinyl, tyrosinyl, asparaginyl, glutaminyl, aspartoyl, glutaroyl, lysinyl, argininyl, histidinyl, β -valinyl, β -leucinyl, β -isoleuccinyl, β -prolinyl, β -phenylalaninyl, β -tryptophanyl, β -methioninyl, β -glycyl, β -serinyl, β -threoninyl, β -cysteinyl, β -tyrosinyl, β -asparaginyl, β -glutaminyl, β -aspartoyl, β -glutaroyl, β -lysinyl, β -argininyl or β -histidinyl. When the term amino acid is used, it is considered to be a specific and independent disclosure of each of the esters of a natural or synthetic amino acid, including but not limited to α , β , γ or δ glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine in the D and L-configurations.

The term "protected" as used herein and unless otherwise defined includes a group that is added to an oxygen, nitrogen, sulfur or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis (see Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, inc., New York, N.Y., 1999).

The term aryl, as used herein, and unless otherwise specified, includes phenyl, biphenyl, or naphthyl, and preferably phenyl. The aryl group is optionally substituted with one or more moieties such as halo, hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene, et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, 3rd Ed., 1999.

The term alkaryl or alkylaryl includes an alkyl group with an aryl substituent. The term aralkyl or arylalkyl includes an aryl group with an alkyl substituent.

The term halo, as used herein, includes chloro, bromo, iodo, and fluoro.

The tetra acyl includes a carboxylic acid ester in which the non-carbonyl moiety of the ester group is selected from straight, branched, or cyclic alkyl or lower alkyl, alkoxyalkyl including methoxymethyl, aralkyl including benzyl, aryloxyalkyl such as phenoxyethyl, aryl including phenyl optionally substituted with halogen, C₁ to C₄ alkyl or C₁ to C₄ alkoxy, sulfonate esters such as alkyl or aralkyl sulfonyl including methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxytrityl, substituted benzyl, trialkylsilyl (e.g. dimethyl-t-butylsilyl) or diphenylmethylsilyl. Aryl groups in the esters optimally comprise a phenyl group. The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

As used herein, the term "substantially free of" or "substantially in the absence of" with respect to enantiomeric purity, refers to a composition that includes at least 85% or 90% by weight, preferably 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the designated enantiomer. In a preferred embodiment, in the methods and compounds of this invention, the compounds are substantially free of other enantiomers.

Similarly, the term "isolated" refers to a compound composition that includes at least 85% or 90% by weight, preferably 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the compound, the remainder comprising other chemical species or enantiomers.

The term "independently" is used herein to indicate that the variable, which is independently applied, varies independently from application to application. Thus, in a compound such as R"XYR", wherein R" is "independently carbon or nitrogen," both R" can be carbon, both R" can be nitrogen, or one R" can be carbon and the other R" nitrogen.

The term "pharmaceutically acceptable salt or prodrug" is used throughout the specification to describe any pharmaceutically acceptable form (such as an ester, phosphate ester, salt of an ester or a related group) of a compound, which, upon administration to a patient, provides the compound. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. Pharmaceutically acceptable prodrugs refer to a compound that is metabolized, for example hydrolyzed or oxidized, in the host to form the compound of the present invention. Typical examples of prodrugs include compounds that have biologically labile protecting groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, dephosphorylated to produce the active compound.

The term "pharmaceutically acceptable esters" as used herein, unless otherwise specified, includes those esters of one or more compounds, which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of hosts without undue toxicity, irritation, allergic response and the like, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.

The term "subject" as used herein refers to an animal, preferably a mammal, most preferably a human. Mammals can include non-human mammals, including, but not limited to, pigs, sheep, goats, cows (bovine), deer, mules, horses, monkeys and other non-human primates, dogs, cats, rats, mice, rabbits or any other known or disclosed herein.

II. In Vivo Efficacy/Dosing Regimens

In another aspect of the present invention, dosing regimens are provided that limit the toxic side effects of TCN and related compounds. In one embodiment, such dosing regimens minimize the following toxic side effects, including, but not limited to, hepatotoxicity, thrombocytopenia, hyperglycemia, vomiting, hypocalcemia, anemia, hypoalbuminemia, myelosuppression, hypertriglyceridemia, hyperamylasemia, diarrhea, stomatitis and/or fever.

In another embodiment, the administration of TCN, TCN-P or related compounds provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with esophageal adenocarcinoma by administering to the subject an effective amount of TCN, TCN-P or a related compound according to a dosing schedule that includes administering the drug one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e. via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the drug is not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m² of TCN, TCN-P or a related compound can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 mg/m² or TCN, TCN-P or a related compound can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat esophageal adenocarcinoma in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P or a related compound one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P or a related compound as disclosed herein can be administered one time per week.

In embodiments of the present invention, the compound disclosed herein can be administered as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the drug via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The drugs can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The drug can be administered according to any combination disclosed herein, for example, the drug can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the drug is not administered.

The TCN, TCN-P and related compounds as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P or related compounds can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a compound disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P or a related compound disclosed herein can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than 20 mg/m² of TCN and related compounds. In one embodiment, less than 20 mg/m² of TCN or related compounds can be administered once a week. In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15

mg/m² of TCN or a related compound can be administered to a subject. In another embodiment, less than 10 mg/m² can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

III. Screening of Patient Populations

In another aspect of the present invention, methods are provided to identify esophageal adenocarcinoma susceptible to the toxic effects of tricitiribine (TCN) and related compounds. In one embodiment, methods are provided to esophageal adenocarcinoma in a mammal by (i) obtaining a biological sample from the tumor; (ii) determining whether the cancer or tumor overexpresses Akt kinase or hyperactivated and phosphorylated Akt kinase, and (iii) treating the cancer or tumor with tricitiribine or a related compound as described herein. In one embodiment, the biological sample can be a biopsy. In other embodiments, the biological sample can be fluid, cells and/or aspirates obtained from the tumor or cancer.

The biological sample can be obtained according to any technique known to one skilled in the art. In one embodiment, a biopsy can be conducted to obtain the biological sample. A biopsy is a procedure performed to remove tissue or cells from the body for examination. Some biopsies can be performed in a physician's office, while others need to be done in a hospital setting. In addition, some biopsies require use of an anesthetic to numb the area, while others do not require any sedation. In certain embodiments, an endoscopic biopsy can be performed. This type of biopsy is performed through a fiberoptic endoscope (a long, thin tube that has a close-focusing telescope on the end for viewing) through a natural body orifice (i.e., mouth) or a small incision (i.e., arthroscopy). The endoscope is used to view the organ in question for abnormal or suspicious areas, in order to obtain a small amount of tissue for study. Endoscopic procedures are named for the organ or body area to be visualized and/or treated. The physician can insert the endoscope into the gastrointestinal tract (alimentary tract endoscopy), bladder (cystoscopy), abdominal cavity (laparoscopy), joint cavity (arthroscopy), mid-portion of the chest (mediastinoscopy), or trachea and bronchial system (laryngoscopy and bronchoscopy).

In another embodiment, a bone marrow biopsy can be performed. This type of biopsy can be performed either from the sternum (breastbone) or the iliac crest hipbone (the bone area on either side of the pelvis on the lower back area). The skin is cleansed and a local anesthetic is given to numb the area. A long, rigid needle is inserted into the marrow, and cells are aspirated for study; this step is occasionally uncomfortable. A core biopsy (removing a small bone 'chip' from the marrow) may follow the aspiration.

In a further embodiment, an excisional or incisional biopsy can be performed on the mammal. This type of biopsy is often used when a wider or deeper portion of the skin is needed. Using a scalpel (surgical knife), a full thickness of skin is removed for further examination, and the wound is sutured (sewed shut with surgical thread). When the entire tumor is removed, it is referred to as an excisional biopsy technique. If only a portion of the tumor is removed, it is referred to as an incisional biopsy technique. Excisional biopsy is often the method usually preferred, for example, when melanoma (a type of skin cancer) is suspected.

In still further embodiments, a fine needle aspiration (FNA) biopsy can be used. This type of biopsy involves using a thin needle to remove very small pieces from a tumor. Local anesthetic is sometimes used to numb the area, but the test rarely causes much discomfort and leaves no scar. FNA is not,

for example, used for diagnosis of a suspicious mole, but may be used, for example, to biopsy large lymph nodes near a melanoma to see if the melanoma has metastasized (spread). A computed-tomography scan (CT or CAT scan) can be used to guide a needle into a tumor in an internal organ such as the lung or liver.

In other embodiments, punch shave and/or skin biopsies can be conducted. Punch biopsies involve taking a deeper sample of skin with a biopsy instrument that removes a short cylinder, or "apple core," of tissue. After a local anesthetic is administered, the instrument is rotated on the surface of the skin until it cuts through all the layers, including the dermis, epidermis, and the most superficial parts of the subcutis (fat). A shave biopsy involves removing the top layers of skin by shaving it off. Shave biopsies are also performed with a local anesthetic. Skin biopsies involve removing a sample of skin for examination under the microscope to determine if, for example, melanoma is present. The biopsy is performed under local anesthesia.

In particular embodiment, methods are provided to determine whether the tumor overexpresses an Akt kinase. Akt kinase overexpression can refer to the phosphorylation state of the kinase. Hyperphosphorylation of Akt can be detected according to the methods described herein. In one embodiment, a tumor biopsy can be compared to a control tissue. The control tissue can be a normal tissue from the mammal in which the biopsy was obtained or a normal tissue from a healthy mammal. Akt kinase overexpression or hyperphosphorylation can be determined if the tumor biopsy contains greater amounts of Akt kinase and/or Akt kinase phosphorylation than the control tissue, such as, for example, at least approximately 1.5, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 5.5, 6, 7, 8, 9, or 10-fold greater amounts of Akt kinase than contained in the control tissue.

In one embodiment, the present invention provides a method to detect aberrant Akt kinase expression in a subject or in a biological sample from the subject by contacting cells, cell extracts, serum or other sample from the subjects or said biological sample with an immunointeractive molecule specific for an Akt kinase or antigenic portion thereof and screening for the level of immunointeractive molecule-Akt kinase complex formation, wherein an elevated presence of the complex relative to a normal cell is indicative of an aberrant cell that expresses or overexpresses Akt. In one example, cells or cell extracts can be screened immunologically for the presence of elevated levels of Akt kinase.

In an alternative embodiment, the aberrant expression of Akt in a cell is detected at the genetic level by screening for the level of expression of a gene encoding an Akt kinase wherein an elevated level of a transcriptional expression product (i.e. mRNA) compared to a normal cell is indicative of an aberrant cell. In certain embodiments, real-time PCR as well as other PCR procedures can be used to determine transcriptional activity. In one embodiment, mRNA can be obtained from cells of a subject or from a biological sample from a subject and cDNA optionally generated. The mRNA or cDNA can then be contacted with a genetic probe capable of hybridizing to and/or amplifying all or part of a nucleotide sequence encoding Akt kinase or its complementary nucleotide sequence and then the level of the mRNA or cDNA can be detected wherein the presence of elevated levels of the mRNA or cDNA compared to normal controls can be assessed.

Yet another embodiment of the present invention contemplates the use of an antibody, monoclonal or polyclonal, to Akt kinase in a quantitative or semi-quantitative diagnostic kit to determine relative levels of Akt kinase in suspected

cancer cells from a patient, which can include all the reagents necessary to perform the assay. In one embodiment, a kit utilizing reagents and materials necessary to perform an ELISA assay is provided. Reagents can include, for example, washing buffer, antibody dilution buffer, blocking buffer, cell staining solution, developing solution, stop solution, anti-phospho-protein specific antibodies, anti-Pan protein specific antibodies, secondary antibodies, and distilled water. The kit can also include instructions for use and can optionally be automated or semi-automated or in a form which is compatible with automated machine or software. In one embodiment, a phosphor-ser-473 Akt antibody that detects the activated form of AKT (Akt phosphorylated at serine 474) can be utilized as the antibody in a diagnostic kit. See, for example, Yuan et al. (2000) "Frequent Activation of AKT2 and induction of apoptosis by inhibition of phosphoinositide-3-OH kinase/Akt pathway in human ovarian cancer," *Oncogene* 19:2324-2330.

Akt Kinases

Akt, also named PKB³, represents a subfamily of the serine/threonine kinase. Three members, AKT1, AKT2, and AKT3, have been identified in this subfamily. Akt is activated by extracellular stimuli in a PI3K-dependent manner (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). Full activation of Akt requires phosphorylation of Thr³⁰⁸ in the activation loop and Ser⁴⁷³ in the C-terminal activation domain. Akt is negatively regulated by PTEN tumor suppressor. Mutations in PTEN have been identified in various tumors, which lead to activation of Akt pathway (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). In addition, amplification, overexpression and/or activation of Akt have been detected in a number of human malignancies (Dana, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999, Cheng, J. Q., and Nicosia, S. V. AKT signal transduction pathway in oncogenesis. In Schwab D, editor. *Encyclopedic Reference of Cancer*. Berlin Heidelberg and New York: Springer, 2001. pp 35-7). Ectopic expression of Akt, especially constitutively active Akt, induces cell survival and malignant transformation whereas inhibition of Akt activity stimulates apoptosis in a range of mammalian cells (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999, Cheng, J. Q., and Nicosia, S. V. AKT signal transduction pathway in oncogenesis. In Schwab D, editor. *Encyclopedic Reference of Cancer*. Berlin Heidelberg and New York: Springer, 2001. pp 35-7, Sun, M., et al. *Am. J. Pathol.*, 159: 431-437, 2001, Cheng, J. Q., et al. *Oncogene*, 14: 2793-2801, 1997). Further, activation of Akt has been shown to associate with tumor invasiveness and chemoresistance (West, K. A., et al. *Drug Resist. Updat.*, 5: 234-248, 2002).

Activation of the Akt pathway plays a pivotal role in malignant transformation and chemoresistance by inducing cell survival, growth, migration, and angiogenesis. The present invention provides methods to determine levels of Akt kinase overexpression and/or hyperactivated and phosphorylated Akt kinase.

The Akt kinase can be any known Akt family kinase, or kinase related thereto, including, but not limited to Akt 1, Akt 2, Akt 3. The mRNA and amino acid sequences of human Akt1, Akt2, and Akt 3 are illustrated in FIGS. 6a-c, 7a-d, and 8a-c, respectively.

Diagnostic Assays

Immunological Assays

In one embodiment, a method is provided for detecting the aberrant expression of an Akt kinase in a cell in a mammal or in a biological sample from the mammal, by contacting cells, cell extracts or serum or other sample from the mammal or biological sample with an immuno interactive molecule specific for an Akt kinase or antigenic portion thereof and screen-

ing for the level of immunointeractive molecule-Akt kinase complex formations and determining whether an elevated presence of the complex relative to a normal cell is present.

The immunointeractive molecule can be a molecule having specificity and binding affinity for an Akt kinase or its antigenic parts or its homologs or derivatives thereof. In one embodiment, the immunointeractive molecule can be an immunoglobulin molecule. In other embodiments, the immunointeractive molecules can be an antibody fragments, single chain antibodies, and/or deimmunized molecules including humanized antibodies and T-cell associated antigen-binding molecules (TABMs). In one particular embodiment, the antibody can be a monoclonal antibody. In another particular embodiment, the antibody can be a polyclonal antibody. The immunointeractive molecule can exhibit specificity for an Akt kinase or more particularly an antigenic determinant or epitope on an Akt kinase. An antigenic determinant or epitope on an Akt kinase includes that part of the molecule to which an immune response is directed. The antigenic determinant or epitope can be a B-cell epitope or where appropriate a T-cell epitope. In one embodiment, the antibody is a phosphor-ser 473 Akt antibody.

One embodiment of the present invention provides a method for diagnosing the presence of cancer or cancer-like growth in a mammal, in which aberrant Akt activity is present, by contacting cells or cell extracts from the mammal or a biological sample from the subject with an Akt kinase-binding effective amount of an antibody having specificity for the Akt kinase or an antigenic determinant or epitope thereon and then quantitatively or qualitatively determining the level of an Akt kinase-antibody complex wherein the presence of elevated levels of said complex compared to a normal cell is determined.

Antibodies can be prepared by any of a number of means known to one skilled in the art. For example, for the detection of human Akt kinase, antibodies can be generally but not necessarily derived from non-human animals such as primates, livestock animals (e.g. sheep, cows, pigs, goats, horses), laboratory test animals (e.g. mice, rats, guinea pigs, rabbits) and/or companion animals (e.g. dogs, cats). Antibodies may also be recombinantly produced in prokaryotic or eukaryotic host cells. Generally, antibody based assays can be conducted in vitro on cell or tissue biopsies. However, if an antibody is suitably deimmunized or, in the case of human use, humanized, then the antibody can be labeled with, for example, a nuclear tag, administered to a patient and the site of nuclear label accumulation determined by radiological techniques. The Akt kinase antibody can be a cancer targeting agent. Accordingly, another embodiment of the present invention provides deimmunized forms of the antibodies for use in cancer imaging in human and non-human patients.

In general, for the generation of antibodies to an Akt kinase, the enzyme is required to be extracted from a biological sample whether this be from animal including human tissue or from cell culture if produced by recombinant means. The Akt kinase can be separated from the biological sample by any suitable means. For example, the separation may take advantage of any one or more of the Akt kinase's surface charge properties, size, density, biological activity and its affinity for another entity (e.g. another protein or chemical compound to which it binds or otherwise associates). Thus, for example, separation of the Akt kinase from the biological fluid can be achieved by any one or more of ultra-centrifugation, ion-exchange chromatography (e.g. anion exchange chromatography, cation exchange chromatography), electrophoresis (e.g. polyacrylamide gel electrophoresis, isoelectric focussing), size separation (e.g., gel filtration, ultra-filtration)

and affinity-mediated separation (e.g. immunoaffinity separation including, but not limited to, magnetic bead separation such as Dynabead™ separation, immunochromatography, immuno-precipitation). The separation of Akt kinase from the biological fluid can preserve conformational epitopes present on the kinase and, thus, suitably avoids techniques that cause denaturation of the enzyme. In a further embodiment, the kinase can be separated from the biological fluid using any one or more of affinity separation, gel filtration and/or ultra-filtration.

Immunization and subsequent production of monoclonal antibodies can be carried out using standard protocols known in the art, such as, for example, described by Kohler and Milstein (Kohler and Milstein, *Nature* 256: 495-499, 1975; Kohler and Milstein, *Eur. J. Immunol.* 6(7): 511-519, 1976), Coligan et al ("Current Protocols in Immunology, John Wiley & Sons, Inc., 1991-1997) or Toyama et al. (Monoclonal Antibody, Experiment Manual", published by Kodansha Scientific, 1987). Essentially, an animal is immunized with an Akt kinase-containing biological fluid or fraction thereof or a recombinant form of Akt kinase by standard methods to produce antibody-producing cells, particularly antibody-producing somatic cells (e.g. B lymphocytes). These cells can then be removed from the immunized animal for immortalization. In certain embodiment, a fragment of an Akt kinase can be used to generate antibodies. The fragment can be associated with a carrier. The carrier can be any substance of typically high molecular weight to which a non- or poorly immunogenic substance (e.g. a hapten) is naturally or artificially linked to enhance its immunogenicity.

Immortalization of antibody-producing cells can be carried out using methods which are well-known in the art. For example, the immortalization may be achieved by the transformation method using Epstein-Barr virus (EBV) (Kozbor et al., *Methods in Enzymology* 121: 140, 1986). In another embodiment, antibody-producing cells are immortalized using the cell fusion method (described in Coligan et al., 1991-1997, supra), which is widely employed for the production of monoclonal antibodies. In this method, somatic antibody-producing cells with the potential to produce antibodies, particularly B cells, are fused with a myeloma cell line. These somatic cells may be derived from the lymph nodes, spleens and peripheral blood of primed animals, preferably rodent animals such as mice and rats. In a particular embodiment, mice spleen cells can be used. In other embodiments, rat, rabbit, sheep or goat cells can also be used. Specialized myeloma cell lines have been developed from lymphocytic tumours for use in hybridoma-producing fusion procedures (Kohler and Milstein, 1976, supra; Shulman et al., *Nature* 276: 269-270, 1978; Volk et al., *J. Virol.* 42(1): 220-227, 1982). Many myeloma cell lines can also be used for the production of fused cell hybrids, including, e.g. P3.times.63-Ag8, P3.times.63-AG8.653, P3/NS1-Ag4-1 (NS-1), Sp2/0-Ag14 and S194/5.XXO.Bu.1. The P3.times.63-Ag8 and NS-1 cell lines have been described by Kohler and Milstein (1976, supra). Shulman et al. (1978, supra) developed the Sp2/0-Ag14 myeloma line. The S194/5.XXO.Bu.1 line was reported by Trowbridge (*J. Exp. Med.* 148(1): 313-323, 1978). Methods for generating hybrids of antibody-producing spleen or lymph node cells and myeloma cells usually involve mixing somatic cells with myeloma cells in a 10:1 proportion (although the proportion may vary from about 20:1 to about 1:1), respectively, in the presence of an agent or agents (chemical, viral or electrical) that promotes the fusion of cell membranes. Fusion methods have been described (Kohler and Milstein, 1975, supra; Kohler and Milstein, 1976, supra; Gefter et al., *Somatic Cell Genet.* 3: 231-236,

1977; Volk et al., 1982, supra). The fusion-promoting agents used by those investigators were Sendai virus and polyethylene glycol (PEG). In certain embodiments, means to select the fused cell hybrids from the remaining unfused cells, particularly the unfused myeloma cells, are provided. Generally, the selection of fused cell hybrids can be accomplished by culturing the cells in media that support the growth of hybridomas but prevent the growth of the unfused myeloma cells, which normally would go on dividing indefinitely. The somatic cells used in the fusion do not maintain long-term viability in in vitro culture and, hence do not pose a problem. Several weeks are required to selectively culture the fused cell hybrids. Early in this time period, it is necessary to identify those hybrids which produce the desired antibody, so that they may subsequently be cloned and propagated. Generally, around 10% of the hybrids obtained produce the desired antibody, although a range of from about 1 to about 30% is not uncommon. The detection of antibody-producing hybrids can be achieved by any one of several standard assay methods, including enzyme-linked immunoassay and radioimmunoassay techniques as, for example, described in Kennet et al. (Monoclonal Antibodies and Hybridomas: A New Dimension in Biological Analyses, pp 376-384, Plenum Press, New York, 1980) and by FACS analysis (O'Reilly et al., *Biotechniques* 25: 824-830, 1998).

Once the desired fused cell hybrids have been selected and cloned into individual antibody-producing cell lines, each cell line may be propagated in either of two standard ways. A suspension of the hybridoma cells can be injected into a histocompatible animal. The injected animal will then develop tumours that secrete the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can be tapped to provide monoclonal antibodies in high concentration. Alternatively, the individual cell lines may be propagated in vitro in laboratory culture vessels. The culture medium containing high concentrations of a single specific monoclonal antibody can be harvested by decantation, filtration or centrifugation, and subsequently purified.

The cell lines can then be tested for their specificity to detect the Akt kinase of interest by any suitable immunodetection means. For example, cell lines can be aliquoted into a number of wells and incubated and the supernatant from each well is analyzed by enzyme-linked immunosorbent assay (ELISA), indirect fluorescent antibody technique, or the like. The cell line(s) producing a monoclonal antibody capable of recognizing the target LIM kinase but which does not recognize non-target epitopes are identified and then directly cultured in vitro or injected into a histocompatible animal to form tumours and to produce, collect and purify the required antibodies.

The present invention provides, therefore, a method of detecting in a sample an Akt kinase or fragment, variant or derivative thereof comprising contacting the sample with an antibody or fragment or derivative thereof and detecting the level of a complex containing the antibody and Akt kinase or fragment, variant or derivative thereof compared to normal controls wherein elevated levels of Akt kinase is determined. Any suitable technique for determining formation of the may be used. For example, an antibody according to the invention, having a reporter molecule associated therewith, may be utilized in immunoassays. Such immunoassays include but are not limited to radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs) immunochromatographic techniques (ICTs), and Western blotting which are well known to those of skill in the art. Immunoassays can also

include competitive assays. The present invention encompasses qualitative and quantitative immunoassays.

Suitable immunoassay techniques are described, for example, in U.S. Pat. Nos. 4,016,043, 4,424,279 and 4,018,653. These include both single-site and two-site assays of the non-competitive types, as well as the traditional competitive binding assays. These assays also include direct binding of a labeled antigen-binding molecule to a target antigen.

The invention further provides methods for quantifying Akt protein expression and activation levels in cells or tissue samples obtained from an animal, such as a human cancer patient or an individual suspected of having cancer. In one embodiment, the invention provides methods for quantifying Akt protein expression or activation levels using an imaging system quantitatively. The imaging system can be used to receive, enhance, and process images of cells or tissue samples, that have been stained with AKT protein-specific stains, in order to determine the amount or activation level of AKT protein expressed in the cells or tissue samples from such an animal. In embodiments of the methods of the invention, a calibration curve of AKT1 and AKT2 protein expression can be generated for at least two cell lines expressing differing amounts of AKT protein. The calibration curve can then be used to quantitatively determine the amount of AKT protein that is expressed in a cell or tissue sample. Analogous calibration curves can be made for activated AKT proteins using reagents specific for the activation features. It can also be used to determine changes in amounts and activation state of AKT before and after clinical cancer treatment.

In one particular embodiment of the methods of the invention, AKT protein expression in a cell or tissue sample can be quantified using an enzyme-linked immunoabsorbent assay (ELISA) to determine the amount of AKT protein in a sample. Such methods are described, for example, in U.S. Patent Publication No. 2002/0015974.

In other embodiments enzyme immunoassays can be used to detect the Akt kinase. In such assays, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. The substrates to be used with the specific enzymes are generally chosen for the production of, upon hydrolysis by the corresponding enzyme, a detectable colour change. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates. The enzyme-labeled antibody can be added to the first antibody-antigen complex, allowed to bind, and then the excess reagent washed away. A solution containing the appropriate substrate can then be added to the complex of antibody-antigen-antibody. The substrate can react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of antigen which was present in the sample. Alternatively, fluorescent compounds, such as fluorescein, rhodamine and the lanthanide, europium (EU), can be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labeled antibody adsorbs the light energy, inducing a state to excitability in the molecule, followed by emission of the light at a characteristic colour visually detectable with a light microscope. The fluorescent-labeled antibody is allowed to bind to the first antibody-antigen complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to light of an appropriate wavelength. The fluorescence observed indicates the presence of the antigen of interest. Immunofluorometric assays (IFMA) are well established in the art and are particularly useful for the present method.

However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules can also be employed.

In a particular embodiment, antibodies to Akt kinase can also be used in ELISA-mediated detection of Akt kinase especially in serum or other circulatory fluid. This can be accomplished by immobilizing anti-Akt kinase antibodies to a solid support and contacting these with a biological extract such as serum, blood, lymph or other bodily fluid, cell extract or cell biopsy. Labeled anti-Akt kinase antibodies can then be used to detect immobilized Akt kinase. This assay can be varied in any number of ways and all variations are encompassed by the present invention and known to one skilled in the art. This approach can enable rapid detection and quantitation of Akt kinase levels using, for example, a serum-based assay.

In one embodiment, an Akt Elisa assay kit may be used in the present invention. For example, a Cellular Activation of Signaling ELISA kit for Akt S473 from SuperArray Bioscience can be utilized in the present invention. In one embodiment, the antibody can be an anti-pan antibody that recognizes Akt S473. Elisa assay kit containing an anti-Akt antibody and additional reagents, including, but not limited to, washing buffer, antibody dilution buffer, blocking buffer, cell staining solution, developing solution, stop solution, secondary antibodies, and distilled water.

Nucleotide Detection

In another embodiment, a method to detect Akt kinases is provided by detecting the level of expression in a cell of a polynucleotide encoding an Akt kinase. Expression of the polynucleotide can be determined using any suitable technique known to one skilled in the art. In one embodiment, a labeled polynucleotide encoding an Akt kinase can be utilized as a probe in a Northern blot of an RNA extract obtained from the cell. In other embodiments, a nucleic acid extract from an animal can be utilized in concert with Oligonucleotide primers corresponding to sense and antisense sequences of a polynucleotide encoding the kinase, or flanking sequences thereof, in a nucleic acid amplification reaction such as RT-PCR. A variety of automated solid-phase detection techniques are also available to one skilled in the art, for example, as described by Fodor et al. (Science 251: 767-777, 1991) and Kazal et al. (Nature Medicine 2: 753-759, 1996).

In other embodiments, methods are provided to detect akt kinase encoding RNA transcripts. The RNA can be isolated from a cellular sample suspected of containing Akt kinase RNA, e.g. total RNA isolated from human cancer tissue. RNA can be isolated by methods known in the art, e.g. using TRIzol reagent (GIBCO-BRL/Life Technologies, Gaithersburg, Md.). Oligo-dT, or random-sequence oligonucleotides, as well as sequence-specific oligonucleotides can be employed as a primer in a reverse transcriptase reaction to prepare first-strand cDNAs from the isolated RNA. Resultant first-strand cDNAs can then be amplified with sequence-specific oligonucleotides in PCR reactions to yield an amplified product.

Polymerase chain reaction or "PCR" refers to a procedure or technique in which amounts of a preselected fragment of nucleic acid, RNA and/or DNA, are amplified as described, for example, in U.S. Pat. No. 4,683,195. Generally, sequence information from the ends of the region of interest or beyond is employed to design oligonucleotide primers. These primers will be identical or similar in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific RNA sequences and cDNA transcribed from total cellular RNA. See generally Mullis et al. (Quant. Biol. 51: 263, 1987; Erlich, eds., PCR Technology, Stockton Press, NY,

1989). Thus, amplification of specific nucleic acid sequences by PCR relies upon oligonucleotides or "primers" having conserved nucleotide sequences wherein the conserved sequences are deduced from alignments of related gene or protein sequences, e.g. a sequence comparison of mammalian Akt kinase genes. For example, one primer is prepared which is predicted to anneal to the antisense strand and another primer prepared which is predicted to anneal to the sense strand of a cDNA molecule which encodes a Akt kinase. To detect the amplified product, the reaction mixture is typically subjected to agarose gel electrophoresis or other convenient separation technique and the relative presence of the Akt kinase specific amplified DNA detected. For example, Akt kinase amplified DNA may be detected using Southern hybridization with a specific oligonucleotide probe or comparing its electrophoretic mobility with DNA standards of known molecular weight. Isolation, purification and characterization of the amplified Akt kinase DNA can be accomplished by excising or eluting the fragment from the gel (for example, see references Lawn et al., *Nucleic Acids Res.* 2: 6103, 1981; Goeddel et al., *Nucleic Acids Res.* 8: 4057-1980), cloning the amplified product into a cloning site of a suitable vector, such as the pCRII vector (Invitrogen), sequencing the cloned insert and comparing the DNA sequence to the known sequence of LIM kinase. The relative amounts of LIM kinase mRNA and cDNA can then be determined.

In one embodiment, real-time PCR can be used to determine transcriptional levels of Akt nucleotides. Determination of transcriptional activity also includes a measure of potential translational activity based on available mRNA transcripts. Real-time PCR as well as other PCR procedures use a number of chemistries for detection of PCR product including the binding of DNA binding fluorophores, the 5' endonuclease, adjacent liner and hairpin oligoprobes and the self-fluorescing amplicons. These chemistries and real-time PCR in general are discussed, for example, in Mackay et al., *Nucleic Acids Res* 30(6): 1292-1305, 2002; Walker, J. *Biochem. Mol. Toxicology.* 15(3): 121-127, 2001; Lewis et al., *J. Pathol.* 195: 66-71, 2001.

In an alternate embodiment, the aberrant expression of Akt can be identified by contacting a nucleotide sequences isolated from a biological sample with an oligonucleotide probe having a sequence complementary to an Akt sequences selected from the nucleotide sequences of FIGS. 6a-c, 7a-d, or 8a-c, or fragment thereof, and then detecting the sequence by hybridizing the probe to the sequence, and comparing the results to a normal sample. The hybridization of the probe to the biological sample can be detected by labeling the probe using any detectable agent. The probe can be labeled for example, with a radioisotope, or with biotin, fluorescent dye, electron-dense reagent, enzyme, hapten or protein for which antibodies are available. The detectable label can be assayed by any desired means, including spectroscopic, photochemical, biochemical, immunochemical, radioisotopic, or chemical means. The probe can also be detected using techniques such as an oligomer restriction technique, a dot blot assay, a reverse dot blot assay, a line probe assay, and a 5' nuclease assay. Alternatively, the probe can be detected using any of the generally applicable DNA array technologies, including microarray, microarray and DNA microchip technologies. The oligonucleotide probe typically includes approximately at least 14, 15, 16, 18, 20, or 28 nucleotides that hybridize to the nucleotides selected from FIGS. 6a-c, 7a-d, and 8a-c, or a fragment thereof. It is generally not preferred to use a probe

that is greater than approximately 25 or 28 nucleotides in length. The oligonucleotide probe is designed to identify an Akt nucleotide sequence.

Kinase Assays

The activity of the Akt kinases can be measured using any suitable kinase assay known in the art. For example, and not by way of limitation, the methods described in Hogg et al (*Oncogene* 1994 9:98-96), Mills et al (*J. Biol. Chem.* 1992 267:16000-006) and Tomizawa et al 2001 (*FEBS Lett.* 2001 492: 221-7), Schmandt et al, (*J. Immunol.* 1994, 152:96-105) can be used. Further serine, threonine and tyrosine kinase assays are described in Ausubel et al. (*Short Protocols in Molecular Biology*, 1999, unit 17.6).

Akt kinase assays can generally use an Akt polypeptide, a labeled donor substrate, and a receptor substrate that is either specific or non-specific for Akt. In such assays Akt transfers a labeled moiety from the donor substrate to the receptor substrate, and kinase activity is measured by the amount of labeled moiety transferred from the donor substrate to the receptor substrate. Ala polypeptide can be produced using various expression systems; can be purified from cells, can be in the form of a cleaved or cleaved recombinant fusion protein and/or can have non-Akt polypeptide sequences, for example a His tag or .beta.-galactosidase at its N- or C-terminus. Akt activity can be assayed in cancerous cells lines if the cancerous cell lines are used as a source of the Ala to be assayed. Suitable donor substrates for Ala assays include any molecule that is susceptible to dephosphorylation by Akt., such as, for example include .gamma.-labeled ATP and ATP analogs, wherein the label is ³³P, ³²P, ³⁵S or any other radioactive isotope or a suitable fluorescent marker. Suitable recipient substrates for Ala assays include any polypeptide or other molecule that is susceptible to phosphorylation by Akt. Recipient substrates can be derived from fragments of in vivo targets of Akt. Recipient substrates fragments can be 8 to 50 amino acids in length, usually 10 to 30 amino acids and particularly of about 10, 12, 15, 18, 20 and 25 amino acids in length. Further recipient substrates can be determined empirically using a set of different polypeptides or other molecules. Targets of Recipient substrates for TTK can be capable of being purified from other components of the reaction once the reaction has been performed. This purification is usually done through a molecular interaction, where the recipient substrates is biotinylated and purified through its interaction with streptavidin, or a specific antibody is available that can specifically recognize the recipient substrates. The reaction can be performed in a variety of conditions, such as on a solid support, in a gel, in solution or in living cells. The choice of detection methods depends on type of label used for the donor molecule and may include, for example, measurement of incorporated radiation or fluorescence by autoradiography, scintillation, scanning or fluorography.

IV. Methods of Treatment

The compounds and pharmaceutical compositions provided herein can be used in the treatment of esophageal adenocarcinoma and other disorders associated with abnormal cell proliferation. In one embodiment, the compounds of the present invention can be used to treat esophageal adenocarcinoma.

Drug Resistant Tumors or Cancers

The invention provides compounds that can be used to treat drug resistant esophageal adenocarcinoma. In one embodiment, the compound, such as TCN, TCN-P or a related compound as disclosed herein, can be co-administered with a second drug.

Multidrug resistance (MDR) occurs in human cancers and can be a significant obstacle to the success of chemotherapy.

35

Multidrug resistance is a phenomenon whereby tumor cells in vitro that have been exposed to one cytotoxic agent develop cross-resistance to a range of structurally and functionally unrelated compounds. In addition, MDR can occur intrinsically in some cancers without previous exposure to chemotherapy agents. Thus, in one embodiment, the present invention provides methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer, by administration of TCN, TCN-P or a related compound as disclosed herein. In certain embodiments, TCN, TCN-P and related compounds can be used to treat cancers that are resistant to taxol, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P or a related compound as disclosed herein can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

Combination Therapy

In one aspect of the present invention, the compounds and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, compounds disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the compounds and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopryl, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, taxol, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimetabolic effects, such as those which target cytoskeletal elements, including microtubule modulators such as taxane drugs (such as taxol, paclitaxel, taxotere, docetaxel), podophylotoxins or ulna alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anticycline drugs adriamycin, doxorubicin, pharvorubicin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyproterone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplavin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other

36

bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed below in Table 1.

TABLE 1

Chemotherapeutic Agents	
13-cis-Retinoic Acid	Neosar
2-Amino-6-Mercaptopurine	Neulasta
2-CdA	Neumega
2-Chlorodeoxyadenosine	Neupogen
5-fluorouracil	Nilandron
5-FU	Nilutamide
6-TG	Nitrogen Mustard
6-Thioguanine	Novaldex
6-Mercaptopurine	Novantrone
6-MP	Octreotide
Accutane	Octreotide acetate
Actinomycin-D	Oncospar
Adriamycin	Oncovin
Adrucil	Ontak
Agrylin	Onxal
Ala-Cort	Opivelkin
Aldesleukin	Orapred
Alemtuzumab	Orasone
Alitretinoin	Oxaliplatin
Alkaban-AQ	Paclitaxel
Alkeran	Pamidronate
All-transretinoic acid	Panretin
Alpha interferon	Paraplatin
Altretamine	Pediapred
Amethopterin	PEG Interferon
Amifostine	Pegaspargase
Aminoglutethimide	Pegfilgrastim
Anagrelide	PEG-INTRON
Anandron	PEG-L-asparaginase
Anastrozole	Phenylalanine Mustard
Arabinosylcytosine	Platinol
Ara-C	Platinol-AQ
Aranesp	Prednisolone
Aredia	Prednisone
Arimidex	Prelone
Aromasin	Procarbazine
Arsenic trioxide	PROCRIT
Asparaginase	Proleukin
ATRA	Proliferospaspan 20 with Carmustine implant
Avastin	Purinethol
BCG	Raloxifene
BCNU	Rheumatrex
Bevacizumab	Rituxan
Bexarotene	Rituximab
Bicalutamide	Roveron-A (interferon alfa-2a)
BiCNU	Rubex
Blenoxane	Rubidomycin hydrochloride
Bleomycin	Sandostatin
Bortezomib	Sandostatin LAR
Busulfan	Sargramostim
Busulfex	Solu-Cortef
C225	Solu-Medrol
Calcium Leucovorin	STI-571
Campath	Streptozocin
Campotosar	Tamoxifen
Campothecin-11	Targretin
Capecitabine	Taxol
Carac	Taxotere
Carboplatin	Temodar
Carmustine	Temozolomide
Carmustine wafer	Teniposide
Casodex	TESPA
CCNU	Thalidomide
CDDP	Thalomid

TABLE 1-continued

Chemotherapeutic Agents	
CeeNU	TheraCys
Cerubidine	Thioguanine
cetuximab	Thioguanine Tabloid
Chlorambucil	Thiophosphoamide
Cisplatin	Thioplex
Citrovorum Factor	Thiotepa
Cladribine	TICE
Cortisone	Toposar
Cosmegen	Topotecan
CPT-11	Toremifene
Cyclophosphamide	Trastuzumab
Cytadren	Tretinoin
Cytarabine	Trexall
Cytarabine liposomal	Trisenox
Cytosar-U	TSPA
Cytoxan	VCR
Dacarbazine	Velban
Dactinomycin	Velcade
Darbepoetin alfa	VePesid
Daunomycin	Vesanoid
Daunorubicin	Viadur
Daunorubicin hydrochloride	Vinblastine
Daunorubicin liposomal	Vinblastine Sulfate
DaunoXome	Vincasar Pfs
Decadron	Vincristine
Delta-Cortef	Vinorelbine
Deltasone	Vinorelbine tartrate
Denileukin diftitox	VLB
DepoCyt	VP-16
Dexamethasone	Vumon
Dexamethasone acetate	Xeloda
dexamethasone sodium phosphate	Zanosar
Dexasone	Zevalin
Dexrazoxane	Zinecard
DHAD	Zoladex
DIC	Zoledronic acid
Diodex	Zometa
Docetaxel	Gliadel wafer
Doxil	Glivec
Doxorubicin	GM-CSF
Doxorubicin liposomal	Goserelin
Droxia	granulocyte - colony stimulating factor
	Granulocyte macrophage colony stimulating factor
DTIC	Halotestin
DTIC-Dome	Herceptin
Duralone	Hexadrol
Efudex	Hexalen
Eligard	Hexamethylmelamine
Ellence	HMM
Eloxatin	Hycamtin
Elspar	Hydrea
Emcyt	Hydrocort Acetate
Epinubicin	Hydrocortisone
Epoetin alfa	Hydrocortisone sodium phosphate
Eribitux	Hydrocortisone sodium succinate
Erwinia L-asparaginase	Hydrocortone phosphate
Estramustine	Hydroxyurea
Ethylol	Ibritumomab
Etopophos	Ibritumomab Tiuxetan
Etoposide	Idamycin
Etoposide phosphate	Idarubicin
Eulexin	Ifex
Evista	IFN-alpha
Exemestane	Ifosfamide
Fareston	IL-2
Faslodex	IL-11
Femara	Imatinib mesylate
Filgrastim	Imidazole Carboxamide
Floxuridine	Interferon alfa
Fludara	Interferon Alfa-2b (PEG conjugate)
Fludarabine	Interleukin-2
Fluoroplex	Interleukin-11
Fluorouracil	Intron A (interferon alfa-2b)
Fluorouracil (cream)	Leucovorin
Fluoxynesterone	Leukeran
Flutamide	Leukine
Folinic Acid	Leuprolide

TABLE 1-continued

Chemotherapeutic Agents	
	Leurocristine
5	FUDR
	Fulvestrant
	G-CSF
	Gefitinib
	Gercitabine
	Gerntuzumab ozogamicin
	Gemzar
10	Gleevec
	Lupron
	Lupron Depot
	Matulane
	Maxidex
15	Mechlorethamine
	Mechlorethamine
	Hydrochlorine
	Medralone
	Medrol
	Megace
	Megestrol
20	Megestrol Acetate
	Melphalan
	Mercaptopurine
	Mesna
	Mesnex
	Methotrexate
25	Methotrexate Sodium
	Methylprednisolone
	Mylocel
	Letrozole
	Leustatin
	Liposomal Ara-C
	Liquid Pred
	Lomustine
	L-PAM
	L-Sarcosylsin
	Meticorten
	Mitomycin
	Mitomycin-C
	Mitoxantrone
	M-Prednisol
	MTC
	MTX
	Mustargen
	Mustine
	Mutamycin
	Myleran
	Iressa
	Irinotecan
	Isotretinoin
	Kidrolase
	Lanacort
	L-asparaginase
	LCR

In certain embodiments, interferons (IFNs) can be used in combinations with the compounds of the present invention. Suitable interferons include: interferon alpha-2a, interferon alpha-2b, pegylated interferon alpha, including interferon alpha-2a and interferon alpha 2b, interferon beta, interferon gamma, interferon tau, interferon omega, INFERGEN (interferon alfacon-1) by InterMune, OMNIFERON (natural interferon) by Viragen, ALBUFERON by Human Genome Sciences, REBIF (interferon beta-1a) by Ares-Serono, Omega Interferon by BioMedicine, Oral Interferon Alpha by Amarillo Biosciences, and interferon gamma, interferon tau, and/or interferon gamma-1b by InterMune.

In one embodiment TCN, TCN-P or a related compound as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 3, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextiguldipine, valspodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

V. Pharmaceutical Compositions

Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with other active ingredients.

Compositions comprising the compounds disclosed herein may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcu-

taneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration.

The compositions may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association one or more compositions of the present invention and one or more pharmaceutical carriers or excipients.

The compounds can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel Introduction to Pharmaceutical Dosage Forms, Fourth Edition 1985, 126).

In the compositions, effective concentrations of one or more compounds or pharmaceutically acceptable derivatives thereof may be mixed with one or more suitable pharmaceutical carriers. The compounds may be derivatized as the corresponding salts, esters, enol ethers or esters, acetals, ketals, orthoesters, hemiacetals, hemiketals, acids, bases, solvates, hydrates or prodrugs prior to formulation. The concentrations of the compounds in the compositions are effective for delivery of an amount, upon administration, that treats, prevents, or ameliorates one or more of the symptoms of the target disease or disorder. In one embodiment, the compositions are formulated for single dosage administration. To formulate a composition, the weight fraction of compound is dissolved, suspended, dispersed or otherwise mixed in a selected carrier at an effective concentration such that the treated condition is relieved, prevented, or one or more symptoms are ameliorated.

Compositions suitable for oral administration may be presented as discrete units such as, but not limited to, tablets, caplets, pills or dragees capsules, or cachets, each containing a predetermined amount of one or more of the compositions; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion or as a bolus, etc.

Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing an active compound as defined above and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, solubilizing agents, pH buffering agents, preservatives, flavoring agents, and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 15th Edition, 1975.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or more of the compositions of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouth-

washes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

The tablets, pills, capsules, troches and the like can contain one or more of the following ingredients, or compounds of a similar nature: a binder; a lubricant; a diluent; a glidant; a disintegrating agent; a coloring agent; a sweetening agent; a flavoring agent; a wetting agent; an emetic coating; and a film coating. Examples of binders include microcrystalline cellulose, gum tragacanth, glucose solution, acacia mucilage, gelatin solution, molasses, polyvinylpyrrolidone, povidone, crospovidones, sucrose and starch paste. Lubricants include talc, starch, magnesium or calcium stearate, lycopodium and stearic acid. Diluents include, for example, lactose, sucrose, starch, kaolin, salt, mannitol and dicalcium phosphate. Glidants include, but are not limited to, colloidal silicon dioxide. Disintegrating agents include crosscarmellose sodium, sodium starch glycolate, alginic acid, corn starch, potato starch, bentonite, methylcellulose, agar and carboxymethylcellulose. Coloring agents include, for example, any of the approved certified water soluble FD and C dyes, mixtures thereof; and water insoluble FD and C dyes suspended on alumina hydrate. Sweetening agents include sucrose, lactose, mannitol and artificial sweetening agents such as saccharin, and any number of spray dried flavors. Flavoring agents include natural flavors extracted from plants such as fruits and synthetic blends of compounds which produce a pleasant sensation, such as, but not limited to peppermint and methyl salicylate. Wetting agents include propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate and polyoxyethylene laural ether. Emetic-coatings include fatty acids, fats, waxes, shellac, ammoniated shellac and cellulose acetate phthalates. Film coatings include hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate.

Compositions suitable for topical administration to the skin may be presented as ointments, creams, gels, and pastes, having one or more of the compositions administered in a pharmaceutical acceptable carrier.

Compositions for rectal administration may be presented as a suppository with a suitable base comprising, for example, cocoa butter or a salicylate.

Compositions suitable for nasal administration, when the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to 500 microns which is administered in the manner in which snuff is taken, (i.e., by rapid inhalation through the nasal passage from a container of the powder held close up to the nose). When the carrier is a liquid (for example, a nasal spray or as nasal drops), one or more of the compositions can be admixed in an aqueous or oily solution, and inhaled or sprayed into the nasal passage.

Compositions suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing one or more of the compositions and appropriate carriers.

Compositions suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The compositions may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspen-

sions may be prepared from sterile powders, granules, and tablets of the kind previously described above.

Pharmaceutical organic or inorganic solid or liquid carrier media suitable for enteral or parenteral administration can be used to fabricate the compositions. Gelatin, lactose, starch, magnesium marine, talc, vegetable and animal fats and oils, gum, polyalkylene glycol, water, or other known carriers may all be suitable as carrier media.

Compositions may be used as the active ingredient in combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

It will be understood, however, that the total daily usage of the compositions will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular host will depend upon a variety of factors, including for example, the disorder being treated and the severity of the disorder, activity of the specific composition employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration; route of administration; rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific composition employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the composition at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

Compositions are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of the administered ingredient. For example, approximately 1-5 mg per day of a compound disclosed herein can reduce the volume of a solid tumor in mice.

The dosage will depend on host factors such as weight, age, surface area, metabolism, tissue distribution, absorption rate and excretion rate. In one embodiment, approximately 0.5 to 7 grams per day of a compound disclosed herein may be administered to humans. Optionally, approximately 1 to 4 grams per day of the compound can be administered to humans. In certain embodiments 0.001-5 mg/day is administered to a human. The therapeutically effective dose level will depend on many factors as noted above. In addition, it is well

within the skill of the art to start doses of the composition at relatively low levels, and increase the dosage until the desired effect is achieved.

Compositions comprising a compound disclosed herein may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho) esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (copolymers of lactic acid and glycolic acid).

The compounds may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The Liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The compounds may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microfilm powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions comprising the compounds disclosed herein may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with one or more compositions of the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

Additional Embodiments

The pharmaceutical compositions of the subject invention can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, *Remington Pharmaceutical Sciences* (Martin E W [1995] Easton Pa., Mack Publishing Company, 19th ed.) describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile

suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The methods of the present invention, for example, for inhibiting the growth of a cancerous cell, can be advantageously combined with at least one additional therapeutic method, including but not limited to chemotherapy, radiation therapy, therapy that selectively inhibits Ras oncogenic signaling, or any other therapy known to those of skill in the art of the treatment and management of cancer, such as administration of an anti-cancer agent.

Administration of API-2 (tricitabine) as a salt may be carried out. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartrate, succinate, benzoate, ascorbate, alpha-ketoglutarate, and alpha-glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

The compounds of the present invention can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the compounds of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added.

When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethyleneglycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the compounds of the invention, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the compounds of the invention may be incorporated into sustained-release preparations and devices.

The active agent (i.e., API-2 or pharmaceutically acceptable salts thereof) may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agent or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating compounds of the invention in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

For topical administration, the compounds of the invention may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer than to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the compounds of

45

the invention can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user. Examples of useful dermatological compositions which can be used to deliver the compounds of the invention to the skin are disclosed in Jacquet et al. (U.S. Pat. No. 4,608,392), Gala (U.S. Pat. No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Woltzman (U.S. Pat. No. 4,820,508).

Useful dosages of the pharmaceutical compositions of the present invention can be determined by comparing their in vitro activity, and in vivo activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art for example, see U.S. Pat. No. 4,938,949.

In one non-limiting embodiment, the concentration of the active agent in a liquid composition, such as a lotion, can be from about 0.1-25 wt-%, or from about 0.5-10 wt-%. In one embodiment, the concentration in a semi-solid or solid composition such as a gel or a powder can be about 0.1-5 wt-%, preferably about 0.5-2.5 wt-%. In one embodiment, single dosages for injection, infusion or ingestion will generally vary between 5-1500 mg, and may be administered, i.e., 1-3 times daily, to yield levels of about 0.1-50 mg/kg, for adults. A non-limiting dosage of the present invention can be between 7.5 to 45 mg per kg, administered orally, with appropriate adjustment for the body weight of an individual.

Accordingly, the present invention includes a pharmaceutical composition comprising API-2, as described herein, or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, comprising an amount of API-2, or a pharmaceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to effect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the active agent in tumor tissue which is known to effect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

Mammalian species which benefit from the disclosed methods for the inhibition of cancer cell growth, include, but are not limited to, primates, such as apes, chimpanzees, orangutans, humans, monkeys; domesticated animals (e.g., pets) such as dogs, cats, guinea pigs, hamsters, Vietnamese pot-bellied pigs, rabbits, and ferrets; domesticated farm animals such as cows, buffalo, bison, horses, donkey, swine, sheep, and goats; exotic animals typically found in zoos, such as bear, lions, tigers, panthers, elephants, hippopotamus, rhinoceros, giraffes, antelopes, sloth, gazelles, zebras, wildebeests,

46

prairie dogs, koala bears, kangaroo, opossums, raccoons, pandas, hyena, seals, sea lions, elephant seals, otters, porpoises, dolphins, and whales. The terms "patient" and "subject" are used herein interchangeably and are intended to include such human and non-human mammalian species. Likewise, in vitro methods of the present invention can be earned out on cells of such mammalian species.

Patients in need of treatment using the methods of the present invention can be identified using standard techniques known to those in the medical profession.

The following examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1

In Vitro Screening

Materials

Cell Lines and NCI Diversity Set. All cell lines used in this study were either purchased from ATCC or described previously (Cheng, J. Q., et al. *Oncogene*, 14: 2793-2801, 1997, West, K. A., et al. *Drug Resist. Updat.*, 5: 234-248, 2002, Satyamoorthy, K., et al. *Cancer Res.* 61: 7318-7324, 2001). The NCI Structural Diversity Set is a library of 1,992 compounds selected from the approximately 140,000-compound NCI drug depository. In-depth data on the selection, structures, and activities of these diversity set compounds can be found on the NCI Developmental Therapeutics Program web site.

Screening for Inhibition of Akt-transformed Cell Growth. AKT2 transformed NIH3T3 cells or LXSN vector-transfected NIH3T3 control cells (Cheng, J. Q., et al. *Oncogene*, 14: 2793-2801, 1997) were plated into 96-well tissue culture plate. Following treatment with 5 μ M of NCI Diversity Set compound, cell growth was detected with CellTier 96 One Solution Cell Proliferation kit (Promega). Compounds that inhibit growth in AKT2-transformed but not LXSN-transfected NIH3T3 cells were considered as candidates of Akt inhibitor and subjected to further analysis.

In vitro Protein Kinase, Cell Survival and Apoptosis Assays. In vitro kinase was performed as previously described (see, for example, Jiang, K., Coppola, et al. *Mol. Cell. Biol.*, 20:139-148, 2000). Cell survival was assayed with MTS (Promega). Apoptosis was detected with annexin V, which was performed as previously described (Jiang, K., Coppola, et al. *Mol. Cell. Biol.*, 20:139-148, 2000). Recombinant Akt and PDK1 were purchased from Upstate Biotechnology Inc.

Results

Identification of Small Molecule Inhibitor of Akt Signaling Pathway, API-2. Frequent alterations of Akt have been detected in human cancer and disruption of Akt pathway induces apoptosis and inhibits tumor growth (Jetzt, A., et al. *Cancer Res.*, 63: 697-706, 2003). Thus, Akt has been considered as an attractive molecular target for development of novel cancer therapeutics. To identify small molecule inhibitor(s) of Akt, a chemical library of 1,992-compounds from the NCI (the NCI Diversity Set) was evaluated for agents capable of inhibition of growth in AKT2-transformed but not empty vector LXSN-transfected NIH3T3 cells as described in "Materials and Methods". Repeated experiments showed that 32 compounds inhibited growth only in AKT2-transformed

cells. The most potent of these compounds, API-2 (NCI identifier. NSC 154020), suppressed cell growth at a concentration of 50 nM. FIG. 1A shows the chemical structure of API-2, which is also known as triciribine (Schweinsberg, P. D., et al. *Biochem Pharmacol.*, 30: 2521-2526, 1981). The fact that API-2 inhibited selectively AKT-2 transformed cells over untransformed parental cells prompted us to determine whether API-2 is an inhibitor of AKT2 kinase. To this end, AKT2 was immunoprecipitated with anti-AKT2 antibody from AKT-2 transformed NIH3T3 cells following treatment with API-2. AKT2 immunoprecipitates were immunoblotted with anti-phospho-Akt antibodies. As shown in FIG. 1B, API-2 significantly inhibited AKT2 phosphorylation at both threonine-309 and serine-474, which are required for full activation of AKT2 (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). As three isoforms of Akt share high homology and similar structure, the effect of API-2 on their kinase activities was evaluated. HEK293 cells were transfected with HA-Akt1, -AKT2 and -AKT3, serum-starved overnight and treated with API-2 for 60 min prior to EGF (50 ng/ml) stimulation. Triple experiments showed that API-2 suppressed EGF-induced kinase activity and phosphorylation of Akt1, AKT2 and AKT3 (FIG. 1C). However, kinase activity of recombinant constitutively active AKT2 (Myr-AKT2) was not inhibited by API-2 in an in vitro kinase reaction (FIG. 1D), suggesting that API-2 does not directly inhibit Akt in vitro and that API-2 neither functions as ATP competitor nor as the substrate competitor that binds to active site of Akt.

API-2 Does Not Inhibit Known Upstream Activators of Akt. It has been well documented that Akt is activated by extracellular stimuli and intracellular signal molecules, such as active Ras and Src, through a PI3K-dependent manner. Therefore, API-2 inhibition of Akt could result from targeting upstream molecule(s) of Akt. As PI3K and PDK1 are direct upstream regulators of Akt (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999), whether API-2 inhibits PI3K and/or PDK1 was examined. HEK293 cells were serum-starved and then treated with API-2 or PI3K inhibitor, wortmannin, for 30 min prior to EGF stimulation. PI3K was immunoprecipitated with anti-p110 α antibody. The immunoprecipitates were subjected to in vitro PI3K kinase assay using PI-4-P as a substrate. As shown in FIG. 2A, the EGF-induced PI3K activity was inhibited by wortmannin but not by API-2. To evaluate the effect of API-2 on PDK1, an assay in which recombinant PDK1 promotes the threonine-309 phosphorylation of AKT2 peptides was used in the presence of lipid vesicles containing phosphatidylinositol. As shown in FIG. 2B, the assay was potentially inhibited by the control PDK1 inhibitor staurosporine (IC₅₀=5 nM). In contrast, API-2 displayed only 21% inhibition of the assay at the highest concentration tested (5.1 μ M). These data demonstrate that API-2 is not a potent inhibitor of PDK1. To further evaluate the effect of API-2 on PDK1 activation, the autophosphorylation level of PDK1 at serine-241, a residue that is phosphorylated by itself and is critical for its activity was examined (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999), following API-2 treatment of HEK293 cells. Triplicate experiments show that phosphorylation levels of PDK1 were not inhibited by API-2 (FIG. 2B). However, PI3K inhibitor wortmannin, as expected, inhibited EGF-stimulated PDK1 (FIG. 2B).

API-2 Is Highly Selective for the Akt over PKC, PKA, SGK, STAT, JNK, p38, and ERK Signaling Pathways. Akt belongs to AGC (PKA/PKG/PKC) kinase family, which also include PICA, PKC, serum- and glucocorticoid-inducible kinase (SGK), p90 ribosomal S6 kinase, p70^{S6k}, mitogen- and stress-activated protein kinase and PKC-related kinase. Among AGC kinase family, protein structures of PKA, PKC and SGK are more close to Akt kinase than other members. Therefore, next examined were the effects of API-2 on the enzymatic activities of these 3 kinases. HEK293 cells were transfected with HA-tagged PKA, PKC α or SGK. In vitro kinase assay and immunoblotting analysis showed that the kinase activities of PKA and PKC α were inhibited by PKAI and Ro 31-8220, a PKC inhibitor, respectively, whereas API-2 exhibited no effect on their activities (FIGS. 2C and 2E). Further, serum-induced SGK kinase activity was attenuated by wortmannin but not by API-2 (FIG. 2D). In addition, it was determined whether API-2 has effect on other oncogenic survival pathways. Western blotting analyses with commercially available anti-phospho-antibodies revealed that phosphorylation levels of Stat3, JNK, p38 and Erk1/2 were not affected by API-2 treatment (FIG. 2F). These data indicate that API-2 specifically inhibits Akt signaling pathway.

API-2 Suppresses Cell Growth and Induces Apoptosis in Akt-overexpressing/activating Human Cancer Cell Lines. The ability of API-2 to selectively inhibit the Akt pathway suggests that it should inhibit proliferation and/or induces apoptosis preferentially in those tumor cells with aberrant expression/activation of Akt. As activation of Akt in human malignancies commonly results from overexpression of Akt or PTEN mutations, API-2 was used to treat the cells that express constitutively active Akt, caused by overexpression of AKT2 (OVCAR3, OVCAR8, PANC1 and AKT2-transformed NIH3T3) or mutations of the PTEN gene (PC-3, LNCaP, MDA-MB-468), and cells that do not (OVCAR5, DU-145, T47D, COLO357 and LXSNIH3T3) as well as melanoma cells that are activated by IGF-1 to activate Akt or do not respond to growth stimulation by IGF-1 (Satyamoorthy, K., et al. *Cancer Res.* 61: 7317-7324, 2001). Immunoblotting analysis showed that phosphorylation levels of Akt were inhibited by API-2 only in the cells expressing elevated Akt or responding to IGF-1 stimulation (FIG. 3A). Accordingly, API-2 inhibited cell growth to a much higher degree in Akt-overexpressing/activating cells as compared to those with low levels of Akt. As shown in FIG. 3B, API-2 treatment inhibited cell proliferation by approximate 50-60% in Akt-overexpressing/activating cell lines, LNCaP, PC-3, OVCAR3, OVCA8, PANC1, MDA-MB-468, and WM35, whereas only by about 10-20% in DU145, OVCAR5, COLO357, T47D and WM852 cells, which exhibit low levels of Akt or do not respond to growth stimulation by IGF-1. Moreover, API-2 induces apoptosis by 8-fold (OVCAR3), 6-fold (OVCAR8), 6-fold (PANC1), and 3-fold (AKT2-NIH3T3). No significant difference of apoptosis was observed between API-2 and vehicle (DMSO) treatment in OVCAR5, COLO357 and LXSNIH3T3 cells. Thus, API-2 inhibits cell growth and induces apoptosis preferentially in cells that express aberrant Akt.

API-2 Inhibits Downstream Targets of Akt. It has been shown that Akt exerts its cellular effects through phosphorylation of a number of proteins (Datta, S. R., et al. *Genes Dev.*

13: 2905-2927, 1999). More than 20 proteins have been identified as Akt substrates, including the members of Forkhead protein family (FKHR, AFX and FKHL1), tuberlin/TSC2, p70^{S6K}, GSK-3 β , p21^{WAF1/Cip1}, p27^{kip1}, MDM2, Bad, ASK1 and IKK α etc. It was next examined whether API-2 inhibits downstream targets of Akt. As anti-phospho-tuberlin, -Bad, -AFX, and -GSK-3 β antibodies are commercially available, therefore, the effect of API-2 on their phosphorylation induced by Akt was determined. Following API-2 (1 \square M) treatment, OVCAR3 cells were lysed and immunoblotted with the individual anti-phospho-antibody. FIG. 4A shows that API-2 considerably inhibited the phosphorylation levels of tuberlin leading to stabilization and upregulation of tuberlin (Dan, H. C., et al. J. Biol. Chem., 277: 35364-35370, 2002). The phosphorylation levels of Bad, GSK-3 β , and AFX were partially attenuated by API-2. These data suggest that API-2 induces cell death and cell growth arrest by inhibiting phosphorylation of its downstream targets. API-2 inhibition of Akt downstream targets at different degrees could be due to the fact that phosphorylation sites of these targets are also regulated by other kinase(s), for instance, Bad serine-136 is phosphorylated by PAK1 in addition to Akt (Schurmann, A., et al. Mol. Cell. Biol., 20: 453-461, 2000).

Example 2

Antitumor Activity in the Nude Mouse Tumor Xenograft Model

Tumor cells were harvested, resuspended in PBS, and injected s.c. into the right and left flanks (2 \times 10⁶ cells/flank) of 8-week-old female nude mice as reported previously (Sun, J., Blaskovic, et al. Cancer Res., 59: 4919-4926, 1999). When tumors reached about 100-150 mm³, animals were randomized and dosed i.p. with 0.2 ml vehicle of drug daily. Control animals were received DMSO (20%) vehicle, whereas treated animals were injected with API-2 (1 mg/kg/day) in 20% DMSO.

API-2 Inhibits the Growth of Tumors in Nude Mice that Overexpress Akt. Frequent overexpression/activation and/or amplification of AKT1 and AKT2 in human ovarian and pancreatic cancer was shown (Cheng, J. Q., and Nicosia, S. V. AKT signal transduction pathway in oncogenesis. In Schwab D, Editor, Encyclopedic Reference of Cancer. Berlin Heidelberg and New York: Springer; 2001. pp 35-7). Inhibition of Akt pathway by inhibitors of Pint., HSP70, Src and farnesyl-transferase resulted in cell growth arrest and induction of apoptosis (Sat, D. B., et al. Cancer Res., 63: 2139-2144, 2003, Xu, W., et al. Cancer Res., 63: 7777-7784, 2003). A recent study showed that the tumor growth of xenografts with elevated Akt was also significantly inhibited by intratumoral injection of adenovirus of dominant negative Akt (Jetzt, A., et al. Cancer Res., 63: 697-706, 2003). Because API-2 inhibits Akt signaling and induces apoptosis and cell growth arrest only in cancer cells with elevated levels of Akt (FIG. 3), the growth of tumors with elevated levels of Akt should be more

sensitive to API-2 than that of tumors with low levels of Akt in nude mice. To this end, s.c. Akt-overexpressing cells (OVCAR3, OVCAR8 and PANC-1) were s.c. implanted into the right flank, and those cell lines that express low levels of Akt (OVCAR5 and COLO357) into the left flank of mice. When the tumors reached an average size of about 100-150 mm³, the animals were randomized and treated i.p. with either vehicle or API-2 (1 mg/kg/day). As illustrated in FIG. 4B, OVCAR-5 and COLO357 tumors treated with vehicle grew to about 800-1,000 mm³ 49 days after tumor implantation. OVCAR3, OVCAR8 and PANC1 tumors treated with vehicle control grew to about 700-900 mm³ 49 days after tumor implantation. API-2 inhibited OVCAR3, OVCAR8 and PANC1 tumor growth by 90%, 88% and 80%, respectively. In contrast, API-2 had little effect on the growth of OVCAR5 and COLO357 cells in nude mice (FIGS. 4B-4D and data not shown). At dose 1 mg/kg/day, API-2 had no effects on blood glucose level, body weight, activity and food intake of mice. In treated tumor samples, Akt activity was inhibited by API-2 without change of total Akt content (FIG. 4E). Taken together, these results indicate that API-2 selectively inhibits the growth of tumors with elevated levels of Akt.

Example 3

TCN Directly Inhibits Wild Type Akt Kinase Activity

API-2 (TCN) can directly inhibit wild type Akt kinase activity induced by PDK1 in vitro (FIG. 1). This result supports that API-2 is a direct Akt inhibitor and that the underlying mechanism may be API-2 binding to PH domain and/or threonine-308 of Akt. An in vitro kinase assay was performed with recombinant of PDK1 and Akt in a kinase buffer containing phosphatidylinositol-3,4,5-P3 (PIP3), API-2 and histone H2B as substrate. After incubation of 30 min, the reactions were separated by SDS-PAGE and exposed in a film.

Example 4

TCN is Effective in Cancer Resistant Cells

The effects of TCN (API-2) were tested in cisplatin, paclitaxel, and tamoxifen resistant A270CP, C-13, OVCAR433 and MCF7/TAM cells. API-2 overcame cisplatin, paclitaxel, and tamoxifen resistance in these cells

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention will be evident to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of this invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1
 <211> LENGTH: 1443
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(1443)

<400> SEQUENCE: 1

atg aac gac gta gcc att gtg aag gag ggc tgg ctg cac aaa cga ggg	48
Met Asn Asp Val Ala Ile Val Lys Glu Gly Trp Leu His Lys Arg Gly	
1 5 10 15	
gaa tat att aaa acc tgg cgg cca cgc tac ttc ctc ctc aag aac gat	96
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Asn Asp	
20 25 30	
ggc acc ttt att ggc tac aag gaa cgg cct cag gat gtg gat cag cga	144
Gly Thr Phe Ile Gly Tyr Lys Glu Arg Pro Gln Asp Val Asp Gln Arg	
35 40 45	
gag tcc cca ctc aac aac ttc tca gtg gca caa tgc cag ctg atg aag	192
Glu Ser Pro Leu Asn Asn Phe Ser Val Ala Gln Cys Gln Leu Met Lys	
50 55 60	
aca gag cgg cca agg ccc aac acc ttt atc atc cgc tgc ctg cag tgg	240
Thr Glu Arg Pro Arg Pro Asn Thr Phe Ile Ile Arg Cys Leu Gln Trp	
65 70 75 80	
acc aca gtc att gag cgc acc ttc cat gta gaa acg cct gag gag cgg	288
Thr Thr Val Ile Glu Arg Thr Phe His Val Glu Thr Pro Glu Glu Arg	
85 90 95	
gaa gaa tgg gcc acc gcc att cag act gtg gcc gat gga ctc aag agg	336
Glu Glu Trp Ala Thr Ala Ile Gln Thr Val Ala Asp Gly Leu Lys Arg	
100 105 110	
cag gaa gaa gag acg atg gac ttc cga tca ggc tca ccc agt gac aac	384
Gln Glu Glu Glu Thr Met Asp Phe Arg Ser Gly Ser Pro Ser Asp Asn	
115 120 125	
tca ggg gct gaa gag atg gag gtg tcc ctg gcc aag ccc aag cac cgt	432
Ser Gly Ala Glu Glu Met Glu Val Ser Leu Ala Lys Pro Lys His Arg	
130 135 140	
gtg acc atg aac gag ttt gag tac ctg aaa cta ctg ggc aag ggc acc	480
Val Thr Met Asn Glu Phe Glu Tyr Leu Lys Leu Leu Gly Lys Gly Thr	
145 150 155 160	
ttt ggg aaa gtg att ctg gta aaa gag aag gcc aca ggc cgc tac tat	528
Phe Gly Lys Val Ile Leu Val Lys Glu Lys Ala Thr Gly Arg Tyr Tyr	
165 170 175	
gcc atg aag atc ctc aag aag gag gtc atc gtc gcc aag gat gag gtt	576
Ala Met Lys Ile Leu Lys Lys Glu Val Ile Val Ala Lys Asp Glu Val	
180 185 190	
gcc cac acg ctt act gaa aac cgt gtc ctg cag aac tct agg cat ccc	624
Ala His Thr Leu Thr Glu Asn Arg Val Leu Gln Asn Ser Arg His Pro	
195 200 205	
ttc ctt acg gcc ctc aag tac tca ttc cag acc cac gac cgc ctc tgc	672
Phe Leu Thr Ala Leu Lys Tyr Ser Phe Gln Thr His Asp Arg Leu Cys	
210 215 220	
ttt gtc atg gag tat gcc aac ggg ggc gag ctc ttc ttc cac ctg tct	720
Phe Val Met Glu Tyr Ala Asn Gly Gly Glu Leu Phe Phe His Leu Ser	
225 230 235 240	
cga gag cgc gtg ttc tcc gag gac cgg gcc cgc ttc tat ggt gcg gag	768
Arg Glu Arg Val Phe Ser Glu Asp Arg Ala Arg Phe Tyr Gly Ala Glu	
245 250 255	

-continued

att gtg tct gcc ctg gac tac ttg cac tcg gag aag aac gtg gtg tac Ile Val Ser Ala Leu Asp Tyr Leu His Ser Glu Lys Asn Val Val Tyr 260 265 270	816
cgg gac ctg aag cta gag aac ctc atg ctg gac aag gac ggg cac atc Arg Asp Leu Lys Leu Glu Asn Leu Met Leu Asp Lys Asp Gly His Ile 275 280 285	864
aag ata acg gac ttc ggg ctg tgc aag gag ggg atc aag gat ggt gcc Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile Lys Asp Gly Ala 290 295 300	912
act atg aag aca ttc tgc gga acg ccg gag tac ctg gcc cct gag gtg Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu Ala Pro Glu Val 305 310 315 320	960
ctg gaa gac aac gac tac ggc cgt gca gtg gac tgg tgg ggg ctg ggc Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp Trp Gly Leu Gly 325 330 335	1008
gtg gtc atg tat gag atg atg tgt ggc cgc ctg ccc ttc tac aac cag Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro Phe Tyr Asn Gln 340 345 350	1056
gac cac gag aag ctg ttc gag ctg atc ctc atg gag gag atc cgc ttc Asp His Glu Lys Leu Phe Glu Leu Ile Leu Met Glu Glu Ile Arg Phe 355 360 365	1104
ccg cgc aca ctc ggc cct gag gcc aag tcc ctg ctc tcc ggg ctg ctc Pro Arg Thr Leu Gly Pro Glu Ala Lys Ser Leu Leu Ser Gly Leu Leu 370 375 380	1152
aag aag gac cct aca cag agg ctc ggt ggg ggc tct gag gat gcc aag Lys Lys Asp Pro Thr Gln Arg Leu Gly Gly Ser Glu Asp Ala Lys 385 390 395 400	1200
gag atc atg cag cac cgg ttc ttt gcc aac atc gtg tgg cag gat gtg Glu Ile Met Gln His Arg Phe Phe Ala Asn Ile Val Trp Gln Asp Val 405 410 415	1248
tat gag aag aag ctg agc cca cct ttc aag ccc cag gtc acc tct gag Tyr Glu Lys Lys Leu Ser Pro Pro Phe Lys Pro Gln Val Thr Ser Glu 420 425 430	1296
act gac acc agg tat ttc gat gag gag ttc aca gct cag atg atc acc Thr Asp Thr Arg Tyr Phe Asp Glu Glu Phe Thr Ala Gln Met Ile Thr 435 440 445	1344
atc acg ccg cct gat caa gat gac agc atg gag tgt gtg gac agt gag Ile Thr Pro Pro Asp Gln Asp Asp Ser Met Glu Cys Val Asp Ser Glu 450 455 460	1392
cgg agg ccg cac ttc ccc cag ttc tcc tac tca gcc agt ggc aca gcc Arg Arg Pro His Phe Pro Gln Phe Ser Tyr Ser Ala Ser Gly Thr Ala 465 470 475 480	1440
tga	1443

<210> SEQ ID NO 2

<211> LENGTH: 480

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Asn Asp Val Ala Ile Val Lys Glu Gly Trp Leu His Lys Arg Gly
1 5 10 15

Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Asn Asp
20 25 30

Gly Thr Phe Ile Gly Tyr Lys Glu Arg Pro Gln Asp Val Asp Gln Arg
35 40 45

Glu Ser Pro Leu Asn Asn Phe Ser Val Ala Gln Cys Gln Leu Met Lys
50 55 60

-continued

Thr	Glu	Arg	Pro	Arg	Pro	Asn	Thr	Phe	Ile	Ile	Arg	Cys	Leu	Gln	Trp	65	70	75	80
Thr	Thr	Val	Ile	Glu	Arg	Thr	Phe	His	Val	Glu	Thr	Pro	Glu	Glu	Arg	85	90	95	
Glu	Glu	Trp	Ala	Thr	Ala	Ile	Gln	Thr	Val	Ala	Asp	Gly	Leu	Lys	Arg	100	105	110	
Gln	Glu	Glu	Glu	Thr	Met	Asp	Phe	Arg	Ser	Gly	Ser	Pro	Ser	Asp	Asn	115	120	125	
Ser	Gly	Ala	Glu	Glu	Met	Glu	Val	Ser	Leu	Ala	Lys	Pro	Lys	His	Arg	130	135	140	
Val	Thr	Met	Asn	Glu	Phe	Glu	Tyr	Leu	Lys	Leu	Leu	Gly	Lys	Gly	Thr	145	150	155	160
Phe	Gly	Lys	Val	Ile	Leu	Val	Lys	Glu	Lys	Ala	Thr	Gly	Arg	Tyr	Tyr	165	170	175	
Ala	Met	Lys	Ile	Leu	Lys	Lys	Glu	Val	Ile	Val	Ala	Lys	Asp	Glu	Val	180	185	190	
Ala	His	Thr	Leu	Thr	Glu	Asn	Arg	Val	Leu	Gln	Asn	Ser	Arg	His	Pro	195	200	205	
Phe	Leu	Thr	Ala	Leu	Lys	Tyr	Ser	Phe	Gln	Thr	His	Asp	Arg	Leu	Cys	210	215	220	
Phe	Val	Met	Glu	Tyr	Ala	Asn	Gly	Gly	Glu	Leu	Phe	Phe	His	Leu	Ser	225	230	235	240
Arg	Glu	Arg	Val	Phe	Ser	Glu	Asp	Arg	Ala	Arg	Phe	Tyr	Gly	Ala	Glu	245	250	255	
Ile	Val	Ser	Ala	Leu	Asp	Tyr	Leu	His	Ser	Glu	Lys	Asn	Val	Val	Tyr	260	265	270	
Arg	Asp	Leu	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	Gly	His	Ile	275	280	285	
Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Lys	Asp	Gly	Ala	290	295	300	
Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	Pro	Glu	Val	305	310	315	320
Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	Gly	Leu	Gly	325	330	335	
Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	Tyr	Asn	Gln	340	345	350	
Asp	His	Glu	Lys	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Glu	Ile	Arg	Phe	355	360	365	
Pro	Arg	Thr	Leu	Gly	Pro	Glu	Ala	Lys	Ser	Leu	Leu	Ser	Gly	Leu	Leu	370	375	380	
Lys	Lys	Asp	Pro	Thr	Gln	Arg	Leu	Gly	Gly	Gly	Ser	Glu	Asp	Ala	Lys	385	390	395	400
Glu	Ile	Met	Gln	His	Arg	Phe	Phe	Ala	Asn	Ile	Val	Trp	Gln	Asp	Val	405	410	415	
Tyr	Glu	Lys	Lys	Leu	Ser	Pro	Pro	Phe	Lys	Pro	Gln	Val	Thr	Ser	Glu	420	425	430	
Thr	Asp	Thr	Arg	Tyr	Phe	Asp	Glu	Glu	Phe	Thr	Ala	Gln	Met	Ile	Thr	435	440	445	
Ile	Thr	Pro	Pro	Asp	Gln	Asp	Asp	Ser	Met	Glu	Cys	Val	Asp	Ser	Glu	450	455	460	
Arg	Arg	Pro	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser	Ala	Ser	Gly	Thr	Ala	465	470	475	480

-continued

```

<211> LENGTH: 1515
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(1446)

<400> SEQUENCE: 3

atg aat gag gtg tct gtc atc aaa gaa ggc tgg ctc cac aag cgt ggt      48
Met Asn Glu Val Ser Val Ile Lys Glu Gly Trp Leu His Lys Arg Gly
1          5          10          15

gaa tac atc aag acc tgg agg cca cgg tac ttc ctg ctg aag agc gac      96
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Ser Asp
          20          25          30

ggc tcc ttc att ggg tac aag gag agg ccc gag gcc cct gat cag act     144
Gly Ser Phe Ile Gly Tyr Lys Glu Arg Pro Glu Ala Pro Asp Gln Thr
          35          40          45

cta ccc ccc tta aac aac ttc tcc gta gca gaa tgc cag ctg atg aag     192
Leu Pro Pro Leu Asn Asn Phe Ser Val Ala Glu Cys Gln Leu Met Lys
          50          55          60

acc gag agg ccg cga ccc aac acc ttt gtc ata cgc tgc ctg cag tgg     240
Thr Glu Arg Pro Arg Pro Asn Thr Phe Val Ile Arg Cys Leu Gln Trp
65          70          75          80

acc aca gtc atc gag agg acc ttc cac gtg gat tct cca gac gag agg     288
Thr Thr Val Ile Glu Arg Thr Phe His Val Asp Ser Pro Asp Glu Arg
          85          90          95

gag gag tgg atg cgg gcc atc cag atg gtc gcc aac agc ctc aag cag     336
Glu Glu Trp Met Arg Ala Ile Gln Met Val Ala Asn Ser Leu Lys Gln
          100          105          110

cgg gcc cca ggc gag gac ccc atg gac tac aag tgt ggc tcc ccc agt     384
Arg Ala Pro Gly Glu Asp Pro Met Asp Tyr Lys Cys Gly Ser Pro Ser
          115          120          125

gac tcc tcc acg act gag gag atg gaa gtg gcg gtc agc aag gca cgg     432
Asp Ser Ser Thr Thr Glu Glu Met Glu Val Ala Val Ser Lys Ala Arg
          130          135          140

gct aaa gtg acc atg aat gac ttc gac tat ctc aaa ctc ctt ggc aag     480
Ala Lys Val Thr Met Asn Asp Phe Asp Tyr Leu Lys Leu Leu Gly Lys
          145          150          155          160

gga acc ttt ggc aaa gtc atc ctg gtg cgg gag aag gcc act ggc cgc     528
Gly Thr Phe Gly Lys Val Ile Leu Val Arg Glu Lys Ala Thr Gly Arg
          165          170          175

tac tac gcc atg aag atc ctg cga aag gaa gtc atc att gcc aag gat     576
Tyr Tyr Ala Met Lys Ile Leu Arg Lys Glu Val Ile Ile Ala Lys Asp
          180          185          190

gaa gtc gct cac aca gtc acc gag agc cgg gtc ctc cag aac acc agg     624
Glu Val Ala His Thr Val Thr Glu Ser Arg Val Leu Gln Asn Thr Arg
          195          200          205

cac ccg ttc ctc act gcg ctg aag tat gcc ttc cag acc cac gac cgc     672
His Pro Phe Leu Thr Ala Leu Lys Tyr Ala Phe Gln Thr His Asp Arg
          210          215          220

ctg tgc ttt gtg atg gag tat gcc aac ggg ggt gag ctg ttc ttc cac     720
Leu Cys Phe Val Met Glu Tyr Ala Asn Gly Gly Glu Leu Phe Phe His
          225          230          235          240

ctg tcc cgg gag cgt gtc ttc aca gag gag cgg gcc cgg ttt tat ggt     768
Leu Ser Arg Glu Arg Val Phe Thr Glu Glu Arg Ala Arg Phe Tyr Gly
          245          250          255

gca gag att gtc tcg gct ctt gag tac ttg cac tcg cgg gac gtg gta     816
Ala Glu Ile Val Ser Ala Leu Glu Tyr Leu His Ser Arg Asp Val Val
          260          265          270

tac cgc gac atc aag ctg gaa aac ctc atg ctg gac aaa gat ggc cac     864
Tyr Arg Asp Ile Lys Leu Glu Asn Leu Met Leu Asp Lys Asp Gly His
          275          280          285

```

-continued

atc aag atc act gac ttt ggc ctc tgc aaa gag ggc atc agt gac ggg Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile Ser Asp Gly 290 295 300	912
gcc acc atg aaa acc ttc tgt ggg acc ccg gag tac ctg gcg cct gag Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu Ala Pro Glu 305 310 315 320	960
gtg ctg gag gac aat gac tat ggc cgg gcc gtg gac tgg tgg ggg ctg Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp Trp Gly Leu 325 330 335	1008
ggg gtg gtc atg tac gag atg atg tgc ggc cgc ctg ccc ttc tac aac Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro Phe Tyr Asn 340 345 350	1056
cag gac cac gag cgc ctc ttc gag ctc atc ctc atg gaa gag atc cgc Gln Asp His Glu Arg Leu Phe Glu Leu Ile Leu Met Glu Glu Ile Arg 355 360 365	1104
ttc ccg cgc acg ctc agc ccc gag gcc aag tcc ctg ctt gct ggg ctg Phe Pro Arg Thr Leu Ser Pro Glu Ala Lys Ser Leu Leu Ala Gly Leu 370 375 380	1152
ctt aag aag gac ccc aag cag agg ctt ggt ggg ggg ccc agc gat gcc Leu Lys Lys Asp Pro Lys Gln Arg Leu Gly Gly Gly Pro Ser Asp Ala 385 390 395 400	1200
aag gag gtc atg gag cac agg ttc ttc ctc agc atc aac tgg cag gac Lys Glu Val Met Glu His Arg Phe Phe Leu Ser Ile Asn Trp Gln Asp 405 410 415	1248
gtg gtc cag aag aag ctc ctg cca ccc ttc aaa cct cag gtc acg tcc Val Val Gln Lys Lys Leu Leu Pro Pro Phe Lys Pro Gln Val Thr Ser 420 425 430	1296
gag gtc gac aca agg tac ttc gat gat gaa ttt acc gcc cag tcc atc Glu Val Asp Thr Arg Tyr Phe Asp Asp Glu Phe Thr Ala Gln Ser Ile 435 440 445	1344
aca atc aca ccc cct gac cgc tat gac agc ctg ggc tta ctg gag ctg Thr Ile Thr Pro Pro Asp Arg Tyr Asp Ser Leu Gly Leu Leu Glu Leu 450 455 460	1392
gac cag cgg acc cac ttc ccc cag ttc tcc tac tcg gcc agc atc cgc Asp Gln Arg Thr His Phe Pro Gln Phe Ser Tyr Ser Ala Ser Ile Arg 465 470 475 480	1440
gag tga gcagtctgcc cagcagagg acgcacgctc gctgccatca ccgctgggtg Glu	1496
gttttttacc cctgcccg	1515

<210> SEQ ID NO 4

<211> LENGTH: 481

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Asn Glu Val Ser Val Ile Lys Glu Gly Trp Leu His Lys Arg Gly
1 5 10 15

Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Ser Asp
20 25 30

Gly Ser Phe Ile Gly Tyr Lys Glu Arg Pro Glu Ala Pro Asp Gln Thr
35 40 45

Leu Pro Pro Leu Asn Asn Phe Ser Val Ala Glu Cys Gln Leu Met Lys
50 55 60

Thr Glu Arg Pro Arg Pro Asn Thr Phe Val Ile Arg Cys Leu Gln Trp
65 70 75 80

-continued

Thr	Thr	Val	Ile	Glu	Arg	Thr	Phe	His	Val	Asp	Ser	Pro	Asp	Glu	Arg
			85						90					95	
Glu	Glu	Trp	Met	Arg	Ala	Ile	Gln	Met	Val	Ala	Asn	Ser	Leu	Lys	Gln
			100					105					110		
Arg	Ala	Pro	Gly	Glu	Asp	Pro	Met	Asp	Tyr	Lys	Cys	Gly	Ser	Pro	Ser
		115						120					125		
Asp	Ser	Ser	Thr	Thr	Glu	Glu	Met	Glu	Val	Ala	Val	Ser	Lys	Ala	Arg
		130					135					140			
Ala	Lys	Val	Thr	Met	Asn	Asp	Phe	Asp	Tyr	Leu	Lys	Leu	Leu	Gly	Lys
145					150					155					160
Gly	Thr	Phe	Gly	Lys	Val	Ile	Leu	Val	Arg	Glu	Lys	Ala	Thr	Gly	Arg
			165						170					175	
Tyr	Tyr	Ala	Met	Lys	Ile	Leu	Arg	Lys	Glu	Val	Ile	Ile	Ala	Lys	Asp
			180					185					190		
Glu	Val	Ala	His	Thr	Val	Thr	Glu	Ser	Arg	Val	Leu	Gln	Asn	Thr	Arg
		195						200					205		
His	Pro	Phe	Leu	Thr	Ala	Leu	Lys	Tyr	Ala	Phe	Gln	Thr	His	Asp	Arg
		210					215				220				
Leu	Cys	Phe	Val	Met	Glu	Tyr	Ala	Asn	Gly	Gly	Glu	Leu	Phe	Phe	His
225					230					235					240
Leu	Ser	Arg	Glu	Arg	Val	Phe	Thr	Glu	Glu	Arg	Ala	Arg	Phe	Tyr	Gly
			245						250					255	
Ala	Glu	Ile	Val	Ser	Ala	Leu	Glu	Tyr	Leu	His	Ser	Arg	Asp	Val	Val
			260					265					270		
Tyr	Arg	Asp	Ile	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	Gly	His
		275						280					285		
Ile	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Ser	Asp	Gly
		290					295				300				
Ala	Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	Pro	Glu
305					310					315					320
Val	Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	Gly	Leu
			325						330					335	
Gly	Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	Tyr	Asn
			340					345					350		
Gln	Asp	His	Glu	Arg	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Glu	Ile	Arg
		355						360					365		
Phe	Pro	Arg	Thr	Leu	Ser	Pro	Glu	Ala	Lys	Ser	Leu	Leu	Ala	Gly	Leu
		370					375				380				
Leu	Lys	Lys	Asp	Pro	Lys	Gln	Arg	Leu	Gly	Gly	Gly	Pro	Ser	Asp	Ala
385					390					395					400
Lys	Glu	Val	Met	Glu	His	Arg	Phe	Phe	Leu	Ser	Ile	Asn	Trp	Gln	Asp
			405						410					415	
Val	Val	Gln	Lys	Lys	Leu	Leu	Pro	Pro	Phe	Lys	Pro	Gln	Val	Thr	Ser
			420					425					430		
Glu	Val	Asp	Thr	Arg	Tyr	Phe	Asp	Asp	Glu	Phe	Thr	Ala	Gln	Ser	Ile
		435						440				445			
Thr	Ile	Thr	Pro	Pro	Asp	Arg	Tyr	Asp	Ser	Leu	Gly	Leu	Leu	Glu	Leu
		450					455				460				
Asp	Gln	Arg	Thr	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser	Ala	Ser	Ile	Arg
465					470					475					480
Glu															

<210> SEQ ID NO 5

<211> LENGTH: 1510

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (3)..(1460)

<400> SEQUENCE: 5

gg gct cag agg gga gtc atc atg agc gat gtt acc att gtg aaa gaa      47
  Ala Gln Arg Gly Val Ile Met Ser Asp Val Thr Ile Val Lys Glu
    1             5             10             15

ggt tgg gtt cag aag agg gga gaa tat ata aaa aac tgg agg cca aga      95
Gly Trp Val Gln Lys Arg Gly Glu Tyr Ile Lys Asn Trp Arg Pro Arg
    20             25             30

tac ttc ctt ttg aag aca gat ggc tca ttc ata gga tat aaa gag aaa     143
Tyr Phe Leu Leu Lys Thr Asp Gly Ser Phe Ile Gly Tyr Lys Glu Lys
    35             40             45

cct caa gat gtg gat tta cct tat ccc ctc aac aac ttt tca gtg gca     191
Pro Gln Asp Val Asp Leu Pro Tyr Pro Leu Asn Asn Phe Ser Val Ala
    50             55             60

aaa tgc cag tta atg aaa aca gaa cga cca aag cca aac aca ttt ata     239
Lys Cys Gln Leu Met Lys Thr Glu Arg Pro Lys Pro Asn Thr Phe Ile
    65             70             75

atc aga tgt ctc cag tgg act act gtt ata gag aga aca ttt cat gta     287
Ile Arg Cys Leu Gln Trp Thr Thr Val Ile Glu Arg Thr Phe His Val
    80             85             90             95

gat act cca gag gaa agg gaa gaa tgg aca gaa gct atc cag gct gta     335
Asp Thr Pro Glu Glu Arg Glu Glu Trp Thr Glu Ala Ile Gln Ala Val
    100            105            110

gca gac aga ctg cag agg caa gaa gag gag aga atg aat tgt agt cca     383
Ala Asp Arg Leu Gln Arg Gln Glu Glu Glu Arg Met Asn Cys Ser Pro
    115            120            125

act tca caa att gat aat ata gga gag gaa gag atg gat gcc tct aca     431
Thr Ser Gln Ile Asp Asn Ile Gly Glu Glu Glu Met Asp Ala Ser Thr
    130            135            140

acc cat cat aaa aga aag aca atg aat gat ttt gac tat ttg aaa cta     479
Thr His His Lys Arg Lys Thr Met Asn Asp Phe Asp Tyr Leu Lys Leu
    145            150            155

cta ggt aaa ggc act ttt ggg aaa gtt att ttg gtt cga gag aag gca     527
Leu Gly Lys Gly Thr Phe Gly Lys Val Ile Leu Val Arg Glu Lys Ala
    160            165            170            175

agt gga aaa tac tat gct atg aag att ctg aag aaa gaa gtc att att     575
Ser Gly Lys Tyr Ala Met Lys Ile Leu Lys Lys Glu Val Ile Ile
    180            185            190

gca aag gat gaa gtg gca cac act cta act gaa agc aga gta tta aag     623
Ala Lys Asp Glu Val Ala His Thr Leu Thr Glu Ser Arg Val Leu Lys
    195            200            205

aac act aga cat ccc ttt tta aca tcc ttg aaa tat tcc ttc cag aca     671
Asn Thr Arg His Pro Phe Leu Thr Ser Leu Lys Tyr Ser Phe Gln Thr
    210            215            220

aaa gac cgt ttg tgt ttt gtg atg gaa tat gtt aat ggg ggc gag ctg     719
Lys Asp Arg Leu Cys Phe Val Met Glu Tyr Val Asn Gly Gly Glu Leu
    225            230            235

ttt ttc cat ttg tcg aga gag cgg gtg ttc tct gag gac cgc aca cgt     767
Phe Phe His Leu Ser Arg Glu Arg Val Phe Ser Glu Asp Arg Thr Arg
    240            245            250            255

ttc tat ggt gca gaa att gtc tct gcc ttg gac tat cta cat tcc gga     815
Phe Tyr Gly Ala Glu Ile Val Ser Ala Leu Asp Tyr Leu His Ser Gly
    260            265            270

```

-continued

aag att gtg tac cgt gat ctc aag ttg gag aat cta atg ctg gac aaa	863
Lys Ile Val Tyr Arg Asp Leu Lys Leu Glu Asn Leu Met Leu Asp Lys	
275 280 285	
gat ggc cac ata aaa att aca gat ttt gga ctt tgc aaa gaa ggg atc	911
Asp Gly His Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile	
290 295 300	
aca gat gca gcc acc atg aag aca ttc tgt ggc act cca gaa tat ctg	959
Thr Asp Ala Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu	
305 310 315	
gca cca gag gtg tta gaa gat aat gac tat ggc cga gca gta gac tgg	1007
Ala Pro Glu Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp	
320 325 330 335	
tgg ggc cta ggg gtt gtc atg tat gaa atg atg tgt ggg agg tta cct	1055
Trp Gly Leu Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro	
340 345 350	
ttc tac aac cag gac cat gag aaa ctt ttt gaa tta ata tta atg gaa	1103
Phe Tyr Asn Gln Asp His Glu Lys Leu Phe Glu Leu Ile Leu Met Glu	
355 360 365	
gac att aaa ttt cct cga aca ctc tct tca gat gca aaa tca ttg ctt	1151
Asp Ile Lys Phe Pro Arg Thr Leu Ser Ser Asp Ala Lys Ser Leu Leu	
370 375 380	
tca ggg ctc ttg ata aag gat cca aat aaa cgc ctt ggt gga gga cca	1199
Ser Gly Leu Leu Ile Lys Asp Pro Asn Lys Arg Leu Gly Gly Gly Pro	
385 390 395	
gat gat gca aaa gaa att atg aga cac agt ttc ttc tct gga gta aac	1247
Asp Asp Ala Lys Glu Ile Met Arg His Ser Phe Phe Ser Gly Val Asn	
400 405 410 415	
tgg caa gat gta tat gat aaa aag ctt gta cct cct ttt aaa cct caa	1295
Trp Gln Asp Val Tyr Asp Lys Lys Leu Val Pro Pro Phe Lys Pro Gln	
420 425 430	
gta aca tct gag aca gat act aga tat ttt gat gaa gaa ttt aca gct	1343
Val Thr Ser Glu Thr Asp Thr Arg Tyr Phe Asp Glu Glu Phe Thr Ala	
435 440 445	
cag act att aca ata aca cca cct gaa aaa tat gat gag gat ggt atg	1391
Gln Thr Ile Thr Ile Thr Pro Pro Glu Lys Tyr Asp Glu Asp Gly Met	
450 455 460	
gac tgc atg gac aat gag agg cgg ccg cat ttc cct caa ttt tcc tac	1439
Asp Cys Met Asp Asn Glu Arg Arg Pro His Phe Pro Gln Phe Ser Tyr	
465 470 475	
tct gca agt gga cga gaa taa gtctctttca ttctgtact tcaactgtcat	1490
Ser Ala Ser Gly Arg Glu	
480 485	
cttcaattta ttactgaaaa	1510

<210> SEQ ID NO 6

<211> LENGTH: 485

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens


<400> SEQUENCE: 6

Ala Gln Arg Gly Val Ile Met Ser Asp Val Thr Ile Val Lys Glu Gly	
1 5 10 15	
Trp Val Gln Lys Arg Gly Glu Tyr Ile Lys Asn Trp Arg Pro Arg Tyr	
20 25 30	
Phe Leu Leu Lys Thr Asp Gly Ser Phe Ile Gly Tyr Lys Glu Lys Pro	
35 40 45	
Gln Asp Val Asp Leu Pro Tyr Pro Leu Asn Asn Phe Ser Val Ala Lys	
50 55 60	
Cys Gln Leu Met Lys Thr Glu Arg Pro Lys Pro Asn Thr Phe Ile Ile	
65 70 75 80	

69

I claim:

1. A method for treating esophageal adenocarcinoma in a mammal comprising treating the mammal with an effective amount of a compound of the formula:



wherein each R_2' , R_3' and R_5' are independently hydrogen, optionally substituted phosphate or phosphonate; acyl; alkyl; amide; sulfonate ester; sulfonyl, methanesulfonyl, benzyisulfonyl, arylsulfonyl, wherein the phenyl group is optionally substituted with one or more substituents; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein R_2' , R_3' or R_5' is independently H or mono-, di- or tri-phosphate;

wherein R^x and R^y are independently hydrogen, optionally substituted phosphate; acyl; amide, alkyl; aromatic, polyoxyalkylene; polyethyleneglycol, optionally substituted arylsulfonyl; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group;

wherein R_1 and R_2 each are independently H, optionally substituted straight chained, branched or cyclic alkyl, alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl,

70


CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxy-alkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl.

2. The method of claim 1, wherein the mammal has been diagnosed with esophageal adenocarcinoma.

3. The method of claim 1, comprising first obtaining a biological sample from the tumor or cancer; then determining whether the tumor or cancer overexpresses an Akt kinase, wherein the level of Akt kinase expression is determined by assaying the cancer for the presence of a phosphorylated Akt kinase.

4. The method of claim 3, wherein the level of Akt kinase expression is determined by assaying the cancer for the presence of a phosphorylated Akt kinase with an antibody.

5. A method of treating esophageal adenocarcinoma in a mammal comprising administering to the mammal an effective amount of a compound of formula:



wherein each R_2' , R_3' and R_5' are independently hydrogen, optionally substituted phosphate or phosphonate; acyl; alkyl; amide; sulfonate ester; sulfonyl, methanesulfonyl, benzyisulfonyl, arylsulfonyl, wherein the phenyl group

71

is optionally substituted with one or more substituents; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein R^{2'}, R^{3'} or R^{5'} is independently H or mono-, di- or tri-phosphate;

wherein R^x and R^y are independently hydrogen, optionally substituted phosphate; acyl; amide, alkyl; aromatic, polyoxyalkylene; polyethyleneglycol, optionally substituted arylsulfonyl; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group;

wherein R₁ and R₂ each are independently H, optionally substituted straight chained, branched or cyclic alkyl, alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl;

wherein the compound is administered one time per week via intravenous infusion for three weeks followed by a one week period wherein the compound is not administered.



6. The method of claim 5, wherein the dosing schedule is repeated at least twice.

7. The method of claim 5, wherein the dosing schedule is repeated at least 4 times.

8. The method of claim 5, wherein at least 10 mg/m² of the compound is administered.


9. The method of claim 5, wherein 10 mg/m² or less of the compound is administered.

10. A method of treating esophageal adenocarcinoma in a mammal comprising administering to the mammal a dose of 10 mg/m² or less of a compound of the formula:



72

-continued



wherein each R^{2'}, R^{3'} and R^{5'} are independently hydrogen, optionally substituted phosphate or phosphonate; acyl; alkyl; amide; sulfonate ester; sulfonyl, methanesulfonyl, benzyulfonyl, arylsulfonyl, wherein the phenyl group is optionally substituted with one or more substituents; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein R^{2'}, R^{3'} or R^{5'} is independently H or mono-, di- or tri-phosphate;

wherein R^x and R^y are independently hydrogen, optionally substituted phosphate; acyl; amide, alkyl; aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group;

wherein R₁ and R₂ each are independently H, optionally substituted straight chained, branched or cyclic alkyl, alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl;

wherein the compound is administered one time per week.

11. The method of claim 10, wherein the dosing regimen is repeated for at least two weeks.

12. The method of claim 10, wherein the dosing regimen is repeated for three weeks, and wherein the three week period is followed by a one week period wherein no drug is administered.

13. The method of claim 12, wherein the dosing regimen represents a dosing cycle.

14. The method of claim 13, wherein the dosing cycle is repeated at least twice.


15. The method of claim 13 wherein the dosing cycle is repeated until regression of the cancer is achieved.

16. The method of claim 10, wherein the drug is administered intravenously.

17. The method of claim 10, wherein the subject has been diagnosed with a carcinoma, sarcoma, lymphoma, leukemia, or myeloma.

18. The method of claim 10 wherein the mammal is a human.

19. The method of claim 10, wherein the compound is the compound of Formula IB:

73


IB

5

10

15

20. The method of claim **10**, wherein the compound is the compound of Formula IIA:

74

IIA

* * * * *