

April 2012

Materials and methods for treatment of inflammatory and cell proliferation disorders

Shyam S. Mohapatra

Follow this and additional works at: https://digitalcommons.usf.edu/usf_patents

Recommended Citation

Mohapatra, Shyam S., "Materials and methods for treatment of inflammatory and cell proliferation disorders" (2012). *USF Patents*. 309.
https://digitalcommons.usf.edu/usf_patents/309

This Patent is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Patents by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.



US008148114B2

(12) **United States Patent**
Mohapatra

(10) **Patent No.:** **US 8,148,114 B2**
(45) **Date of Patent:** **Apr. 3, 2012**

(54) **MATERIALS AND METHODS FOR
TREATMENT OF INFLAMMATORY AND
CELL PROLIFERATION DISORDERS**

(75) Inventor: **Shyam S Mohapatra**, Tampa, FL (US)

(73) Assignee: **University of South Florida**, Tampa, FL
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 266 days.

(21) Appl. No.: **12/259,110**

(22) Filed: **Oct. 27, 2008**

(65) **Prior Publication Data**

US 2009/0176706 A1 Jul. 9, 2009

Related U.S. Application Data

(63) Continuation of application No. 11/059,814, filed on
Feb. 17, 2005, now abandoned.

(60) Provisional application No. 60/521,072, filed on Feb.
17, 2004.

(51) **Int. Cl.**

A61P 35/00 (2006.01)

A61K 31/7088 (2006.01)

C07H 21/04 (2006.01)

C12N 15/09 (2006.01)

C12N 15/74 (2006.01)

(52) **U.S. Cl.** **435/91.1; 435/69.1; 435/320.1;**
435/375; 435/455; 536/23.1

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,753,945 A 6/1988 Gilbard et al.
4,957,735 A 9/1990 Huang
5,013,556 A 5/1991 Woodle et al.
5,043,164 A 8/1991 Huang et al.
5,087,617 A 2/1992 Smith
5,093,246 A 3/1992 Cech et al.
5,116,742 A 5/1992 Cech et al.
5,135,917 A 8/1992 Burch
5,144,019 A 9/1992 Rossi et al.
5,168,053 A 12/1992 Altman et al.
5,176,996 A 1/1993 Hogan et al.
5,180,818 A 1/1993 Cech et al.
5,190,931 A 3/1993 Inouye
5,272,262 A 12/1993 Rossi et al.
5,545,614 A 8/1996 Stamler et al.
5,595,756 A * 1/1997 Bally et al. 424/450
5,602,143 A 2/1997 Krauss
5,625,056 A 4/1997 Genieser et al.
5,646,032 A 7/1997 Volker ter Meulen et al.
5,665,861 A 9/1997 Forssmann et al.
5,686,101 A 11/1997 Tagawa et al.
5,691,310 A 11/1997 Vesely
5,705,187 A 1/1998 Unger
5,817,856 A 10/1998 Tirosh et al.
5,820,873 A 10/1998 Choi et al.
5,840,341 A 11/1998 Watts et al.
5,858,694 A 1/1999 Piazza et al.

6,013,630 A 1/2000 Shimkets
6,028,055 A 2/2000 Lowe et al.
6,184,037 B1 2/2001 Rolland et al.
6,506,559 B1 1/2003 Fire et al.
6,528,245 B2 3/2003 Sanchez-Ramos et al.
6,943,147 B2 9/2005 Vesely
7,022,828 B2 4/2006 McSwiggen
7,354,908 B2 4/2008 Mohapatra et al.
7,488,713 B2 2/2009 Vesely
7,595,303 B1 9/2009 Mohapatra et al.
7,655,772 B2 * 2/2010 Mohapatra 536/23.1
7,825,092 B2 11/2010 Vesely
7,846,900 B2 12/2010 Vesely
2001/0027181 A1 10/2001 Kitakaze et al.
2002/0094326 A1 7/2002 Donahue et al.
2002/0146821 A1 10/2002 Sanchez-Ramos et al.
2002/0193579 A1 12/2002 Usman et al.
2003/0069186 A1 4/2003 Burnett et al.
2003/0073628 A1 4/2003 Shailubhai et al.
2003/0105000 A1 6/2003 Pero et al.
2003/0138793 A1 7/2003 Su et al.
2003/0147943 A1 8/2003 Luo et al.
2003/0204063 A1 10/2003 Gravel et al.
2003/0215528 A1 11/2003 Graham et al.
2004/0002458 A1 1/2004 Seilhamer et al.
2004/0067889 A1 4/2004 Belenky et al.
2004/0138134 A1 7/2004 Golembo et al.
2004/0146549 A1 7/2004 Ben-Sasson et al.
2004/0171550 A1 9/2004 Backstrom et al.
2004/0203081 A1 10/2004 James et al.
2004/0213782 A1 10/2004 Wax et al.

(Continued)

FOREIGN PATENT DOCUMENTS

WO WO 99/32619 A1 7/1999
WO WO 00/07156 A2 11/2000
WO WO 01/68836 A3 9/2001
WO WO 01/75164 A3 10/2001
WO WO 01/92513 A1 12/2001
WO WO 2004/011498 A3 2/2004
WO WO 2004/022003 A2 3/2004
WO WO 2004/022579 A3 3/2004
WO WO 2004/083236 A3 9/2004
WO WO 2005/094420 A3 10/2005
WO WO 2006/026536 A3 3/2006
WO WO 2007/127487 A2 11/2007
WO WO 2007/130672 A3 11/2007
WO WO 2009/073527 A2 6/2009

OTHER PUBLICATIONS

Sporn MB, Suh, N, "Chemoprevention of cancer," Carcinogenesis,
2000, 21(3): 525-530.*

(Continued)

Primary Examiner — Julie Ha

(74) *Attorney, Agent, or Firm* — Saliwanchik, Lloyd &
Eisenschenk

(57) **ABSTRACT**

Compositions and methods for treatment of inflammatory
and cell proliferation disorders using natriuretic hormone
peptide (NP), polynucleotides encoding NP, or agents that
reduce the activity of atrial natriuretic peptide receptor A
(NPR-A).

20 Claims, 11 Drawing Sheets

U.S. PATENT DOCUMENTS

2004/0224903	A1	11/2004	Berry et al.
2004/0229784	A1	11/2004	Vesely
2004/0258687	A1	12/2004	Waldman et al.
2004/0266673	A1	12/2004	Bakis et al.
2005/0008617	A1	1/2005	Chen et al.
2005/0014287	A1	1/2005	Friese et al.
2005/0014289	A1	1/2005	Parsons et al.
2005/0176641	A1	8/2005	Bakis et al.
2005/0209139	A1	9/2005	Vesely
2005/0266093	A1	12/2005	Mohapatra
2005/0272650	A1	12/2005	Mohapatra
2006/0014689	A1	1/2006	Vesely
2006/0019256	A1	1/2006	Clarke et al.
2006/0110359	A1	5/2006	Sanchez-Ramos et al.
2006/0205642	A1	9/2006	Vesely
2006/0276382	A1	12/2006	Mohapatra
2007/0036867	A1	2/2007	Mohapatra et al.
2007/0116767	A1	5/2007	Mohapatra
2007/0265204	A1	11/2007	Mohapatra et al.
2008/0039394	A1	2/2008	Vesely
2008/0070858	A1	3/2008	Mohapatra
2008/0214437	A1	9/2008	Mohapatra et al.
2009/0062206	A1	3/2009	Vesely
2009/0170196	A1	7/2009	Vesely
2009/0176706	A1	7/2009	Mohapatra
2011/0034386	A1	2/2011	Vesely
2011/0039777	A1	2/2011	Vesely

OTHER PUBLICATIONS

- Cancer Drug Design and Discovery Neidle, Stephen, ed. (Elsevier/Academic Press, 2008) pp. 427-431.*
- Rudinger J, "Characteristics of the amino acids as components of a peptide hormone sequence," Peptide Hormones, JA Parsons Edition, University Park Press, Jun. 1976, pp. 1-7.*
- "Designing Custom Peptides," from SIGMA Genosys, pp. 1-2. Accessed Dec. 16, 2004.*
- Schinzl R, Drucekes P, "The phosphate recognition site of *Escherichia coli* maltodextrin phosphorylase," FEBS, Jul. 1991, 286(1,2): 125-128.*
- Berendsen HJC, "A Glimpse of the Holy Grail?" Science, 1998, 282: 642-643.*
- Voet D, Voet JG, Biochemistry, Second Edition, John Wiley & Sons, Inc., 1995, pp. 235-241.*
- Ngo JT, Marks J, Karplus M, "Computational Complexity, Protein Structure Prediction, and the Levinthal Paradox," The Protein Folding Problem and Tertiary Structure Prediction, K. Merc Jr. and S. Le Grand Edition, 1994, pp. 491-495.*
- Bradley CM, Barrick D, "Limits of Cooperativity in a Structurally Modular Protein: Response of the Notch Ankyrin Domain to Analogous Alanine Substitutions in Each Repeat," J. Mol. Biol., 2002, 324: 373-386.*
- U.S. Appl. No. 11/799,225, filed Apr. 30, 2007, Mohapatra.
- U.S. Appl. No. 10/526,584, filed Mar. 3, 2005, Mohapatra.
- Abbey, S. et al. "Lysophosphatidic acid inhibits C-type natriuretic peptide activation of guanylyl cyclase-B" *Endocrinology*, 2003, 144:240-246.
- Angus, R.M. et al. "Effect of inhaled atrial natriuretic peptide on methacholine induced bronchoconstriction in asthma" *Clin Exp Allergy* 1994, 24:784-788.
- Auerbach, R. et al. "Angiogenesis assays, Problems and Pitfalls" *Cancer and Metastasis Reviews* 2000, 19: 167-172.
- Bliss, D. et al. "Expression of the atrial natriuretic factor gene in small cell lung cancer tumors and tumor cell lines" *J Natl Can Inst*, 1990, 82:305-310.
- Bryan, P.M. et al. "The Atrial Natriuretic Peptide Receptor (NPR-A/GC-1) is Dephosphorylated by Distinct Microcystin-sensitive and Magnesium-dependent Protein Phosphatase" *J. Biol. Chem.* 2002, 277(18): 16041-16047.
- Chen, S. et al. "1, 25 dihydroxyvitamin D amplifies type A natriuretic peptide receptor expression and activity in target cells" *J. Am. Soc. Nephrol.*, 2005, 16:329-339.
- Chen, X. et al. "Human bone marrow stromal cell cultures" *J Neurosci Res*, 2002, 69:687-691.
- Chen, X. et al. "Ischemic rat brain extracts induce human marrow stromal cell growth factor production" *Neuropathology*, 2002, 22:275-279.
- Chen, J. et al. "Therapeutic benefit of intravenous administration of bone marrow stromal cells after cerebral ischemia in rats" *Stroke*, 2001, 32:1005-1011.
- Chen, J. et al. "Therapeutic benefit of intracerebral transplantation of bone marrow stromal cells after cerebral ischemia in rats" *J. Neurological Sci.*, 2001, 189:49-57.
- Chopp, M. et al. "Spinal cord injury in rat: treatment with bone marrow stromal cell transplantation" *Neuroreport*, 2000, 11:3001-3005.
- Delporte, C. et al. "Discovery of a potent atrial natriuretic peptide antagonist for ANP_A receptors in the human neuroblastoma NB-OK-1 cell line" *Eur. J. Pharmacol.*, 1992, 224(2-3):183-188.
- Doczi, T.P. et al. "Atrial natriuretic peptide (ANP) attenuates brain oedema accompanying experimental subarachnoid haemorrhage" *Acta Neurochir (Wien)*, 1995, 132:87-91.
- Drewett, J.G. and Garbers, D.L. "The family of guanylyl cyclase receptors and their ligands" *Endocrine Reviews*, 1994, 15(2):135-162.
- El-Ayoubi, R. et al. "Urinary responses to acute moxonidine are inhibited by natriuretic peptide receptor antagonist" *Br. J. Pharmacol.*, 2005, 145:50-56.
- Ernst, P. "Review article: the role of inflammation in the pathogenesis of gastric cancer" *Aliment Pharmacol Ther.*, 1999, 13(1):13-18.
- Fonarow, G.C. et al. "Combining natriuretic peptides and necrosis markers in determining prognosis in heart failure" *Rev. Cardiovasc. Med.*, 2003, 4(suppl 4):S20-S28.
- Fujiseki, Y. et al. "Natriuretic peptide receptors, NPR-A and NPR-B, in cultured rabbit retinal pigment epithelium cells" *Jpn. J. Pharmacol.*, 1999, 79(3):359-368.
- Fürst, R. et al. "Atrial natriuretic peptide induces mitogen-activated protein kinase phosphatase-1 in human endothelial cells via Rac1 and NAD(P)H oxidase/Nox2-activation" *Circ. Res.*, 2005, 96:43-53.
- Gower, W.R. et al. "Regulation of atrial natriuretic peptide secretion by cholinergic and PACAP neurons of the gastric antrum" *Am. J. Physiol. Gastrointest. Liver Physiol.*, 2003, 284:G68-G74.
- Greten, F.R. et al. "IKK β links inflammation and tumorigenesis in a mouse model of colitis-associated cancer" *Cell*, 2004, 118:285-296.
- Gura T. "Cancer Models: Systems for Identifying New Drugs are Often Faulty" *Science* 1997, 278(5340): 1041-1042, 1-5.
- Izumi, T. et al. "Blockade of the natriuretic peptide receptor guanylyl cyclase-A inhibits NF- κ B activation and alleviates myocardial ischemia/reperfusion injury" *J Clin Invest* 2001, 108:203-213.
- Jain, R.K. "Barriers to Drug Delivery in Solid Tumors" *Scientific American* Jul. 1994, 58-65.
- Jensen, K.T. et al. "A new, fast and reliable radioimmunoassay of brain natriuretic peptide in human plasma. Reference values in healthy subjects and in patients with different diseases" *Scand J Clin Lab Invest*, 1997, 57:529-540.
- Jin, H. et al. "Novel analog of atrial natriuretic peptide selective for receptor-A produces increased diuresis and natriuresis in rats" *J. Clin. Invest.*, 1996, 98:969-976.
- Kanwal, S. et al. "Intracellular fragments of the natriuretic peptide receptor-C (NPR-C) attenuate dopamine efflux" *Endocrinology*, 1999, 140(3):1118-1124.
- Kaneko, T. et al. "C-type natriuretic peptide (CNP) is the major natriuretic peptide in human cerebrospinal fluid" *Brain Res*, 1993, 612:104-109.
- Kanwal, S. et al. "Intracellular fragments of the natriuretic peptide receptor-C (NPR-C) attenuate dopamine efflux" *Endocrinology*, 1999, 140(3):1118-1124.
- Kojima, M. et al. "Cloning and sequence analysis of cDNA encoding a precursor for rat brain natriuretic peptide" *Biochem. Biophys. Res. Commun.*, 1989, 159(3):1420-1426.
- Kelly, R. and Struthers, A. "Are natriuretic peptides clinically useful as markers of heart failure?" *Ann. Clin. Biochem.*, 2001, 38:94-102.
- Khurana, M.L. and Pandey, K.N. "Receptor-mediated stimulatory effect of atrial natriuretic factor, brain natriuretic peptide, and C-type natriuretic peptide on testosterone production in purified mouse Leydig cells: Activation of cholesterol side-chain cleavage enzyme" *Endocrinology*, 1993, 133:2141-2149.

- Kiemer, A. and Vollmar, A. "Autocrine regulation of inducible nitric-oxide synthase in macrophages by atrial natriuretic peptide" *J Biol Chem*, 1998, 273:13444-13451.
- Kiemer, A. et al. "cGMP-mediated inhibition of TNF- α production by the atrial natriuretic peptide in murine macrophages" *J Immunol*, 2000, 165:175-181.
- Kumar, R. et al. "Expression of guanylyl cyclase-A/atrial natriuretic peptide receptor blocks the activation of protein kinase C in vascular smooth muscle cells" *Hypertension*, 1997, 29(part 2):414-421.
- Kurihara, M. et al. "Lower number of atrial natriuretic peptide receptors in thymocytes and spleen cells of spontaneously hypertensive rats" *Biochem Biophys Res Commun* 1987, 149:1132-1140.
- Liang, F. et al. "Sp1 dependence of natriuretic peptide receptor A gene transcription in rat aortic smooth muscle cells" *Endocrinology*, 1999, 140(4):1695-1701.
- Li, Y. et al. "Intrastriatal transplantation of bone marrow nonhematopoietic cells improves functional recovery after stroke in adult mice" *Journal of Cerebral Blood Flow & Metabolism*, 2000, 20:1311-1319.
- Li, Y. et al. "Human marrow stromal cell therapy for stroke in rat: Neurotrophins and functional recovery" *Neurology*, 2002, 59:514-523.
- Lin, K-F. et al. "Human atrial natriuretic peptide gene delivery reduces blood pressure in hypertensive rats" *Hypertension*, 1995, 26:847-853.
- Lin, K-F. et al. "Atrial natriuretic peptide gene delivery attenuates hypertension, cardiac hypertrophy, and renal injury in salt-sensitive rats" *Human Gene Therapy*, 1998, 9:1429-1438.
- Lu, D. et al. "Adult bone marrow stromal cells administered intravenously to rats after traumatic brain injury migrate into brain and improve neurological outcome" *Neuroreport*, 2001, 12:559-563.
- Mahmood, A. et al. "Treatment of traumatic brain injury in female rats with intravenous administration of bone marrow stromal cells" *Neurosurgery*, 2001, 49:1196-1204.
- Mahmood, A. et al. "Intracranial bone marrow transplantation after traumatic brain injury improving functional outcome in adult rats" *Journal of Neurosurgery*, 2001, 94:589-595.
- Mahmood, A. et al. "Intracerebral transplantation of marrow stromal cells cultured with neurotrophic factors promotes functional recovery in adult rats subjected to traumatic brain injury" *J Neurotrauma*, 2002, 19:1609-1617.
- Maisel, A.S. et al. "Cardiac natriuretic peptides: A proteomic window to cardiac function and clinical management" *Rev. Cardiovasc. Med.*, 2003, 4(suppl 4):S3-S12.
- Martin, J. et al. "Modulation by biologic response modifiers of hepatitis C virus antigen-independent cytokine secretion in blood mononuclear cells" *Cytokine*, 1999, 11:267-273.
- Matanić, D. et al. "Cytokines in patients with lung cancer" *Scand J Immunol*, 2003, 57:173-178.
- Matsukawa, N. et al. "The natriuretic peptide clearance receptor locally modulates the physiological effects of the natriuretic peptide system" *Proc. Natl. Acad. Sci. USA*, 1999, 96:7403-7408.
- Mohapatra, S. et al. "Natriuretic peptides and genesis of asthma: An emerging paradigm?" *J Allergy Clin Immunol*, 2004, 114:520-526.
- Motohashi, S. et al. "Preserved IFN- α production of circulating V α 24 NKT cells in primary lung cancer patients" *Int J Cancer*, 2002, 102:159-165.
- Mueller, C. and Buser, P. "B-type natriuretic peptide (BNP): can it improve our management of patients with congestive heart failure?" *Swiss Med Wkly*, 2002, 132:618-622.
- Nakagawa, K. et al. "Plasma concentrations of atrial and brain natriuretic peptides in a case with hypertensive encephalopathy" *Neurol. Res.*, 2002, 24:627-630.
- Nakao, N. et al. "Effect of atrial natriuretic peptide on ischemic brain edema: Changes in brain water and electrolytes" *Neurosurgery*, 1990, 27:39-44.
- Naruse, S. et al. "Effects of atrial natriuretic peptide on brain oedema: The change of water, sodium, and potassium contents in the brain" *Acta Neurochir Suppl (Wien)*, 1990, 51:118-121.
- Nazario, B. et al. "Atrial and brain natriuretic peptides stimulate the production and secretion of C-type natriuretic peptide from bovine aortic endothelial cells" *J. Clin. Invest.*, 1995, 95:1151-1157.
- Nocera, R. et al. "Novel strategies of neuroprotection against pathologic consequences of stroke in the aged brain" *Society for Neurosci. Abstracts*, 2001, 27(2):2302, Meeting date Nov. 10-15, 2001.
- Ogawa, Y. et al. "Molecular cloning of the complementary DNA and gene that encode mouse brain natriuretic peptide and generation of transgenic mice that overexpress the brain natriuretic peptide gene" *J. Clin. Invest.*, 1994, 93(5):1911-1921.
- Ohsaki, Y. et al. "Human small cell lung cancer cell lines express functional atrial natriuretic peptide receptors" *Cancer Res*, 1993, 53:3165-3171.
- Ohsaki, Y. et al. "Human small cell lung cancer cells produce brain natriuretic peptide" *Oncology*, 1999, 56:155-159.
- Ohyama, Y. et al. "Stable expression of natriuretic peptide receptors: Effects of HS-142-1, a non-peptide ANP antagonist" *Biochem. Biophys. Res. Commun.*, 1992, 189(1):336-342.
- Pandey, K.N. et al. "Natriuretic peptide receptor-A negatively regulates mitogen-activated protein kinase and proliferation of mesangial cells: Role of cGMP-dependent protein kinase" *Biochem. Biophys. Res. Commun.*, 2000, 271(2):374-379.
- Pandey, K.N. et al. "Internalization and trafficking of guanylyl (guanylate) cyclase/natriuretic peptide receptor A is regulated by an acidic tyrosine-based cytoplasmic motif GDAY" *Biochem. J.*, 2005, 388:103-113.
- Pandey, K.N. et al. "Ligand-regulated internalization, trafficking, and down-regulation of guanylyl cyclase/atrial natriuretic peptide receptor-A in human embryonic kidney 293 cells" *J. Biol. Chem.*, 2002, 277(7):4618-4627.
- Pandey, K.N. et al. "Functional domains and expression of truncated atrial natriuretic peptide receptor-A: The carboxyl-terminal regions direct the receptor internalization and sequestration in COS-7 cells" *Molecular Pharmacology*, 2000, 57:259-267.
- Pandey, K.N. "Dynamics of internalization and sequestration of guanylyl cyclase/atrial natriuretic peptide receptor-A" *Can. J. Physiol. Pharmacol.*, 2001, 79(8):631-639.
- Pandey, K.N. "Intracellular trafficking and metabolic turnover of ligand-bound guanylyl cyclase/atrial natriuretic peptide receptor-A into subcellular compartments" *Mol. Cell. Biochem.*, 2002, 230(1-2):61-72.
- Pikarsky, E. et al. "NF- κ B functions as a tumour promoter in inflammation-associated cancer" *Nature*, 2004, 431: 461-466.
- Porter, J.G. et al. "Cloning of a cDNA encoding porcine brain natriuretic peptide" *J. Biol. Chem.*, 1989, 264(12):6689-6692.
- Prins, B.A. et al. "Atrial natriuretic peptide inhibits mitogen-activated protein kinase through the clearance receptor" *J. Biol. Chem.*, 1996, 271(24):14156-14162.
- Quan, H. et al. "Inducible regulation of human brain natriuretic peptide promoter in transgenic mice" *Am. J. Physiol. Heart Circ. Physiol.*, 2001, 280:H368-H376.
- Roy, K. et al. "Oral gene delivery with chitosan-DNA nanoparticles generates immunologic protection in a murine model of peanut allergy" *Nat Med*, 1999, 5:387-391.
- Roy, R.N. and Flynn, T.G. "Organization of the gene for iso-rANP, a rat B-type natriuretic peptide" *Biochem. Biophys. Res. Commun.*, 1990, 171(1):416-423.
- Rouleau, N. et al. "Development of a Non-radioactive Homogenous HTS Platform to Measure the Activity of Guanylate Cyclase", Poster #P10144, Presented at 10th Annual SBS Conference and Exhibition, Orlando, FL, Sep. 11-15, 1004, Perkinelmer Biosignal Inc., Canada.
- Rutherford, R. et al. "Identification of renal natriuretic peptide receptor subpopulations by use of the non-peptide antagonist, HS-142-1" *Br. J. Pharmacol.*, 1994, 113:931-939.
- Sanchez-Ramos, J.R. "Neural cells derived from adult bone marrow and umbilical cord blood" *J. Neurosci. Res.*, 2002, 69:880-893.
- Seidman, C. et al. "Nucleotide sequences of the human and mouse atrial natriuretic factor genes" *Science*, 1984, 226:1206-1209.
- Seilhamer, J.J. et al. "Human and canine gene homologs of porcine brain natriuretic peptide" *Biochem. Biophys. Res. Commun.*, 1989, 165(2):650-658.
- Sekiguchi, T. et al. "Molecular cloning of natriuretic peptide receptor A from bullfrog (*Rana catesbeiana*) brain and its functional expression" *Gene*, 2001, 273:251-257.

- Sharma, G.D. et al. "Expression of atrial natriuretic peptide receptor-A antagonizes the mitogen-activated protein kinases (Erk2 and P38^{MAPK}) in cultured human vascular smooth muscle cells" *Mol. Cell. Biochem.*, 2002, 233(1-2):pp. 165-173.
- Shi, S.-J. et al. "Natriuretic peptide receptor A mediates renal sodium excretory responses to blood volume expansion" *Am. J. Physiol. Renal Physiol.*, 2003, 285:F694-F702.
- Shimizu, K. et al. "Ectopic atrial natriuretic peptide production in small cell lung cancer with the syndrome of inappropriate antidiuretic hormone secretion" *Cancer* 1991, 68:2284-2288.
- Silberbach, M. and Roberts Jr., C. "Natriuretic peptide signaling molecular and cellular pathways to growth regulation" *Cell Signal*, 2001, 13:221-231.
- Simkins, J. "Nesiritide (Natrekor®) for Decompensated CHF" in The University of Montana's School of Pharmacy and Allied Health Sciences Drug Information Service, Apr. 2002, vol. 6 No. 4.
- Song, S. et al. "Nerve growth factor and retinoic acid induce development of neuronal cells from bone marrow stromal cells of both young and old mice" *Society for Neurosci. Abstracts*, 2001, 27(1):940, Meeting date Nov. 10-15, 2001.
- Song, S. et al. "Expression of brain natriuretic peptide by human bone marrow stromal cells" *Society for Neurosci. Abstracts*, 2002, Abstract No. 824.3, Meeting date Nov. 2-7, 2002.
- Song, S. and Sanchez-Ramos, J. "Preparation of Neural Progenitors from Bone Marrow and Umbilical Cord Blood" in *Protocols for Neural Stem Cell Methods*, Zigova, T. et al., Eds., 2002, pp. 79-88.
- Song, S. et al. "Expression of brain natriuretic peptide by human bone marrow stromal cells" *Experimental Neurology*, 2004, 185:191-197.
- Steinhilber, M.E. "Structure, expression, and genomic mapping of the mouse natriuretic peptide type-B gene" *Circ. Res.*, 1993, 72(5):984-992.
- Sudoh, T. et al. "Brain natriuretic peptide-32: N-terminal six amino acid extended form of brain natriuretic peptide identified in porcine brain" *Biochem Biophys Res Commun*, 1988, 155:726-732.
- Suenobu, N. et al. "Natriuretic peptides and nitric oxide induce endothelial apoptosis via a cGMP-dependent mechanism" *Arterioscler Thromb Vasc Biol.*, 1999, 19:140-146.
- Surić-Lambić, L. et al. "Vasoactive natriuretic peptides and kidney" *Facta Universitatis: Medicine and Biology*, 1998, 5(1):6-11.
- True, D. et al. "Comparison of Kinase Assay Technologies for High Throughput Screening" poster presented at Society for Biomolecular Screening (SBS), 8th Annual Conference, Sep. 22-26, 2002.
- Vesely, B.A. et al. "Four peptides decrease the number of human pancreatic adenocarcinoma cells" *Eur. J. Clin. Invest.*, 2003, 33:998-1005.
- Vesely, D.L. "Atrial natriuretic peptides in pathophysiological diseases" *Cardiovascular Res*, 2001, 51:647-658.
- Vlasuk, G.P. et al. "Structure and analysis of the bovine atrial natriuretic peptide precursor gene" *Biochem. Biophys. Res. Commun.*, 1986, 136(1):396-403.
- Wang, W. et al. "AlbBNP, a recombinant B-type natriuretic peptide and human serum albumin fusion hormone, as a long-term therapy of congestive heart failure" *Pharm. Res.*, 2004, 21(11):2105-2111.
- Zivin, R.A. et al. "Molecular cloning and characterization of DNA sequences encoding rat and human atrial natriuretic factors" *Proc. Natl. Acad. Sci. USA*, 1984, 81(20):6325-6329.
- Boiteau, R. et al. "Increase in atrial natriuretic factor (ANF) in acute severe asthma (ASA)" *Am Rev Res Dis.*, 1988, 137:A484.
- Chanez, P. et al. "Atrial natriuretic factor (ANF) is a potent bronchodilator in asthma" *J. Allergy Clin. Immunol.*, 1990, 86:321-324.
- Greenberg, B.D. et al. "Nucleotide sequence of the gene encoding human atrial natriuretic factor precursor" *Nature*, 1984, 312(5995):656-658.
- Hamet, P. et al. "Aspects physiologiques et physiopathologiques du facteur natriuretique auriculaire" *Nephrologie*, 1987, 8:7-12, abstract.
- Hulks, G. et al. "Effect of atrial natriuretic factor on bronchomotor tone in the normal human airway" *Clin Sci*, 1990, 79:51-55.
- Hulks, G. et al. "Inhaled atrial natriuretic peptide and asthmatic airways" *Br. Med. J.*, 1992, 304:1156.
- Hulks, G. et al. High dose inhaled atrial natriuretic peptide is a bronchodilator in asthmatic subjects *Eur Respir J.*, 1994, 7:1593-1597.
- Ishii, Y. et al. "Effects of atrial natriuretic peptide on Type II alveolar epithelial cells of the rat lung. Autoradiographic and morphometric studies" *J. Anat.*, 1989, 166:85-95.
- Kumar, M. et al. "Intranasal IFN- γ gene transfer protects BALB/c mice against respiratory syncytial virus infection" *Vaccine*, 2000, 18:558-567.
- Kumar, M. et al. "Atrial natriuretic peptide gene transfer by means of intranasal administration attenuates airway reactivity in a mouse model of allergic sensitization" *J Allergy Clin Immunol.*, 2002, 110:879-882.
- Louzier, V. et al. "Adenovirus-mediated atrial natriuretic protein expression in the lung protects rats from hypoxia-induced pulmonary hypertension" *Hum Gene Ther*, 2001, 12:503-513.
- Matsuse, H. et al. "Recurrent Respiratory Syncytial Virus Infections in Allergen-Sensitized Mice Lead to Persistent Airway Inflammation and Hyperresponsiveness" *The Journal of Immunology*, 2000, 164:6583-6592.
- Needleman, P. et al. "Atriopeptin: a cardiac hormone intimately involved in fluid, electrolyte, and blood-pressure homeostasis" *N Engl J Med*, 1986, 314:828-834.
- Ohbayashi, H. et al. "Compared effects of natriuretic peptides on ovalbumin-induced asthmatic model" *Eur. J. Pharmacol.*, 1998, 346:55-64.
- Seidman, C.E. et al. "The structure of rat preproatrial natriuretic factor as defined by a complementary DNA clone" *Science*, 1984, 225:324-326.
- Winquist, R. et al. "Atrial natriuretic factor elicits an endothelium-independent relaxation and activates particulate guanylate cyclase in vascular smooth muscle" *Proc. Natl. Acad. Sci. USA*, 1984, 81:7661-7664.
- "Atrial natriuretic factor receptor A", MeSH results, accessed <http://www.ncbi.nlm.nih.gov/mesh> on Dec. 3, 2009, pp. 1-2.
- Ahn, K.S. et al. "Simvastatin Potentiates TNF- α -Induced Apoptosis through the Down-Regulation of NF- κ B-Dependent Antiapoptotic Gene Products: Role of I κ B α Kinase and TGF- β -Activated Kinase-1" *Journal of Immunology*, 2007, 178:2507-2516.
- Allen, T.M. et al. "Large unilamellar liposomes with low uptake into the reticuloendothelial system" *FEBS Letters*, 1987, 223:42-46.
- Arenberg, D. "Chemokines in the Biology of Lung Cancer" *Journal of Thoracic Oncology*, 2006, 1(4):287-288.
- Ashworth, T. et al. "Cutting Edge: TFII-I Controls B Cell Proliferation via Regulating NF- κ B" *Journal of Immunology*, 2007, 178:2631-2635.
- Baldini, P.M. et al. "Decrease of polyamine levels and enhancement of transglutaminase activity in selective reduction of B16-F10 melanoma cell proliferation induced by atrial natriuretic peptide" *Melanoma Research*, 2006, 16:501-507.
- Bass, B.L. "RNA interference: The short answer" *Nature*, 2001, 411:428-429.
- Bernstein, E. et al. "The rest is silence" *RNA*, 2001, 7:1509-1521.
- Bernstein, E. et al. "Role for a bidentate ribonuclease in the initiation step of RNA interference" *Nature*, Jan. 2001, 409:363-366.
- Cane, A. et al. "The Endogenous Oxindoles 5-Hydroxyoxindole and Isatin Are Antiproliferative and Proapoptotic" *Biochemical and Biophysical Research Communications*, 2000, 276:379-384.
- Carthew, R.W. "Gene silencing by double-stranded RNA" *Current Opinion in Cell Biology*, 2001, 13:244-248.
- Chin, L. et al. "Malignant melanoma: modern black plague and genetic black box" *Genes & Development*, 1998, 12(22):3467-3481.
- Chiu, Y.L. et al. "RNAi in Human Cells: Basic Structural and Functional Features of Small Interfering RNA" *Molecular Cell*, Sep. 2002, 10(3):549-561.
- Clark, A.R. "Mechanisms of steroid action and resistance in inflammation: MAP kinase phosphatase 1: a novel mediator of biological effects of glucocorticoids" *Journal of Endocrinology*, 2003, 178:5-12.
- Clemens, J.C. et al. "Use of double-stranded RNA interference in Drosophila cell lines to dissect signal transduction pathways" *PNAS*, 2000, 97:6499-6503.

- Clusel, C. et al. "Ex vivo regulation of specific gene expression by nanomolar concentration of double-stranded dumbbell oligonucleotides" *Nucleic Acids Research* 1993, 21(15):3405-3411.
- Collins, F.S. et al. "Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences" *Proc Natl Acad Sci*, 2002;99:16899-16903.
- Dorn, G. et al. "siRNA relieves chronic neuropathic pain" *Nucleic Acids Research*, 2004, 32:e49.
- Eichelbaum, E.J. et al. "Cardiac and kidney hormones cure up to 86% of human small-cell lung cancers in mice" *Eur J Clin Invest*, 2008, 38(8):562-570.
- Elbashir, S.M. et al. "Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells" *Nature*, May 24, 2001, 411:494-498.
- Elbashir, S.M. et al. "RNA interference is mediated by 21- and 22-nucleotide RNAs" *Genes and Development*, 2001, 15:188-200.
- Fattal, E. et al. "Biodegradable polyalkylcyanoacrylate nanoparticles for the delivery of oligonucleotides" *Journal of Controlled Release*, 1998, 53:137-143.
- Filomeni, G. et al. "Pro-apoptotic Activity of Novel Isatin-Schiff Base Copper(II) Complexes Depends on Oxidative Stress Induction and Organelle-selective Damage" *Journal of Biological Chemistry*, 2007, 282(16):12010-12021.
- Fiscus, R.R. "Involvement of Cyclic GMP and Protein Kinase G in the Regulation of Apoptosis and Survival in Neural Cells" *NeuroSignals*, 2002, 11:175-190.
- Forssmann, W.G. et al. "The renal urodilatin system: clinical implications" *Cardiovascular Research*, 2001, 51:450-462.
- Gabellini, C. et al. "Involvement of RB gene family in tumor angiogenesis" *Oncogene*, 2006, 25:5326-5332.
- Glover, V. et al. "Isatin: Identity with the Purified Endogenous Monamine Oxidase Inhibitor Tribulin" *Journal of Neurochemistry*, 1988, 51(2):656-659.
- Gopalakrishnan, M. et al. "Stable expression and pharmacological properties of the human α_7 nicotinic acetylcholine receptor" *European Journal of Pharmacology: Molecular Pharmacology Section*, 1995, 290(3):237-246.
- Halder, J. et al. "Focal Adhesion Kinase Targeting Using In vivo Short Interfering RNA Delivery in Neutral Liposomes for Ovarian Carcinoma Therapy" *Clinical Cancer Research*, 2006, 12:4916-4924.
- Hammond, S.M. et al. "Argonaute2, a Link Between Genetic and Biochemical Analyses of RNAi" *Science*, Aug. 10, 2001, 293:1146-1150.
- Harborth, J. et al. "Identification of essential genes in cultured mammalian cells using small interfering RNAs" *Journal of Cell Science*, 2001, 114:4457-4565.
- Haseloff, J. et al. "Simple RNA enzymes with new and highly specific endoribonuclease activities" *Nature*, 1988, 334(6183):585-591.
- Helene, C. et al. "Control of Gene Expression by Triple Helix-Forming Oligonucleotides: The Antisense Strategy" *Annals of the New York Academy of Sciences*, 1992, 660(1):27-36.
- Hellerman, G. et al. "Mechanism of bronchoprotective effects of a novel natriuretic hormone peptide" *Journal of Allergy and Clinical Immunology*, 2004, 113:79-85.
- Hirata, Y. "Heterologous Down-Regulation of Vascular Atrial Natriuretic Peptide Receptors by Phorbol Esters" *Biochemical and Biophysical Research Communications*, May 16, 1988, 152(3):1097-1103.
- Ho, R.J.Y. et al. "Target-sensitive immunoliposomes: preparation and characterization" *Biochemistry*, 1986, 25:5500-5506.
- Ho, R.J.Y. et al. "Interactions of Target-sensitive Immunoliposomes with Herpes Simplex Virus" *The Journal of Biological Chemistry*, 1987, 262(29):13979-13984.
- Ho, R.J.Y. et al. "Target-sensitive immunoliposomes as an efficient drug carrier for antiviral activity" *The Journal of Biological Chemistry*, 1987, 262(29):13973-13978.
- Howard, K.A. et al. "RNA Interference in Vivo and in Vitro Using a Chitosan/siRNA Nanoparticle System" *Molecular Therapy*, 2006, 14(4):476-484.
- Hutvagner, G. et al. "RNAi: nature abhors a double-strand" *Current Opinion in Genetics & Development*, 2002, 12:225-232.
- Igosheva, N. et al. "Isatin, an endogenous monamine oxidase inhibitor, triggers a dose- and time-dependent switch from apoptosis to necrosis in human neuroblastoma cells" *Neurochemistry International*, 2005, 47(3):216-224.
- Inoue, J.I. et al. "NFkB activation in development and progression of cancer" *Cancer Science*, 2007, 98:268-274.
- Ivanova, K. et al. "Differential Expression of Functional Guanylyl Cyclases in Melanocytes: Absence of Nitric-Oxide-Sensitive Isoform in Metastatic Cells" *Journal of Investigative Dermatology*, Mar. 2001, 116(3):409-416.
- Jacque, J.M. et al. "Modulation of HIV-1 replication by RNA interference" *Nature*, Jul. 25, 2002, 418(6896):435-438.
- Karin, M. "Mitogen activated protein kinases as targets for development of novel anti-inflammatory drugs" *Annals of the Rheumatic Diseases*, 2004, 63(Suppl. 2):ii62-ii64.
- Katas, H. et al. "Development and characterisation of chitosan nanoparticles for siRNA delivery" *Journal of Controlled Release*, 2006, 115(2):216-225.
- Kim, J.W. et al. "Effect of phosphorylation and S—S bond-induced dimerization on DNA binding and transcriptional activation by C/EBP β " *Proc Natl Acad Sci USA*, 2007, 104:1913-1918.
- Kisilew, M. et al. "Isoform-specific knockdown and expression of adaptor protein ShcA using small interfering RNA" *Biochem J*, 2002, 363(1):1-5.
- Klibanov, A.L. et al. "Amphipathic polyethyleneglycols effectively prolong the circulation time of liposomes" *FEBS Letters*, 1990, 268:235-237.
- Knaapen, A.M. et al. "Inhaled particles and lung cancer. Part A: Mechanisms" *International Journal of Cancer*, 2004, 109:799-809.
- Kong, X. et al. "Mice Deficient in Atrial Natriuretic Peptide Receptor A (NPRA) Exhibit Decreased Lung Inflammation: Implication of NPRA Signaling in Asthma Pathogenesis" *Journal of Allergy and Clinical Immunology*, Jan. 2007, 119(1):5127, abstract 501.
- Kong, X. et al. "Natriuretic Peptide Receptor A as a Novel Anticancer Agent" *Cancer Research*, Jan. 2008, 68(1):249-256.
- Lambert, G. et al. Nanoparticulate systems for the delivery of antisense oligonucleotides *Advanced Drug Delivery Reviews*, Mar. 2001, 47:99-112.
- Landen, C.N. et al. "Intraperitoneal Delivery of Liposomal siRNA for Therapy of Advanced Ovarian Cancer" *Cancer Biology & Therapy*, 2006, 5(12):1708-1713.
- Lee, N.S. et al. "Expression of small interfering RNAs targeted against HIV-1 *rev* transcripts in human cells" *Nature Biotechnology*, May 2002, 20:500-505.
- Lenz, A. et al. "Cardiac Hormones Eliminate some Human Squamous Lung Carcinomas in Athymic Mice" *European Journal of Clinical Investigation*, 2010 in press, p. 1-13.
- Levin, E.R. et al. "Mechanisms of Disease: Natriuretic Peptides" *New England Journal of Medicine*, 1998, 339:321-328.
- Lieberman, J. et al. "Interfering with disease: opportunities and roadblocks to harnessing RNA interference" *Trends in Molecular Medicine*, Sep. 2003, 9(9):397-403.
- Liu, Y. et al. "Discovery of Inhibitors that Elucidate the Role of UCH-L1 Activity in the H1299 Lung Cancer Cell Line" *Chemistry & Biology*, 2003, 10(9):837-846.
- Liu, X. et al. "The influence of polymeric properties on chitosan/siRNA nanoparticle formulation and gene silencing" *Biomaterials*, 2007, 28:1280-1288.
- Mailand, N. et al. "Deregulated human Cdc14a phosphatase disrupts centrosome separation and chromosome segregation" *Nature Cell Biology*, 2002, 4(4):317-322.
- Martey, C.A. et al. "Cigarette smoke induces cyclooxygenase-2 and microsomal prostaglandin E_2 synthase in human lung fibroblasts: implications for lung inflammation and cancer" *Am J Physiol Lung Cell Mol Physiol*, 2004, 287:L981-991.
- Massion, P.P. et al. "The molecular basis of lung cancer: molecular abnormalities and therapeutic implications" *Respiratory Research*, 2003, 4(1):12.1-12.15.
- Matzke, M. et al. "RNA: Guiding Gene Silencing" *Science*, 2001, 293:1080-1083.
- McCaffrey, A.P. et al. "Gene expression: RNA interference in adult mice" *Nature*, Jul. 2002, 418(6893):38-39.

- Misono, K.S. "Natriuretic peptide receptor: Structure and signaling" *Molecular and Cellular Biochemistry*, 2002, 230:49-60.
- Mizuguchi, M. et al. "Bronchoprotective effects of atrial natriuretic peptide against propranolol-induced bronchoconstriction after allergic reaction in guinea pigs" *Clinical and Experimental Allergy*, 2000, 30:439-444.
- Mohapatra, S.S. et al. "Role of natriuretic peptide signaling in modulating asthma and inflammation" *Can J Physiol Pharmacol*, 2007, 85:754-759.
- Morita, R. et al. "Atrial Natriuretic Peptide Polarizes Human Dendritic Cells Toward a Th2-Promoting Phenotype Through Its Receptor Guanylyl Cyclase-Coupled Receptor A1" *The Journal of Immunology*, 2003, 170:5869-5875.
- Nafee, N. et al. "Chitosan-coated PLGA nanoparticles for DNA/RNA delivery: effect of the formulation parameters on complexation and transfection of antisense oligonucleotides" *Nanomedicine: Nanotechnology, Biology, and Medicine*, 2007, 3(3):173-183.
- Novina, C.D. et al. "The RNAi revolution" *Nature*, 2004, 430:161-164.
- Nuglozeh, E. et al. "Gene expression of natriuretic peptide receptors in rats with DOCA-salt hypertension" *Am J Physiol Cell Physiol*, 1997, 273:1427-1434.
- Nykanen, A. et al. "ATP Requirements and small Interfering RNA Structure" *Cell*, Nov. 2001, 107:300-321.
- Nyormoi, O. et al. "Transcriptional regulation of metastasis-related genes in human melanoma" *Clinical and Experimental Metastasis*, 2003, 20:251-263.
- Ogata, A. et al. "Isatin, an endogenous MAO inhibitor, improves bradykinesia and dopamine levels in a rat model of Parkinson's disease induced by Japanese encephalitis virus" *Journal of the Neurological Sciences*, 2003, 206(1):79-83.
- Oka, D. et al. "Sesquiterpene lactone parthenolide suppresses tumor growth in a xenograft model of renal cell carcinoma by inhibiting the activation of NF- κ B" *International Journal of Cancer*, 2007, 120:2576-2581.
- Oliveira, A.M. et al. "Tumor Suppressor Genes in Breast Cancer: The Gatekeepers and the Caretakers" *American Society for Clinical Pathology*, 2005, 124:S16-S28.
- Paddison, P.J. et al. "Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells" *Genes & Development*, 2002, 15:948-958.
- Palaparti, A. et al. "Inhibition of atrial natriuretic peptide (ANP) C receptor expression by antisense oligodeoxynucleotides in A10 vascular smooth-muscle cells is associated with attenuation of ANP-C-receptor-mediated inhibition of adenylyl cyclase" *Biochem J*, 2000, 346:313-320.
- Pandey, K.N. et al. "Molecular Cloning and Expression of Murine Guanylate Cyclase/Atrial Natriuretic Factor Receptor cDNA" *The Journal of Biological Chemistry*, Jul. 25, 1990, 265(21):12342-12348.
- Paul, C.P. et al. "Effective expression of small interfering RNA in human cells" *Nature Biotechnology*, May 2002, 20:505-508.
- Pedram, A. et al. "Natriuretic Peptides Inhibit G Protein Activation. Mediation Through Cross-Talk Between Cyclic GMP-Dependent Protein Kinase and Regulators of G Protein-Signaling Proteins" *J. Biol. Chem.*, 2000, 275:7365-7372.
- Plasterk, R.H.A., "RNA Silencing: The Genome's Immune System" *Science*, 2002, 296:1263-1265.
- Popp, F.D. "Synthesis of Potential Antineoplastic Agents. XX. Compounds Related to the 3-o-Nitrophenylhydrazones of Isatin" *Journal of Medicinal Chemistry*, 1969, 12(1):182-184.
- Reich, S.J. et al. "Small interfering RNA (siRNA) targeting VEGF effectively inhibits ocular neovascularization in a mouse model" *Molecular Vision*, 2003, 9:210-216.
- Saccani, A. et al. "p50 Nuclear Factor- κ B Overexpression in Tumor-Associated Macrophages Inhibits M1 Inflammatory Responses and Antitumor Resistance" *Cancer Research*, Dec. 2006, 66(23):11432-11440.
- Scadden, A.D.J. "RNAi is antagonized by A \rightarrow I hyper-editing" *EMBO Reports*, 2001, 11(2):1107-1111.
- Scherr, M. et al. "Inhibition of GM-CSF Receptor Function by Stable RNA Interference in a NOD/SCID Mouse Hematopoietic Stem Cell Transplantation Model" *Oligonucleotides*, Oct. 2003, 13(5): 353-363.
- Schipper, N.G.M. et al. "Chitosans as Absorption Enhancers for Poorly Absorbable Drugs. 1: Influence of Molecular Weight and Degree of Acetylation on Drug Transport Across Human Intestinal Epithelial (Caco-2) Cells" *Pharmaceutical Research*, 1996, 13(11):1686-1692.
- Schmidt, D. et al. "Critical role for NF κ B-induced JunB in VEGF regulation and tumor angiogenesis" *The EMBO Journal*, 2007, 26:710-729.
- Schwartz, A.G. et al. "The molecular epidemiology of lung cancer" *Carcinogenesis*, 2007, 28:507-518.
- Sharp, P.A. "RNAi and double-strand RNA" *Genes & Dev.*, 1999, 13:139-141.
- Sharp, P.A. "RNA interference—2001" *Genes & Development*, 2001, 15:485-490.
- Song, E. et al. "RNA interference targeting Fas protects mice from fulminant hepatitis" *Nature Medicine*, 2003, 9(3):347-351.
- Soutschek, J. et al. "Therapeutic silencing of an endogenous gene by systemic administration of modified siRNAs" *Nature*, Nov. 2004, 432:173-178.
- Sumpter, W.C. "The Chemistry of Isatin" *Chemical Reviews*, 1944, 34(3):393-434.
- Svoboda, P. et al. "Selective reduction of dormant maternal mRNAs in mouse oocytes by RNA interference" *Development*, Oct. 2000, 127:4147-4156.
- Tolentino, M.J. et al. "Intravitreal Injection of Vascular Endothelial Growth Factor Small Interfering RNA Inhibits Growth and Leakage in a Nonhuman Primate, Laser-Induced Model of Choroidal Neovascularization" *Retina*, 2004, 24:132-138.
- Tremblay, J. et al. "Biochemistry and physiology of the natriuretic peptide receptor guanylyl cyclases" *Molecular and Cellular Biochemistry*, 2002, 230:31-47.
- Tunny, T.J. et al. "Association of Restriction Fragment Length Polymorphism at the Atrial Natriuretic Peptide Gene Locus with Aldosterone Responsiveness to Angiotensin in Aldosterone-Producing Adenoma" *Biochemical and Biophysical Research Communications*, 1994, 204:1312-1317.
- Tuschl, T. et al. "Targeted mRNA degradation by double-stranded RNA in vitro" *Genes & Development*, Dec. 1999, 13:3191-3197.
- Tuschl, T. "RNA Interference and Small Interfering RNAs" *ChemBiochem*, 2001, 2:239-245.
- Tuschl, T. "Expanding small RNA interference" *Nature Biotechnology*, 2002, 20:446-448.
- Vellaichamy, E. et al. "Reduced cGMP signaling activates NF- κ B in hypertrophied hearts of mice lacking natriuretic peptide receptor-A" *Biochemical and Biophysical Research Communications*, 2005, 327:106-111.
- Verma, I.M. et al. "Gene Therapy—promises, problems and prospects" *Nature*, 1997, 389:239-242.
- Vesely, D.L. et al. "Atrial natriuretic peptides negatively and positively modulate circulating endothelin in humans" *Metabolism*, 1996, 45:315-319.
- Vesely, D.L. et al. "Vessel dilator, long acting natriuretic peptide, and kaliuretic peptide increase circulating prostaglandin E₂" *Life Sciences*, 2000, 66:095-913.
- Vesely, D.L. et al. "Atrial Natriuretic Hormone, Vessel Dilator, Long-Acting Natriuretic Hormone, and Kaliuretic Hormone Decrease the Circulating Concentrations of CRH, Corticotropin, and Cortisol" *J Clin Endocrinol Metab*, 2001, 86:4244-4249.
- Vesely, D.L. et al. "Atrial Natriuretic Hormone, Vessel Dilator, Long Acting Natriuretic Hormone, and Kaliuretic Hormone Decrease Circulating Prolactin Concentrations" *Norm Metab Res*, 2002, 34:245-249.
- Vesely, D.L. "Atrial Natriuretic Peptide Prohormone Gene Expression: Hormones and Diseases That Upregulate its Expression" *IUBMB Life*, 2002, 53(3):153-159.
- Vesely, D.L. et al. "Four Cardiac Hormones Eliminate up to Two-Thirds of Human Breast Cancers in Athymic Mice" *In Vivo*, 2007, 21:973-978.

- Vesely, D.L. et al. "Elimination of Up to 80% of Human Pancreatic Adenocarcinomas in Athymic Mice by Cardiac Hormones" *In Vivo*, 2007, 21:445-452.
- Vesely, B.A. et al. "Four peptide hormones decrease the number of human breast adenocarcinoma cells" *European Journal of Clinical Investigation*, 2005, 35:60-69.
- Vesely, B.A. et al. "Four Cardiac Hormones Cause Cell Death of Melanoma Cells and Inhibit Their DNA Synthesis" *Am J Med Sci*, 2007, 334(5):342-349.
- Vesely, D.L. "Atrial natriuretic peptides: anticancer agents" *J Investig Med.*, 2005, 53(7):360-365.
- Vilimas, T. et al. "Targeting the NF- κ B signaling pathway in Notch1-induced T-cell leukemia" *Nature Medicine*, 2007, 13:70-77.
- Vine, K.L. et al. "In vitro cytotoxicity evaluation of some substituted isatin derivatives" *Bioorganic & Medicinal Chemistry*, 2007, 15(2):931-938.
- Wang, X. et al. "siRNA Targeting the Natriuretic Peptide Receptor-A Prevents Airway Inflammation in a Mouse Model of Allergic Asthma" *Journal of Allergy and Clinical Immunology*, Jan. 2007, 119(1):S131, abstract 515.
- Wang, X. et al. "Modulation of lung inflammation by vessel dilator in a mouse model of allergic asthma" *Respiratory Research*, Jul. 2009, 10(66):1-8.
- Wang, X. et al. "Prevention of airway inflammation with topical cream containing imiquimod and small interfering RNA for natriuretic peptide receptor" *Genetic Vaccines and Therapy*, Feb. 2008, 6(7):1-9.
- Woodle, M.C. et al. "Sterically stabilized liposomes" *Biochimica et Biophysica Acta: Reviews on Biomembranes*, 1992, 1113:171-199.
- Xia, H. et al. "siRNA-mediated gene silencing in vitro and in vivo" *Nature Biotechnology*, Oct. 2002, 20(10):1006-1010.
- Yu, J.Y. et al. "RNA interference by expression of short-interfering RNAs and hairpin RNAs in mammalian cells" *PNAS USA*, Apr. 2002, 99(9):6047-6052.
- Yuhas, J.M. et al. "Specific and Nonspecific Stimulation of Resistance to the Growth and Metastasis of the Line 1 Lung Carcinoma" *Cancer Research*, 1975, 35:242-244.
- Zamore, P.D. "Ancient Pathways Programmed by Small RNAs" *Science*, 2002, 296(5571):1265-1269.
- Zamore, P.D. "RNA interference: listening to the sound of silence" *Nature Structural Biology*, Sep. 2001, 8(9):746-750.
- Zamore, P.D. et al. "RNAi: Double-Stranded RNA Directs the ATP-Dependent Cleavage of mRNA at 21-23 Nucleotide Intervals" *Cell*, Mar. 31, 2000, 101(1):25-33.
- Zhang, X. et al. "Small Interfering RNA Targeting Heme Oxygenase-1 Enhances Ischemia-Reperfusion-induced Lung Apoptosis" *The Journal of Biological Chemistry*, Mar. 2004, 279(11):10677-10684.
- Zeng, Y. et al. "Both Natural and Designed Micro RNAs Can Inhibit the Expression of Cognate mRNAs When Expressed in Human Cells" *Molecular Cell*, Jun. 2002, 9:1327-1333.
- Zimmerman, T.S. et al. "RNAi-mediated gene silencing in non-human primates" *Nature*, May 2006, 441:111-114.
- Abadi, A.H. et al. "Synthesis of 3-substituted-2-oxindole analogues and their evaluation as kinase inhibitors, anticancer and antiangiogenic agents" *European Journal of Medicinal Chemistry*, 2006, 41(3):296-305.
- Office Action dated Aug. 9, 2007 in U.S. Appl. No. 11/059,814, filed Feb. 17, 2005.
- Office Action dated Apr. 25, 2008 in U.S. Appl. No. 11/059,814, filed Feb. 17, 2005.
- Advisory Action dated Oct. 15, 2008 in U.S. Appl. No. 11/059,814, filed Feb. 17, 2005.
- Office Action dated Sep. 2, 2009 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Office Action dated Apr. 1, 2010 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Office Action dated Aug. 20, 2010 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Office Action dated Jun. 1, 2009 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.
- Office Action dated Dec. 11, 2009 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.
- Office Action dated May 18, 2010 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.
- Office Action dated Oct. 13, 2010 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.
- Benson, J.D. et al. "Validating cancer drug targets" *Nature*, May 2006, 441:451-456.
- Brafford, P. et al. "Gene expression profiling of melanoma cells—searching the haystack" *Journal of Translational Medicine*, 2005, 3:2, pp. 1-2.
- Carr, K.M. et al. "Gene-expression profiling in human cutaneous melanoma" *Oncogene*, 2003, 22:3076-3080.
- Chen, J.H. "Application of cationic polymer vector for gene delivery systems" Yao Xue Xue Bao, Apr. 2003, 38(4):316-20, abstract.
- Chengalvala, M.V. et al. "Gene Expression Profiling and its Practice in Drug Development" *Current Genomics*, 2007, 8(4):262-270.
- Chin, L. et al. "Malignant melanoma: genetics and therapeutics in the genomic era" *Genes & Development*, 2006, 20:2149-2182.
- De Wit, N.J.W. et al. "Analysis of differential gene expression in human melanocytic tumour lesions by custom made oligonucleotide arrays" *British Journal of Cancer*, 2005, 92:2249-2261.
- Fluge, T. et al. "Bronchodilation using combined urotilatin—albuterol administration in asthma: a randomized, double-blind, placebo-controlled trial" *European Journal of Medical Research*, 1999, 4(10):411-415.
- Gogas, H. et al. "Biomarkers in melanoma" *Annals of Oncology*, Aug. 2009, 20(Supplement 6):vi8-vi13.
- Haberman, A.B. "Strategies to Move Beyond Target Validation" *Genetic Engineering News*, Dec. 2005, 25(21):36.
- Hulks, G. et al. "Bronchodilator effect of atrial natriuretic peptide in asthma" *Br Med J*, Oct. 28, 1989, 299:1081-1082.
- Levy, J.A. et al. "Inactivation of Murine RNA Tumor Viruses by Isatin Beta-Thiosemicarbazone" *Virology*, Oct. 15, 1976, 74(2):426-431.
- Lobbezoo, M.W. et al. "Signal Transduction Modulators for Cancer Therapy: From Promise to Practice?" *The Oncologist*, 2003, 8:210-213.
- Martinez, S.R. et al. "Molecular Markers in Malignant Cutaneous Melanoma: Gift Horse or One-Trick Pony?" *Journal of Cellular Biochemistry*, 2005, 96:473-483.
- Rahmanto, Y.S. et al. "Identification of distinct changes in gene expression after modulation of melanoma tumor antigen p97 (melanotransferrin) in multiple models in vitro and in vivo" *Carcinogenesis*, 2007, 28(10):2172-2183.
- Schwab, G. et al. "An approach for new anticancer drugs: Oncogene-targeted antisense DNA" *Ann Oncol*, 1994, 5(Supplement 4):S55-S58.
- Schwarze, J. et al. "Respiratory Syncytial Virus Infection Results in Airway Hyperresponsiveness and Enhanced Airway Sensitization to Allergen" *J Clin Invest*, 1997, 99:226-233.
- Seftor, E.A. et al. "Expression of multiple molecular phenotypes by aggressive melanoma tumor cells: role in vasculogenic mimicry" *Critical Reviews in Oncology/Hematology*, 2002, 44:17-27.
- Skubitz, A.P.N. et al. "Differential gene expression identifies subgroups of ovarian carcinoma" *Translational Research*, Nov. 2006, 148(5):223-248.
- Sprenger, H. et al. "The lack of receptors for atrial natriuretic peptides on human monocytes prevents a rise of cGMP and induction of tumor necrosis factor- α synthesis" *Immunobiology*, Sep. 1991, 183(1-2):94-101.
- Sun, Y. et al. "Atrial Natriuretic Peptide and Long Acting Natriuretic Peptide Inhibit ERK 1/2 in Prostate Cancer Cells" *Anticancer Research*, 2006, 26:4143-4148.
- Sun, Y. et al. "Vessel Dilator and Kaliuretic Peptide Inhibit ERK 1/2 Activation in Human Prostate Cancer Cells" *Anticancer Research*, 2006, 26:3217-3222.
- Vesely, D.L. et al. "The N-Terminus of the Atrial Natriuretic Factor Prohormone in the Pleural Fluid of Congestive Heart Failure Patients" *Chest*, 1990, 97(6):1295-1298.
- Vesely, D.L. "Aprotinin blocks the binding of pro atrial natriuretic peptides 1 to 30, 31 to 67, and 99-126 to human placental membranes" *Am J Obstet Gynecol*, 1991, 165(3):567-573.

- Vesely, D.L. "Atrial Natriuretic Hormones Originating from the N-Terminus of the Atrial Natriuretic Factor Prohormone" *Clin Exp Pharmacol Physiol*, 1995, 22(2):108-114.
- Vesely, D.L. et al. "Long-Acting Natriuretic Peptide, Vessel Dilator, and Kaliuretic Peptide Enhance the Urinary Excretion Rate of β_2 -Microglobulin" *Metabolism*, Dec. 2000, 49(12):1592-1597.
- Vita, M. et al. "The Myc oncoprotein as a therapeutic target for human cancer" *Seminars in Cancer Biology*, 2006, 16:318-330.
- Winnepennickx, V. et al. "Gene Expression Profiling of Primary Cutaneous Melanoma and Clinical Outcome" *Journal of the National Cancer Institute*, Apr. 5, 2006, 98(7):472-482.
- Wong, H.H. et al. "Pancreatic cancer: molecular pathogenesis and new therapeutic targets" *Nat Rev Gastroenterol Hepatol*, 2009, 6:412-422.
- Yang, J. et al. "Conditional ablation of *Ikkb* inhibits melanoma tumor development in mice" *The Journal of Clinical Investigation*, Jul. 2010, 120(7):2563-2574.
- ATCC No. CCL-248 (T84, 1984), 2 pages.
- Bellone, M. et al. "Cancer immunotherapy: synthetic and natural peptides in the balance" *Immunology Today*, Oct. 1999, 20(10):457-462.
- Benjamin, B.A. et al. "Effect of proANF-(31-67) on sodium excretion in conscious monkeys" *Am J Physiol Regul Integr Comp Physiol*, 1995, 269:R1351-R1355, Abstract.
- Buskins, C. et al. "Adenocarcinomas of the Gastro-Esophageal Junction: A Comparative Study of the Gastric Cardia and the Esophagus with Respect to Cyclooxygenase-2 Expression" *Digestive Disease Week Abstracts and Itinerary Planner*, 2003, Abstract No. 850.
- Capizzi, R.L. "Molecular and Cellular Biology of Cancer" In: Internal Medicine, 4th Edition, Ed. Jay Stein, Elsevier Science, 1994, pp. 707-729.
- Carter, S.K. et al. "Chemotherapy of Cancer" Second Edition, John Wiley & Sons: New York, 1981, Appendix C, 5 pages.
- Chow, W.H. et al. "Rising Incidence of Renal Cell Cancer in the United States" *JAMA*, May 5, 1999, 281(17):1628-1631.
- Clark, J.I. et al. "Adjuvant High-Dose Bolus Interleukin-2 for Patients With High-Risk Renal Cell Carcinoma: A Cytokine Working Group Randomized Trial" *J Clin Oncol*, Aug. 15, 2003, 21(16):3133-3140.
- Cohen, H.T. et al. "Medical Progress: Renal-Cell Carcinoma" *N. Engl J Med*, Dec. 8, 2005, 353(23):2477-2490.
- Daggubati, S. et al. "Adrenomedullin, endothelin, neuropeptide Y, atrial, brain, and C-natriuretic prohormone peptides compared as early heart failure indicators" *Cardiovascular Research*, 1997, 36:246-255.
- De Palo, E.F. et al. "Circulating Immunoreactive proANP(1-30) and proANP(31-67) in Sedentary Subjects and Athletes" *Clinical Chemistry*, 2000, 46(6):843-847.
- Dermer, G.B. "The Last Word: Another Anniversary for the War on Cancer" *Bio/Technology*, Mar. 1994, 12:320.
- Dietz, J.R. et al. "Evidence supporting a physiological role for proANP(1-30) in the regulation of renal excretion" *Am J Physiol Regul Integr Comp Physiol*, 2001, 280:R1510-R1517.
- Fisher, E. et al. "Comparative Histopathologic, Histochemical, Electron Microscopic and Tissue Culture Studies of Bronchial Carcinoids and Oat Cell Carcinomas of Lung" *Am J Clin Pathol*, Feb. 1978, 69(2):165-172.
- Franz, M. et al. "N-terminal fragments of the proatrial natriuretic peptide in patients before and after hemodialysis treatment" *Kidney International*, 2000, 58:374-383.
- Franz, M. et al. "Plasma concentration and urinary excretion of N-terminal proatrial natriuretic peptides in patients with kidney diseases" *Kidney International*, 2001, 59:1928-1934.
- Freshney, R.I. "Culture of Animal Cells: A Manual of Basic Technique" Alan R. Liss, Inc.: New York, 1983, pp. 3-4.
- Friedman, H.S. et al. "Glioblastoma Multiforme and the Epidermal Growth Factor Receptor" *N. Engl J Med*, Nov. 10, 2005, 353(19):1997-1999.
- Gaiger, A. et al. "Immunity to WT1 in the animal model and in patients with acute myeloid leukemia" *Blood*, Aug. 15, 2000, 96(4):1480-1489.
- Garrett, M.D. et al. "Discovering Novel Chemotherapeutic Drugs for the Third Millennium" *European Journal of Cancer*, 1999, 35(14):2010-2030.
- Gower, W.R. et al. "Four Peptides Decrease Human Colon Adenocarcinoma Cell Number and DNA Synthesis via Cyclic GMP" *International Journal of Gastrointestinal Cancer*, 2005, 36(2):77-88.
- Gratzner, H.G., "Monoclonal Antibody to 5-Bromo- and 5-Iododeoxyuridine: A New Reagent for Detection of DNA Replication" *Science*, Oct. 29, 1982, 218:474-475.
- Gunning, M.E. et al. "Atrial Natriuretic Peptide₍₃₁₋₆₇₎Inhibits Na⁺Transport in Rabbit Inner Medullary Collecting Duct Cells: Role of Prostaglandin E₂" *Journal of Clinical Investigation*, May 1992, 89:1411-1417.
- Heim, J.M. et al. "Urodilatin and β -ANF: Binding Properties and Activation of Particulate Guanylate Cyclase" *Biochemical and Biophysical Research Communications*, Aug. 30, 1989, 163(1):37-41.
- Hunter, E.F.M. et al. "Analysis of peptides derived from Pro Atrial Natriuretic Peptide that circulate in man and increase in heart disease" *Scand J Clin Lab Invest*, 1998, 58:205-216.
- Jemal, A. et al. "Cancer Statistics, 2005" *CA: A Cancer Journal for Clinicians*, 2005, 55:10-30.
- Kaiser, J. et al. "CANCER: First Pass at Cancer Genome Reveals Complex Landscape" *Science*, Sep. 8, 2006, 313:1370.
- Krontiris, T.G. "Molecular and Cellular Biology of Cancer" In: Internal Medicine, 4th Edition, Ed. Jay Stein, Elsevier Science, 1994, pp. 699-707.
- Kumar, R. et al. "Stimulation of atrial natriuretic peptide receptor/guanylyl cyclase- A signaling pathway antagonizes the activation of protein kinase C- α in murine Leydig cells" *Biochimica et Biophysica Acta*, 1997, 1356:221-228.
- La Vecchia, C. et al. "Smoking and Renal Cell Carcinoma" *Cancer Research*, Sep. 1, 1990, 50:5231-5233.
- Lelièvre, V. et al. "Proliferative Actions of Natriuretic Peptides on Neuroblastoma Cells" *The Journal of Biological Chemistry*, Nov. 23, 2001, 276(47):43668-43676.
- Linder, M.W. et al. "Pharmacogenetics: a laboratory tool for optimizing therapeutic efficiency" *Clinical Chemistry*, 1997, 43(2):254-266.
- Martin, D.R. et al. "Three peptides from the ANF prohormone NH(2)-terminus are natriuretic and/or kaliuretic" *Am J Physiol Renal Physiol*, Dec. 1990, 258(5):F1401-F1408, Abstract.
- McLaughlin, J.K. et al. "A Population-Based Case-Control Study of Renal Cell Carcinoma" *J Natl Cancer Inst*, Feb. 1984, 72:275-284.
- Mellinghoff, I.K. et al. "Molecular Determinants of the Response of Glioblastomas to EGFR Kinase Inhibitors" *N Engl J Med*, Nov. 10, 2005, 353(19):2012-2024.
- Morstyn, G. et al. "Immunohistochemical Identification of Proliferating Cells in Organ Culture Using Bromodeoxyuridine and a Monoclonal Antibody" *The Journal of Histochemistry and Cytochemistry*, 1986, 34(6):697-701.
- Nguyen, T.D. et al. "Citrus Flavonoids Stimulate Secretion by Human Colonic T₈₄ Cells^{1,2}" *The Journal of Nutrition*, 1993, 123:259-268.
- Pan, E. et al. "Central Nervous System: Primary Neoplasms of the Central Nervous System" In: Kufe, D.W., Pollock, R.F., Weichselbaum, R.R., Bast, R.C., Jr., Gansler, I.S., Holland, J.F., and Frei, E., III (Eds.), *Cancer Medicine*, 6th Edition, 2003, pp. 1193-1226, London: BC Decker.
- Pitari, G.M. et al. "Guanylyl cyclase C agonists regulate progression through the cell cycle of human colon carcinoma cells" *PNAS*, Jul. 3, 2001, 98(14):7846-7851.
- Rashed, H.M. et al. "Atrial Natriuretic Peptide Inhibits Growth of Hepatoblastoma (HEP G2) Cells by Means of Activation of Clearance Receptors" *Hepatology*, Apr. 1993, 17(4):677-684.
- Ropper, A.H. and Brown, R.H. (Eds.) "Intracranial Neoplasms and Paraneoplastic Disorders" In: Adams and Victor's Principles of Neurology, 8th Edition, 2005, New York: McGraw-Hill, pp. 546-591.
- Rosenzweig, A. et al. "Atrial Natriuretic Factor and Related Peptide Hormones" *Annu. Rev. Biochem.*, 1991, 60:229-255.
- Saba, S.R. et al. "Immunocytochemical Localization of Atrial Natriuretic Peptide, Vessel Dilator, Long-acting Natriuretic Peptide, and Kaliuretic Peptide in Human Pancreatic Adenocarcinomas" *Journal of Histochemistry & Cytochemistry*, 2005, 53(8):989-995.

- Salani, D. et al. "Endothelin-1 Induces an Angiogenic Phenotype in Cultured Endothelial Cells and Stimulates Neovascularization *In Vivo*" *American Journal of Pathology*, Nov. 2000, 157(5):1703-1711.
- Saxenhofer, H. et al. "Urodilatin: binding properties and stimulation of cGMP generation in rat kidney cells" *Am J Physiol Renal Physiol*, 1993, 264:F267-F273, Abstract.
- Schulz-Knappe, P. et al. "Isolation and Structural Analysis of "Urodilatin", a New Peptide of the Cardiodilatin-(ANP)-Family, Extracted from Human Urine" *Klin Wochenschr*, 1988, 66:752-759.
- Schwede, F. et al. "Cyclic nucleotide analogs as biochemical tools and prospective drugs" *Pharmacology & Therapeutics*, 2000, 87:199-226.
- Schweitz, H. et al. "A New Member of the Natriuretic Peptide Family Is Present in the Venom of the Green Mamba (*Dendroaspis angusticeps*)" *The Journal of Biological Chemistry*, Jul. 15, 1992, 267(20):13928-13932.
- Scott, D.A. et al. "The Pendred syndrome gene encodes a chloride-iodide transport protein" *Nature Genetics*, Apr. 1999, 21:440-443.
- Shapiro, J.A. et al. "Body Mass Index and Risk of Renal Cell Carcinoma" *Epidemiology*, Mar. 1999, 10(2):188-191.
- Senger, D. et al. "Long-Term Survivors of Glioblastoma: Statistical Aberration or Important Unrecognized Molecular Subtype?" *The Cancer Journal*, May/Jun. 2003, 9(3):214-221.
- Sudoh, T. et al. "C-Type Natriuretic Peptide (CNP): A New Member of Natriuretic Peptide Family Identified in Porcine Brain" *Biochemical and Biophysical Research Communications*, Apr. 30, 1990, 168(2):863-870.
- Turner, G.A. et al. "Urine cyclic nucleotide concentrations in cancer and other conditions; cyclic GMP: a potential marker for cancer treatment" *J Clin Pathol*, 1982, 35:800-806.
- Valentin, J.P. et al. "Urodilatin Binds to and Activates Renal Receptors for Atrial Natriuretic Peptide" *Hypertension*, Apr. 1993, 21(4):432-438.
- Van Meir, E.G. et al. "Release of an inhibitor of angiogenesis upon induction of wild type p35 expression in glioblastoma cells" *Nature Genetics*, Oct. 1994, 8:171-176.
- Vesely, B.A. et al. "Five cardiac hormones decrease the number of human small-cell lung cancer cells" *Eur J Clin Invest*, 2005, 35:388-398.
- Vesely, B.A. et al. "Four cardiac hormones cause cell death in 81% of human ovarian adenocarcinoma cells" *Cancer Therapy*, 2007, 5:97-104.
- Vesely, B.A. et al. "Four cardiac hormones eliminate 4-fold more human glioblastoma cells than the green mamba snake peptide" *Cancer Letters*, 2007, 254:94-101.
- Vesely, B.A. et al. "Four peptide hormones' specific decrease (up to 97%) of human prostate carcinoma cells" *Eur J Clin Invest*, 2005, 35(11):700-710.
- Vesely, B.A. et al. "Primary Malignant Tumors of the Heart: Four Cardiovascular Hormones Decrease the Number and DNA Synthesis of Human Angiosarcoma Cells" *Cardiology*, 2006, 105:226-233.
- Vesely, B.A. et al. "Vessel dilator: Most potent of the atrial natriuretic peptides in decreasing the number of DNA synthesis of human squamous lung cancer cells" *Cancer Letters*, 2006, 233:226-231.
- Vesely, B.A. et al. "Four peptide hormones decrease the number of human breast adenocarcinoma cells" *Eur J Clin Invest*, 2005, 35(1):60-69.
- Vesely, B.A. et al. "Urodilatin and four cardiac hormones decrease human renal carcinoma cell numbers" *Eur J Clin Invest*, 2006, 36(11):810-819.
- Vesely, D.L. et al. "Atrial Natriuretic Peptide Increases Urodilatin in the Circulation" *American Journal of Nephrology*, 1998, 18(3):204-213, Abstract.
- Vesely, D.L. et al. "Atrial Natriuretic Prohormone Peptides 1-30, 31-67, and 79-98 Vasodilate the Aorta" *Biochemical and Biophysical Research Communications*, Nov. 13, 1987, 148(3):1540-1548.
- Vesely, D.L. et al. "Increased Release of the N-Terminal and C-Terminal Portions of the Prohormone of Atrial Natriuretic Factor During Immersion-Induced Central Hypervolemia in Normal Humans¹" *Proc Soc Exp Biol Med*, 1989, 192:230-235.
- Vesely, D.L. et al. "Negative Feedback of Atrial Natriuretic Peptides" *Journal of Clinical Endocrinology and Metabolism*, 1994, 78(5):1128-1134.
- Vesely, D.L. et al. "Novel therapeutic approach for cancer using four cardiovascular hormones" *Eur J Clin Invest*, 2004, 34(10):674-682.
- Vesely, D.L. et al. "Three Peptides From the Atrial Natriuretic Factor Prohormone Amino Terminus Lower Blood Pressure and Produce Diuresis, Natriuresis, and/or Kaliuresis in Humans" *Circulation*, Sep. 1994, 90(3):1129-1140.
- Vesely, D.L. et al. "Vessel Dilator Enhances Sodium and Water Excretion and Has Beneficial Hemodynamic Effects in Persons With Congestive Heart Failure" *Circulation*, Jul. 28, 1998, 98:323-329.
- Vesely, D.L. "Natriuretic peptides and acute renal failure" *Am J Physiol Renal Physiol*, 2003, 285:F167-F177.
- Villarreal, D. et al. "Hemodynamic and Renal Effects of ProANF₃₁₋₆₇ in Hypertensive Rats₍₄₄₃₉₉₎" *Proceedings of the Society for Experimental Biology and Medicine*, 1999, 221(3):166-170.
- Weller, M. et al. "Predicting Chemoresistance in Human Malignant Glioma Cells: The Role of Molecular Genetic Analysis" *Int. J. Cancer (Pred. Oncol.)*, 1998, 79:640-644.
- White, R.E. et al. "Potassium channel stimulation by natriuretic peptides through cGMP-dependent dephosphorylation" *Nature*, Jan. 21, 1993, 361:263-266.
- Wigle, D.A. et al. "ANP secretion from small cell lung cancer lines: a potential model of ANP release" *Am J Physiol Heart Circ Physiol*, 1995, 268(5):H1869-H1874, Abstract.
- Winters, C.J. et al. "The N-Terminus and a 4,000-MW Peptide From the Midportion of the N-Terminus of the Atrial Natriuretic Factor Prohormone Each Circulate in Humans and Increase in Congestive Heart Failure" *Circulation*, Sep. 1989, 80(3):438-449.
- Yagoda, A. et al. "Chemotherapy for Advanced Renal-Cell Carcinoma: 1983-1993" *Seminars in Oncology*, Feb. 1995, 22(1):42-60.
- Yang, J.C. et al. "Randomized Study of High-Dose and Low-Dose Interleukin-2 in Patients With Metastatic Renal Cancer" *J Clin Oncol*, Aug. 15, 2003, 21(16):3127-3132.
- Yu, C.C.W. et al. "The assessment of cellular proliferation by immunohistochemistry: A review of currently available methods and their applications" *The Histochemical Journal*, Mar. 1992, 24(3):121-131.
- Yu, M.C. et al. "Cigarette Smoking, Obesity, Diuretic Use, and Coffee Consumption as Risk Factors for Renal Cell Carcinoma" *J Natl Cancer Inst*, Aug. 1986, 77:351-356.
- Zeidel, M.L. "Regulation of Collecting Duct Na⁺ Reabsorption by ANP 31-67" *Clinical and Experimental Pharmacology and Physiology*, 1995, 22(2):121-124, Abstract.
- Zellner, A. et al. "Disparity in Expression of Protein Kinase C α in Human Glioma versus Glioma-derived Primary Cell Lines: Therapeutic Implications" *Clinical Cancer Research*, Jul. 1998, 4:1797-1802.
- Zips, D. et al. "New Anticancer Agents: *In Vitro* and *In Vivo* Evaluation" *In Vivo*, Jan./Feb. 2005, 19(1):1-7.
- Office Action dated Jan. 26, 2011 in Canadian Application Serial No. 2,518,550, filed Mar. 19, 2004.
- Office Action dated Apr. 2, 2009 in U.S. Appl. No. 11/833,757, filed Aug. 3, 2007.
- Office Action dated Oct. 19, 2009 in U.S. Appl. No. 11/833,757, filed Aug. 3, 2007.
- Response to Office Action dated Apr. 2, 2009 in U.S. Appl. No. 11/833,757, filed Aug. 3, 2007.
- Response to Office Action dated Oct. 19, 2009 in U.S. Appl. No. 11/833,757, filed Aug. 3, 2007.
- Office Action dated Apr. 27, 2011 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Response to Office Action dated Sep. 2, 2009 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Response to Office Action dated Apr. 1, 2010 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Response to Office Action dated Aug. 20, 2010 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Response to Office Action dated Apr. 27, 2011 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.
- Response to Office Action dated Jun. 1, 2009 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.
- Response to Office Action dated Dec. 11, 2009 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.

Response to Office Action dated May 18, 2010 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.

Response to Office Action dated Oct. 13, 2010 in U.S. Appl. No. 11/998,792, filed Nov. 30, 2007.

Notice of Allowance dated Jul. 27, 2011 in U.S. Appl. No. 11/799,225, filed Apr. 30, 2007.

Office Action dated Jun. 27, 2011 in U.S. Appl. No. 12/908,806, filed Oct. 20, 2010.

Gower, W.R. et al. "Four Peptides Decrease Human Colon Adenocarcinoma Cell Number and DNA Synthesis via Cyclic GMP" *International Journal of Gastrointestinal Cancer*, 2005, 36(2):77-88.
Keimer, A.K. et al. "Effects of Different Natriuretic Peptides on Nitric Oxide Synthesis in Macrophages" *Endocrinology*, 1997, 138(10):4282-4290.

* cited by examiner




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4

FIG. 5A

DAPI



FIG. 5B

Control




FIG. 5C


 α -FLAG

FIG. 5D




FIG. 5E





FIG. 6A

**Control**

FIG. 6B

**Vehicle**

FIG. 6C

**NP₇₃₋₁₀₂**




FIG. 6D




FIG. 7




FIG. 8

1

MATERIALS AND METHODS FOR TREATMENT OF INFLAMMATORY AND CELL PROLIFERATION DISORDERS

CROSS-REFERENCE TO RELATED APPLICATION

The present application is a continuation of U.S. application Ser. No. 11/059,814, filed Feb. 17, 2005, now abandoned which claims the benefit of U.S. Provisional Application Ser. No. 60/521,072, filed Feb. 17, 2004, each of which is hereby incorporated by reference herein in its entirety, including any figures, tables, nucleic acid sequences, amino acid sequences, or drawings.

BACKGROUND OF THE INVENTION

The vast majority of cancers of the lung, breast and colon are adenocarcinomas, which arise from pre-existing adenomatous polyps that develop in the normal colonic mucosa. This adenoma-carcinoma sequence is a well-characterized clinical and histopathologic series of events with which discrete molecular genetic alterations have been associated. Lung tumor development and metastasis are complex processes that include transformation, proliferation, resistance to apoptosis, neovascularization, and metastatic spread. A number of gene products have been identified that play critical roles in these processes. It has been suggested that the development of epithelial-derived tumors, the most common class of cancers, involves mutations of tumor suppressors and proto-oncogenes or epigenetic alterations of signaling pathways affecting cell proliferation and/or survival, which in turn may be caused by inflammation induced by infections and reactive oxygen species (ROS) (Ernst, P. *Aliment Pharmacol Ther.*, 1999, 13(1):13-18).

A group of four peptide hormones, originating from the 126-amino acid atrial natriuretic factor (ANF) prohormone, have become known for their vasodilator activity. These four peptide hormones, consisting of amino acids 1-30, 31-67, 79-98, and 99-126 of this prohormone, have been named long acting natriuretic peptide (LANP), vessel dilator (VD), kaliuretic peptide (KP), and atrial natriuretic peptide (ANP), respectively, for their most prominent effects (Angus R. M. et al., *Clin Exp Allergy* 1994, 24:784-788). The ANP sequence, particularly the C-terminal portion, is highly conserved among species (Seidman et al., *Science*, 1984, 226:1206-1209). ANP has been proposed to be useful for treatment of various cardiovascular, respiratory, and renal diseases (Vesely, D. L. *Cardiovascular*, 2001, 51:647-658), but also causes inflammation. The family of natriuretic hormone peptides has been shown to have broad physiologic effects, including vasodilation and inhibition of aldosterone secretion and cardiovascular homeostasis.

As indicated above, ANF, the 126 amino acid prohormone, gives rise to four peptides: LANP (amino acids 1-30), VD (amino acids 31-67), KP (amino acids 79-98) and ANP (amino acids 99-126, also referred to herein as NP₉₉₋₁₂₆) (Angus R. M. et al., *Clin Exp Allergy*, 1994, 24:784-788). The ANP sequence particularly the C-terminal portion is highly conserved among species (Seidman et al., *Science*, 1984, 226:1206-1209). The natriuretic peptide receptors (NPRs), NPR-A and NPR-B, are expressed in many different tissues of various organs systems, and are coupled to guanylyl cyclase. ANP and BNP are thought to signal primarily through NPR-A by increasing cGMP and activating cGMP-dependent protein kinase (PKG). NPR-A is the primary receptor for ANP while NPR-B seems to bind CNP most

2

effectively. PKG activation in turn activates ion transporters and transcription factors, which together affect cell growth and proliferation, apoptosis and inflammation. NPR-C is a clearance receptor for ANP removal, but also appears to signal phospholipase C activation and a decrease in adenylyl cyclase activity through a cGMP-independent pathway (Abbey and Potter, *Endocrinology*, 2003, 144: 240-246; Silberbach and Roberts, *Cell Signal*, 2001, 13:221-231). The signaling mechanisms underlying ANP's growth regulatory effects are poorly understood, although a number of reports suggest that ANP acts through mitogen-activated protein kinases (Silberbach and Roberts, *Cell Signal*, 2001, 13:221-231).

Most cells of the mucosal immune system have ANP receptors (NPRs) and there is evidence that natriuretic peptides regulate the immune response and inflammation (Kurihara et al., *Biochem Biophys Res Commun* 1987, 149:1132-1140). ANP stimulates migration of human neutrophils (Izumi et al., *J Clin Invest* 2001, 108:203-213), and inhibit nitric oxide and TNF- α production by murine macrophages (Kierner and Vollmar, *J Biol Chem* 1998, 273:13444-13451; Kierner et al., *J Immunol* 2000, 165:175-81). It has been suggested that the ANP system may be a critical component of the immune response through its actions on both immune and non-immune cells. In patients with lung tumors, the immune response plays a large part in the progression of the disease and, consequently, the NPR system may potentially be involved. The alveolar macrophages in lung cancer patients secrete more pro-inflammatory cytokines, such as IL-6 and IL-1 β , after LPS stimulation than in persons with non-malignant disease (Matanic et al., *Scand J Immunol* 2003, 57:173-178). Increased IL-6 in lung cancer patients enhances the acute phase response, and is correlated with poor nutritional status and lowered survival (Martin et al., *Cytokine* 1999, 11: 267-273). The cells of the immune system, such as natural killer (NK) cells, V α 24 NKT, which are necessary for cancer surveillance may also be reduced in lung tumor patients (Motohashi et al., *Int J Cancer* 2002, 102:159-165). The most common clinical paraneoplastic syndrome in patients with small-cell lung cancer (SCLC) is hyponatremia, which is believed to be caused by tumor secretion of vasopressin. Tumor biopsies from patients with SCLC and hyponatremia expressed the gene for ANP (Shimizu et al., *Cancer* 1991, 68: 2284-2288; Bliss et al., *J Natl Can Inst*, 1990, 82: 305-310). Thus, the reduced sodium levels seen in SCLC patients may be attributed to the secretion of ANP (Bliss et al., *J Natl Can Inst*, 1990, 82: 305-310). Human SCLC cell lines express functional ANP receptors (Ohsaki et al., *Cancer Res* 1993, 53: 3165-3171). A majority of SCLC cell lines produce ANP and some produce BNP as well (Ohsaki et al., *Oncology* 1999, 56:155-159). In contrast, in NSCLC cell lines, which are derived mostly from adenocarcinomas that comprise about two-thirds of all lung cancers, little is known about their growth regulation in response to ANP cascade.

The present inventor has found that the N-terminal natriuretic peptides, such as pNP73-1102, are capable of inhibiting NF κ B activation (Mohapatra, international application WO 2004/022003, published Mar. 18, 2004, which is incorporated herein by reference in its entirety), and that the ANP cascade plays a critical role in cell proliferation and inflammation. NF κ B, a transcription factor and a key player in inflammatory processes, has been implicated in the development of cancer in liver and mammary tissues (Greten F. R. et al. *Cell*, 2004, 118: 285-296; Pikarsky F. et al. *Nature*, 2004, 431: 461-466). Activation of the NF- κ B pathway enhances tumor development and may act primarily in the late stages of tumorigenesis. Inhibition of NF- κ B signaling uniformly sup-

pressed tumor development; however, depending upon the model studied, this salutary effect was attributed to an increase in tumor cell apoptosis, reduced expression of tumor cell growth factors supplied by surrounding stromal cells, or abrogation of a tumor cell dedifferentiation program that is critical for tumor invasion/metastasis.

BRIEF SUMMARY OF THE INVENTION

The present invention pertains to methods for reducing natriuretic peptide receptor-A (also known in the art as NPRA, NPR-A, and guanylate cyclase A) activity in vitro or in vivo. The method of the invention may be used for treating inflammatory and cell proliferation disorders, such as cancer.

In another aspect, the present invention concerns methods for identifying agents useful for treating inflammatory and cell proliferation disorders by determining whether the candidate agent reduces activity of the natriuretic peptide receptor-A (also known in the art as NPRA, NPR-A, and guanylate cyclase A) in vitro and/or in vivo (also referred to herein as the diagnostic method or assay of the invention).

In another aspect, the method of the present invention may be used for reducing the growth of cancer cells in vitro or in vivo. In one aspect, the method is useful for treating cancers, such as adenocarcinomas of lung, breast, ovary and melanomas, which may be caused by cell proliferation and inflammation induced by the atrial natriuretic peptide (ANP) cascade.

In one embodiment, the method of the present invention comprises administering a therapeutically effective amount of an agent that reduces NPR-A activity. In another embodiment, the method of the present invention comprises administering a therapeutically effective amount of an N-terminal natriuretic peptide (referred to herein as NP or NP peptide), or a polynucleotide encoding NP and an operably-linked promoter sequence, to a patient in need of such treatment. As used herein, NP refers to peptides derived from atrial natriuretic factor (ANF) hormone, or a biologically active fragment, homolog, or variant thereof. In another embodiment, the method of the present invention comprises administering an effective amount of NP, or a polynucleotide encoding NP and an operably-linked promoter, to one or more cancer cells, wherein the NP is capable of reducing cell proliferation and/or tumor growth. The effect of the NP or a biologically active fragment, homolog, or variant thereof, is capable of reducing cancer cell growth in vitro or in vivo.

Specifically exemplified NPs comprise an amino acid sequence selected from the group consisting of amino acids 1-30 of ANF (also known as "long acting natriuretic peptide" and referred to herein as NP₁₋₃₀ or SEQ ID NO:1), amino acids 31-67 of ANF (also known as "vessel dilator" and referred to herein as NP₃₁₋₆₇ or SEQ ID NO:2), and amino acids 79-98 of ANF (also known as "kaliuretic peptide" and referred to herein as NP₇₉₋₉₈ or SEQ ID NO:3), or biologically active fragments or homologs of any of the foregoing. Other exemplified NPs comprise amino acids 73-102 of proANF (referred to herein as NP₇₃₋₁₀₂ or SEQ ID NO:5), or SEQ ID NO:6, or biologically active fragment(s) or homolog(s) of the foregoing. In one embodiment, the NP administered to the patient does not consist of NP₉₉₋₁₂₆ (SEQ ID NO:4).

In another embodiment, the method of the present invention comprises administering an effective amount of at least one nucleic acid molecule encoding an NP to a patient in need of such treatment. The present inventor has determined that introduction of a nucleic acid molecule encoding NP is capable of inhibiting tumor growth and tumor metastasis. The

gene delivery method of the present invention permits long-term expression of NP-encoding nucleic acid sequences in vivo, thereby conferring anti-cancer effects. In one embodiment, a therapeutically effective amount of at least one nucleic acid molecule encoding a peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5 or biologically active fragments or homologs of any of the foregoing, are administered to the patient.

In another aspect, the present invention concerns an isolated peptide comprising the amino acid sequence NP₇₃₋₁₀₂ (SEQ ID NO:5) or SEQ ID NO:6, or a biologically active fragment or homolog of the foregoing. In another aspect, the present invention concerns an isolated nucleic acid molecule encoding the amino acid sequence of NP₇₃₋₁₀₂ (SEQ ID NO:5) or encoding the amino acid sequence of SEQ ID NO:6, or a biologically active fragment or homolog thereof.

In another aspect, the present invention concerns an expression vector comprising a nucleic acid sequence encoding an NP, and a promoter sequence that is operably linked to the NP-encoding nucleic acid sequence. In one embodiment, the expression vector is a DNA plasmid or virus. In another aspect, the present invention concerns a pharmaceutical composition comprising a nucleic acid sequence encoding an NP, and a pharmaceutically acceptable carrier.

BRIEF DESCRIPTION OF THE DRAWINGS

For a fuller understanding of the nature and objects of the invention, reference should be made to the following detailed description, taken in connection with the accompanying drawings, in which:

FIG. 1 shows pNP 73-102 inhibits NPRA expression. Pregnant (12 days) mice were injected i.p. with pVAX (vector), or pNP73-102. After 1 day, mice were sacrificed, thymi removed from the embryo, and homogenized. Cells were centrifuged and erythrocytes were lysed and incubated with anti-NPR-Ab or anti-NPR-C for 1 hour, washed, and incubated with PE-conjugated secondary antibodies. Levels of NPRA and NPRC were determined by flow cytometry.

FIGS. 2A-2D show NPRA deficiency decreases pulmonary inflammation. Groups (n=3) of wild type DBA/2 (wt) (FIG. 2A) and NPR-C deficient (NPRC^{-/-}) (FIG. 2B) mice and wild type C57/BL6 (wt) (FIG. 2C) and NPR-A (NPR^{-/-}) (FIG. 2D) were sensitized with OVA (20 mg/mouse) and after 2 weeks challenged i.n. with OVA (20 mg/mouse). One day later mice were sacrificed and lung sections were stained with H & E to examine inflammation.

FIGS. 3A-3D demonstrate that A549 cells transfected with pNP₇₃₋₁₀₂ show a significantly higher level of apoptosis compared to control and pANP or pVAX (FIGS. 3A-3C). Cells were transfected with pNP73-102, pANP and pVAX (as control) and cells were stained with PI and annexin and quantified by flow cytometry (FIG. 3D). The proteins were isolated and an equal amount of the cell lysates were western-blotted using an antibody to poly-ADP ribose polymerase (PARP). The results demonstrate that pNP73-102 shows a higher accumulation of apoptotic cells compared to cells transfected with pANP and pVAX controls.

FIG. 4 shows that pNP73-102 decreases tumorigenesis in a colony formation assay by A549. Six centimeter tissue culture plates were covered with 4 ml of 0.5% soft agar. A549 cells were transfected with pANP, pNP73-102 and pVAX plasmid DNA (V) or nothing (C). After 40 h of transfection, cells were suspended in 2 ml of 0.3% soft agar and added to each plate. Cells were plated in duplicate at a density of 2×10^4

5

cells/dish and incubated for two weeks. Plates were photographed under a microscope. Cell colonies were counted and plotted. The results of one representative experiment of two is shown.

FIGS. 5A-5E show expression of NP₇₃₋₁₀₂-FLAG in the BAL cells after i.n. administration of chitosan encapsulated plasmid pNP₇₃₋₁₀₂-FLAG construct. BAL was performed in mice (n=3) after 24 hours and BAL cells were stained with either the second antibody control or anti-FLAG antibody (SIGMA) and then with DAPI. A representative staining is shown (FIGS. 5A-5C). FIG. 5D shows lungs removed from mice treated with chitosan nanoparticles carrying pNP₇₃₋₁₀₂ (CPNP73-102) (Rx) or empty plasmid pVAX (control). The lungs of control mice showed several lung nodules in contrast to mice treated with CPNP73-102, which showed very few tumors. Intranasal CPNP73-102 administration abrogated tumor formation in A549 injected nude mice. Nude mice were given 5×10⁶ cells intravenously (tail vein) and weekly injections of nanoparticle carrying either empty plasmid (control) or pNP73-102 (Rx). Three weeks later, mice were sacrificed and lung sections were stained with H & E to examine the lung nodules (FIG. 5D). Control shows nodules and tumor cell mass, whereas the treated group had no tumors. Sections were also stained with antibodies to cyclinB and to phospho-Bad (FIG. 5E). The results show that mice treated with CPNP73-102 had no tumors in the lung and did not show any staining for pro-mitotic Cyclin-B and anti-apoptotic marker phospho-Bad.

FIGS. 6A-6D demonstrate that treatment with chitosan nanoparticles carrying pNP₇₃₋₁₀₂ (CPNP73-102) decreases the tumor burden in a spontaneous tumorigenesis model of immunocompetent BALB/c mice. Two groups of mice (n=4) were administered with the Line-1 tumor cells (100,000 cells/mouse) at the flanks. One group was administered with CPNP73-102 the same day, whereas another group was administered with vehicle alone (nanoparticle carrying a plasmid without NP73-102) and the third group was given the saline. Treatment was continued with CPNP73-102 or control at weekly intervals for 5 weeks. The tumors were dissected out from the mice of each group (FIGS. 6A-6C) and the tumor burden was calculated by weighing them on a balance and expressed as tumor mass per g lung weight. Results are shown in FIG. 6D.

FIG. 7 shows that CPNP73-102 induces apoptosis in chemoresistant ovarian cancer cells. C-13 and OV2008 ovarian cancer cells were transfected with pNP73-102. Forty-eight hours later, cells were processed for TUNEL assay to examine apoptosis. The results of one of two representative experiments are shown.

FIG. 8 shows breast cancer MCF-7 cell counts. The cells were transfected with pVAX, pANP, and pANP₇₃₋₁₀₂ and counted at 24 and 48 hours after transfection. 30 ml of Trypan Blue was mixed with 30 ml for measuring the cell viability. The results of one of two representative experiments are shown.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO: 1 is the amino acid sequence of human “long acting natriuretic peptide” or NP₁₋₃₀:

¹NPMYN AVSNADLMDF KNLLDHLEEK MPLED³⁰. (SEQ ID NO: 1)

6

SEQ ID NO:2 is the amino acid sequence of human “vessel dilator” or NP₃₁₋₆₇:

³¹EVVPP QVLSEPNEEA GAALSPLPEV PPWTGEVSPA QR⁶⁷. (SEQ ID NO: 2)

SEQ ID NO:3 is the amino acid sequence of human “kali-uretic peptide” or NP₇₉₋₉₈:

⁷⁹SSDRSAL LKSKLRALLT APR⁹⁸. (SEQ ID NO: 3)

SEQ ID NO:4 is the amino acid sequence of human “atrial natriuretic peptide” (ANP) or NP₉₉₋₁₂₆:

⁹⁹SLRRSSC FGGRMDRIGA QSGLCNSFR Y¹²⁶. (SEQ ID NO: 4)

SEQ ID NO:5 is the amino acid sequence of cloned mouse pNP₇₃₋₁₀₂:

⁷³GSPWDPSDRS ALLKSKLRAL LAGPRSLRRS¹⁰². (SEQ ID NO: 5)

SEQ ID NO:6 is the amino acid sequence of cloned mouse NP fragment:

^{VSNTDLMDFK NLLDHLEEK PVEDEVMPQ ALSEQTE}. (SEQ ID NO: 6)

SEQ ID NO:7 is the amino acid sequence for the human preproANP (NCBI ACCESSION # NM_006172) wherein the underlined amino acids represent the signal sequence which is cleaved off to form the mature peptide:

¹MSSRSTTVS FLLLAFOLL GOTRANPMYN AVSNADLMDF
KNLLDHLEEK MPLEDEVVPP QVLSEPNEEA GAALSPLPEV
PPWTGEVSPA QRDGGALGRG PWDSSDRSAL LKSKLRALLT
APRSLRRSSC FGGRMDRIGA QSGLCNSFR Y¹⁵¹. (SEQ ID NO: 7)

SEQ ID NO:8 is a forward primer for the cDNA sequence encoding mouse prepro ANF protein:

^{5'-gac ggc aag ctt act atg ggc agc ccc tgg}
^{gac cc-3'.} (SEQ ID NO: 8)

SEQ ID NO:9 is a reverse primer for the cDNA sequence encoding mouse pre-proANF protein:

^{5'-acc ccc ctc gag tta tta tct tcg tag gct ccg-3'.} (SEQ ID NO: 9)

SEQ ID NO:10 is a forward primer for the cDNA sequence encoding mouse NP fragment:

^{5'-aat cct aag ctt agt atg gtc tcc aac aca gat-3'.} (SEQ ID NO: 10)

US 8,148,114 B2

7

SEQ ID NO:11 is a reverse primer for the cDNA sequence encoding mouse NP fragment:

(SEQ ID NO: 11)
5'-tgc gaa ctc gag tta ctc agt ctg ctc act cag
ggc ctg cg-3'.

SEQ ID NO:12 is the nucleotide sequence encoding cloned mouse pNP₇₃₋₁₀₂:

(SEQ ID NO: 12)
ggc agc ccc tgg gac ccc tcc gat aga tct gcc ctc
ttg aaa agc aaa ctg agg gct ctg ctc gct ggc
cct cgg agc cta cga aga tcc.

8

SEQ ID NO:13 is the nucleotide sequence encoding cloned mouse pNP fragment:

(SEQ ID NO: 13)
5 atg gtg tcc aac aca gat ctg atg gat ttc aag aac
ctg cta gac cac ctg gag gag aag atg ccg gta gaa
10 gat gag gtc atg ccc ccg cag gcc ctg agt gag cag
act gag taa.

SEQ ID NO:14 is the mRNA nucleotide sequence encoding human ANP (NCBI Accession # NM_006172:

(SEQ ID NO: 14)
1 tggcgaggga cagacgtagg ccaagagagg ggaaccagag aggaaccaga ggggagagac
61 agagcagcaa gcagtggatt gctccttgac gacgccagca tgagctcctt ctccaccacc
121 accgtgagct tctccttttt actggcattc cagctcctag gtcagaccag agctaatecc
181 atgtacaatg ccgtgtccaa cgcagacctg atggatttca agaatttctt ggaccatttg
241 gaagaaaaga tgccctttaga agatgaggtc gtgccccac aagtgtctag tgagccgaat
301 gaagaagcgg gggctgtctt cagccccctc cctgaggtgc ctccctggac cggggaagtc
361 agcccagccc agagagatgg aggtgccctc gggcggggcc cctgggactc ctctgatcga
421 tctgccctcc taaaaagcaa gctgagggcg ctgctcactg cccctcggag cctgcggaga
481 tccagctgct tcgggggcag gatggacagg attggagccc agagcggact gggctgtaac
541 agcttccggt actgaagata acagccaggg aggacaagca gggctgggcc tagggacaga
601 ctgcaagagg ctctgtgtcc ctggggtctc tgtgtcattt gtgtcatctt gttgccatgg
661 agttgtgate atcccatcta agctgcagct tctgtcaac acttctcaca tcttatgcta
721 actgtagata aagtggtttg atggtgactt cctgcctct cccaccccat gcattaaatt
781 ttaaggtaga acctcacctg ttactgaaag tggtttgaaa gtgaataaac ttcagcacca
841 tggac.

SEQ ID NO:15 is the human gene for atrial natriuretic factor propeptide (coding sequence includes-join (570 . . . 692, 815 . . . 1141, 2235 . . . 2240); sig. peptide=570 . . . 644; mat. peptide=join (645 . . . 692, 815 . . . 1141, 2235 . . . 2237), (NCBI ACCESSION NO: X01471; Greenberg, B. D. et al., *Nature*, 1984, 312(5995):656-658):

(SEQ ID NO: 15)
1 ggatccattt gtctegggct gctggtgcc tgccatttcc tctctccac ccttatttgg
61 aggccttgac agctgagcca caaacaacc aggggagctg ggcaccagca agcgtcaccc
121 tctgtttccc cgcacggtag cagcgtcgag gagaaagaat cctgaggcac ggcggtgaga
181 taaccaagga ctctttttta ctcttctcac acctttgaag tgggagcctc ttgagtcaaa
241 tcagtaagaa tgccgctctt gcagctgagg gtctgggggg ctgttggggc tgcccaaggc
301 agagaggggc tgtgacaagc cctgcggatg ataactttaa aagggcatct cctgctggct
361 tctcacttgg cagctttatc actgcaagtg acagaatggg gagggttctg tctctcctgc
421 gtgcttgagg agctgggggg ctataaaaag aggcggcact gggcagctgg gagacaggga
481 cagacgtagg ccaagagagg ggaaccagag aggaaccaga ggggagagac agagcagcaa

-continued

541 gcagtgagatt gctccttgac gacgccagca tgagctcctt ctccaccacc accgtgagct
601 tctcctttt actggcattc cagctcctag gtcagaccag agctaataccc atgtacaatg
661 ccgtgtccaa cgcagacctg atggatttca aggtagggcc aggaaagcgg gtgcagtctg
721 gggccagggg gctttctgat gctgtgctca ctctcttga tttcctccaa gtcagtgagg
781 tttatccctt tcctgtatt ttcttttct aaagaatttg ctggaccatt tggaagaaaa
841 gatgccttta gaagatgagg tcgtgcccc acaagtgtct agtgagccga atgaagaagc
901 gggggctgct ctcagccccc tcctgaggt gcctccctgg accggggaag tcagcccagc
961 ccagagagat ggaggtgcc tcgggggggg cccctgggac tcctctgate gatctgccct
1021 cctaaaaaagc aagctgaggg cgctgctcac tgcccctcgg agcctgcgga gatccagctg
1081 cttcgggggc aggatggaca ggattggagc ccagagcggga ctgggctgta acagcttcgg
1141 ggtaagagga actggggatg gaaatgggat gggatggaca ctactgggag acaccttcag
1201 caggaaaggg accaatgcag aagctcattc cctctcaagt ttctgcccc acaccagag
1261 tgccccatgg gtgtcaggac atgccatcta ttgtccttag ctagtctgct gagaaaatgc
1321 ttaaaaaaaa aagggggggg gctgggcacg gtcgtcacgc ctgtaataccc agcacttttg
1381 gaggccaggc agcggatcat gaggtcaaga gatcaagact atcctggcca acatggtgaa
1441 accccagctc tactaaaaat acaaaaatta gctgggtgtg tggcgggcac ctgtactctc
1501 agctacttgg gaggtgagg caggagaatc acttgaaccc aggaggcaga ggttcagtg
1561 agcagagatc acgccactgc agtccagcct aggtgataga gcgagactgt ctcaaaaaa
1621 aaaaaaaaag gccaggcgcg gtggctcacg cctgtaatcc cagcgctttg ggaggccaag
1681 gcgggtgat cagcaggtca ggagatggag accatcctgg ctaacacggt gaaacccgt
1741 ctctactaaa aatacaaaaa attagccagg cgtggtggca ggcgctgta agtcctagct
1801 actccggagg ctgaggcagg agaattggct gaacccggga ggaggagctt gcagtgagca
1861 gagatggcac cactgcactc cagcctgggc gacagagcaa gactccgtct caaaaaaaa
1921 aaaaaaaa gcaactgcca ctagcactgg gaaattaaaa tattcataga gccaagttat
1981 ctttgcattg ctgattagca gttcatattc ctcccagaa ttgcaagatc ctgaagggt
2041 taagtgaat ttactctgat gagtaacttg cttatcaatt catgaagctc agagggtcat
2101 caggctgggg tgggggccgg tgggaagcag gtggtcagta atcaagtcca gaggatgggc
2161 acactcatac atgaagctga cttttccagg acagccaggt caccaagcca gatatgtctg
2221 tgttctcttt gcagtactga agataacagc caggaggagc aagcagggtt gggcctaggg
2281 acagactgca agaggtcctt gtcctcggg gtctctgctg catttggtgc atcttggtgc
2341 catggagttg tgatcatccc atctaagctg cagcttccctg tcaaacacttc tcacatctta
2401 tgctaactgt agataaagtg gtttgatggt gacttctcgc cctctccac cccatgcatt
2461 aaattttaag gtagaacctc acctgttact gaaagtgtt tgaaagtga taaacttcag
2521 caccatggac agaagacaaa tgctgcgtt ggtgtgcttt ctttctctt gggaagagaa
2581 ttc.

SEQ ID NO:16 is the amino acid sequence for the mouse
preproANP peptide:

(SEQ ID NO: 16)
MGFSITLGF FLVLAFWLPG HIGANPVYSA VSNTDLMDFK

NLLDHLEEKM PVEDEVMPQ ALSEQTEEAG AALSSLPEVP

-continued

60 PWTGEVNPPL RDGSALGRSP WDPSDRSALL KSKLRALLAG

PRSLRRSSCF GGRIDRIGAQ SGLGCNSFRY RR.

65 SEQ ID NO: 17 is the genetic sequence for the mouse pre-
proANP peptide wherein the coding sequence starts at
nucleic acid molecule position 81 and ends at nucleic acid
molecule position 539:

(SEQ ID NO: 17)

```

1  caaaagctga gagagagaga gaaagaaacc agagtgggca gagacagcaa acatcagatc
61  gtgccccgac ccacgccagc atgggctcct tctccatcac cctgggcttc ttcctcgtct
121  tggccttttg gcttccaggc catattggag caaatcctgt gtacagtgcg gtgtccaaca
181  cagatctgat ggatttcaag aacctgctag accacctgga ggagaagatg ccggtagaag
241  atgaggtcat gccccgcagc gccctgagtg agcagactga ggaagcaggg gccgcactta
301  gctccctccc cgaggtgcct ccttgactg gggaggtcaa cccacctctg agagacggca
361  gtgctctagg gcgcagcccc tgggacctct ccgatagatc tgcctctctg aaaagcaaac
421  tgagggtctt gctcgtggc cctcggagcc tacgaagatc cagctgcttc ggggtagga
481  ttgacaggat tggagcccag agtggactag gctgcaacag cttccggtac cgaagataac
541  agccaaggag gaaaaggcag tcgattctgc ttgagcagat cgcaaaagat cctaagccct
601  tgtggtgtgt caccgacgtt ggtcacattg ccactgtggc gtggtgaaca cctcctgga
661  gctgcggtct cctgccttca tctatcacga tcgatgttaa atgtagatga gtggtctagt
721  ggggtcttgc ctctccact ctgcatatta aggtagatcc tcacctttt cagaaagcag
781  ttggaaaaaa aaaaaagaa taaacttcag caccaaggac agacgccgag gccctgatgt
841  gcttcttttg cttctgcct cagttctttg ctctcccc.

```

SEQ ID NO:18 is amino acid sequence of human natriuretic peptide receptor-A (NPR-A):

(SEQ ID NO: 18)

```

MPGPRRPAGSRLRLLLLLLPLLLLLLRGSHAGNLTAVVVLPLANTSYFW
SWARVGPAPVELALAQVKARPDLLPGWTVRTVLGSSSENALGVCSDTAAPLA
AVDLKWEHNPVFLGPGCVYAAAPVGRFTAHRVPLLTAGAPALGFQVKD
EYALTTRAGPSYAKLGFVAALHRRLGWERQALMLYAYRPGDEEHCFVLV
EGLFMRVRDLNITVDHLEFAEDDLSHYTRLLRTMPRKGRVYIICSSPDA
FRTLMLLALAEAGLCGEDYVFFHLDFGQSLQGGQGPAPRRPWERGDGQDV
SARQAFQAAKIIITYKDPDNPEYLEFLKQLKHLAYEQFNFTMEDVLVNTIP
ASFHDGLLLYIQAVTETLAHGTVTDGENITQRMWNRSFQGTGYLKIDS
SGDRETDPSLWMDPENGAFRVVLNYNGTSQELVAVSGRKLNWPLGYPPP
DIPKCGFDNEDPACNQDHLSTLEVLALVGSLSLLGILIVSFFIYRKMLE
KELASELWRVRWEDVEPSSLERHLRSAGSRLTSLGRGSNYGSLLTTEGQF
QVFAKTAYYKGNLVAVKRVNRKRIELTRKVLFEKHMARDVQNEHLTRFVG
ACTDPNIIICILTEYCPRGSLQDILENESITLDWMFRYSLTNDIVKGMFLF

```

- continued

```

HNGAICSHGNLKSSNCVVDGRFVLKITYGLESFRDLDPQGHQHTVYAKKL
30  WTAPELLRMASPPVRGSQAGDVYSFGIILQEIALRSGVFHVEGLDLSPE
IIERVTRGEQPPFRPSLALQSHLEELGLLMQRCWAEDPQERPPFQQIRLT
LRKFNRENSSNILDNLLSRMEQYANNLEELVEERTQAYLEEKRAEALLY
35  QILPHSVAEQKRGETVQAEAFDSVTIYFSDIVGTALSASTPMQVVTL
LNDLYTCFDAVIDNFDVYKVETIGDAYMVVSGLPVRNRLHACEVARMAL
ALLDAVRSFRIRHRPQEQRLRLRIGIHTGPVCAGVGLKMPRYCLFGDVTN
40  TASRMESNGEALKIHLSETKAVLEEFGGFELELRGDVEMKKGKVRTYV
LLGERGSSTRG.

```

(NCBI ACCESSION NO. NM_000906; Airhart N. et al. *J. Biol. Chem.*, 2003, 278(40):38693-38698; Pitzalis M. V. et al., *J. Hypertens.*, 2003, 21(8):1491-1496; Mokentin J. D. *J. Clin. Invest.*, 2003, 111(9):1275-1277; De L. et al., *J. Biol. Chem.*, 2003, 278(13):11159-11166; Knowles J. W. et al., *Hum. Genet.* 2003, 12(1):62-70; Pandey K. N. et al., *J. Biol. Chem.* 2002, 277(7):4618-4627).

SEQ ID NO:19 is the nucleotide coding sequence for human natriuretic peptide receptor-A (NPR-A):

(SEQ ID NO: 19)

```

ggttccctcc ggatagccgg agacttgggc cggccggacg ccccttctgg cacactccct
61  ggggcaggcg ctacgcacg ctacaaacac acactcctct ttctccctc gcgcgcctc
121  tctcctctt cttcacgaag cgctcactcg caccctttct ctctctctct ctctctctaa
181  caccgcacga cactccagtg tgttcacact cgggtcctct ccagcccgac gttctcctgg
241  caccacactg ctccgcggcg cctcgcgcg cccctcgggt cgcgccctt gcgctctcgg
301  ccagaccgt cgcagctaca gggggcctcg agccccggg tgagcgtccc cgtcccgtc
361  ctgctccttc ccatagggac gcgcctgatg cctgggaccg gccgctgagc ccaaggggac

```

-continued

421 cgaggaggcc atggtaggag cgctcgctg ctgcgggtgcc cgtgaggcc atgccggggc
 481 cccggcgccc cgtggtctcc cgcctgcgcc tgcctctgct cctgctgctg ccgcctgctg
 541 tgctgctgct ccggggcagc caccggggca acctgacggt agccgtggta ctgccgctgg
 601 ccaatacctc gtaccctctg tcgtggggcg gcgtggggacc cgcctgggag ctggccctgg
 661 cccaggtgaa ggcgcgcccc gacttgctgc cgggctggac ggtccgcacg gtgctgggca
 721 gcagcgaaaa cgcgctgggc gtctgctccg acaccgcagc gcccctggcc gcggtggacc
 781 tcaagtggga gcacaacccc gctgtgttcc tgggccccgg ctgcgtgtac gccgcccccc
 841 cagtggggcg cttcacgcgc cactggcggg tcccgtgctg gaccgcccgc gccccggcgc
 901 tgggcttcgg tgtcaaggac gagtatgcgc tgaccacccg cgcggggccc agctacgcca
 961 agctggggga cttcgtggcg gcgctgcacc gacggctggg ctgggagcgc caagcgctca
 1021 tgctctacgc ctaccggcgc ggtgacgaag agcactgctt cttcctctg gaggggctgt
 1081 tcatgcgggt ccgcgaccgc ctcaatatta cggtggaaca cctggagttc gccgaggacg
 1141 acctcagcca ctacaccagg ctgctgcgga ccatgccgcg caaaggccga gttatctaca
 1201 tctgcagetc ccctgatgcc ttcagaaccc tcctgctcct ggccctggaa gctggttgt
 1261 gtggggagga ctacgttttc ttccacctgg atatctttgg gcaaagcctg caaggtggac
 1321 agggccctgc tccccgcagg ccctgggaga gaggggatgg gcaggatgtc agtcccgcgc
 1381 aggcctttca ggctgccaaa atcattacat ataaagaccc agataatccc gactacttgg
 1441 aattcctgaa gcagttaaaa cacctggcct atgagcagtt caacttcacc atggaggatg
 1501 tcctggtgaa caccatccca gcctccttcc acgacgggct cctgctctat atccaggcag
 1561 tgacggagac tctggcacat gggggaactg ttactgatgg ggagaacatc actcagcgga
 1621 tgtggaaccc aagctttcaa ggtgtgacag gatacctgaa aattgatagc agtggcgatc
 1681 gggaaacaga cttctccctc tgggatatgg atcccagaa tgggtgcctc agggttgtac
 1741 tgaactacaa tgggacttcc caagagctgg tggctgtgtc ggggcgcaaa ctgaactggc
 1801 ccctggggta ccctcctcct gacatcccca aatgtggcct tgacaacgaa gaccagcat
 1861 gcaaccaaga tcacctttcc accctggagg tgctggcttt ggtgggcagc ctctccttgc
 1921 tcggcattct gattgtctcc ttcttcatat acaggaagat gcagctggag aaggaaactgg
 1981 cctcggaact gtggcggggt cgctgggagg acgttgagcc cagtagcctt gagaggcacc
 2041 tgcggagtgc aggcagccgg ctgacccctga gcgggagagg ctccaattac ggctccctgc
 2101 taaccacaga gggccagttc caagtctttg ccaagacagc atattataag ggcaacctcg
 2161 tggctgtgaa acgtgtgaac cgtaaacgca ttgagctgac acgaaaagtc ctgtttgaac
 2221 tgaagcatat gcgggatgtg cagaatgaac acctgaccag gtttgtggga gcttcacccg
 2281 acccccccaa tatctgcac ctcacagagt actgtccccg tgggagcctg caggacattc
 2341 tggagaatga gacatcacc ctggactgga tgttcggta ctcactcacc aatgacatcg
 2401 tcaagggcat gctgtttcta cacaatgggg ctatctgttc ccatgggaac ctcaagtcac
 2461 ccaactgcgt ggtagatggg cgctttgtgc tcaagatcac cgactatggg ctggagagct
 2521 tcagggacct ggaccagag caaggacaca ccgtttatgc caaaaagctg tggacggccc
 2581 ctgagctcct gcgaatggct tcaccccctg tgcggggctc ccaggctggg gacgtataca
 2641 gctttgggat catccttcag gagattgccc tgaggagtgg ggtcttcac gtggaaggtt
 2701 tggacctgag ccccaagag atcatcgagc gggtgactcg gggtagagc ccccccttcc
 2761 ggccctccct ggccctgcag agtcacctgg aggagtggg gctgctcatg cagcgggtgct
 2821 gggctgagga cccacaggag aggccacat tccagcagat ccgcctgacg ttgcgcaaat

-continued

2881 ttaacagggga gaacagcagc aacatcctgg acaacctgct gtcccgcagt gagcagtagc
 2941 cgaacaatct ggaggaaactg gtggaggagc ggaccaggc atacctggag gagaagcgca
 3001 aggtcgaggc cctgctctac cagatcctgc ctactcagt ggctgagcag ctgaagcgtg
 3061 ggggagcgtg gcaggccgaa gcctttgaca gtgttaccat ctacttcagt gacattgtgg
 3121 gtttcacagc gctgtcggcg gagagcagc ccatgcaggt ggtgacctg ctcaatgacc
 3181 tgtacacttg ctttgatgct gtcatagaca actttgatgt gtacaagggt gagacaattg
 3241 gcgatgccta catggtggtg tcagggtcc ctgtgcggaa cgggcggcta cagcctgcg
 3301 aggtagcccg catggccctg gcaactgctg atgctgtgcg ctccctccga atccgccacc
 3361 ggccccagga gcagctgcgc ttgctcattg gcaccacac aggacctgtg tgtgctggag
 3421 tgggtgggact gaagatgccc cgttactgtc tctttgggga tacagtcaac acagcctcaa
 3481 gaatggagtc taatggggaa gccctgaaga tccactgtc ttctgagacc aaggctgtcc
 3541 tggaggagtt tgggtggttc gagctggagc ttcgaggga tgtagaaatg aagggcaaag
 3601 gcaaggttcg gacctactgg ctccctgggg agagggggag tagcaccga ggctgacctg
 3661 cctcctctcc tatccctcca caccctccct accctgtgcc agaagcaaca gaggtgccag
 3721 gcctcagcct caccacagc agcccatcg ccaaaggatg gaagtaattt gaatagctca
 3781 ggtgtgctta cccagtgaa gacaccagat aggacctctg agaggggact ggcatggggg
 3841 gatctcagag cttacaggct gagccaagcc caggccatg cacagggaca ctcacacagg
 3901 cacacgcacc tgctctccac ctggactcag gccgggctgg gctgtggatt cctgateccc
 3961 tccccctccc atgctctcct ccctcagcct tgetacctg tgacttactg ggaggagaaa
 4021 gagtccactg aaggggaaca tgaaaagaga ctaggatgaag agagggcagg ggagccaca
 4081 tctggggctg gccacaata cctgctccc cgacccctc caccagcag tagacacagt
 4141 gcacagggga gaagaggggt ggccgagaag ggttgggggc ctgtatgcct tgctctacc
 4201 atgagcagag acaattaaaa tctttattcc aaaaaaaaa aaaaaa

(NCBI ACCESSION NO. NM_000906; Airhart N. et al., *J. Biol. Chem.*, 2003, 278(40):38693-38698; Pitzalis M. V. et al., *J. Hypertens.*, 2003, 21(8):1491-1496; Mokentin J. D. *J. Clin. Invest.*, 2003, 111(9):1275-1277; De L. et al., *J. Biol. Chem.*, 2003, 278(13):11159-11166; Knowles J. W. et al., *Hum. Genet.*, 2003, 12(1):62-70; Pandey K. N. et al. *J. Biol. Chem.*, 2002, 277(7):4618-4627).

SEQ ID NO:20 is amino acid sequence of the human atrial natriuretic peptide clearance receptor precursor (ANP-C; also referred to as NPR-C, NPRC, and atrial natriuretic peptide C-type receptor):

(SEQ ID NO: 20)

MPSLLVLTFS PCVLLGWALL AGTGGGGVG GGGGAGIGG
 GRQEREALPP QKIEVLVLLP QDDSYLFSLT RVRPAIEYAL
 RSVEGNGTGR RLLPPGTRFQ VAYEDSDCGN RALFSLVDRV
 AAARGAKPDL ILGPVCEYAA APVARLASHW DLPMLSAGAL
 AAGFQHKDSE YSHLTRVAPA YAKMGEMMLA LFRHHWSRA
 ALVYSDDKLE RNCYFTLEGV HEVFQEEGLH TSIYSFDETK
 DLDLEDIVRN IQASERVVIM CASSDTIRSI MLVAHRHGMT
 SGDYAFFNIE LFNSSSYGDG SWKRGDKHDF EAKQAYSSLQ
 TVTLLRTPVK EFEKFSMEVK SSVEKQGLNM EDVNMFVEG

-continued

FHDAILYVL ALHEVLRAGY SKKDGKIIQ QTWNRTFEGI
 AGQVSIDANG DRYGDFSVIA MTDVEAGTQE VIGDYFGKEG
 RFEMRPNVKY PWGPKLRID ENRIVEHTNS SPCKSSGGLE
 ESAVTGIVVG ALLGAGLLMA FYFRKKYRI TIERRTQEE
 SNLGKHREL R EDSIRSHFSV A

(NCBI ACCESSION NO. P17342; Lowe D. G. et al., *Nucleic Acids Res.*, 1990, 18(11):3412; Porter J. G. et al., *Biochem. Biophys. Res. Commun.*, 1990, 171(2):796-803; Stults J. T. et al., *Biochemistry*, 1994, 33(37):11372-11381).

SEQ ID NO: 21 is an siRNA specific for NPR-A (human).
 tat tac ggt gga cca cct gtt caa gag aca ggt ggt
 cca ccg taa tat ttttt
 SEQ ID NO: 22 is an siRNA specific for NPR-A (human).
 aga att cca gaa acg cag ctt caa gag agc tgc gtt
 tct gga att ctt ttttt

DETAILED DISCLOSURE

The present invention pertains to methods for reducing natriuretic peptide receptor-A (also known in the art as

NPRA, NPR-A, and guanylate cyclase A) activity in vitro or in vivo. In one aspect, the method of the invention may be used for treating inflammatory and cell proliferation disorders, such as cancer. In another aspect, the present invention concerns methods for identifying agents useful for treating inflammatory and cell proliferation disorders by determining whether the candidate agent reduces activity of the natriuretic peptide receptor-A (also known in the art as NPRA, NPR-A, and guanylate cyclase A) in vitro and/or in vivo (also referred to herein as the diagnostic method or assay of the invention).

As used herein, an "inflammatory disorder" includes those conditions characterized by an aberrant increase in one or more of the following: IL-6, IL-1 beta, TNF-alpha, IL-8, eosinophil production, neutrophil production, release of histamines, proliferants, hyperplasia, and cell adhesion molecule expression. As used herein, a "cell proliferation disorder" is characterized by one or more of the following: uncontrolled proliferation, a high mitogenic index, over-expression of cyclin D1, cyclin B1, expression of an oncogene such as c-jun and/or c-fos, aberrant activation of NFkB and/or ERK (extracellular receptor kinase), and matrix metalloproteinase expression (such as MMP-2 and/or MMP-9).

In one embodiment, the inflammatory disorder and cell proliferation disorder is not one that is amenable to effective treatment by administration of a vasodilator. In one embodiment, the inflammatory disorder and cell proliferation disorder is not a cardiovascular disorder (such as hypertension or stroke). In another embodiment, the inflammatory disorder and cell proliferation disorder is not a disorder of the central nervous system (such as Alzheimer's disease or other dementia). In another embodiment, the inflammatory disorder and cell proliferation disorder is not kidney failure or other kidney disorder.

The agent used to reduce NPR-A activity in vitro or in vivo can be virtually any substance and can encompass numerous chemical classes, including organic compounds or inorganic compounds. Preferably, an effective amount of the agent is administered to the cells with a pharmaceutically acceptable carrier. The agent may be a substance such as genetic material, protein, lipid, carbohydrate, small molecules, a combination of any of two or more of foregoing, or other compositions. The agent may be naturally occurring or synthetic, and may be a single substance or a mixture. The agent can be obtained from a wide variety of sources including libraries of compounds. The agent can be or include, for example, a polypeptide, peptidomimetic, amino acid(s), amino acid analog(s), function-blocking antibody, polynucleotide(s), polynucleotide analog(s), nucleotide(s), nucleotide analog(s), or other small molecule(s). A polynucleotide may encode a polypeptide that potentially reduces NPR-A activity within the cell, or the polynucleotide may be a short interfering RNA (siRNA), a hairpin RNA (shRNA), antisense oligonucleotide, ribozyme, or other polynucleotide that targets an endogenous or exogenous gene for silencing of gene expression and potentially NPR-A activity within the cell.

In one embodiment, the agent used to reduce NPR-A activity is an interfering RNA specific for NPR-A mRNA, preferably human NPR-A mRNA. Interfering RNA is capable of hybridizing with the mRNA of a target gene and reduce and/or eliminate translation through the mechanism of RNA interference. Examples of such interfering RNA include SEQ ID NO:21 and SEQ ID NO:22, which were determined to have a relatively high probability of reducing NPR-A activity using an siRNA Target Finder program (AMBIION) and in accordance with published guidelines (Tuschl T., *Nature Biotechnol.*, 2002, 20:446448). As used herein, the term "RNA interference" ("RNAi") refers to a selective intracellular deg-

radation of RNA. RNAi occurs in cells naturally to remove foreign RNAs (e.g., viral RNAs). Natural RNAi proceeds via fragments cleaved from free dsRNA which direct the degradative mechanism to other similar RNA sequences. Alternatively, RNAi can be initiated by the hand of man, for example, to silence the expression of target genes.

As used herein, the term "small interfering RNA" ("siRNA") (also referred to in the art as "short interfering RNAs") refers to an RNA (or RNA analog) that is capable of directing or mediating RNA interference. In one embodiment, the siRNA is between about 10-50 nucleotides (or nucleotide analogs). Optionally, a polynucleotide (e.g., DNA) encoding the siRNA may be administered to cells in vitro or in vivo, such as in a vector, wherein the DNA is transcribed.

As used herein, a siRNA having a "sequence sufficiently complementary to a target mRNA sequence to direct target-specific RNA interference (RNAi)" means that the siRNA has a sequence sufficient to trigger the destruction of the target mRNA by the RNAi machinery or process. "mRNA" or "messenger RNA" or "transcript" is single-stranded RNA that specifies the amino acid sequence of one or more polypeptides. This information is translated during protein synthesis when ribosomes bind to the mRNA.

The scientific literature is replete with reports of endogenous and exogenous gene expression silencing using siRNA, highlighting their therapeutic potential (Gupta, S. et al. *PNAS*, 2004, 101:1927-1932; Takaku, H. *Antivir Chem. Chemother*, 2004, 15:57-65; Pardridge, W. M. *Expert Opin. Biol. Ther.*, 2004, 4:1103-1113; Zheng, B. J. *Antivir. Ther.*, 2004, 9:365-374; Shen, W. G. *Chin., Med. J. (Engl)*, 2004, 117:1084-1091; Fuchs, U. et al. *Curr. Mol. Med.*, 2004, 4:507-517; Wadhwa, R. et al. *Mutat. Res.*, 2004, 567:71-84; Ichim, T. E. et al. *Am. J. Transplant*, 2004, 4:1227-1236; Jana, S. et al. *Appl. Microbiol. Biotechnol.*, 2004, 65:649-657; Ryther, R. C. et al. *Gene Ther.*, 2005, 12:5-11; Chae, S-S. et al., *J. Clin. Invest.*, 2004, 114:1082-1089; Fougerolles, A. et al., *Methods Enzymol.*, 2005, 392:278-296), each of which is incorporated herein by reference in its entirety. Therapeutic silencing of endogenous genes by systemic administration of siRNAs has been described in the literature (Kim B. et al., *American Journal of Pathology*, 2004, 165:2177-2185; Soutschek J. et al., *Nature*, 2004, 432:173-178; Pardridge W. M., *Expert Opin. Biol. Ther.*, 2004, July, 4(7):1103-1113), each of which is incorporated herein by reference in its entirety.

In another embodiment, the decrease in NPR-A activity (e.g., a reduction in NPR-A expression) may be achieved by administering an analogue of ANP (e.g., ANP4-23) or non-peptide antagonists (e.g., HS-142-1; Rutherford et al., *Br. J. Pharmacol.*, 1994, 113:931-939; El-Ayoubi et al., *Br. J. Pharmacol.*, 2005, Feb. 7, Epub ahead of print; Delpont C. et al., *Eur. J. Pharmacol.*, 1992, 224(2-3):183-188; Ohyama Y. et al., *Biochem. Biophys. Res. Commun.*, 1992, 189(1):336-342). In another embodiment, the agent is an anti-human NPR-A function-blocking antibody (preferably, humanized), or soluble NPR-A or NPR-C (as a receptor decoy). Other examples of agents include NPR-A antagonists that specifically inhibit cGMP-dependent protein kinase (PKG) such as A71915 and KT5823 (Pandey K. N. et al., *Biochemical and Biophysical Research Communications*, 2000, 271:374-379).

The methods of the invention may include further steps. In some embodiments, a subject with the relevant inflammatory disorder and/or cell proliferation disorder is identified or a patient at risk for the disorder is identified. A patient may be someone who has not been diagnosed with the disease or condition (diagnosis, prognosis, and/or staging) or someone

diagnosed with disease or condition (diagnosis, prognosis, monitoring, and/or staging), including someone treated for the disease or condition (prognosis, staging, and/or monitoring). Alternatively, the person may not have been diagnosed with the disease or condition but suspected of having the disease or condition based either on patient history or family history, or the exhibition or observation of characteristic symptoms.

In one aspect, the therapeutic method of the invention involves administering a natriuretic hormone peptide (NP), or a fragment, homolog or variant thereof, or a nucleic acid sequence encoding an NP, or a fragment, homolog, or variant thereof, to a patient. The present inventor has demonstrated that a prolonged, substantial reduction of tumor burden in lungs can be achieved by intranasal delivery of pDNA-encoding a peptide comprising amino acid residues 73 to 102 (NP73-102). Without being bound by theory, the NP decreased viability due to the induction of apoptosis in a lung adenocarcinoma cell line A549 cell, and can reduce tumorigenesis and metastasis in a number of cancers.

In specific embodiments, the peptides used in the subject invention comprise at least one amino acid sequence selected from the group consisting of NP₁₋₃₀, NP₃₁₋₆₇, NP₇₉₋₉₈, and NP₇₃₋₁₀₂, (SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, and SEQ ID NO:5, respectively), SEQ ID NO:6, or a biologically active fragment or homolog thereof. In some embodiments, a combination of NP or NP-encoding nucleic acid sequences is utilized. In one embodiment, the peptide utilized does not consist of the amino acid sequence of NP₉₉₋₁₂₆ (SEQ ID NO:4).

In another aspect, the therapeutic method of the invention involves administering an agent that reduces activity of the natriuretic peptide receptor-A (also known in the art as NPR-A, NPR-A, and guanylate cyclase A) to a patient, wherein the agent is administered in an amount effective to reduce receptor (NPR-A) activity. NPR-A activity can be determined, for example, by one or more of the following biological parameters: production/accumulation of cGMP, expression of the NPR-A (transcription or translation), and/or cellular internalization of the NPR-A.

According to the gene therapy method of the present invention, the NP-encoding nucleic acid sequence is administered locally at the target site (e.g., at the site of cancer or pre-cancer), or systemically to the patient. In order to treat cancer of the lung, for example, the NP-encoding nucleic acid sequence is preferably administered to the airways of the patient, e.g., nose, sinus, throat and lung, for example, as nose drops, by nebulization, vaporization, or other methods known in the art. More preferably, the nucleic acid sequence encoding NP is administered to the patient orally or intranasally, or otherwise intratracheally. For example, the nucleic acid sequence can be inhaled by the patient through the oral or intranasal routes, or injected directly into tracheal or bronchial tissue.

In specific embodiments, the nucleic acid sequences used in the subject invention encode at least one amino acid sequence selected from the group consisting of NP₁₋₃₀, NP₃₁₋₆₇, NP₇₉₋₉₈, NP₉₉₋₁₂₆, and NP₇₃₋₁₀₂, (SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5, respectively), SEQ ID NO:6, or a biologically active fragment or homolog of any of the foregoing.

Preferably, the nucleic acid sequence encoding the NP is administered with a nucleic acid sequence that is operatively linked with the NP-encoding nucleic acid sequence and operates as a regulatory sequence. For example, the regulatory sequence can be a promoter sequence that controls transcription and drives expression of the NP-encoding nucleic acid

sequence at the desired site, such as at, or adjacent to, the patient's respiratory epithelial cells. The promoter can be a constitutive or inducible promoter to allow selective transcription. The promoter can be a vertebrate or viral promoter. Optionally, enhancers may be used to obtain desired transcription levels. An enhancer is generally any non-translated nucleic acid sequence that works contiguously with the coding sequence (in cis) to change the basal transcription level dictated by the promoter.

The NP-encoding nucleic acid sequences used in the methods, expression vectors, and pharmaceutical compositions of the present invention are preferably isolated. According to the present invention, an isolated nucleic acid molecule or nucleic acid sequence, is a nucleic acid molecule or sequence that has been removed from its natural milieu. As such, "isolated" does not necessarily reflect the extent to which the nucleic acid molecule has been purified. An isolated nucleic acid molecule or sequence useful in the present composition can include DNA, RNA, or any derivatives of either DNA or RNA. An isolated nucleic acid molecule or sequence can be double stranded (i.e., containing both a coding strand and a complementary strand) or single stranded.

A nucleic acid molecule can be isolated from a natural source, or it can be produced using recombinant DNA technology (e.g., polymerase chain reaction (PCR) amplification, cloning) or chemical synthesis. Nucleic acid molecules can be generated or modified using a variety of techniques including, but not limited to, classic mutagenesis techniques and recombinant DNA techniques, such as site-directed mutagenesis, chemical treatment of a nucleic acid molecule to induce mutations, restriction enzyme cleavage of a nucleic acid fragment, ligation of nucleic acid fragments, polymerase chain reaction (PCR) amplification and/or mutagenesis of selected regions of a nucleic acid sequence, synthesis of oligonucleotide mixtures and ligation of mixture groups to "build" a mixture of nucleic acid molecules, and combinations thereof.

Although the phrase "nucleic acid molecule" primarily refers to the physical nucleic acid molecule and the phrase "nucleic acid sequence" primarily refers to the sequence of nucleotides on the nucleic acid molecule, the two phrases are used interchangeably herein. As used herein, a "coding" nucleic acid sequence refers to a nucleic acid sequence that encodes at least a portion of a peptide or protein (e.g., a portion of an open reading frame), and can more particularly refer to a nucleic acid sequence encoding a peptide or protein which, when operatively linked to a transcription control sequence (e.g., a promoter sequence), can express the peptide or protein.

The nucleotide sequences encoding NP used in the subject invention include "homologous" or "modified" nucleotide sequences. Modified nucleic acid sequences will be understood to mean any nucleotide sequence obtained by mutagenesis according to techniques well known to persons skilled in the art, and exhibiting modifications in relation to the normal sequences. For example, mutations in the regulatory and/or promoter sequences for the expression of a polypeptide that result in a modification of the level of expression of a polypeptide according to the invention provide for a "modified nucleotide sequence". Likewise, substitutions, deletions, or additions of nucleic acids to the polynucleotides of the invention provide for "homologous" or "modified" nucleotide sequences. In various embodiments, "homologous" or "modified" nucleic acid sequences have substantially the same biological or serological activity as the native (naturally occurring) natriuretic peptide. A "homologous" or "modified" nucleotide sequence will also be understood to mean a

splice variant of the polynucleotides of the instant invention or any nucleotide sequence encoding a "modified polypeptide" as defined below.

A homologous nucleotide sequence, for the purposes of the present invention, encompasses a nucleotide sequence having a percentage identity with the bases of the nucleotide sequences of between at least (or at least about) 20.00% to 99.99% (inclusive).

The aforementioned range of percent identity is to be taken as including, and providing written description and support for, any fractional percentage, in intervals of 0.01%, between 20.00% and 99.99%. These percentages are purely statistical and differences between two nucleic acid sequences can be distributed randomly and over the entire sequence length.

In various embodiments, homologous sequences exhibiting a percentage identity with the bases of the nucleotide sequences of the present invention can have 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 percent identity with the polynucleotide sequences of the instant invention. Homologous nucleotide and amino acid sequences include mammalian homologs of the human NP sequences.

The NP homologs include peptides containing, as a primary amino acid sequence, all or part of an exemplified NP polypeptide sequence. The NP homologs thus include NP polypeptides having conservative substitutions, i.e., altered sequences in which functionally equivalent amino acid residues are substituted for residues within the sequence resulting in a peptide which is biologically active. For example, one or more amino acid residues within the sequence can be substituted by another amino acid of a similar polarity which acts as a functional equivalent, resulting in a silent alteration. In one aspect of the present invention, conservative substitutions for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs (see Table 1). Conservative substitutions also include substitutions by amino acids having chemically modified side chains which do not eliminate the biological activity of the resulting NP homolog.

TABLE 1

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

Both protein and nucleic acid sequence homologies may be evaluated using any of the variety of sequence comparison algorithms and programs known in the art. Such algorithms and programs include, but are by no means limited to, TBLASTN, BLASTP, FASTA, TFASTA, and CLUSTALW (Pearson and Lipman *Proc. Natl. Acad. Sci. USA*, 1988, 85(8):2444-2448; Altschul et al. *J. Mol. Biol.*, 1990, 215(3): 403-410; Thompson et al. *Nucleic Acids Res.*, 1994, 22(2): 4673-4680; Higgins et al. *Methods Enzymol.*, 1996, 266:383-402; Altschul et al. *J. Mol. Biol.*, 1990, 215(3):403-410; Altschul et al. *Nature Genetics*, 1993, 3:266-272).

Identity and similarity of related nucleic acid molecules and polypeptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; York (1988);

Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; York (1993); Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Jersey (1994); Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Press (1987); Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; York (1991); and Carillo et al., SIAM J. Applied Math., 48:1073 (1988).

The methods, pharmaceutical compositions, and vectors of the present invention can utilize biologically active fragments of nucleic acid sequences encoding the 126-amino acid atrial natriuretic factor (ANF) prohormone, such as nucleic acid sequences encoding NP₁₋₃₀, NP₃₁₋₆₇, NP₇₉₋₉₈, NP₉₉₋₁₂₆, and NP₇₃₋₁₀₂ (SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5, respectively), SEQ ID NO:6, and including biologically active fragments of the nucleic acid sequences encoding SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6.

Representative fragments of the nucleotide sequences according to the invention will be understood to mean any polynucleotide fragment having at least 8 or 9 consecutive nucleotides, preferably at least 12 consecutive nucleotides, and still more preferably at least 15 or at least 20 consecutive nucleotides of the sequence from which it is derived, with retention of biological activity as described herein. The upper limit for such fragments is one nucleotide less than the total number of nucleotides found in the full-length sequence (or, in certain embodiments, of the full length open reading frame (ORF) identified herein).

In other embodiments, fragments can comprise consecutive nucleotides of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, and up to one nucleotide less than the full length ANF prohormone. In some embodiments, fragments comprise biologically active fragments of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, or SEQ ID NO:6.

It is also well known in the art that restriction enzymes can be used to obtain biologically active fragments of the nucleic acid sequences, such as those encoding SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6. For example, Bal31 exonuclease can be conveniently used for time-controlled limited digestion of DNA (commonly referred to as "erase-a-base" procedures). See, for example, Maniatis et al. [1982] *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York; Wei et al., *J. Biol. Chem.*, 1983, 258:13006-13512.

The methods and pharmaceutical compositions of the present invention can utilize amino acid sequences that are biologically active fragments of the 126-amino acid atrial natriuretic factor (ANF) prohormone, such as NP₁₋₃₀, NP₃₁₋₆₇, NP₇₉₋₉₈, NP₉₉₋₁₂₆, and NP₇₃₋₁₀₂ (SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and SEQ ID NO:5, respectively), SEQ ID NO:6, and including biologically active fragments of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6.

Representative fragments of the polypeptides according to the invention will be understood to mean any polypeptide fragment having at least 8 or 9 consecutive amino acids, preferably at least 12 amino acids, and still more preferably at

least 15 or at least 20 consecutive amino acids of the polypeptide sequence from which it is derived, with retention of biological activity as described herein. The upper limit for such fragments is one amino acid less than the total number of amino acids found in the full-length sequence.

In other embodiments, fragments of the polypeptides can comprise consecutive amino acids of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, and up to one amino acid less than the full-length ANF prohormone. Fragments of polypeptides can be any portion of the full-length ANF prohormone amino acid sequence (including human or non-human mammalian homologs of the ANF prohormone) that exhibit biological activity as described herein, e.g., a C-terminally or N-terminally truncated version of the ANF prohormone, or an intervening portion of the ANF prohormone. In some embodiments, fragments comprise biologically active fragments of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6.

The present invention can be practiced using other biologically equivalent forms of ANF fragments or homologs thereof as can be appreciated by the sequence comparison below. Sequence similarities between mouse and human forms of ANP are shown where areas of conservation are clearly seen. NCBI BLAST Comparison of mouse (Query) to human (Subject) ANP a.a. sequences.

```
Query: 1MGFSIT-LGFFLVLAFLPGHIGANPVYSAVSNLDLMDFKNLLDHLEEKMPVEDEVMP
      M SFS T + F L-LAF L G ANP+Y+AVSN DLMDFKNLLDHLEEKMP+EDEV+PP
Sbjct: 1MSFSSTTTVSFLLLLAFQLLGQTRANPMYNAVSNADLMDFKNLLDHLEEKMPLEDEVVPP

Query: 60QALSEQTEEAGAALSSLEVPWPWTGEVNPPLRDGSA LGRSPWDPDSDXXXXXXXXXXXXX
      Q LSE EEAGAALS LPEVPPWTGEV+P RDG ALGR PWD SD
Sbjct: 61QVLSPEPNEEAGAALSPLPEVPPWTGEVSPAQRDGGALGRGPWDDSDRSALLKSKLRALLT

Query: 120GPRSLRRSSCFGGRIDRIGAQSGLGCNSFRY      150
      PRSLRRSSCFGGR+DRIGAQSGLGCNSFRY
Sbjct: 121APRSLRRSSCFGGRMDRIGAQSGLGCNSFRY      151
```

The NP utilized in the subject invention can be peptide derivatives, such as those disclosed in U.S. Patent Publication No. 2004/0266673 (Bakis et al.), which is incorporated herein by reference in its entirety. These NP derivatives include an NP and a reactive entity coupled to the NP peptide. The reactive entity is able to covalently bond with a functionality on a blood component. Such NP derivatives are reported to have an extended half-life in vivo. The NP utilized in the subject invention can be a modified NP, such as those described in U.S. Patent Publication No. 2004/0002458 (Seilhamer et al.) and U.S. Patent Publication No. 2003/0204063 (Gravel et al.), which are incorporated herein by reference in their entirety.

The NP utilized may be a fusion polypeptide comprising an NP, or Fragment or homolog thereof, and one or more additional polypeptides, such as another NP or a carrier protein, including those described in U.S. Patent Publication No. 2004/0138134 (Golembo et al.), which is incorporated herein by reference in its entirety. The NP utilized may be a chimeric polypeptide, such as those described in U.S. Patent Publication No. 2003/0069186 (Burnett et al.), which is incorporated herein by reference in its entirety. The fusion polypeptide or chimeric polypeptide may be administered to cells in vitro or

in vivo directly (i.e., as a polypeptide), or the fusion polypeptide may be administered as a polynucleotide encoding the fusion polypeptide with an operably linked promoter sequence. See, for example, Wang W. et al., "Albubnp, a Recombinant B-type Natriuretic Peptide and Human Serum Albumin Fusion Hormone, as a Long-Term Therapy of Congestive Heart Failure", Pharmaceutical Research, Springer Science and Business Media B. V., Formerly Kluwer Academic Publishers B. V., ISSN:0724-8741, volume 21, Number 11, November, 2004, pages 2105-2111.

The NP includes all hydrates and salts of natriuretic peptides that can be prepared by those of skill in the art. Under conditions where the compounds of the present invention are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compounds as salts may be appropriate. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids that form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, alpha-ketoglutarate, and alpha-glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts of NP may be obtained using standard procedures well known in the art, for example, by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

The NP of the invention can be prepared by well-known synthetic procedures. For example, the polypeptides can be

prepared by the well-known Merrifield solid support method. See Merrifield, *J. Amer. Chem. Soc.*, 1963, 85:2149-2154 and Merrifield (1965) *Science* 150:178-185. This procedure, using many of the same chemical reactions and blocking groups of classical peptide synthesis, provides a growing peptide chain anchored by its carboxyl terminus to a solid support, usually cross-linked polystyrene or styrenedivinylbenzene copolymer. This method conveniently simplifies the number of procedural manipulations since removal of the excess reagents at each step is effected simply by washing of the polymer.

Alternatively, these peptides can be prepared by use of well-known molecular biology procedures. Polynucleotides, such as DNA sequences, encoding the NP of the invention can be readily synthesized. Such polynucleotides are a further aspect of the present invention. These polynucleotides can be used to genetically engineer eukaryotic or prokaryotic cells, for example, bacteria cells, insect cells, algae cells, plant cells, mammalian cells, yeast cells or fungi cells for synthesis of the peptides of the invention.

For purposes of the present invention, the biological activity attributable to the homologs and fragments of NP and NP-encoding nucleic acid sequences means the capability to

prevent or alleviate symptoms associated with inflammatory and/or cell proliferation disorders such as cancer. This biological activity can be mediated by one or more of the following mechanisms: increased production of intracellular Calf concentration (e.g., in epithelial cells), increased production of nitric oxide (O), and decreased activation of transcription factors such as NFkB, ERK1,2 and/or AP1.

The methods of the subject invention also contemplate the administration of cells that have been genetically modified to produce NP, or biologically active fragments, variants, or homologs thereof. Such genetically modified cells can be administered alone or in combinations with different types of cells. Thus, genetically modified cells of the invention can be co-administered with other cells, which can include genetically modified cells or non-genetically modified cells. Genetically modified cells may serve to support the survival and function of the co-administered cells, for example.

The term "genetic modification" as used herein refers to the stable or transient alteration of the genotype of a cell of the subject invention by intentional introduction of exogenous nucleic acids by any means known in the art (including for example, direct transmission of a polynucleotide sequence from a cell or virus particle, transmission of infective virus particles, and transmission by any known polynucleotide-bearing substance) resulting in a permanent or temporary alteration of genotype. The nucleic acids may be synthetic, or naturally derived, and may contain genes, portions of genes, or other useful polynucleotides in addition to those encoding NP. A translation initiation codon can be inserted as necessary, making methionine the first amino acid in the sequence. The term "genetic modification" is not intended to include naturally occurring alterations such as that which occurs through natural viral activity, natural genetic recombination, or the like. The genetic modification may confer the ability to produce NP, wherein the cell did not previously have the capability, or the modification may increase the amount of NP endogenously produced by the cell, e.g., through increased expression.

Exogenous nucleic acids and/or vectors encoding NP can be introduced into a cell by viral vectors (retrovirus, modified herpes virus, herpes virus, adenovirus, adeno-associated virus, lentivirus, and the like) or direct DNA transfection (lipofection, chitosan-nanoparticle mediated transfection, calcium phosphate transfection, DEAE-dextran, electroporation, and the like), microinjection, cationic lipid-mediated transfection, transduction, scrape loading, ballistic introduction and infection (see, for example, Sambrook et al. [1989] *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.).

Preferably, the exogenous polynucleotide encoding the NP is operably linked to a promoter sequence that permits expression of the polynucleotide in a desired tissue within the patient. The promoters can be inducible, tissue-specific, or event-specific, as necessary.

The genetically modified cell may be chosen from eukaryotic or prokaryotic systems, for example, bacterial cells (Gram negative or Gram positive), yeast cells, animal cells, plant cells, and/or insect cells using baculovirus vectors, for example. In some embodiments, the genetically modified cell for expression of the nucleic acid sequences encoding NP, are human or non-human mammal cells.

According to the methods of the present invention, NP or polynucleotides encoding NP can be administered to a patient in order to alleviate (e.g., reduce or eliminate) a variety of symptoms associated with cancers, in various stages of pathological development. Treatment with NP or nucleic acid sequences encoding NP is intended to include prophylactic

intervention to prevent or reduce cancer cell growth (e.g., tumor growth) and onset of the symptoms associated with cancer cell growth (e.g. tumor growth), such as pain. The nucleic acid sequences and pharmaceutical compositions of the invention can be co-administered (concurrently or consecutively) to a patient with other therapeutic agents useful for treating cancers of the lung, ovarian, breast, as well as melanomas.

Suitable expression vectors for NP include any that are known in the art or yet to be identified that will cause expression of NP-encoding nucleic acid sequences in mammalian cells. Suitable promoters and other regulatory sequences can be selected as is desirable for a particular application. The promoters can be inducible, tissue-specific, or event-specific, as necessary. For example, the cytomegalovirus (CMV) promoter (Boshart et al., *Cell*, 1985, 41:521-530) and SV40 promoter (Subramani et al., *Mol. Cell. Biol.*, 1981, 1:854-864) have been found to be suitable, but others can be used as well. Optionally, the NP-encoding nucleic acid sequences used in the subject invention include a sequence encoding a signal peptide upstream of the NP-encoding sequence, thereby permitting secretion of the NP from a host cell. Also, various promoters may be used to limit the expression of the peptide in specific cells or tissues, such as lung cells.

A tissue-specific and/or event-specific promoter or transcription element that responds to the target microenvironment and physiology can also be utilized for increased transgene expression at the desired site. There has been an immense amount of research activity directed at strategies for enhancing the transcriptional activity of weak tissue-specific promoters or otherwise increasing transgene expression with viral vectors. It is possible for such strategies to provide enhancement of gene expression equal to one or two orders of magnitude, for example (see Nettelbeck et al., *Gene Ther.*, 1998, 5(12):1656-1664 and Qin et al., *Hum. Gene Ther.*, 1997, 8(17):2019-2019, the abstracts of which are submitted herewith for the Examiner's convenience). Examples of cardiac-specific promoters are the ventricular form of MLC-2v promoter (see, Zhu et al., *Mol. Cell Biol.*, 1993, 13:4432-4444, Navankasattusas et al., *Mol. Cell Biol.*, 1992, 12:1469-1479, 1992) and myosin light chain-2 promoter (Franz et al., *Circ. Res.*, 1993, 73:629-638). The E-cadherin promoter directs expression specific to epithelial cells (Behrens et al., *PNAS*, 1991, 88:11495-11499), while the estrogen receptor (ER) 3 gene promoter directs expression specifically to the breast epithelium (Hopp et al., *J. Mammary Gland Biol. Neoplasia*, 1998, 3:73-83). The human C-reactive protein (CRP) gene promoter (Ruther et al., *Oncogene* 8:87-93, 1993) is a liver-specific promoter. An example of a muscle-specific gene promoter is human enolase (ENO3) (Peshavaria et al., *Biochem. J.* 1993, 292(Pt 3):701-704). A number of brain-specific promoters are available such as the thy-1 antigen and gamma-enolase promoters (Vibert et al., *Eur. J. Biochem.* 181:33-39, 1989). The prostate-specific antigen promoter provides prostate tissue specificity (Pang et al., *Gene Ther.*, 1995, 6(11):1417-1426; Lee et al., *Anticancer Res.*, 1996, 16(4A):1805-1811). The surfactant protein B promoter provides lung specificity (Strayer et al., *Am. J. Respir. Cell Mol. Biol.*, 1998, 18(1):1-11). Any of the aforementioned promoters may be selected for targeted or regulated expression of the NP-encoding polynucleotide.

Various viral or non-viral vectors may be used to deliver polynucleotides encoding NP to cells in vitro or in vivo, resulting in expression and production of NP. Tissue-specific promoters or event-specific promoters may be utilized with polynucleotides encoding NP to further optimize and localize expression at target sites, such as within diseased tissues (e.g.,

cancer cells or tissues containing cancer cells). Robson et al. review various methodologies and vectors available for delivering and expressing a polynucleotide in vivo for the purpose of treating cancer (Robson, T. Hirst, D. G., *J. Biomed. and Biotechnol.*, 2003, 2003(2):110-137). Among the various targeting techniques available, transcriptional targeting using tissue-specific and event-specific transcriptional control elements is discussed. For example, Table 1 at page 112 of the Robson et al. publication lists several tissue-specific promoters useful in cancer therapy. Tables 2-4 of the Robson et al. publication list tumor-specific promoters, tumor environment-specific promoters, and exogenously controlled inducible promoters, many of which were available at the time the patent application was filed. The successful delivery and expression of the p53 tumor suppressor gene in vivo has been documented (Horowitz, J. *Curr. Opin. Mol. Ther.*, 1999, 1(4): 500-509; Von Gruenigen, V. F. et al. *Int. J. Gynecol. Cancer*, 1999, 9(5):365-372; Fujiwara, T. et al., *Mol. Urol*, 2000, 4(2):51-54, respectively).

Many techniques for delivery of drugs and proteins are available in the art to reduce the effects of enzymatic degradation, to facilitate cell uptake, and to reduce any potential toxicity to normal (undiseased) cells, etc. Such methods and reagents can be utilized for administration of NP to cells in vitro or in vivo. For example, peptides known as "cell penetrating peptides" (CPP) or "protein transduction domains" (PTD) have an ability to cross the cell membrane and enter the cell. PTDs can be linked to a cargo moiety such as a drug, peptide, or full-length protein, and can transport the moiety across the cell membrane. One well characterized PTD is the human immunodeficient virus (HIV)-1 Tat peptide (see, for example, Frankel et al., U.S. Pat. Nos. 5,804,604; 5,747,641; 6,674,980; 5,670,617; and 5,652,122; Fawell, S. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1994, 91:664-668). Peptides such as the homeodomain of *Drosophila antennapedia* (ANTp) and arginine-rich peptides display similar properties (Derossi, D. et al., *J. Biol. Chem.*, 1994, 269:10444-10450; Derossi, D. et al., *Trends Cell Biol.*, 1998, 8:84-87; Rojas, M. et al., *Nat. Biotechnol.*, 1998, 16:370-375; Futaki, S. et al., *J. Biol. Chem.*, 2001, 276:5836-5840). VP22, a tegument protein from Herpes simplex virus type 1 (HISV-1), also has the ability to transport proteins across a cell membrane (Elliot et al., *Cell*, 1997, 88:223-233; Schwarze S. R. et al., *Trends Pharmacol. Sci.*, 2000, 21:45-48). A common feature of these carriers is that they are highly basic and hydrophilic (Schwarze S. R. et al., *Trends Cell Biol.*, 2000, 10:290-295). Coupling of these carriers to marker proteins such as beta-galactosidase has been shown to confer efficient internalization of the marker protein into cells. More recently, chimeric, in-frame fusion proteins containing these carriers have been used to deliver proteins to a wide spectrum of cell types both in vitro and in vivo. For example, VP22-p53 chimeric protein retained its ability to spread between cells and its pro-apoptotic activity, and had a widespread cytotoxic effect in p53 negative human osteosarcoma cells in vitro (Phelan, A. et al., *Nature Biotechnol.*, 1998, 16:440-443). Intraperitoneal injection of the beta-galactosidase protein fused to the HIV-1 Tat peptide resulted in delivery of the biologically active fusion protein to all tissues in mice, including the brain (Schwarze S. R. et al., *Science*, 1999, 285:1569-1572).

Liposomes of various compositions can also be used for site-specific delivery of proteins and drugs (Witschi, C. et al., *Pharm. Res.*, 1999, 16:382-390; Yell, M. K. et al., *Pharm. Res.*, 1996, 16:1693-1698). The interaction between the liposomes and the protein cargo usually relies on hydrophobic interactions or charge attractions, particularly in the case of cationic lipid delivery systems (Zelphati, O. et al., *J. Biol.*

Chem., 2001, 276:35103-35110). Tat peptide-bearing liposomes have also been constructed and used to deliver cargo directly into the cytoplasm, bypassing the endocytotic pathway (Torchilin V. P. et al., *Biochim. Biophys. Acta-Biomembranes*, 2001, 1511:397-411; Torchilin V. P. et al., *Proc. Natl. Acad. Sci. USA*, 2001, 98:8786-8791). When encapsulated in sugar-grafted liposomes, pentamidine isethionate and a derivative have been found to be more potent in comparison to normal liposome-encapsulated drug or to the free drug (Banerjee, G. et al., *J. Antimicrob. Chemother.*, 1996, 38(1):145-150). A thermo-sensitive liposomal taxol formulation (heat-mediated targeted drug delivery) has been administered in vivo to tumor-bearing mice in combination with local hyperthermia, and a significant reduction in tumor volume and an increase in survival time was observed compared to the equivalent dose of free taxol with or without hyperthermia (Sharma, D. et al., *Melanoma Res.*, 1998, 8(3):240-244). Topical application of liposome preparations for delivery of insulin, IFN-alpha, IFN-gamma, and prostaglandin E1 have met with some success (Cevc G. et al., *Biochim. Biophys. Acta*, 1998, 1368:201-215; Foldvari M. et al., *J. Liposome Res.*, 1997, 7:115-126; Short S. M. et al., *Pharm. Res.*, 1996, 13:1020-1027; Foldvari M. et al., *Urology*, 1998, 52(5):838-843; U.S. Pat. No. 5,853,755).

Antibodies represent another targeting device that may make liposome uptake tissue-specific or cell-specific (Mastrobattista, E. et al., *Biochim. Biophys. Acta*, 1999, 1419(2): 353-363; Mastrobattista, E. et al., *Adv. Drug Deliv. Rev.*, 1999, 40(1-2):103-127). The liposome approach offers several advantages, including the ability to slowly release encapsulated drugs and proteins, the capability of evading the immune system and proteolytic enzymes, and the ability to target tumors and cause preferentially accumulation in tumor tissues and their metastases by extravasation through their leaky neovasculature. Other carriers have also been used to deliver anti-cancer drugs to neoplastic cells, such as polyvinylpyrrolidone nanoparticles and maleylated bovine serum albumin (Sharma, D. et al., *Oncol. Res.*, 1996, 8(7-8):281-286; Mukhopadhyay, A. et al., *FEBS Lett.*, 1995, 376(1-2): 95-98). Thus, using targeting and encapsulation technologies, which are very versatile and amenable to rational design and modification, delivery of NP to desired cells can be facilitated. Furthermore, because many liposome compositions are also viable delivery vehicles for genetic material, many of the advantages of liposomes are equally applicable to polynucleotides encoding NP.

As indicated above, the pharmaceutical composition of the present invention can include a liposome component. According to the present invention, a liposome comprises a lipid composition that is capable of fusing with the plasma membrane of a cell, thereby allowing the liposome to deliver a nucleic acid molecule and/or a protein composition into a cell. Some preferred liposomes include those liposomes commonly used in gene delivery methods known to those of skill in the art. Some preferred liposome delivery vehicles comprise multilamellar vesicle (MLV) lipids and extruded lipids, although the invention is not limited to such liposomes. Methods for preparation of MLVs are well known in the art. "Extruded lipids" are also contemplated. Extruded lipids are lipids that are prepared similarly to MLV lipids, but which are subsequently extruded through filters of decreasing size, as described in Templeton et al., *Nature Biotech.*, 1997, 15:647-652, which is incorporated herein by reference in its entirety. Small unilamellar vesicle (SUV) lipids can also be used in the compositions and methods of the present invention. Other preferred liposome delivery vehicles comprise liposomes having a polycationic lipid composition (i.e. cationic lipo-

somes). For example, cationic liposome compositions include, but are not limited to, any cationic liposome complexed with cholesterol, and without limitation, include DOTMA and cholesterol, DOTAP and cholesterol, DOTIM and cholesterol, and DDAB and cholesterol. Liposomes utilized in the present invention can be any size, including from about 10 to 1000 nanometers (nm), or any size in between.

A liposome delivery vehicle can be modified to target a particular site in a mammal, thereby targeting and making use of an NP-encoding nucleic acid molecule of the present invention at that site. Suitable modifications include manipulating the chemical formula of the lipid portion of the delivery vehicle. Manipulating the chemical formula of the lipid portion of the delivery vehicle can elicit the extracellular or intracellular targeting of the delivery vehicle. For example, a chemical can be added to the lipid formula of a liposome that alters the charge of the lipid bilayer of the liposome so that the liposome fuses with particular cells having particular charge characteristics. In one embodiment, other targeting mechanisms, such as targeting by addition of exogenous targeting molecules to a liposome (i.e., antibodies) may not be a necessary component of the liposome of the present invention, since effective immune activation at immunologically active organs can already be provided by the composition when the route of delivery is intravenous or intraperitoneal, without the aid of additional targeting mechanisms. However, in some embodiments, a liposome can be directed to a particular target cell or tissue by using a targeting agent, such as an antibody, soluble receptor or ligand, incorporated with the liposome, to target a particular cell or tissue to which the targeting molecule can bind. Targeting liposomes are described, for example, in Ho et al., *Biochemistry*, 1986, 25: 5500-6; Ho et al., *J Biol Chem*, 1987a, 262: 13979-84; Ho et al., *J Biol Chem*, 1987b, 262: 13973-8; and U.S. Pat. No. 4,957,735 to Huang et al., each of which is incorporated herein by reference in its entirety. In one embodiment, if avoidance of the efficient uptake of injected liposomes by reticuloendothelial system cells due to opsonization of liposomes by plasma proteins or other factors is desired, hydrophilic lipids, such as gangliosides (Allen et al., *FEBS Lett*, 1987, 223: 42-6) or polyethylene glycol (PEG)-derived lipids (Klibanov et al., *FEBS Lett*, 1990, 268: 235-7), can be incorporated into the bilayer of a conventional liposome to form the so-called sterically-stabilized or "stealth" liposomes (Woodle et al., *Biochim Biophys Acta*, 1992, 1113: 171-99). Variations of such liposomes are described, for example, in U.S. Pat. No. 5,705,187 to Unger et al., U.S. Pat. No. 5,820,873 to Choi et al., U.S. Pat. No. 5,817,856 to Tirosh et al.; U.S. Pat. No. 5,686,101 to Tagawa et al.; U.S. Pat. No. 5,043,164 to Huang et al., and U.S. Pat. No. 5,013,556 to Woodle et al., all of which are incorporated herein by reference in their entireties).

The NP-encoding nucleic acid sequences of the present invention can be conjugated with chitosan. For example, DNA chitosan nanospheres can be generated, as described by Roy, K. et al. (1999, *Nat Wed* 5:387). Chitosan allows increased bioavailability of the NP-encoding nucleic acid sequences because of protection from degradation by serum nucleases in the matrix and thus has great potential as a mucosal gene delivery system. Chitosan also has many beneficial effects, including anticoagulant activity, wound-healing properties, and immunostimulatory activity, and is capable of modulating immunity of the mucosa and bronchus-associated lymphoid tissue.

Mammalian species which benefit from the disclosed methods of treatment include, and are not limited to, apes, chimpanzees, orangutans, humans, monkeys; domesticated animals (e.g., pets) such as dogs, cats, guinea pigs, hamsters,

Vietnamese pot-bellied pigs, rabbits, and ferrets; domesticated farm animals such as cows, buffalo, bison, horses, donkey, swine, sheep, and goats; exotic animals typically found in zoos, such as bear, lions, tigers, panthers, elephants, hippopotamus, rhinoceros, giraffes, antelopes, sloth, gazelles, zebras, wildebeests, prairie dogs, koala bears, kangaroo, opossums, raccoons, pandas, hyena, seals, sea lions, elephant seals, otters, porpoises, dolphins, and whales. The terms "patient" and "subject" are used interchangeably herein are intended to include such human and non-human mammalian species. According to the method of the present invention, human or non-human mammalian NP (or nucleic acid sequences encoding human or non-human mammalian NP) can be administered to the patient. The NP may be naturally occurring within the patient's species or a different mammalian species. The expression vectors used in the subject invention can comprise nucleic acid sequences encoding any human or non-human mammalian NP. In instances where genetically modified cells are administered to a patient, the cells may be autogenic, allogeneic, or xenogeneic, for example.

In another aspect, the present invention concerns pharmaceutical compositions containing a therapeutically effective amount of agent that reduces NPR-A activity, such as an NP, or polynucleotides encoding NP, and a pharmaceutically acceptable carrier. Preferably, if the agent is a polynucleotide, such as an NP-encoding nucleic acid sequence, the polynucleotide is contained within an expression vector, such as plasmid DNA or a virus. Pharmaceutical compositions including a therapeutically effective amount of an agent that reduces NPR-A activity such as NP, or nucleic acid sequences encoding NP, and a pharmaceutically acceptable carrier, can be administered to a patient by any effective route, including local or systemic delivery. Administration can be continuous or at distinct intervals as can be determined by a person skilled in the art.

The agent that reduces NPR-A activity, such as NP or polynucleotides encoding NP (and pharmaceutical compositions containing them), can be administered to a patient by any route that results in prevention (or reduction of onset) or alleviation of symptoms associated with cancer, such as pain. For example, the agent (e.g., NP or NP-encoding nucleic acid molecule) can be administered parenterally, intravenously (I.V.), intramuscularly (I.M.), subcutaneously (S.C.), intradermally (I.D.), topically, transdermally, orally, intranasally, etc.

If desired, the pharmaceutical composition can be adapted for administration to the airways of the patient, e.g., nose, sinus, throat and lung, for example, as nose drops, as nasal drops, by nebulization as an inhalant, vaporization, or other methods known in the art. Examples of intranasal administration can be by means of a spray, drops, powder or gel and also described in U.S. Pat. No. 6,489,306, which is incorporated herein by reference in its entirety. One embodiment of the present invention is the administration of the invention as a nasal spray. Alternate embodiments include administration through any oral or mucosal routes, sublingual administration and even eye drops. However, other means of drug administrations are well within the scope of the present invention.

The pharmaceutical compositions of the subject invention can be formulated according to known methods for preparing pharmaceutically useful compositions. Furthermore, as used herein, the phrase "pharmaceutically acceptable carrier" includes any of the standard pharmaceutically acceptable carriers. The pharmaceutically acceptable carrier can include diluents, adjuvants, and vehicles, as well as implant carriers, and inert, non-toxic solid or liquid fillers, diluents, or encaps-

sulating material that does not react with the active ingredients of the invention. Examples include, but are not limited to, phosphate buffered saline, physiological saline, water, and emulsions, such as oil/water emulsions. The carrier can be a solvent or dispersing medium containing, for example, ethanol, polyol (for example, glycerol, propylene glycol, liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. Formulations containing pharmaceutically acceptable carriers are described in a number of sources which are well known and readily available to those skilled in the art. For example, *Remington's Pharmaceutical Sciences* (Martin E. W., 1995, Easton Pa., Mack Publishing Company, 19th ed.), which is incorporated herein by reference in its entirety, describes formulations that can be used in connection with the subject invention.

Pharmaceutical compositions of the present invention useful for parenteral injection can include pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents, or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. Formulations suitable for parenteral administration include, for example, aqueous injectable solutions that may contain antioxidants, buffers, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions, which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that, in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The pharmaceutical compositions used in the methods of the present invention may also contain adjuvants such as preservatives, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption, such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the active agent (e.g., NP), it is desirable to slow the absorption from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the NP or NP-encoding polynucleotide then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered NP or NP-

encoding polynucleotide is accomplished by dissolving or suspending the NP in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the agent (e.g. NP or NP-encoding polynucleotide) in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of active agent (e.g., NP or NP-encoding polynucleotide) to polymer and the nature of the particular polymer employed, the rate of release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active agents (NP or NP-encoding polynucleotide) are mixed with at least one pharmaceutically acceptable excipient or carrier such as sodium nitrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid; b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay; and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hardfilled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. Optionally, the solid dosage forms contain opacifying agents, and can be of a composition that releases the NP or NP-encoding polynucleotide only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

The active agents (NP or NP-encoding polynucleotide) can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the NP or NP-encoding polynucleotide, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth, and mixtures thereof.

Topical administration includes administration to the skin or mucosa, including surfaces of the lung and eye. Compositions for topical administration, including those for inhalation, may be prepared as a dry powder, which may be pressurized or non-pressurized. In nonpressurized powder compositions, the active ingredients in finely divided form may be used in admixture with a larger-sized pharmaceutically acceptable inert carrier comprising particles having a size, for example, of up to 100 μm in diameter. Suitable inert carriers include sugars such as lactose. Desirably, at least 95% by weight of the particles of the active ingredient have an effective particle size in the range of 0.01 to 10 μm .

Alternatively, the pharmaceutical composition may be pressurized and contain a compressed gas, such as nitrogen or a liquefied gas propellant. The liquefied propellant medium or the entire composition is preferably such that the active ingredients do not dissolve therein to any substantial extent. The pressurized composition may also contain a surface active agent. The surface active agent may be a liquid or solid non-ionic surface active agent or may be a solid anionic surface active agent. It is preferred to use the solid anionic surface active agent in the form of a sodium salt.

The compositions and methods of the invention can further incorporate permeation enhancers, such as those described in U.S. Patent Publication No. 2003/0147943 (Luo et al.), penetrating peptides capable of facilitating penetration of an NP, or an NP-encoding polynucleotide, across a biological barrier, such as those described in U.S. Patent Publication No. 2004/0146549 (Ben-Sasson et al.), enhancer compounds that enhance the absorption of a polypeptide in the respiratory tract, such as those described in U.S. Patent Publication No. 2004/0171550 (Backstrom et al.), and suspension vehicles, such as those described in U.S. Patent Publication No. 2004/0224903 (Berry et al.), each of which are incorporated herein by reference in their entirety.

The agent that reduces NPR-A activity (such as NP or NP-encoding polynucleotide) is administered and dosed in accordance with good medical practice, taking into account the clinical condition of the individual patient, the site and method of administration, scheduling of administration, patient age, sex, body weight, and other factors known to medical practitioners. The pharmaceutically "effective amount" for purposes herein is thus determined by such considerations as are known in the art. For example, an effective amount of NP-encoding polynucleotide is that amount necessary to provide an effective amount of NP, when expressed *in vivo* or *in vitro*. The amount of the agent (e.g., NP or NP-encoding nucleic acid molecule) must be effective to achieve some improvement including, but not limited to, improved survival rate, more rapid recovery, total prevention of symptoms associated with an inflammatory or cell proliferation disorder, such as cancer, or improvement or elimination of symptoms associated with an inflammatory or cell proliferation disorder, such as cancer, and other indicators as are selected as appropriate measures by those skilled in the art. In accordance with the present invention, a suitable single dose size is a dose that is capable of preventing or alleviating (reducing or eliminating) a symptom in a patient when

administered one or more times over a suitable time period. One of skill in the art can readily determine appropriate single dose sizes for local or systemic administration based on the size of a mammal and the route of administration.

In accordance with the invention, a mammal (such as a human) that is predisposed to or suffering from a physical disorder may be treated by administering to the mammal an effective amount of an agent that reduces NPR-A activity (such as NP or NP-encoding polynucleotide), in combination with a pharmaceutically acceptable carrier or excipient therefore (as described below). Physical disorders treatable with the compositions and methods of the present invention include any physical disorder that may be delayed, prevented cured or otherwise treated by administration of an agent that reduces NPR-A activity (such as NP or NP-encoding polynucleotide) in a mammal suffering from or predisposed to the physical disorder. Such physical disorders include, but are not limited to, a variety of carcinomas and other cancers, such as skin cancers (including melanomas and Kaposi's Sarcoma), oral cavity cancers, lung cancers, breast cancers, prostatic cancers, bladder cancers, liver cancers, pancreatic cancers, cervical cancers, ovarian cancers, head and neck cancers, colon cancers, germ cell cancers (including teratocarcinomas) and leukemias. Other physical disorders treatable by the methods of the present invention include inflammatory disorders such as rheumatoid arthritis, multiple sclerosis, systemic lupus erythematosus, pelvic inflammatory disease, and Crohn's disease. The methods of the invention may also be used to treat a mammal suffering from or predisposed to a fibrotic disorder, including pulmonary fibrosis, cystic fibrosis, endomyocardial fibrosis, hepatic fibrosis (particularly hepatic cirrhosis), myelofibrosis, scleroderma, and systemic sclerosis. Other physical disorders treatable by the methods of the invention include osteoporosis, atherosclerosis, and ocular disorders such as corneal ulceration and diabetic retinopathy. The methods of the present invention may also be used in the prevention of disease progression, such as in chemoprevention of the progression of a premalignant lesion to a malignant lesion, and to treat a mammal suffering from, or predisposed to, other physical disorders that respond to treatment with compositions that differentially modulate gene expression.

Cell proliferation disorders include but are not limited to solid tumors, such as cancers of the breast, respiratory tract, brain, reproductive organs, digestive tract, urinary tract, eye, liver, skin, head and neck, thyroid, parathyroid and their distant metastases. Those disorders also include lymphomas, sarcomas, and leukemias.

Cancers of any organ can be treated, including cancers of, but are not limited to, e.g., colon, pancreas, breast, prostate, bone, liver, kidney, lung, testes, skin, pancreas, stomach, colorectal cancer, renal cell carcinoma, hepatocellular carcinoma, melanoma, etc.

Examples of breast cancer include, but are not limited to, invasive ductal carcinoma, invasive lobular carcinoma, ductal carcinoma *in situ*, and lobular carcinoma *in situ*. Examples of cancers of the respiratory tract include, but are not limited to, small-cell and non-small-cell lung carcinoma, as well as bronchial adenoma and pleuropulmonary blastoma. Examples of brain cancers include, but are not limited to, brain stem and hypothalamic glioma, cerebellar and cerebral astrocytoma, medulloblastoma, ependymoma, as well as neuroectodermal and pineal tumor. Tumors of the male reproductive organs include, but are not limited to, prostate and testicular cancer. Tumors of the female reproductive organs include, but are not limited to, endometrial, cervical, ovarian, vaginal, and vulvar cancer, as well as sarcoma of the uterus.

Tumors of the digestive tract include, but are not limited to, anal, colon, colorectal, esophageal, gallbladder, gastric, pancreatic, rectal, small-intestine, and salivary gland cancers. Tumors of the urinary tract include, but are not limited to, bladder, penile, kidney, renal pelvis, ureter, and urethral cancers. Eye cancers include, but are not limited to, intraocular melanoma and retinoblastoma. Examples of liver cancers include, but are not limited to, hepatocellular carcinoma (liver cell carcinomas with or without fibrolamellar variant), cholangiocarcinoma (intrahepatic bile duct carcinoma), and mixed hepatocellular cholangiocarcinoma. Skin cancers include, but are not limited to, squamous cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell skin cancer, and non-melanoma skin cancer. Head-and-neck cancers include, but are not limited to, laryngeal, hypopharyngeal, nasopharyngeal, and/or oropharyngeal cancers, and lip and oral cavity cancer. Lymphomas include, but are not limited to, AIDS-related lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, Hodgkin's disease, and lymphoma of the central nervous system. Sarcomas include, but are not limited to, sarcoma of the soft tissue, osteosarcoma, malignant fibrous histiocytoma, lymphosarcoma, and rhabdomyosarcoma. Leukemias include, but are not limited to, acute myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, and hairy cell leukemia. In addition to reducing the proliferation of tumor cells, agents that reduce NPR-A activity can also cause tumor regression, e.g., a decrease in the size of a tumor, or in the extent of cancer in the body.

In addition to chemotherapeutic agents, the methods and compositions of the subject invention can incorporate treatments and agents utilizing, for example, angiogenesis inhibitors (Thalidomide, Bevacizumab), Bcl-2 antisense oligonucleotides (G3139), a PSA based vaccine, a PDGF receptor inhibitor (Gleevec), microtubule stabilizers (Epothilones), and a pro-apoptotic agent (Perifosine). Thus, an NP or NP-encoding polynucleotide can be administered to a patient in combination (simultaneously or consecutively) with other agents for useful for treating inflammatory disorders and/or cell proliferation disorders. Likewise, the pharmaceutical compositions of the subject invention can include such agents.

The term "gene therapy", as used herein, refers to the transfer of genetic material (a polynucleotide, e.g., DNA or RNA) of interest into a host to treat or prevent a genetic or acquired disease or condition phenotype. The genetic material of interest encodes a product (e.g., a protein, polypeptide, peptide, or functional RNA) whose production in vivo is desired, such as NP. In addition to one or more NP, the genetic material of interest can encode a hormone, receptor, enzyme, polypeptide or peptide of therapeutic and/or diagnostic value. For a review see, in general, the text "Gene Therapy" (Advances in Pharmacology 40, Academic Press, 1997).

Two basic approaches to gene therapy have evolved: (1) ex vivo and (2) in vivo gene therapy. In ex vivo gene therapy, cells are removed from a patient and, while being cultured, are treated in vitro. Generally, a functional replacement gene is introduced into the cell via an appropriate gene delivery vehicle/method (transfection, transduction, homologous recombination, etc.) and an expression system as needed and then the modified cells are expanded in culture and returned to the host/patient. These genetically reimplanted cells have been shown to produce the transfected gene product in situ.

In vivo gene therapy, target cells are not removed from the subject, rather the gene to be transferred is introduced into the cells of the recipient organism in situ, that is within the recipient. Alternatively, if the host gene is defective, the gene

is repaired in situ. Thus, these genetically altered cells produce the transfected gene product (e.g. NP) in situ.

The gene expression vector is capable of delivery/transfer of heterologous nucleic acid sequences (e.g. NP-encoding nucleic acid sequences) into a host cell. The expression vector may include elements to control targeting, expression and transcription of the nucleic acid sequence in a cell selective manner as is known in the art. It should be noted that often the 5'UTR and/or 3'UTR of the gene may be replaced by the 5'UTR and/or 3'UTR of the expression vehicle.

The expression vector can include a promoter for controlling transcription of the heterologous material and can be either a constitutive or inducible promoter to allow selective transcription. The expression vector can also include a selection gene.

Vectors can be introduced into cells or tissues by any one of a variety of known methods within the art. Such methods can be found generally described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory, New York (1989, 1992), in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Md. (1989), Chang et al., *Somatic Gene Therapy*, CRC Press, Ann Arbor, Mich. (1995), Vega et al., *Gene Targeting*, CRC Press, Ann Arbor, Mich. (1995), *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, Butterworths, Boston Mass. (1988) and include, for example, stable or transient transfection, lipofection, electroporation, and infection with recombinant viral vectors.

Introduction of nucleic acids by infection offers several advantages over the other listed methods. Higher efficiency can be obtained due to their infectious nature. Moreover, viruses are very specialized and typically infect and propagate in specific cell types. Thus, their natural specificity can be used to target the vectors to specific cell types in vivo or within a tissue or mixed culture of cells. Viral vectors can also be modified with specific receptors or ligands to alter target specificity through receptor mediated events.

A specific example of a DNA viral vector for introducing and expressing recombinant sequences is the adenovirus derived vector Adenov3TK. This vector expresses a herpes virus thymidine kinase (TK) gene for either positive or negative selection and an expression cassette for desired recombinant sequences. This vector can be used to infect cells that have an adenovirus receptor which includes most cancers of epithelial origin as well as others. This vector as well as others that exhibit similar desired functions can be used to treat a mixed population of cells and can include, for example, an in vitro or ex vivo culture of cells, a tissue or a human subject.

Additional features can be added to the vector to ensure its safety and/or enhance its therapeutic efficacy. Such features include, for example, markers that can be used to negatively select against cells infected with the recombinant virus. An example of such a negative selection marker is the TK gene described above that confers sensitivity to the antibiotic ganciclovir. Negative selection is therefore a means by which infection can be controlled because it provides inducible suicide through the addition of antibiotic. Such protection ensures that if, for example, mutations arise that produce altered forms of the viral vector or recombinant sequence, cellular transformation will not occur. Features that limit expression to particular cell types can also be included. Such features include, for example, promoter and regulatory elements that are specific for the desired cell type.

In addition, recombinant viral vectors are useful for in vivo expression of a desired nucleic acid because they offer advantages such as lateral infection and targeting specificity. Lateral infection is inherent in the life cycle of, for example,

retrovirus and is the process by which a single infected cell produces many progeny virions that bud off and infect neighboring cells. The result is that a large area becomes rapidly infected, most of which was not initially infected by the original viral particles. This is in contrast to vertical-type of infection in which the infectious agent spreads only through daughter progeny. Viral vectors can also be produced that are unable to spread laterally. This characteristic can be useful if the desired purpose is to introduce a specified gene into only a localized number of targeted cells.

In another aspect, the present invention concerns an isolated peptide comprising the amino acid sequence NP₇₃₋₁₀₂ (SEQ ID NO:5), or a biologically active fragment or homolog thereof. NP₇₃₋₁₀₂ is amino acids 73-102 of the 151-amino acid long human atrial natriuretic factor (ANF). In another aspect, the present invention concerns an isolated peptide comprising the amino acid sequence of SEQ ID NO:6, or a biologically active fragment or homolog thereof. SEQ ID NO:6 is a biologically active fragment of the human ANF. In another aspect, the present invention concerns an isolated nucleic acid molecule encoding the amino acid sequence of NP₇₃₋₁₀₂ (SEQ ID NO:5), or a biologically active fragment or homolog thereof. In another aspect, the present invention concerns an isolated nucleic acid molecule (SEQ ID NO:13) encoding the amino acid sequence of SEQ ID NO:6, or a biologically active fragment or homolog thereof.

As used herein, the terms "peptide", "polypeptide", and "protein" refer to amino acid sequences of any length unless otherwise specified.

Assays for Identifying Agents that Reduce Natriuretic Peptide Receptor-A Activity

The present invention also includes methods for identifying agents that reduce the activity of natriuretic peptide receptor-A (also known in the art as NPR-A, NPR-A, and guanylate cyclase A) in vitro or in vivo (also referred to herein as the diagnostic method or screening assay of the invention). Such agents are potentially useful for treating inflammatory or cell proliferation disorders in a patient. In the therapeutic methods and assays of the invention, agents that reduce NPR-A activity include those that, for example, reduce ANP-NPR-A induced c-GMP production, reduce expression of NPR-A, reduce cellular internalization of NPR-A, reduce recycling of NPR-A to the cell membrane, or otherwise interfere with the activity of the receptor.

Production of ANP-NPR-A induced cGMP production can be assayed and used as a high-throughput method for screening agents for anti-proliferative (e.g., anti-cancer) and anti-inflammatory activity. This assay can be carried out using a cell line that transiently or stably expresses the receptor for ANP, NPR-A (Pandey et al., *J Biol. Chem.* 2002, 277:4618-4627) and libraries of agents, such as peptide and compound libraries, which can be novel or obtained commercially. An assay for cGMP can be performed to select agents that are inhibitors of cGMP. Alternatively, ANP peptide can be linked with a moiety that can antagonize cGMP following internalization, which can be checked using a transiently or stably transfected cell line expressing NPR-A.

In the context of the screening assay of the invention, the terms "recombinant host cells", "host cells", "genetically modified host cells", "cells", "cell lines", "cell cultures", and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be, or have been, used as recipients for recombinant vectors or other transfer DNA, immaterial of the method by which the DNA is introduced into the cell or the subsequent disposition of the cell. The terms include the progeny of the original cell that has been transfected. Cells in primary

culture can also be used as recipients. Host cells can range in plasticity and proliferation potential. Host cells can be differentiated cells, progenitor cells, or stem cells, for example.

Host cells can be genetically modified with vectors to express (e.g., overexpress) the NPR-A receptor, or a mutant, isoform, or other variant thereof, which may be a cloning vector or an expression vector, for example. The vector may be in the form of a plasmid, a virus, (e.g. a retrovirus or other virus), a viral particle, a phage, etc. The genetically modified host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants/transfectants or amplifying the receptor-encoding polynucleotide.

In one embodiment, the host cell is a human cell. In another embodiment, the host cell is a non-human mammalian cell. Both prokaryotic and eukaryotic host cells may be used for expression of desired coding sequences when appropriate control sequences (e.g., promoter sequences) that are compatible with the designated host are used. For example, among prokaryotic hosts, *Escherichia coli* may be used. Also, for example, expression control sequences for prokaryotes include but are not limited to promoters, optionally containing operator portions, and ribosome binding sites. Eukaryotic hosts include yeast and mammalian cells in culture systems. *Pichia pastoris*, *Saccharomyces cerevisiae* and *S. carlsbergensis* are commonly used yeast hosts. Yeast-compatible vectors carry markers that permit selection of successful transformants by conferring protrophy to auxotrophic mutants or resistance to heavy metals on wild-type strains. Yeast compatible vectors may employ the 2-p origin of replication (Broach et al. *Meth. Enzymol.* 101:307, 1983), the combination of CEN3 and ARS1 or other means for assuring replication, such as sequences that will result in incorporation of an appropriate fragment into the host cell genome. Control sequences for yeast vectors are known in the art and include but are not limited to promoters for the synthesis of glycolytic enzymes, including the promoter for 3-phosphoglycerate kinase. (See, for example, Hess et al. *J. Adv. Enzyme Reg.* 7:149, 1968; Holland et al. *Biochemistry* 17:4900, 1978; and Hitzeman *J. Biol. Chem.* 255:2073, 1980). For example, some useful control systems are those that comprise the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) promoter or alcohol dehydrogenase (ADH) regulatable promoter, terminators also derived from GAPDH, and, if secretion is desired, leader sequences from yeast alpha factor. In addition, the transcriptional regulatory region and the transcriptional initiation region which are operably linked may be such that they are not naturally associated in the wild-type organism.

Host cells useful for expression of polynucleotides encoding the NPR-A receptor may be primary cells or cells of cell lines. The host cells may be tumor cells (transformed cells) or non-tumor cells. Mammalian cell lines available as hosts for expression are known in the art and are available from depositories such as the American Type Culture Collection. These include but are not limited to HeLa cells, human embryonic kidney (HEK) cells, Chinese hamster ovary (CHO) cells, baby hamster kidney (BHK) cells, and others.

The number of host cells used in a particular assay will vary with the objectives of the assay, the solid support used to support or contain the cell(s), if one is utilized etc. Thus, in some protocols, the host cell may be a single cell. In other protocols, a plurality of host cells will be used.

In accordance with the screening assay of the invention, the polynucleotide encoding the NPR-A is operably linked to a promoter sequence. Suitable promoter sequences for mammalian cells also are known in the art and include viral promoters such as that from Simian Virus 40 (SV40), Rous

sarcoma virus (RSV), adenovirus (ADV), bovine papilloma virus (BPV) and cytomegalovirus (CMV). Mammalian cells also may require terminator sequences and poly A addition sequences; enhancer sequences which increase expression also may be included, and sequences which cause amplification of the gene also may be desirable. These sequences are known in the art. Vectors suitable for replication in mammalian cells may include viral replicons, or sequences which ensure integration of the appropriate sequences including the NPR-A receptor into the host genome. An example of such a mammalian expression system is described in Gopalakrishnan et al. *Eur. J. Pharmacol.-Mol. Pharmacol.* 290: 237-246, 1995).

Candidate agents (and treatments) that may be tested by the screening assays of the present invention include polypeptides, non-peptide small molecules, biological agents, and any other source of candidate agents potentially having the ability to modulate (e.g., reduce) NPR-A activity. Candidate agents and treatments may be useful for the treatment of inflammatory and/or cell proliferation disorders, such as cancer. Candidate agents can be virtually any substance and can encompass numerous chemical classes, including organic compounds or inorganic compounds. A candidate agent may be a substance such as genetic material, protein, lipid, carbohydrate, small molecules, a combination of any of two or more of foregoing, or other compositions. Candidate agents may be naturally occurring or synthetic, and may be a single substance or a mixture. Candidate agents can be obtained from a wide variety of sources including libraries of compounds. A candidate agent can be or include, for example, a polypeptide, peptidomimetic, amino acid(s), amino acid analog(s), polynucleotide(s), polynucleotide analog(s), nucleotide(s), nucleotide analog(s), or other small molecule(s). A polynucleotide may encode a polypeptide that potentially reduces NPR-A activity within the cell, or the polynucleotide may be a short interfering RNA (siRNA), a hairpin RNA (shRNA), antisense oligonucleotide, ribozyme, or other polynucleotide that targets an endogenous or exogenous gene for silencing of gene expression and potentially NPR-A activity within the cell. Candidate treatments may include exposure of the host cells to any conditions that potentially reduce NPR-A activity within the host cells. The treatment may involve exposing the cells to an energy source, for example.

According to the screening assay of the invention, the method for identifying agents (which is intended to be inclusive of treatments) that reduce NPR-A activity is used to identify an agent that is therapeutic for treating an inflammation disorder and/or cell proliferation disorder, such as cancer. In aspect, the screening assay comprising contacting a host cell with a candidate agent, wherein the host cell expresses NPR-A, or an active fragment or variant thereof, and determining whether activity of the receptor is reduced, wherein a decrease in receptor activity is indicative of a potentially therapeutic agent. The method can optionally include an additional step of comparing NPR-A activity in the presence of the candidate agent, with NPR-A activity in the absence of the candidate agent (e.g., or other positive or negative control). The determination of NPR-A activity may be quantitative, semi-quantitative, or qualitative.

Known methods for overexpressing NPR-A in host cells and determining intracellular cGMP may be utilized to determine whether NPR-A activity is reduced (Kumar et al., *Hypertension*, 1997, 29(part 2):414-421; Khurana M. L. and Pandey K. N., *Endocrinology*, 1993, 133:2141-2149; Delport C. et al., *Eur. J. Pharmacol.*, 1992, 224(2-3):183-188; Ohyama Y. et al., *Biochem. Biophys. Res. Commun.*, 1992, 189(1):336-342; Sharma G. D. et al., Expression of Atrial

Natriuretic Peptide Receptor-A Antagonizes the Mitogen-Activated Protein Kinases (erk2 and P38^{MAPK}) in cultured human vascular Smooth Muscle Cells", in *Molecular and Cellular Biochemistry*, Springer Science+Business Media B. V., ISSN:0300-8177, Vol. 233, Number. 1-2, April 2002, pages 165-173; Pandey K. N. et al., *Biochem. Biophys. Res. Commun.*, 2000, 271(2):374-379; Fujiseki Y. et al., *Jpn. J. Pharmacol.*, 1999, 79(3):359-368; Pandey K. N., *Can. J. Physiol. Pharmacol.*, 2001, 79(8):631-639; Pandey K. N., *Mol. Cell. Biochem.*, 2002, 230(1-2):61-72; Sekiguchi T. et al., *Gene*, 2001, 273:251-257; Chen S. et al., *J. Am. Soc. Nephrol.*, 2005, 16:329-339; Pandey K. N. et al., *J. Biol. Chem.*, 2002, 277(7):4618-4627; Pandey K. N. et al., *Biochem. J.*, 2004, Dec. 1, Epub ahead of print; Roueau N. et al., Poster #P10144, "Development of a Non-radioactive Homogenous HTS Platform to Measure the Activity of Guanylate Cyclase", Presented at 10th Annual SBS Conference and Exhibition, Orlando, Fla., Sep. 11-15, 2004, PERKINELMER BIOSIGNAL Inc., Canada) each of which is incorporated herein by reference in its entirety). Functional truncations of NPR-A may also be used in the method of the invention (Pandey K. N. et al., *Molecular Pharmacology*, 2000, 57:259-267, which is incorporated herein by reference in its entirety). For example, using the AlphaScreen, a very sensitive assay platform capable of detecting fmol levels of non-acetylated cGMP has been developed (Rouveau et al., 2004). A biotinylated derivative of cGMP can be used as a tracer in a competitive immunoassay format involving rabbit anti cGMP antibodies. The AlphaScreen signal is generated when streptavidin coated Donor beads and protein A coated Acceptor beads are brought into proximity by the formation of the biotin-cGMP/anti-cGMP IgG complex. Production of cGMP by either particulate or soluble forms of guanylate cyclase leads to a decrease of the AlphaScreen signal by inhibiting the formation of the biotin-cGMP/anti-cGMP IgG complex. Using this assay, the activity of the atrial natriuretic peptide receptor (NPR-A, particulate guanylate cyclase) overexpressed in CHO cells has been characterized as well as that of soluble guanylate cyclase. Pharmacological parameters and Z' values obtained indicate that the assay platform is amenable to HTS.

In addition to determining whether an agent reduces NPR-A activity in vitro (e.g., in a cellular or acellular assay) and/or in vivo (in a human or non-human patient, or an animal model), the method may further comprise determining whether the agent reduces the physiological effects or symptoms associated with an inflammatory disorder and/or cell proliferation disorder, such as cancer, in vitro and/or in vivo (e.g., in an animal model). For example, the method may further comprise determining whether the agent has an apoptotic effect on cancer cells in vitro. These steps may be carried out before, during, or after NPR-A activity is assayed.

Contacting steps in the assays (methods) of the invention can involve combining or mixing the candidate agent and the cell in a suitable receptacle, such as a reaction vessel, microvessel, tube, microtube, well, or other solid support. Host cells and/or candidate agents may be arrayed on a solid support, such as a multi-well plate. "Arraying" refers to the act of organizing or arranging members of a library, or other collection, into a logical or physical array. Thus, an "array" refers to a physical or logical arrangement of, e.g., library members (candidate agent libraries). A physical array can be any "spatial format" or physically gridded format" in which physical manifestations of corresponding library members are arranged in an ordered manner, lending itself to combinatorial screening. For example, samples corresponding to individual or pooled members of a candidate agent library can

be arranged in a series of numbered rows and columns, e.g., on a multiwell plate. Similarly, host cells can be plated or otherwise deposited in microtitered, e.g., 96-well, 384-well, or -1536 well, plates (or trays). Optionally, host cells may be immobilized on the solid support.

A "solid support" (also referred to herein as a "solid substrate") has a fixed organizational support matrix that preferably functions as an organization matrix, such as a microtiter tray. Solid support materials include, but are not limited to, glass, polycryloylmorpholide, silica, controlled pore glass (CPG), polystyrene, polystyrene/latex, polyethylene, polyamide, carboxyl modified teflon, nylon and nitrocellulose and metals and alloys such as gold, platinum and palladium. The solid support can be biological, non-biological, organic, inorganic, or a combination of any of these, existing as particles, strands, precipitates, gels, sheets, tubing, spheres, containers, capillaries, pads, slices, films, plates, slides, etc., depending upon the particular application. Other suitable solid substrate materials will be readily apparent to those of skill in the art. The surface of the solid substrate may contain reactive groups, such as carboxyl, amino, hydroxyl, thiol, or the like for the attachment of nucleic acids, proteins, etc. Surfaces on the solid substrate will sometimes, though not always, be composed of the same material as the substrate. Thus, the surface can be composed of any of a wide variety of materials, for example, polymers, plastics, resins, polysaccharides-silica or silica-based materials, carbon, metals, inorganic glasses, membranes, or any of the above-listed substrate materials.

Measurement of NPR-A gene expression can be carried out using RT-PCR, for example. Screening of candidate agents or treatments (e.g., determination of NPR-A receptor activity) can be performed in a high-throughput format using combinatorial libraries, expression libraries, and the like. Other assays can be carried out on the host cells before, during, and/or after detection of NPR-A activity, and any or all assays may be carried out in an automated fashion, in a high-throughput format.

Alternatively, the aforementioned methods can be modified through the use of a cell-free assay. For example, instead of determining whether NPR-A activity in host cells is reduced by a candidate agent, extracts from host cells may be utilized and a fluorochrome or other detectable moiety can be associated with a nanoparticle or bead.

Once an agent has been determined to be one which reduces NPR-A activity, the agent can be combined with a pharmaceutically acceptable carrier. The method may further include a step of manufacturing the agent. The method may further include the step of packaging the agent.

Various methods of the present invention can include a step that involves comparing a value, level, feature, characteristic, property, etc. to a "suitable control", referred to interchangeably herein as an "appropriate control". A "suitable control" or "appropriate control" is any control or standard familiar to one of ordinary skill in the art useful for comparison purposes. In one embodiment, a "suitable control" or "appropriate control" is a value, level, feature, characteristic, property, etc. determined before, during, or after contacting an NPR-A receptor with a candidate agent, as described herein. For example, a transcription rate, mRNA level, translation rate, protein level, biological activity, cellular characteristic or property, genotype, phenotype, etc. can be determined prior to introducing a candidate into a cell or organism. In another embodiment, a "suitable control" or "appropriate control" is a value, level, feature, characteristic, property, etc. determined in a cell or organism, e.g., a control or normal cell or organism, exhibiting, for example, normal traits. In yet

another embodiment, a "suitable control" or "appropriate control" is a predefined value, level, feature, characteristic, property, etc.

Measuring expression includes determining or detecting the amount of the polypeptide present in a cell or shed by it, as well as measuring the underlying mRNA, where the quantity of mRNA present is considered to reflect the quantity of polypeptide manufactured by the cell. Furthermore, the gene for the NPR-A can be analyzed to determine whether there is a gene defect responsible for aberrant expression or polypeptide activity.

Polypeptide detection can be carried out by any available method, e.g., by Western blots, ELISA, dot blot, immunoprecipitation, RIA, immunohistochemistry, etc. For instance, a tissue section can be prepared and labeled with a specific antibody (indirect or direct and visualized with a microscope. Amount of a polypeptide can be quantitated without visualization, e.g., by preparing a lysate of a sample of interest, and then determining by ELISA or Western the amount of polypeptide per quantity of tissue. Antibodies and other specific binding agents can be used. There is no limitation on how detection of NPR-A activity is performed.

Assays can be utilized which permit quantification and/or presence/absence detection of a target nucleic acid (e.g., NPR-A) in a sample. Assays can be performed at the single-cell level, or in a sample comprising many cells, where the assay is "averaging" expression over the entire collection of cells and tissue present in the sample. Any suitable assay format can be used, including, but not limited to, e.g., Southern blot analysis, Northern blot analysis, polymerase chain reaction ("PCR") (e.g. Saiki et al., *Science* 1988, 241, 53; U.S. Pat. Nos. 4,683,195, 4,683,202, and 6,040,166; PCR Protocols: A Guide to Methods and Applications, Innis et al., eds., Academic Press, New York, 1990), reverse transcriptase polymerase chain reaction ("RT-PCR"), anchored PCR, rapid amplification of cDNA ends ("RACE") (e.g., Schaefer in *Gene Cloning and Analysis: Current Innovations*, Pages 99-115, 1997), ligase chain reaction ("LCR") (EP 320 308), one-sided PCR (Ohara et al., *Proc. Natl. Acad. Sci.* 1989, 86, 5673-5677), indexing methods (e.g., U.S. Pat. No. 5,508,169), in situ hybridization, differential display (e.g., Liang et al., *Nucl. Acid. Res.* 1993, 21, 3269-3275; U.S. Pat. Nos. 5,262,311, 5,599,672 and 5,965,409; WO97/18454; Prashar and Weissman, *Proc. Natl. Acad. Sci.*, 93:659-663, and U.S. Pat. Nos. 6,010,850 and 5,712,126; Welsh et al., *Nucleic Acid Res.*, 20:4965-4970, 1992, and U.S. Pat. No. 5,487,985) and other RNA fingerprinting techniques, nucleic acid sequence based amplification ("NASBA") and other transcription based amplification systems (e.g., U.S. Pat. Nos. 5,409,818 and 5,554,527; WO 88/10315), polynucleotide arrays (e.g., U.S. Pat. Nos. 5,143,854, 5,424,186; 5,700,637, 5,874,219, and 6,054,270; PCT WO 92/10092; PCT WO 90/15070), Qbeta Replicase (PCT/US87/0880), Strand Displacement Amplification ("SDA"), Repair Chain Reaction ("RCR"), nuclease protection assays, subtraction-based methods, Rapid-Scan, etc. Additional useful methods include, but are not limited to, e.g., template-based amplification methods, competitive PCR (e.g., U.S. Pat. No. 5,747,251), redox-based assays (e.g., U.S. Pat. No. 5,871,918), Taqman-based assays (e.g., Holland et al., *Proc. Natl. Acad. Sci.* 1991, 88, 7276-7280; U.S. Pat. Nos. 5,210,015 and 5,994,063), real-time fluorescence-based monitoring (e.g., U.S. Pat. No. 5,928,907), molecular energy transfer labels (e.g., U.S. Pat. Nos. 5,348,853, 5,532,129, 5,565,322, 6,030,787, and 6,177,635; Tyagi and Kramer, *Nature Biotech.*, 14:303-309, 1996). Any method suitable for single cell analysis of gene or protein expression can be used, including in situ hybridization,

immunocytochemistry, MACS, FACS, flow cytometry, etc. For single cell assays, expression products can be measured using antibodies, PCR, or other types of nucleic acid amplification (e.g., Brady et al., *Methods Mol. & Cell. Biol.* 1990, 2, 17-25; Eberwine et al., *Proc. Natl. Acad. Sci.* 1992, 89, 3010-3014; U.S. Pat. No. 5,723,290). These and other methods can be carried out conventionally, e.g. as described in the mentioned publications.

The terms “transfection”, “transformation”, and “introduction”, and grammatical variations thereof, are used interchangeably herein to refer to the insertion of an exogenous polynucleotide (e.g., a nucleic acid sequence encoding an NP, or fragment, homolog, or variant thereof, or a nucleic acid sequence encoding an NPR-A, or fragment, homolog, or variant thereof, into a host cell, irrespective of the method used for the insertion, the molecular form of the polynucleotide that is inserted, or the nature of the cell (e.g., prokaryotic or eukaryotic). The insertion of a polynucleotide per se and the insertion of a plasmid or vector comprised of the exogenous polynucleotide are included. The exogenous polynucleotide may be directly transcribed and translated by the cell, maintained as a nonintegrated vector, for example, a plasmid, or alternatively, may be stably integrated into the host genome. Thus, host cells of the invention include those that have been transfected with polynucleotides encoding an NP, or fragment, variant, or homolog thereof, and those that have been transfected with polynucleotides encoding an NPR-A, or fragment, variant, or homolog thereof.

The phrases “isolated” or “biologically pure” refer to material that is substantially or essentially free from components which normally accompany the material as it is found in its native state.

An “isolated polynucleotide” that encodes a particular polypeptide refers to a polynucleotide that is substantially free of other nucleic acid molecules that do not encode the subject polypeptide; however, the molecule may include functionally and/or structurally conservative mutations as defined herein.

The terms “cell” and “cells” are used interchangeably herein to refer to a single cell or plurality of cells (i.e., at least one cell). Typically, host cells used in the methods of the invention are isolated. However, tissues, and genetically modified or transgenic animals may also be utilized.

The terms “comprising”, “consisting of” and “consisting essentially of” are defined according to their standard meaning. The terms may be substituted for one another throughout the instant application in order to attach the specific meaning associated with each term.

As used in this specification, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to “a cell” includes more than one such cell. Reference to “a receptor” includes more than one such receptor. Reference to “a polynucleotide” includes more than one such polynucleotide. Reference to “a polypeptide” or “agent” includes more than one such polypeptide or agent, and the like.

The practice of the present invention can employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA technology, electrophysiology, and pharmacology, that are within the skill of the art. Such techniques are explained fully in the literature (see, e.g. Sambrook, Fritsch & Maniatis. *Molecular Cloning: A Laboratory Manual*, Second Edition (1989); DNA Cloning, Vols. I and II (D. N. Glover ed. 1985); Perbal, B., *A Practical Guide to Molecular Cloning* (1984); the series, *Methods In Enzymology* (S. Colowick and N. Kaplan eds., Academic Press, Inc.); *Transcription and Translation* (Hames et al. eds.

1984); *Gene Transfer Vectors For Mammalian Cells* (J. H. Miller et al. eds. (1987) Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.); *Scopes, Protein Purification Principles and Practice* (2nd ed., Springer-Verlag); and *PCR: A Practical Approach* (McPherson et al. eds. (1991) IRL Press)).

Example 1

pNP 73-102 Inhibits NPRA Expression

The structures of ANP and ANP like molecules with their ring-structure and receptors associated with it are well characterized. However, the N-terminal peptides do not have this structure. Neither KP nor NP73-102 was shown to bind ANP receptor NPRA (Mohapatra et al., *J Allergy Clin Immunol*, 2004, 114:520-526). The receptors for NP-73-102 are not known.

The highest expression of the ANP and ANP receptors is found in neonatal thymus. To test whether the peptide NP73-102 inhibits in vivo the ANP cascade, pregnant (12 days) mice were injected i.p. with pVAX (vector), or pNP73-102. After 1 day, mice were sacrificed and thymi removed from embryo, were homogenized. Cells were centrifuged and erythrocytes lysed by treating the suspension with ACK buffer. Cells were incubated with anti-NPRA or anti-NPRC antibodies for 1 hour, washed and incubated with PE-conjugated 20 Ab. Levels of NPR's were determined by flow cytometry. The results are shown in FIG. 1. The results demonstrate that pNP73-102 inhibited expression of NPRA in thymocytes. Although the mechanism is not clear, this may be due to feedback inhibition at the level intracellular signaling occurring via NPRA.

Example 2

NPRA Deficiency Decreases Pulmonary Inflammation

Development and chronicity of cancers has been attributed to the chronic inflammation in the affected organs. ANP was reported to have anti-inflammatory activity, although signaling through NPRA is known to cause a number of different biological activity including cell proliferation, immune activation, inflammation and apoptosis. To determine the role of NPRA signaling in the lung inflammation, groups (n=3) of wild type DBA/2 (wt) and NPR-C (ko) deficient mice and wild type C57/BL6 (wt) and NPR-A (ko) were sensitized with ovalbumin (20 mg/mouse) and after 2 weeks challenged i.n. with ovalbumin (20 mg/mouse). One day later, mice were sacrificed and lung sections were stained with H & E to examine inflammation. As shown in FIGS. 2A-2D, there was no significant difference in pulmonary inflammation between the wild-type and NPRC deficient mice. In sharp contrast, a comparison between wild-type C57BL6 and NPRA deficient mice showed that NPRA deficient mice had substantially reduced inflammation compared to wild type. These results indicate that ANP-NPRA signaling is involved in increasing inflammation in the lung.

Example 3

A549 Cells Transfected with pNP₇₃₋₁₀₂ Show a Significantly Higher Level of Apoptosis Compared Control and pANP or pVAX

To determine the effect of overexpression of NP73-102 on proliferation of A549 lung epithelial cells, cells were transfected with either pNP73-102 or vector, pVAX. Cell cycle

45

analysis was performed using propidium iodide (PI) staining and flow cytometry 48 h after transfection. No significant difference was observed between control and pNP73-102-transfected cells in S1, Go-G1 and G2-M stages of cell cycle (data not shown). However, an analysis of apoptosis using flow-cytometry with PI and annexin V, showed that cells transfected with pNP73-102 exhibited significantly higher apoptosis compared to cells transfected with either the control plasmid or a plasmid encoding ANP (FIGS. 3A-3C). This result was confirmed by (i) staining by TUNEL of A549 cells cultured in 8-chamber slide following a 48-hour transfection with either pANP or pNP73-102 (not shown), (ii) by analysis of PARP cleavage in these cells 48 hours after transfection, which was significantly more prominent in pNP73-102 transfected cells (FIG. 3D). The results show that pNP73-102 shows a higher accumulation of apoptotic cells compared to cells transfected with pANP and pVAX controls. Thus, pNP73-102 induces apoptosis of lung adenocarcinoma cells.

In an effort to identify and characterize molecules participating in early signaling pathways, differential gene expression was analyzed using a microarray (AFFYMETRIX). Altered expression of a large number of genes was found, including genes related to cell growth, cell cycle, and apoptosis. These genes included, among others more than, 6- to 8-fold up-regulation of genes such as Caspase (Casp)-8 and FADD like apoptosis regulator, cyclin E binding protein, CDK inhibitor 1A, CDK7, casp4, casp-10, casp-1, apoptosis facilitator BCL2-like 13 and annexin 43 (data not shown). Together, these studies indicate that pNP73-102 is an inducer of apoptosis in A549 lung adenocarcinoma cells.

Example 4

pNP73-102 Decreases Tumorigenesis in a Colony Formation Assay by A549

To test the anti-cancer activity of the pNP73-102 construct, a colony forming assay was undertaken. Thus, six cm tissue culture plates were covered with 4 ml of 0.5% soft agar. A549 cells were transfected with pANP, pNP₇₃₋₁₀₂ and pVAX plasmid DNA. After 40 hours of transfection, equal number of cells were suspended in 2 ml of 0.3% soft agar and added to each plate. Cells were plated in duplicate at a density of 2×10^4 cells/dish and incubated for two weeks. Plates were observed and photographed under a microscope. Cell colonies were counted and plotted. The results of one representative experiment of two experiments performed is shown in FIGS. 5A-5D. The results show that plasmid vector alone caused some reduction in colony formation compared to untransfected control. However, both ANP and pNP₇₃₋₁₀₂ showed substantial reductions in the number of colonies produced compared to vehicle control.

Example 5

Chitosan Nanoparticle Containing pNP₇₃₋₁₀₂ Substantially Decrease Tumor Development in the Lung

To test the effect of de novo expression of pNP₇₃₋₁₀₂, the plasmid was coacervated with chitosan nanoparticles, referred to as CPNP73-102. To examine expression of NP73-102 from CPNP73-102, a construct was developed that carried a C-terminal fusion of marker peptide of FLAG. BALB/c mice were given intranasally the NP73-102-FLAG and the expression of NP73-102-FLAG in the BAL cells after i.n. administration of CPNP73-102-FLAG peptide. A bronchial

46

lavage was performed after 24 hours and lavage cells were stained with either the second antibody control or anti-FLAG antibody (Sigma) and then with DAPI. The results show that intranasal administration induces significant expression of the peptide in the lung cells.

To test whether CPNP73-102 is capable of decreasing tumor formation in the lung, BALB/c nude mice were injected i.v. with 5×10^6 A549 cells, then treated one day afterwards and at weekly intervals with CPNP73-102 or control plasmid. After 4 weeks, mice were examined for lung histology. The control animals showed tumors, whereas no tumors were observed in the CPNP73-102-treated group. Sections were also stained with antibodies to cyclinB and to phospho-Bad. The results show that mice treated with CPNP73-102 had no tumors in the lung and did not show any staining for pro-mitotic Cyclin-B and anti-apoptotic marker phospho-Bad. These results indicate that CPNP73-102 has the potential to decrease tumor formation in the lung.

Example 6

Treatment with CPNP73-102 Decreases the Tumor Burden in a Spontaneous Tumorigenesis Model of Immunocompetent BALB/c Mice

The nude mouse model is deemed to be of less predictive value in terms of translating to human cancer, as mice used are immunodeficient. Therefore, to confirm the results obtained on the potential role of pNP73-102, a syngeneic immunocompetent mouse model of human lung carcinoma was used. For this purpose, Line-1 cell line derived from a bronchioalveolar cell carcinoma (a subtype of lung adenocarcinoma that spontaneously arose in BALB/C mouse (Yuhás et al., *Cancer Research*, 1975, 35:242-244). The cell line forms subcutaneous tumors within 2 to 3 weeks of injection and spontaneously metastasizes to the lung.

To examine whether de novo synthesis of NP73-102 affects tumor development, two groups of BALB/c mice ($n=4$) were administered with the Line-1 tumor cells (100,000 cells/mouse) at the flanks. One group was administered intranasally with CPNP73-102 the same day, whereas another group was administered with vehicle alone (nanoparticle carrying a plasmid without NP73-102), and the third group was given the saline. Treatment was continued with NP73-102 or controls at weekly intervals for 5 weeks. The tumors were dissected out from each group of mice and photographed (FIGS. 6A-6C) and the tumor burden was calculated by weighing them on a balance (FIG. 6D). The results show that mice administered with CPNP73-102 had significantly decreased tumor burden ($P<0.05$).

Example 7

pNP73-102 Induces Apoptosis in Chemoresistant Ovarian Cancer Cells

The adenocarcinomas of various tissues such as lung, ovary, and breasts have many characteristics that are similar. Chemoresistance is a major therapeutic problem in many of the cancers and the current knowledge on cellular mechanisms involved is incomplete. Since A549 cells showed differential sensitivity to apoptosis with pVAX and pNP₇₃₋₁₀₂ the effects of pNP73-102 was tested using chemosensitive (OV2008) and chemoresistant (C13) ovarian cancer cells. C-13 and OV2008 ovarian cancer cells were transfected with pNP73-102 or with pVAX as control. Forty-eight hours later, cells were processed to examine apoptosis by TUNEL assay

(FIG. 7). The results showed that either of the cells when transfected with pVAX did not exhibit any apoptosis. In contrast, both cell lines exhibited apoptosis as evident from TUNEL positive cells. These results indicate that pNP73-102 may induce apoptosis of epithelial adenocarcinomas irrespective of their degree of chemo-sensitivity.

Example 8

MCF-7 Breast Cancer Cells are Also Affected by NP73-102

The effects of de novo synthesis of NP₇₃₋₁₀₂ was examined on the proliferation of the MCF-7 breast cancer cells. Cells were transfected with pVAX, pANP, or pANP₇₃₋₁₀₂. The cells were counted 24 and 48 hours after transfection and their viability was examined by trypan blue staining. The results shown in FIG. 8 indicate that there was a substantial reduction of viable cell numbers in cells transfected with pNP₇₃₋₁₀₂ compared to cells transfected with pANP or control empty vector. To further verify whether this is due to a defect in cell cycle or induction of apoptosis, a cell cycle analysis was undertaken. MCF-7 cells were transfected with pVAX or

pANP₇₃₋₁₀₂ and DNA analysis was undertaken by PI staining 48 hours after transfection. Cells transfected with empty vector plasmid as control showed 37.99% cells in G0-G1, 11.28% in G2-M and 50.73% cells in G2-G1 phase. In contrast, cells transfected with pANP₇₃₋₁₀₂ showed 66.01% cells in G0-G1, 7.07% in G2-M, and 26.91% cells in G2-G1 phase. Transfection with pANP showed results similar to the pNP₇₃₋₁₀₂. These results indicate that both pANP and pNP₇₃₋₁₀₂ expression arrests cells in G0-G1 and blocks progression to S phase, suggesting that treatment with pANP and pNP₇₃₋₁₀₂ or the corresponding peptides may be useful in breast cancer patients.

All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety, including all figures, tables, and sequences, to the extent they are not inconsistent with the explicit teachings of this specification.

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 30

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Asn Pro Met Tyr Asn Ala Val Ser Asn Ala Asp Leu Met Asp Phe Lys
1 5 10 15

Asn Leu Leu Asp His Leu Glu Glu Lys Met Pro Leu Glu Asp
20 25 30

<210> SEQ ID NO 2

<211> LENGTH: 37

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Glu Val Val Pro Pro Gln Val Leu Ser Glu Pro Asn Glu Glu Ala Gly
1 5 10 15

Ala Ala Leu Ser Pro Leu Pro Glu Val Pro Pro Trp Thr Gly Glu Val
20 25 30

Ser Pro Ala Gln Arg
35

<210> SEQ ID NO 3

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Ser Ser Asp Arg Ser Ala Leu Leu Lys Ser Lys Leu Arg Ala Leu Leu
1 5 10 15

Thr Ala Pro Arg
20

-continued

<210> SEQ ID NO 4
 <211> LENGTH: 28
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Ser Leu Arg Arg Ser Ser Cys Phe Gly Gly Arg Met Asp Arg Ile Gly
 1 5 10 15

Ala Gln Ser Gly Leu Gly Cys Asn Ser Phe Arg Tyr
 20 25

<210> SEQ ID NO 5
 <211> LENGTH: 30
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

Gly Ser Pro Trp Asp Pro Ser Asp Arg Ser Ala Leu Leu Lys Ser Lys
 1 5 10 15

Leu Arg Ala Leu Leu Ala Gly Pro Arg Ser Leu Arg Arg Ser
 20 25 30

<210> SEQ ID NO 6
 <211> LENGTH: 37
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Val Ser Asn Thr Asp Leu Met Asp Phe Lys Asn Leu Leu Asp His Leu
 1 5 10 15

Glu Glu Lys Met Pro Val Glu Asp Glu Val Met Pro Pro Gln Ala Leu
 20 25 30

Ser Glu Gln Thr Glu
 35

<210> SEQ ID NO 7
 <211> LENGTH: 151
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Met Ser Ser Phe Ser Thr Thr Thr Val Ser Phe Leu Leu Leu Leu Ala
 1 5 10 15

Phe Gln Leu Leu Gly Gln Thr Arg Ala Asn Pro Met Tyr Asn Ala Val
 20 25 30

Ser Asn Ala Asp Leu Met Asp Phe Lys Asn Leu Leu Asp His Leu Glu
 35 40 45

Glu Lys Met Pro Leu Glu Asp Glu Val Val Pro Pro Gln Val Leu Ser
 50 55 60

Glu Pro Asn Glu Glu Ala Gly Ala Ala Leu Ser Pro Leu Pro Glu Val
 65 70 75 80

Pro Pro Trp Thr Gly Glu Val Ser Pro Ala Gln Arg Asp Gly Gly Ala
 85 90 95

Leu Gly Arg Gly Pro Trp Asp Ser Ser Asp Arg Ser Ala Leu Leu Lys
 100 105 110

Ser Lys Leu Arg Ala Leu Leu Thr Ala Pro Arg Ser Leu Arg Arg Ser
 115 120 125

-continued

Ser Cys Phe Gly Gly Arg Met Asp Arg Ile Gly Ala Gln Ser Gly Leu
 130 135 140

Gly Cys Asn Ser Phe Arg Tyr
 145 150

<210> SEQ ID NO 8

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 8

gacggcaagc ttactatggg cagcccctgg gaccc 35

<210> SEQ ID NO 9

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 9

acccccctcg agttattatc ttcgtaggct ccg 33

<210> SEQ ID NO 10

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 10

aatcctaagc ttagtatggt gtccaacaca gat 33

<210> SEQ ID NO 11

<211> LENGTH: 41

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 11

tgcgaactcg agttactcag tctgctcact cagggcctgc g 41

<210> SEQ ID NO 12

<211> LENGTH: 90

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 12

ggcagcccct gggacccctc cgatagatct gccctcttga aaagcaaaact gagggctctg 60

ctcgtctggcc ctcgagacct acgaagatcc 90

<210> SEQ ID NO 13

<211> LENGTH: 117

<212> TYPE: DNA

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 13

atggtgtcca acacagatct gatggatttc aagaacctgc tagaccacct ggaggagaag 60

atgccggtag aagatgaggt catgcccccg caggccctga gtgagcagac tgagtaa 117

<210> SEQ ID NO 14

<211> LENGTH: 845

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 14

tggcgaggga cagacgtagg ccaagagagg ggaaccagag aggaaccaga ggggagagac	60
agagcagcaa gcagtggatt gtccttgac gacgccagca tgagctcctt ctccaccacc	120
accgtgagct tcctcctttt actggcattc cagctcctag gtcagaccag agctaataccc	180
atgtacaatg ccgtgtccaa cgcagacctg atggatttca agaatttgct ggaccatttg	240
gaagaaaaga tgcctttaga agatgaggtc gtgccccac aagtgtcag tgagccgaat	300
gaagaagcgg gggctgtctc cagccccctc cctgagggtc ctccctggac cggggaagtc	360
agcccagccc agagagatgg aggtgccctc gggcggggcc cctgggactc ctctgatcga	420
tctgccctcc taaaaagcaa gctgagggcg ctgctcactg cccctcggag cctgcggaga	480
tccagctgct tcgggggcag gatggacagg attggagccc agagcggact gggctgtaac	540
agcttcctgg actgaagata acagccaggg aggacaagca gggctgggcc tagggacaga	600
ctgcaagagg ctctgtctcc ctgggggtct tgctgcattt gtgtcatctt gttgccatgg	660
agtttgtgat atcccatcta agctgcagct tcctgtcaac acttctcaca tcttatgcta	720
actgtagata aagtggtttg atgggtgactt cctcgctct cccaccccat gcattaaatt	780
ttaaggtaga acctcacctg ttactgaaag tggtttgaaa gtgaataaac ttcagcacca	840
tggac	845

<210> SEQ ID NO 15

<211> LENGTH: 2583

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

ggatccattt gtctcgggct gctggctgcc tgccatttcc tcctctccac ccttatttgg	60
aggccctgac agctgagcca caaacaacc aggggagctg ggcaccagca agcgtcacc	120
tctgtttccc cgcacggtac cagcgtcag gagaaagaat cctgaggcac ggcggtgaga	180
taaccaagga ctctttttta ctcttctcac acctttgaag tgggagcctc ttgagtcaaa	240
tcagtaagaa tgcggctctt gcagctgagg gtctgggggg ctgttggggc tgcccaaggc	300
agagaggggc tgtgacaagc cctgcggatg ataactttaa aagggcatct cctgctggct	360
tctcacttgg cagctttatc actgcaagt acagaatggg gagggttctg tctctcctgc	420
gtgcttgag agctgggggg ctataaaaag aggcggcact gggcagctgg gagacaggga	480
cagacgtagg ccaagagagg ggaaccagag aggaaccaga ggggagagac agagcagcaa	540
gcagtggatt gtccttgac gacgccagca tgagctcctt ctccaccacc accgtgagct	600
tcctcctttt actggcattc cagctcctag gtcagaccag agctaataccc atgtacaatg	660
ccgtgtccaa cgcagacctg atggatttca aggtagggcc aggaaagcgg gtgcagtctg	720
gggccagggg gctttctgat gctgtgctca ctctcttga tttcctccaa gtcagtggag	780
tttatccctt tcctgtatt ttcttttct aaagaatttg ctggaccatt tggaagaaaa	840
gatgccttta gaagatgagg tcgtgcccc acaagtgtc agtgagccga atgaagaagc	900
gggggctgct ctacgcccc tcctgaggt gcctccctgg accggggaag tcagcccagc	960
ccagagagat ggagggtccc tcgggcgggg cccctgggac tcctctgac gatctgccct	1020
cctaaaaagc aagctgaggg cgctgctcac tgccctcgg agcctgcgga gatccagctg	1080
cttcgggggc aggatggaca ggattggagc ccagagcggg ctgggctgta acagcttccg	1140
ggtaagagga actggggatg gaaatgggat gggatggaca ctactgggag acaccttcag	1200

-continued

caggaaggg accaatgcag aagctcatc cctctcaagt ttctgcccc acaccagag	1260
tgccccatgg gtgtcaggac atgccatcta ttgtccttag ctagtctgct gagaaatgc	1320
ttaaaaaaaaa aagggggggg gctgggcacg gtcgtcacgc ctgtaatccc agcactttgg	1380
gaggccaggc agcggatcat gaggtcaaga gatcaagact atcctggcca acatggtgaa	1440
acccagctc tactaaaaat acaaaaatta gctgggtgtg tggcgggcac ctgtactctc	1500
agctacttgg gaggtgagg caggagaatc acttgaacct aggaggcaga ggttgcagtg	1560
agcagagatc acgccactgc agtccagcct aggtgataga gcgagactgt ctcaaaaaa	1620
aaaaaaaaag gccaggcgcg gtgggtcacg cctgtaatcc cagcgctttg ggaggccaag	1680
gcgggtggat cacgaggtca ggagatggag accatcctgg ctaacacggt gaaaccccg	1740
ctctactaaa aatacaaaaa attagccagg cgtggtggca ggccctgta agtcctagct	1800
actccggagg ctgaggcagg agaatggcgt gaacccggga ggccgagctt gcagtgagca	1860
gagatggcac cactgcactc cagcctgggc gacagagcaa gactccgtct caaaaaaaaa	1920
aaaaaaaaa gcaactgcc ctagcactgg gaaattaaaa tattcataga gccaagtatt	1980
ctttgcatgg ctgattagca gttcatattc ctccccagaa ttgcaagatc ctgaagggt	2040
taagtgaat ttactctgat gagtaacttg cttatcaatt catgaagctc agagggtcat	2100
caggctgggg tgggggcccgg tgggaagcag gtggtcagta atcaagttca gaggatggc	2160
acactcatc atgaagctga cttttccagg acagccagg caccaagcca gatatgtctg	2220
tgttctcttt gcagtactga agataacagc cagggaggac aagcagggtt gggcctagg	2280
acagactgca agaggctcct gtcccctggg gtctctgctg catttggtgc atctgtgtg	2340
catggagttg tgatcatccc atctaagctg cagcttctcg tcaacacttc tcacatctta	2400
tgctaactgt agataaagtg gtttgatggt gacttctcgc cctctcccac cccatgcatt	2460
aaattttaag gtagaacctc acctgttact gaaagtgggt tgaaagtga taaacttcag	2520
caccatggac agaagacaaa tgctgcgtt ggtgtgcttt ctttctctt gggaagagaa	2580
ttc	2583

<210> SEQ ID NO 16

<211> LENGTH: 152

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 16

Met Gly Ser Phe Ser Ile Thr Leu Gly Phe Phe Leu Val Leu Ala Phe	
1 5 10 15	
Trp Leu Pro Gly His Ile Gly Ala Asn Pro Val Tyr Ser Ala Val Ser	
20 25 30	
Asn Thr Asp Leu Met Asp Phe Lys Asn Leu Leu Asp His Leu Glu Glu	
35 40 45	
Lys Met Pro Val Glu Asp Glu Val Met Pro Pro Gln Ala Leu Ser Glu	
50 55 60	
Gln Thr Glu Glu Ala Gly Ala Ala Leu Ser Ser Leu Pro Glu Val Pro	
65 70 75 80	
Pro Trp Thr Gly Glu Val Asn Pro Pro Leu Arg Asp Gly Ser Ala Leu	
85 90 95	
Gly Arg Ser Pro Trp Asp Pro Ser Asp Arg Ser Ala Leu Leu Lys Ser	
100 105 110	
Lys Leu Arg Ala Leu Leu Ala Gly Pro Arg Ser Leu Arg Arg Ser Ser	
115 120 125	

-continued

Cys Phe Gly Gly Arg Ile Asp Arg Ile Gly Ala Gln Ser Gly Leu Gly
 130 135 140

Cys Asn Ser Phe Arg Tyr Arg Arg
 145 150

<210> SEQ ID NO 17
 <211> LENGTH: 878
 <212> TYPE: DNA
 <213> ORGANISM: Mus musculus
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (81)..(539)
 <223> OTHER INFORMATION: coding sequence for preproANP

<400> SEQUENCE: 17

```

caaaagctga gagagagaga gaaagaaacc agagtgggca gagacagcaa acatcagatc      60
gtgccccgac ccacgccagc atgggctcct tctccatcac cctgggcttc ttctcgtct      120
tggccttttg gcttcaggc catattggag caaatcctgt gtacagtgcg gtgtccaaca      180
cagatctgat ggatttcaag aacctgctag accacctgga ggagaagatg ccggtagaag      240
atgaggtcat gcccccgag gccctgagtg agcagactga ggaagcaggg gccgcactta      300
gtccctccc cgaggtgctt ccttgactg gggaggtcaa cccacctctg agagacggca      360
gtgctctagg gcgcagcccc tgggaccctt ccgatagatc tgccctcttg aaaagcaaac      420
tgagggctct gctcgtggc cctcgagacc tacgaagatc cagctgcttc gggggtagga      480
ttgacaggat tggagcccag agtgactag gctgcaacag ctcccggtac cgaagataac      540
agccaaggag gaaaaggcag tcgattctgc ttgagcagat cgcaaaagat cctaagccct      600
tgtggtgtgt cagcagctt ggtcacattg ccaactgtgc gtggtgaaca cctcctgga      660
gtgcggtt cctgccttca tctatcacga tcgatgttaa atgtagatga gtggtctagt      720
ggggtcttgc ctctccact ctgcatatta aggtagatcc tcaccctttt cagaaagcag      780
ttgaaaaaaa aaaaaagaa taaacttcag caccaaggac agacgccgag gccctgatgt      840
gttcttttgg cttctgcctt cagttctttg ctctcccc      878

```

<210> SEQ ID NO 18
 <211> LENGTH: 1061
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

```

Met Pro Gly Pro Arg Arg Pro Ala Gly Ser Arg Leu Arg Leu Leu Leu
1      5      10      15
Leu Leu Leu Leu Pro Pro Leu Leu Leu Leu Arg Gly Ser His Ala
20     25     30
Gly Asn Leu Thr Val Ala Val Val Leu Pro Leu Ala Asn Thr Ser Tyr
35     40     45
Pro Trp Ser Trp Ala Arg Val Gly Pro Ala Val Glu Leu Ala Leu Ala
50     55     60
Gln Val Lys Ala Arg Pro Asp Leu Leu Pro Gly Trp Thr Val Arg Thr
65     70     75     80
Val Leu Gly Ser Ser Glu Asn Ala Leu Gly Val Cys Ser Asp Thr Ala
85     90     95
Ala Pro Leu Ala Ala Val Asp Leu Lys Trp Glu His Asn Pro Ala Val
100    105    110
Phe Leu Gly Pro Gly Cys Val Tyr Ala Ala Ala Pro Val Gly Arg Phe
115    120    125

```

-continued

Thr	Ala	His	Trp	Arg	Val	Pro	Leu	Leu	Thr	Ala	Gly	Ala	Pro	Ala	Leu
130						135					140				
Gly	Phe	Gly	Val	Lys	Asp	Glu	Tyr	Ala	Leu	Thr	Thr	Arg	Ala	Gly	Pro
145					150					155					160
Ser	Tyr	Ala	Lys	Leu	Gly	Asp	Phe	Val	Ala	Ala	Leu	His	Arg	Arg	Leu
				165					170					175	
Gly	Trp	Glu	Arg	Gln	Ala	Leu	Met	Leu	Tyr	Ala	Tyr	Arg	Pro	Gly	Asp
			180					185					190		
Glu	Glu	His	Cys	Phe	Phe	Leu	Val	Glu	Gly	Leu	Phe	Met	Arg	Val	Arg
		195					200					205			
Asp	Arg	Leu	Asn	Ile	Thr	Val	Asp	His	Leu	Glu	Phe	Ala	Glu	Asp	Asp
210						215					220				
Leu	Ser	His	Tyr	Thr	Arg	Leu	Leu	Arg	Thr	Met	Pro	Arg	Lys	Gly	Arg
225					230					235					240
Val	Ile	Tyr	Ile	Cys	Ser	Ser	Pro	Asp	Ala	Phe	Arg	Thr	Leu	Met	Leu
				245					250					255	
Leu	Ala	Leu	Glu	Ala	Gly	Leu	Cys	Gly	Glu	Asp	Tyr	Val	Phe	Phe	His
			260					265					270		
Leu	Asp	Ile	Phe	Gly	Gln	Ser	Leu	Gln	Gly	Gly	Gln	Gly	Pro	Ala	Pro
		275					280					285			
Arg	Arg	Pro	Trp	Glu	Arg	Gly	Asp	Gly	Gln	Asp	Val	Ser	Ala	Arg	Gln
290						295					300				
Ala	Phe	Gln	Ala	Ala	Lys	Ile	Ile	Thr	Tyr	Lys	Asp	Pro	Asp	Asn	Pro
305					310					315					320
Glu	Tyr	Leu	Glu	Phe	Leu	Lys	Gln	Leu	Lys	His	Leu	Ala	Tyr	Glu	Gln
			325					330						335	
Phe	Asn	Phe	Thr	Met	Glu	Asp	Val	Leu	Val	Asn	Thr	Ile	Pro	Ala	Ser
			340					345					350		
Phe	His	Asp	Gly	Leu	Leu	Leu	Tyr	Ile	Gln	Ala	Val	Thr	Glu	Thr	Leu
		355					360					365			
Ala	His	Gly	Gly	Thr	Val	Thr	Asp	Gly	Glu	Asn	Ile	Thr	Gln	Arg	Met
370						375					380				
Trp	Asn	Arg	Ser	Phe	Gln	Gly	Val	Thr	Gly	Tyr	Leu	Lys	Ile	Asp	Ser
385					390					395					400
Ser	Gly	Asp	Arg	Glu	Thr	Asp	Phe	Ser	Leu	Trp	Asp	Met	Asp	Pro	Glu
			405						410					415	
Asn	Gly	Ala	Phe	Arg	Val	Val	Leu	Asn	Tyr	Asn	Gly	Thr	Ser	Gln	Glu
			420					425					430		
Leu	Val	Ala	Val	Ser	Gly	Arg	Lys	Leu	Asn	Trp	Pro	Leu	Gly	Tyr	Pro
			435					440					445		
Pro	Pro	Asp	Ile	Pro	Lys	Cys	Gly	Phe	Asp	Asn	Glu	Asp	Pro	Ala	Cys
450						455					460				
Asn	Gln	Asp	His	Leu	Ser	Thr	Leu	Glu	Val	Leu	Ala	Leu	Val	Gly	Ser
465					470					475					480
Leu	Ser	Leu	Leu	Gly	Ile	Leu	Ile	Val	Ser	Phe	Phe	Ile	Tyr	Arg	Lys
			485					490						495	
Met	Gln	Leu	Glu	Lys	Glu	Leu	Ala	Ser	Glu	Leu	Trp	Arg	Val	Arg	Trp
			500					505					510		
Glu	Asp	Val	Glu	Pro	Ser	Ser	Leu	Glu	Arg	His	Leu	Arg	Ser	Ala	Gly
		515						520				525			
Ser	Arg	Leu	Thr	Leu	Ser	Gly	Arg	Gly	Ser	Asn	Tyr	Gly	Ser	Leu	Leu
530						535					540				
Thr	Thr	Glu	Gly	Gln	Phe	Gln	Val	Phe	Ala	Lys	Thr	Ala	Tyr	Tyr	Lys
545					550					555					560

-continued

Gly	Asn	Leu	Val	Ala	Val	Lys	Arg	Val	Asn	Arg	Lys	Arg	Ile	Glu	Leu
				565					570					575	
Thr	Arg	Lys	Val	Leu	Phe	Glu	Leu	Lys	His	Met	Arg	Asp	Val	Gln	Asn
			580					585					590		
Glu	His	Leu	Thr	Arg	Phe	Val	Gly	Ala	Cys	Thr	Asp	Pro	Pro	Asn	Ile
		595					600					605			
Cys	Ile	Leu	Thr	Glu	Tyr	Cys	Pro	Arg	Gly	Ser	Leu	Gln	Asp	Ile	Leu
	610					615					620				
Glu	Asn	Glu	Ser	Ile	Thr	Leu	Asp	Trp	Met	Phe	Arg	Tyr	Ser	Leu	Thr
625					630					635					640
Asn	Asp	Ile	Val	Lys	Gly	Met	Leu	Phe	Leu	His	Asn	Gly	Ala	Ile	Cys
				645					650					655	
Ser	His	Gly	Asn	Leu	Lys	Ser	Ser	Asn	Cys	Val	Val	Asp	Gly	Arg	Phe
			660					665					670		
Val	Leu	Lys	Ile	Thr	Asp	Tyr	Gly	Leu	Glu	Ser	Phe	Arg	Asp	Leu	Asp
		675					680					685			
Pro	Glu	Gln	Gly	His	Thr	Val	Tyr	Ala	Lys	Lys	Leu	Trp	Thr	Ala	Pro
	690					695					700				
Glu	Leu	Leu	Arg	Met	Ala	Ser	Pro	Pro	Val	Arg	Gly	Ser	Gln	Ala	Gly
705					710					715					720
Asp	Val	Tyr	Ser	Phe	Gly	Ile	Ile	Leu	Gln	Glu	Ile	Ala	Leu	Arg	Ser
				725					730					735	
Gly	Val	Phe	His	Val	Glu	Gly	Leu	Asp	Leu	Ser	Pro	Lys	Glu	Ile	Ile
			740					745					750		
Glu	Arg	Val	Thr	Arg	Gly	Glu	Gln	Pro	Pro	Phe	Arg	Pro	Ser	Leu	Ala
		755					760					765			
Leu	Gln	Ser	His	Leu	Glu	Glu	Leu	Gly	Leu	Leu	Met	Gln	Arg	Cys	Trp
	770					775					780				
Ala	Glu	Asp	Pro	Gln	Glu	Arg	Pro	Pro	Phe	Gln	Gln	Ile	Arg	Leu	Thr
785					790					795					800
Leu	Arg	Lys	Phe	Asn	Arg	Glu	Asn	Ser	Ser	Asn	Ile	Leu	Asp	Asn	Leu
				805				810						815	
Leu	Ser	Arg	Met	Glu	Gln	Tyr	Ala	Asn	Asn	Leu	Glu	Glu	Leu	Val	Glu
			820					825					830		
Glu	Arg	Thr	Gln	Ala	Tyr	Leu	Glu	Glu	Lys	Arg	Lys	Ala	Glu	Ala	Leu
		835					840					845			
Leu	Tyr	Gln	Ile	Leu	Pro	His	Ser	Val	Ala	Glu	Gln	Leu	Lys	Arg	Gly
	850					855					860				
Glu	Thr	Val	Gln	Ala	Glu	Ala	Phe	Asp	Ser	Val	Thr	Ile	Tyr	Phe	Ser
865					870					875					880
Asp	Ile	Val	Gly	Phe	Thr	Ala	Leu	Ser	Ala	Glu	Ser	Thr	Pro	Met	Gln
				885					890					895	
Val	Val	Thr	Leu	Leu	Asn	Asp	Leu	Tyr	Thr	Cys	Phe	Asp	Ala	Val	Ile
			900					905					910		
Asp	Asn	Phe	Asp	Val	Tyr	Lys	Val	Glu	Thr	Ile	Gly	Asp	Ala	Tyr	Met
		915					920					925			
Val	Val	Ser	Gly	Leu	Pro	Val	Arg	Asn	Gly	Arg	Leu	His	Ala	Cys	Glu
		930				935					940				
Val	Ala	Arg	Met	Ala	Leu	Ala	Leu	Leu	Asp	Ala	Val	Arg	Ser	Phe	Arg
945					950					955					960
Ile	Arg	His	Arg	Pro	Gln	Glu	Gln	Leu	Arg	Leu	Arg	Ile	Gly	Ile	His
				965					970					975	
Thr	Gly	Pro	Val	Cys	Ala	Gly	Val	Val	Gly	Leu	Lys	Met	Pro	Arg	Tyr
			980					985					990		

-continued

Cys	Leu	Phe	Gly	Asp	Thr	Val	Asn	Thr	Ala	Ser	Arg	Met	Glu	Ser	Asn
	995						1000					1005			
Gly	Glu	Ala	Leu	Lys	Ile	His	Leu	Ser	Ser	Glu	Thr	Lys	Ala	Val	
	1010					1015					1020				
Leu	Glu	Glu	Phe	Gly	Gly	Phe	Glu	Leu	Glu	Leu	Arg	Gly	Asp	Val	
	1025					1030					1035				
Glu	Met	Lys	Gly	Lys	Gly	Lys	Val	Arg	Thr	Tyr	Trp	Leu	Leu	Gly	
	1040					1045					1050				
Glu	Arg	Gly	Ser	Ser	Thr	Arg	Gly								
	1055					1060									

<210> SEQ ID NO 19

<211> LENGTH: 4246

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

```

ggttccctcc ggatagccgg agacttgggc cggccggacg ccccttcttg cacactccct      60
ggggcaggcg ctcacgcacg ctacaaacac acaactcctt ttctccctc gcgcgccctc      120
tctcatcctt cttcacgaag cgctcactcg caccctttct ctctctctct ctctctctaa      180
cacgcacgca cactcccagt tggtcacact cgggtcctct ccagcccgac gttctcctgg      240
caccacactg ctccgcggcg ccttgccgcg cccctcgggt cgcgccctt gcgtctctcg      300
cccagaccgt cgcagctaca gggggcctcg agccccggg tgagcgtccc cgtcccgtc      360
ctgctccttc ccatagggac gcgcctgatg cctgggacgg gccgctgagc ccaaggggac      420
cgaggaggcc atggtaggag cgctcgctcg ctgcggtgcc cgctgaggcc atgccggggc      480
cccggcgccc cgctggctcc cgctgcgcgc tgctcctgct cctgctgctg ccgccgctgc      540
tgctgctgct ccggggcagc cagcggggca acctgacggt agccgtggta ctgccgctgg      600
ccaatacctc gtacccttgg tcgtgggcgc gcgtgggacc cgcctgggag ctggccctgg      660
cccagggtgaa ggcgcgcccc gacttgctgc cgggctggac ggtccgcacg gtgctgggca      720
gcagcgaaaa cgcgctgggc gtctgtctcg acaccgcagc gcccttgccc gcggtggacc      780
tcaagtggga gcacaacccc gctgtgttcc tgggccccgg ctgcgtgtac gccgcgccc      840
cagtggggcg cttaccgcg cactggcggg tcccgctgct gaccgcggc gccccggcgc      900
tgggcttcgg tgtcaaggac gagtatgcgc tgaccaccgg cgcggggccc agtacgcca      960
agctggggga cttcgtggcg gcgctgcacc gacggtgggg ctgggagcgc caagcgtca      1020
tgctctacgc ctaccggcgg ggtgacgaag agcactgctt cttcctcgtg gaggggctgt      1080
tcatgcgggt ccgcgaccgc ctcaatatta cgggtggacca cctggagttc gccgaggacg      1140
acctcagcca ctaccaggg ctgctgcgga ccatgcccg caaaggccga gttatctaca      1200
tctgcagctc ccctgatgcc ttcagaaccc tcatgctcct ggccctggaa gctggcttgt      1260
gtggggagga ctacgttttc ttccacctgg atatctttgg gcaaagcctg caaggtggac      1320
agggccctgc tccccgcagg ccttgggaga gaggggatgg gcaggatgtc agtgcccgcc      1380
aggcctttca ggctgccaaa atcattacat ataaagaccc agataatccc gagtacttgg      1440
aattcctgaa gcagttaaaa cacctggcct atgagcagtt caacttcacc atggaggatg      1500
tcctgggtgaa caccatccca gcacccctc acgacgggct cctgctctat atccaggcag      1560
tgacggagac tctggcacat gggggaactg ttactgatgg ggagaacatc actcagcgga      1620
tgtggaaccg aagctttcaa ggtgtgacag gatacctgaa aattgatagc agtggcgatc      1680
gggaaacaga cttctccctc tgggatatgg atcccagaaa tgggtgccttc aggggtgtac      1740

```

-continued

tgaactacaa	tgggacttcc	caagagctgg	tggctgtgtc	ggggcgcaaa	ctgaactggc	1800
cctctgggta	ccctcctcct	gacatcccca	aatgtggctt	tgacaacgaa	gacccagcat	1860
gcaaccaaga	tcacctttcc	accctggagg	tgctggcttt	ggtgggcagc	ctctccttgc	1920
tcggcattct	gattgtctcc	ttcttcatat	acaggaagat	gcagctggag	aaggaactgg	1980
cctcggagct	gtggcgggtg	cgtcgggagg	acgttgagcc	cagtagcctt	gagaggcacc	2040
tgcgagagtgc	aggcagccgg	ctgaccctga	gcgggagagg	ctccaattac	ggctccctgc	2100
taaccacaga	gggccagtcc	caagtctttg	ccaagacagc	atattataag	ggcaacctcg	2160
tggctgtgaa	acgtgtgaac	cgtaaacgca	ttgagctgac	acgaaaagtc	ctgtttgaac	2220
tgaagcatat	gcgggatgtg	cagaatgaac	acctgaccag	gtttgtggga	gcctgcaccg	2280
acccccccaa	tatctgcctc	ctcacagagt	actgtccccc	tgggagcctg	caggacattc	2340
tggagaatga	gagcatcacc	ctggactgga	tgctccggta	ctcactcacc	aatgacatcg	2400
tcaagggcat	gctgtttcta	cacaatgggg	ctatctgttc	ccatgggaac	ctcaagtcac	2460
ccaactgcgt	ggtagatggg	cgtcttctgc	tcaagatcac	cgactatggg	ctggagagct	2520
tcagggaacct	ggacccagag	caaggacaca	cgttttatgc	caaaaagctg	tggacggccc	2580
ctgagctcct	gcgaatggct	tcacccctcg	tgccggggctc	ccaggctggt	gacgtataca	2640
gctttgggat	cctccttcag	gagattgccc	tgaggagtgg	ggctctccac	gtggaagggt	2700
tggacctgag	ccccaaagag	atcatcgagc	gggtgactcg	gggtgagcag	cccccttccc	2760
ggccctccct	ggccctgcag	agtcacctgg	aggagtgtgg	gctgctcatg	cagcgggtgct	2820
gggctgagga	cccacaggag	aggccaccat	tccagcagat	ccgcctgacg	ttgcgcaaat	2880
ttaacaggga	gaacagcagc	aacatcctgg	acaacctgct	gtcccgcctg	gagcagtcag	2940
cgaacaatct	ggaggaactg	gtggaggagg	ggacccaggc	atacctggag	gagaagcgca	3000
aggctgaggc	cctgctctac	cagatcctgc	ctcactcagt	ggctgagcag	ctgaagcgtg	3060
gggagacggg	gcaggccgaa	gcctttgaca	gtgttaccat	ctacttcagt	gacattgtgg	3120
gtttcacagc	gctgtcggcg	gagagcacgc	ccatgcaggg	ggtgacctcg	ctcaatgacc	3180
tgtacacttg	ctttgatgct	gtcatagaca	actttgatgt	gtacaagggtg	gagacaattg	3240
gcgatgccta	catggtgggtg	tcagggtctc	ctgtgcggaa	cgggcgggcta	cacgcctgcg	3300
aggtagcccg	catggccctg	gcactgctgg	atgctgtgcg	ctccttccga	atccgccacc	3360
ggccccagga	gcagctgcgc	ttgcgcattg	gcatccacac	aggacctgtg	tgtgctggag	3420
tgggtgggact	gaagatgccc	cgttactgtc	tctttgggga	tacagtcaac	acagcctcaa	3480
gaatggagtc	taatggggaa	gcctgaaga	tccacttgtc	ttctgagacc	aaggctgtcc	3540
tggaggagtt	tgggtggttc	gagctggagc	ttcgagggga	tgtagaaatg	aagggcacaa	3600
gcaagggtcg	gacctactgg	ctccttgggg	agagggggag	tagcaccoga	ggctgacctg	3660
cctcctctcc	tatccctcca	cacctccctc	acctgtgccc	agaagcaaca	gagggtgccag	3720
gcctcagcct	cacccacagc	agcccatcgc	ccaaggatg	gaagtaattt	gaatagctca	3780
ggtgtgctta	ccccagtga	gacaccagat	aggacctctg	agaggggact	ggcatggggg	3840
gatctcagag	cttacaggct	gagccaagcc	cacggccatg	cacagggaca	ctcacacagg	3900
cacacgcacc	tgctctccac	ctggactcag	gccgggctgg	gctgtggatt	cctgatcccc	3960
tccccctccc	atgetctcct	ccctcagcct	tgctaccttg	tgacttactg	ggaggagaaa	4020
gagtcacctg	aaggggaaca	tgaaaagaga	ctaggtgaag	agagggcagg	ggagcccaca	4080
tctggggctg	gcccaacaata	cctgctcccc	cgacccctcc	caccagcag	tagacacagt	4140

-continued

gcacagggga gaagaggggt ggcgcagaag ggttgggggc ctgtatgcct tgcttctacc 4200

atgagcagag acaattaaaa tctttattcc aaaaaaaaaa aaaaaa 4246

<210> SEQ ID NO 20

<211> LENGTH: 541

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Met Pro Ser Leu Leu Val Leu Thr Phe Ser Pro Cys Val Leu Leu Gly
1 5 10 15

Trp Ala Leu Leu Ala Gly Gly Thr Gly Gly Gly Gly Val Gly Gly Gly
20 25 30

Gly Gly Gly Ala Gly Ile Gly Gly Gly Arg Gln Glu Arg Glu Ala Leu
35 40 45

Pro Pro Gln Lys Ile Glu Val Leu Val Leu Leu Pro Gln Asp Asp Ser
50 55 60

Tyr Leu Phe Ser Leu Thr Arg Val Arg Pro Ala Ile Glu Tyr Ala Leu
65 70 75 80

Arg Ser Val Glu Gly Asn Gly Thr Gly Arg Arg Leu Leu Pro Pro Gly
85 90 95

Thr Arg Phe Gln Val Ala Tyr Glu Asp Ser Asp Cys Gly Asn Arg Ala
100 105 110

Leu Phe Ser Leu Val Asp Arg Val Ala Ala Ala Arg Gly Ala Lys Pro
115 120 125

Asp Leu Ile Leu Gly Pro Val Cys Glu Tyr Ala Ala Ala Pro Val Ala
130 135 140

Arg Leu Ala Ser His Trp Asp Leu Pro Met Leu Ser Ala Gly Ala Leu
145 150 155 160

Ala Ala Gly Phe Gln His Lys Asp Ser Glu Tyr Ser His Leu Thr Arg
165 170 175

Val Ala Pro Ala Tyr Ala Lys Met Gly Glu Met Met Leu Ala Leu Phe
180 185 190

Arg His His His Trp Ser Arg Ala Ala Leu Val Tyr Ser Asp Asp Lys
195 200 205

Leu Glu Arg Asn Cys Tyr Phe Thr Leu Glu Gly Val His Glu Val Phe
210 215 220

Gln Glu Glu Gly Leu His Thr Ser Ile Tyr Ser Phe Asp Glu Thr Lys
225 230 235 240

Asp Leu Asp Leu Glu Asp Ile Val Arg Asn Ile Gln Ala Ser Glu Arg
245 250 255

Val Val Ile Met Cys Ala Ser Ser Asp Thr Ile Arg Ser Ile Met Leu
260 265 270

Val Ala His Arg His Gly Met Thr Ser Gly Asp Tyr Ala Phe Phe Asn
275 280 285

Ile Glu Leu Phe Asn Ser Ser Ser Tyr Gly Asp Gly Ser Trp Lys Arg
290 295 300

Gly Asp Lys His Asp Phe Glu Ala Lys Gln Ala Tyr Ser Ser Leu Gln
305 310 315 320

Thr Val Thr Leu Leu Arg Thr Val Lys Pro Glu Phe Glu Lys Phe Ser
325 330 335

Met Glu Val Lys Ser Ser Val Glu Lys Gln Gly Leu Asn Met Glu Asp
340 345 350

Tyr Val Asn Met Phe Val Glu Gly Phe His Asp Ala Ile Leu Leu Tyr
355 360 365

-continued

Val	Leu	Ala	Leu	His	Glu	Val	Leu	Arg	Ala	Gly	Tyr	Ser	Lys	Lys	Asp
370						375					380				
Gly	Gly	Lys	Ile	Ile	Gln	Gln	Thr	Trp	Asn	Arg	Thr	Phe	Glu	Gly	Ile
385					390					395					400
Ala	Gly	Gln	Val	Ser	Ile	Asp	Ala	Asn	Gly	Asp	Arg	Tyr	Gly	Asp	Phe
				405					410					415	
Ser	Val	Ile	Ala	Met	Thr	Asp	Val	Glu	Ala	Gly	Thr	Gln	Glu	Val	Ile
			420					425					430		
Gly	Asp	Tyr	Phe	Gly	Lys	Glu	Gly	Arg	Phe	Glu	Met	Arg	Pro	Asn	Val
		435						440				445			
Lys	Tyr	Pro	Trp	Gly	Pro	Leu	Lys	Leu	Arg	Ile	Asp	Glu	Asn	Arg	Ile
	450					455					460				
Val	Glu	His	Thr	Asn	Ser	Ser	Pro	Cys	Lys	Ser	Ser	Gly	Gly	Leu	Glu
465					470					475					480
Glu	Ser	Ala	Val	Thr	Gly	Ile	Val	Val	Gly	Ala	Leu	Leu	Gly	Ala	Gly
				485					490					495	
Leu	Leu	Met	Ala	Phe	Tyr	Phe	Phe	Arg	Lys	Lys	Tyr	Arg	Ile	Thr	Ile
			500					505					510		
Glu	Arg	Arg	Thr	Gln	Gln	Glu	Glu	Ser	Asn	Leu	Gly	Lys	His	Arg	Glu
		515						520				525			
Leu	Arg	Glu	Asp	Ser	Ile	Arg	Ser	His	Phe	Ser	Val	Ala			
	530					535					540				

<210> SEQ ID NO 21

<211> LENGTH: 53

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: siRNA specific for NPR-A

<400> SEQUENCE: 21

tattacgggtg gaccacctgt tcaagagaca ggtggtccac cgtaatatatt ttt 53

<210> SEQ ID NO 22

<211> LENGTH: 53

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: siRNA specific for NPR-A

<400> SEQUENCE: 22

agaattccag aaacgcagct tcaagagagc tgcgtttctg gaattctttt ttt 53

What is claimed is:

1. A method of inhibiting the growth of cancer cells in a patient, comprising administering a polynucleotide encoding a natriuretic hormone peptide (NP) and an operably-linked promoter sequence, to the patient, wherein the cancer cells comprise lung cancer cells, breast cancer cells, ovarian cancer cells, or melanoma cells.

2. The method of claim 1, wherein the NP comprises the amino acid sequence of SEQ ID NO: 5 or a homolog of SEQ ID NO: 5 having at least one conservative amino acid substitution of SEQ ID NO: 5.

3. The method of claim 1, wherein said administering is by a route selected from the group consisting of oral, intramuscular, parenteral, intravenous, and intranasal.

4. The method of claim 1, wherein the polynucleotide is administered with a pharmaceutically acceptable carrier.

5. The method of claim 1, wherein the polynucleotide is contained within an expression vector.

6. The method of claim 5, wherein the expression vector is a DNA plasmid or virus.

7. The method of claim 1, wherein the polynucleotide sequence is administered with chitosan.

8. The method of claim 1, wherein the patient is suffering from one or more tumors, and wherein the NP causes regression of one or more of the tumors in the patient.

9. The method of claim 1, wherein the NP reduces tumor growth and metastasis in the patient.

10. The method of claim 1, wherein the patient is human.

11. The method of claim 1, wherein the cancer cells comprise lung cancer cells, and wherein the polynucleotide is administered to the airways of the patient.

12. The method of claim 2, wherein the cancer cells comprise lung cancer cells, and wherein the polynucleotide is administered to the airways of the patient.

71

- 13. The method of claim 12, wherein the patient is suffering from one or more tumors, and wherein the NP causes regression of one or more of the tumors in the patient.
- 14. The method of claim 1, wherein the NP comprises the amino acid sequence of SEQ ID NO: 5.
- 15. The method of claim 1, wherein the cancer cells comprise breast cancer cells.
- 16. The method of claim 2, wherein the cancer cells comprise breast cancer cells.

72

- 17. The method of claim 1, wherein the cancer cells comprise ovarian cancer cells.
- 18. The method of claim 2, wherein the cancer cells comprise ovarian cancer cells.
- 19. The method of claim 1, wherein the cancer cells comprise melanoma cells.
- 20. The method of claim 2, wherein the cancer cells comprise melanoma cells.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE

CERTIFICATE OF CORRECTION

PATENT NO. : 8,148,114 B2
APPLICATION NO. : 12/259110
DATED : April 3, 2012
INVENTOR(S) : Shyam S. Mohapatra

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 2,

Line 56, “such as pNP73-1102” should read --such as pNP73-102--.

Column 8,

Line 42, “natritiretic factor” should read --natriuretic factor--.

Column 22,

Line 14, “encoding NP₁₋₀” should read --encoding NP₁₋₃₀--.

Column 25,

Line 5, “Calf concentration” should read --Ca⁺⁺ concentration--.

Line 6, “nitric oxide (O)” should read --nitric oxide (NO)--.

Column 27,

Line 41, “Herpes simplex virus
type 1 (HISV-1)” should read --Herpes simplex virus
type 1 (HSV-1)--.

Line 67, “caLionic lipid delivery” should read --cationic lipid delivery--.

Column 30,

Line 25, “polynucieotides” should read --polynucleotides--.

Column 35,

Lines 22-23, “rhabdomvosarcoma” should read --rhabdomyosarcoma--.

Column 38,

Line 30, “the 2-p origin” should read --the 2-μ origin--.

Line 55, “Hela cells” should read --HeLa cells--.

Signed and Sealed this
Seventeenth Day of July, 2012

David J. Kyros

David J. Kappos
Director of the United States Patent and Trademark Office

CERTIFICATE OF CORRECTION (continued)

Page 2 of 2

U.S. Pat. No. 8,148,114 B2

Column 43,

Line 25, “polyniucleotides” should read --polynucleotides--.