
USF Patents

June 2011

Sphingosine 1-phosphate receptor gene, SPPR

Thomas P. Loughran

Ravi Kothapalli

Follow this and additional works at: https://digitalcommons.usf.edu/usf_patents

Recommended Citation

Loughran, Thomas P. and Kothapalli, Ravi, "Sphingosine 1-phosphate receptor gene, SPPR" (2011). *USF Patents*. 409.

https://digitalcommons.usf.edu/usf_patents/409

This Patent is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Patents by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.



US007964358B2

(12) **United States Patent**
Loughran et al.

(10) **Patent No.:** **US 7,964,358 B2**
(45) **Date of Patent:** **Jun. 21, 2011**

- (54) **SPHINGOSINE 1-PHOSPHATE RECEPTOR GENE, SPPR**
(75) Inventors: **Thomas P. Loughran**, Hummelstown, PA (US); **Ravi Kothapalli**, Wesley Chapel, FL (US)
(73) Assignee: **University of South Florida**, Tampa, FL (US)
(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/879,918**

(22) Filed: **Sep. 10, 2010**

(65) **Prior Publication Data**

US 2011/0021375 A1 Jan. 27, 2011

Related U.S. Application Data

- (63) Continuation of application No. 12/193,428, filed on Aug. 18, 2008, now abandoned, which is a continuation of application No. 11/653,811, filed on Jan. 16, 2007, now abandoned, which is a continuation of application No. 10/024,019, filed on Dec. 21, 2001, now Pat. No. 7,220,580.
(60) Provisional application No. 60/257,119, filed on Dec. 22, 2000.
(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G01N 33/53 (2006.01)
(52) **U.S. Cl.** **435/6; 435/7.1**
(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,220,580 B2	5/2007	Loughran et al.	
2003/0119111 A1*	6/2003	Lal et al.	435/69.1
2007/0105148 A1	5/2007	Loughran et al.	
2009/0181382 A1	7/2009	Loughran et al.	

FOREIGN PATENT DOCUMENTS

EP	1 149 907 A1	10/2001
WO	WO 97/00952 A2	1/1997
WO	WO 99/33972 A1	7/1999
WO	WO 99/46277 A1	9/1999
WO	WO 00/11166 A1	3/2000
WO	WO 00/31258 A2	6/2000
WO	WO 01/66742 A2	9/2001
WO	WO 02/57311 A3	7/2002

OTHER PUBLICATIONS

An et al., "Characterization of a Novel Subtype of Human G Protein-Coupled Receptor for Lysophosphatidic Acid*", *The Journal of Biological Chemistry* (1998), 273(14): 7906-7910.

An et al., "Signaling Mechanisms and Molecular Characteristics of G Protein-Coupled Receptors for Lysophosphatidic Acid and Sphingosine 1-Phosphate", *Journal of Cellular Biochemistry Supplements* (1998), 31:147-157.

Chen et al. "Discordant protein and mRNA expression in lung adenocarcinomas" *Molecular and Cellular Proteomics*, 2002, 1:304-313.

Database EMBL "ONLINE! Dec. 6, 2000, retrieved from EBI Database accession No. af317676 XP002236901 abstract.
Database EMBL "ONLINE! Oct. 8, 1999, retrieved from EBI Database accession No. ACO11461 XP002236899 abstract.
Database EMBL "ONLINE! Sep. 28, 2001, retrieved from EBI Database accession No. AX244600 XP002236900 abstract (Sequence 26 from Patent WO 01 66742).

Epstein and Butow, "Microarray technology—enhanced versatility, persistent challenge", *Current Opinion in Biotechnology* (2000), 11: 36-41.

Glickman et al., "Molecular Cloning, Tissue-Specific Expression, and Chromosomal Localization of a Novel Nerve Growth Factor-Regulated G-Protein-Coupled Receptor, nrg-1", *Molecular and Cellular Neuroscience* (1999), 14:141-152.

Haynes et al. "Proteome analysis: biological assay or data archive?" *Electrophoresis*, 1998, 19:1862-1871.

HLA et al., "Sphingosine-1-Phosphate Signaling via the EDG-1 Family of G-Protein-Coupled Receptors", *Annals New York Academy of Sciences* (2000), 905:16-24.

Hu et al. "Analysis of genomic and proteomic data using advanced literature mining" *J. Proteome Res.*, 2003, 2:405-412.

Im et al., "Characterization of a Novel Sphingosine 1-Phosphate Receptor, Edg-8*", *The Journal of Biological Chemistry* (2000), 275(19):14281-14286.

Im, D.S. et al. "Characterization of the human and mouse sphingosine 1-phosphate receptor, SIP5(Edg-8): Structure-activity relationship of sphingosine 1-phosphate receptors" *Biochem.*, 2001, 40:14053-060.
IM, D.S. et al. as cited in: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=nucleotide&list_uids=18141314&dopt=GenBank "1:NM_030760.Homo sapiens endo . . .", Jan. 13, 2002, p. 1-3.

IM, D.S. et al. as cited in: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=nucleotide&list_uids=18141314&dopt=GenBank "1:NM_030760[gi:13540516]", Dec. 10, 2001, p. 1-2.

Kothapalli, et al., "Characterization of Human Sphingosine-1-Phosphate Receptor Gene (SIP5) and its Differential Expression in LGL Leukemia", *Biochemica et Biophysica Acta* (2002), 117-123..

Kothapalli, R. et al. "Characterization of a novel human sphingosine-1-phosphate receptor gene (sppr) and its differential expression in LGL leukemia" *Blood: Gene Expression in Lymphoid Malignancy*, 2001, 98:3029. Kothapalli, R. et al. as cited in: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=nucleotide&list_uids=18033256&dopt=GenBank "1:AF331840.Homo sapiens SPPR . . .", Jan. 2, 2002, p. 1-2.

Kothapalli, R. et al. as cited in: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=nucleotide&list_uids=18033258&dopt=GenBank "1:AF331841.Homo sapiens SPPR . . .", Jan. 2, 2002, p. 1-2.

Kothapalli, R. et al. as cited in: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=nucleotide&list_uids=18033260&dopt=GenBank "1:AF331842.Homo sapiens SPPR . . .", Jan. 2, 2002, p. 1-2.

Malek, R. et al. "Nrg-1 belongs to the endothelial differentiation gene family of G protein-coupled sphingosine-1-phosphate receptors" *J. Biol. Chem.*, 2001, 276(8):5692-99.

Semenzato et al., "The Lymphoproliferative Disease of Granular Lymphocytes: Updated Criteria for Diagnosis, *Blood* (1997), 89(1): 256-260.

* cited by examiner

Primary Examiner — Christine J Saoud

Assistant Examiner — Jon M Lockard

(74) *Attorney, Agent, or Firm* — Saliwanchik, Lloyd & Eisenschenk


(57) **ABSTRACT**

A novel sphingosine 1-phosphate receptor gene, herein termed sppr, and its splice variants. Sppr is up-regulated in LGL and is useful, for example, in the diagnosis and treatment of certain lymphoproliferative, neurodegenerative and autoimmune diseases.

7 Claims, 16 Drawing Sheets

Microarray

FIG. 1A LGL FIG. 1B Normal



Expression of EST-1 in LGL Leukemia




FIG. 2

Northern Blot:

N=Normal, NA=Normal Activated

LGL = LGL Leukemia

EST-1 = Human Sphingosine 1-Phosphate receptor

BDE: 3.0 (544/182)

FIG. 3

Human sphingosine 1-Phosphate receptor
LOCUS tmpseq_1 2336 bp 4-DEC-2000
SOURCE PEMCs (LGL)
ORGANISM Human
Unclassified.
FEATURES Location/Qualifiers
source 1..2336
CDS 10..1206,10..1206
/note="predicted coding region"
/translation="

MESGLLRPAFVSEVIVLHYNYTGKLRGARYQFGAGLRADAVVCLAVCAFIVLENDVLLV
LGRHPRFHAPMFLLLGSLTSLDLAGAAYAANTLLSGPLTLKLSPALWFAREGGVFVALT
ASVLSLLALERSLTMAROPAPVSSRGRTLAMAAAAGWSLLGLLPALGWNCIGRLD
ACSTVLPLYAKAVVLCVLAFFVGLAATCALYARIYCCVRANARRLPARPGTAGTTSTRA
RRKPRSLALLRLTSLVLLAFVACWGPLFLLLLLDVACPARTCPVLLQADPFLGLAMANSI
LMPHIIYTLNRLRLHALLRLVCCGRHSCGRDPSGSQASASAAEASGGLRLRCLPPLDGSF
SGSERSSPQRDGLDTSSTGSPGAPTAARTLVSEPAAD"

BASE COUNT 461 a 679 c 701 g 495 t
ORIGIN
1 gccggggccca tggagtcggg gctgctggg cggggcggg tgagcgagg catcgctctg
61 cattacaact acacgggcaa gctccgcggt ggcgcgtacc agccgggtgc cggcctgggc
121 gccgacggcg tgggtgtgct ggcggtgagc gccttcctcg tgctagagaa tctagccgtg
181 ttgtttgggc tcggacggca ccggtgcttc caccgtccca tgttcttgct cctggggcagc
241 ctcacgttgt cggatctgct ggcagggccc gctacggcg ccaacatctt actgtcgggg
301 ccgctccagg tgaactgttc ccccgcgctc tggttcgcac gggagggagg cgtcttcgtg
361 gcactcactg cgtccgtgct gaggccctcg gccatcgccc tggagcgccg cctcaccatg
421 gcgcgcgggg ggcggcgccc cgtctccagt cggggggcca cgtctggcgt ggcagccggc
481 gccggggggc tgcgctgct cctcgggctc ctgcacggcg tgggctggaa ttgcctgggt
541 cgcctggcag cttgctccac tgtcttgcgc ctctacggca agggccacgt gctcttctgc
601 gtgctcgctt cgtggggcat cctggcgccg atctgtgcac tctacggcg catctaccgc
661 caggtacggc ccaacggcgc ggcctcgccg gccggggccc ggaactgggg gaccacctcg
721 accggggcgc gtgcgaagcc gcgctcgctg gccttgcctg gcacgctcag cgtggctgct
781 ctggcctttg tggcatgttg gggcccccct tctctgtctg tgttgcctga cgtggcgtgc
841 cggggcgcca ccttctctgt actctgcag gccgatccct tcttgggact gggcatgggc
901 aactcacttc tgaaccccat catctacacg ctcaaccaac gcgacttgcg ccaacggctc
961 ctgcggcctg tctgctgcgc agccactcc tggggcagag acccgagtgg ctcccagcag
1021 tcggcgagcg cggctgaggg ttcggggggc ctggcgcgct gcctggcccc gggccttgat
1081 gggagcttca cgggctcggg cgcctcatcg cccagcgcg acgggctgga caccagcggc
1141 cccacaggca gccccgggtg acccacagcc gcccgggactc tgggtatcga accggctgca
1201 gactgacacc ctccggccac gactgtcttc ccaagtttta cagacttgtt ctttttacc
1261 aaaggaatbt gtaggaatg cagccaaagg tgcagtccga aaagatgcg gggaaatgta
1321 tctatgcagc gacacccac natgtgaaca aacagacaaa aaatctgtgc cctcgtggaa
1381 ttgacgttct gcttgggaac acagaaaaga actcgggtgat gaaataatgg agatgatcc
1441 agtgacaaac gacagagatg gcatggggg tcagggaaga cctctcttga gaggtagga
1501 cctgtgatgt gactgagac cctgttcttg ggaagaccaa aagaaaagca ttccaggatg
1561 agggatggc atgcgcacaag gccctgagggc tgaatgtgc ccattgtgtc taagaaatgc
1621 agcgatgctg cctgtgctgg agcagggagc gagggggaga atgggaggag acaaggagct
1681 gaaggagtag ttcccgaaagg accttgtggg tgatatagag gaacttcgct ttgctctgag
1741 tgaggcggga gccatagaag cttctaaagc gaagagggac ttgccctaat ccaggctgac
1801 acaggtgctt tgtggcctcc atggggaggt gaaaaccaca gaaggtgaag gggggctgca
1861 ctgagccaca ggaacaatga tggagatcc agctaagccc agaccccgta gattctagat
1921 agatcttaga ggcagcagac agaattactg aggaattgag tgtaaggagt gaataaagt
1981 atcaaggaca atgccaaggg tggggtaacc ccaaatctga ctttgggaga ctacgcccac
2041 tctctatcgg taataaaatt tcttttttat tttcttttct tttcttttct tcttttttct
2101 tttttttttt ttgagtttg gatcttctgc tctgtacccc aggtcgaggt gcaatgggca
2161 caatttatgc tcaatgcagc ctggaaactcc tgggatcaag cctggagtcc ctgcttcagc
2221 ctccttagta gctgggacta caggcatgca ccacatgccc cagttaataa aattctctca
2281 aatgcacaaa aaaaaaaaaa aaaaaactcg aggggggggc cggtaaccaa ttogcc

Alignment of deduced Amino acid sequence with Nrg-1 and Edg-8 (rat genes)

Nrg-1
EDG-8
SPPR
MESGLRPAFVSEVIVIHNTYTKRGARYQFGAGLRADAACVAVCAFI VLENI AVLIV
MESGLRPAFVSEVIVIHNTYTKRGARYQFGAGLRADAACVAVCAFI VLENI AVLIV
MESGLRPAFVSEVIVIHNTYTKRGARYQFGAGLRADAACVAVCAFI VLENI AVLIV

LGRHPRFAPMFLLLGSLTSLDLAGAAYATHLLSGPLTRLSPALWFAREGGVFAVA
LGRHPRFAPMFLLLGSLTSLDLAGAAYATHLLSGPLTRLSPALWFAREGGVFAVA
LGRHPRFAPMFLLLGSLTSLDLAGAAYATHLLSGPLTRLSPALWFAREGGVFAVA

ASVLSILAIAIERHATHARGPAPAAARATLMAVAAGLLTLGLLPALGWICLGRLE
ASVLSILAIALERHATHARGPAPAAARATLMAVAAGLLTLGLLPALGWICLGRLE
ASVLSILAIALERHATHARGPAPAAARATLMAVAAGLLTLGLLPALGWICLGRLE

ACSTVLVYAKAYVLCVLAFLGILAAICAIYARIYQVRANARRIRAGPSRRATSSSR
ACSTVLVYAKAYVLCVLAFLGILAAICAIYARIYQVRANARRIRAGPSRRATSSSR
ACSTVLVYAKAYVLCVLAFLGILAAICAIYARIYQVRANARRIRAGPSRRATSSSR

SRHTPRSLALLRTL SVVLLAFVACWGLFL LLLLDVACPARCPVLQADPFLGLAMMS
SRHTPRSLALLRTL SVVLLAFVACWGLFL LLLLDVACPARCPVLQADPFLGLAMMS
ARREPRSLALLRTL SVVLLAFVACWGLFL LLLLDVACPARCPVLQADPFLGLAMMS

LNPILYTTTNRDLRHALLRLCCGRCNQNDSNSLQSPSAVGPSCGGIRRCCLPPTLD
LNPILYTTTNRDLRHALLRLCCGRCNQNDSNSLQSPSAVGPSCGGIRRCCLPPTLD
LNPILYTTTNRDLRHALLRLCCGRCNQNDSNSLQSPSAVGPSCGGIRRCCLPPTLD

RSSPSEHSCPQRDGMDSCTSGPGATANTLTVPDNTD
RSSPSEHSCPQRDGMDSCTSGPGATANTLTVPDNTD
GSFSGSERSSPQRDGLDTSGTSGEGAPTANTLTSEPAAD

SPPR: Nrg -85%
SPPR: EDG -86%
* - single, fully conserved residue
: - conservation of strong groups
. - conservation of weak groups
- no consensus

FIG. 4

FIG. 5

Sphingosine-1- phosphate receptor.1
LOCUS tmpseq_1 1698 bp 30-OCT-2000
DEFINITION No definition line found.
ACCESSION tmpseq_1
VERSION
KEYWORDS
SOURCE Unknown.
ORGANISM Unknown.
Unclassified.
FEATURES Location/Qualifiers
source 1..1698
CDS 11..775,11..775
/note="predicted coding region"
/translation="MESGLLRPAVSEVIVLHYNYTGKLRGARYQPGAGLRADAVVCL
AVCAFIVLENLAVLLVLRHPRFHPMFLLLGSLTSLVSPARPGTAGTTSTRARKPRS
LALLRTL SVLLAFVACWGPLFLDLLLDVACPARTCPVLLQADPFLGLAMANSLLNP
IYTLNRLRLHALLRLVCCGRHSCGRDPSGSQQSAJAEASGGRLRCLPPLDGSFSG
SERSSPQRDLDTSGSTGSPGAPTAARTLVSEPAD"
BASE COUNT 352 a 462 c 516 g 368 t
ORIGIN
1 cgcgcggggccc atggagtgagg ggctgctgcg gccggcgccg gtgagcgagg tcctcgtcct
61 gcattacaac tacacgggca agctccggcg tgcgcgtac cagccgggtg ccggcctgcg
121 cgcgcgagccc gtgggtgtgc tggcgggtgt cgccttcata gtgctagaga acctagccgt
181 gttgttggtg ctgggagccc acccggcgtt ccacgctccc atgttccctg cctggggcag
241 cctcagcttg tgggtgcggg caccggcccg gactgcgggg accacctcga cccgggcggg
301 tggcaagccg cgtctgctgg ccttgcctgg cagcctcagc ggggtgctcc tggcctttgt
361 ggcctgttgg ggcctctctt tcttgcctgc gttgctcgac gtggcgtgcc cggcgccgac
421 ctgtcctgtg ctcttgcagg ccgatccctt cctgggactg gccatggcca actcacttct
481 gaacccccat atctacacgc tcaccaaccg cgacctgcgc cagcgtctcc tgcgcctggt
541 ctgctgcggg cgcactcctt gcggcagaga ccgagtggtc tccagcagt cggcgagcgc
601 ggcctgaggc tccggggggc tggcgcgtg cctgcccccg ggcttctgat ggagcttcag
661 cggctcggag cgtctatcgc cccagcgcca cgggctggac accagtggtt ccacaggcag
721 ccccggtgca cccacagcgc cccggactct ggtatcagaa cgggctgcag actgacccc
781 tgggcccacg actgtctctc caagtctttac agactgttct tttttacata aaggatcttg
841 taggaaatgc agccaaaggc gcagtcggaa aagatgcagg ggaatgtat ttatgcagcg
901 atacccccca atgtgaacaa acagacaaaa aatctgtgct ctcgtggaat tgactgtctg
961 ctggggaaca cagaaaaaga ctccgtgacg aaataatgga gatgattcca gtgacaaacg
1021 acagagatgg tgatgggtgg caggggaagc ctctctgcag aggtagtgcg ctgtgatgtg
1081 agctcgagac tctgtcctgg gaagaccaaa agaaaaagca ttcaggatga gggatggcca
1141 tgcgcaaaag ccttgagggt gaaatgtgct catgtgtctt aagaaatgca gcgatgctgg
1201 tctgtcctgga gcagggaagg agggggagaa tgggaggaga caaggagctg aaggagtagt
1261 tcccgaaagg ccttctgggt gatatagagg acttgccttt tgccttgagt gaggtgggag
1321 ccatagaagg ttctaaagcg aagagggact tgccttaatt cagggtgatca cagggtgtct
1381 gtggcctcca tgggaggttg aaaaccagag aagggtgaag ggggctgcac tgagccacag
1441 gaacaaatgat ggagattcca gctaagccca gacccgtgg attctagata gatttttagag
1501 gcagcagaca gaattactga ggaattgagt gtaagagtgg aataaagtta tcaaggacaa
1561 tgccaagggt ggggcacccc caaatctgac tctgggagac tcagccaaat cctatctggt
1621 aataaaaattt cttttttatt tctctctctt tctctctctt cttctctctt tttctgaggt
1681 gggatctctt gctctgtt

FIG. 6

Sphingosine -1-Phosphate receptor 2

LOCUS tmpseq_1 1245 bp 30-OCT-2000
DEFINITION No definition line found.
ACCESSION tmpseq_1
VERSION
KEYWORDS .
SOURCE Unknown.
ORGANISM Unknown.
Unclassified.


FEATURES Location/Qualifiers
source 1..1245
CDS 11..322,11..322
/note="predicted coding region"
/translation="MESGLLRPAPVSEVIVLHYNMTGKLRGARYQPGAGLRADAVVCL
AVCAFIIVLENLAVLLVLGRHPRFHAPMFLLLGSLTSLDLLAGAAAYAAAARTLVSEPAAD"
D"

BASE COUNT 296 a 284 c 372 g 291 t

ORIGIN
1 cggcgggccc atggagtcgg ggcgcgcgcg gccggcgccg gtgagcgagg tcctcgctct
61 gcattacaac tacaccggca agctccgcgg tgcgcgcctac cagccggggtg ccggccttgcg
121 cggcgagccc gtgggtgtgc tggcggtgtg cgccttcctc gtgctagaga atctagccgt
181 gttggttggtg ctgggacgcc acccgcgctt ccacgctccc atgttccctgc tccctgggcag
241 cctcacgttg tgggactctgc tggcaggcgc cgcctacgca gccgcgcgcc ggactctggt
301 atcagaaccg gcgcgcgact gacaccctcg gccacgact gtcttcccaa gttttacaga
361 cttgttcttt ttacataaag gaatttctag gaaatgcagc caaagggtgca gtgggaaaag
421 atgcagggga aatgtattta tgcagcgaca cccacaaatg tgaacaaaca gacaaaaaat
481 ctgtgcccct gtggaattga cgttctgctt gggaacacag aaaagaactc ggtgatgaaa
541 taatggagat gattccagtg acaaacgaca gagatggtga tgggtggtcag ggaagacctc
601 tctgcagagg tagtgacttg tgatgtgagc tgagacctct gtccctgggaa gacaaaaga
661 aaagcatttc aggatgaggg aatggcatgc gcaaaaggccc tgaggctgaa atgtgcccat
721 gtgttctaaag aaatgcagcg atgctggtgt gcctggagca gggacggagg gggagaatgg
781 gaggagacaa ggagctgaag gagtagtccc cgaaggacct tgtgggtgat atagaggact
841 tgcctcttgc tctgagttag gtgggagcca tagaagcttc taagcagaag agggacttgc
901 cctaattcag gtagtcacag gtgtcttgtg gcctccatgg gaggttgaaa accagagaag
961 gtgaaggggg gctgcactga gccacaggaa caatgatgga gattccagct aagcccagac
1021 ccggtggatt ctagatagat tctagaggca gcagacagaa ttactgagga attgagtgtg
1081 agagtggaaat aaagttatca aggacaatgc caagggtggg gcacccccaa atttgactct
1141 gggagactca gccaaatctt atctggtaat aaaattctct ttttattttt cttttctttc
1201 tttctttctt tttttttttt ttgagttggg atcttctgct ctgtc

FIG. 7

Expression of *Sppr* in Human Tissues



1 gcgcggcccatcgagtcggggctgctgcccggcgccgggtgagcgaggtcctcctg
M E S G L L R P A P V S E V I V L 17
61 cattacaactacacgggcaagctccggggtgcgcgtaccagccgggtgcggcctgcgc
H Y N Y T G K L R G A R Y Q P G A G L R 37
121 gcgcagcccgctggtgtgctcggggtgtgcgccttcctcgtgctagagaatctagccgtg
A D A V V C L A V C A F I V L E N L A V 57
181 ttgtttggtgctcggagccaccccgcttccacgctcccatgttctcgtcctcgggcagc
L L V L G R H P R F H A P M F L L L G S 77
241 ctccagttgtcggatctgctggcaggcgccgcctacgcgcgaacatcctactgtcgggg
L T L S D L L A G A A Y A A N I L L S G 97
301 ccgctcacgctgaaaactgtcccccgctctgttctgcacgggaggaggcgtctctcgtg
P L T L K L S P A L W F A R E G G V F V 117
361 gcaactcaatgcgtccgtgctgagcctcctggccatcgcgctggagcgcagcctcaccatg
A L T A S V L S L L A I A L E R S L T M 137
421 gcgcgcaggggggccgcgcgcgtctccagtcggggcgccacgctggcgatggcagccgcg
A R R G P A P V S S R G R T L A M A A A 157
481 gccggggcgctgtcgtgctcctcgggctcctgcagcgtgggctggaattgcctgggt
A W G V S L L L G L L P A L G W N C L G 177
541 cgcctggagccttgctccactgtcttgccgtctacgcgaaggcctacgtgctctctcgtc
R L D A C S T V L P L Y A K A Y V L F C 197
601 gtgctcgccttcgtgggcacctcgtgcgcgatctgtgcaactcaccgcgcgatctactgc
V L A F V G I L A A I C A L Y A R I Y C 217
661 caggtaacgcgcaacgcgcggcgccgtgcgggcacggccgggactgcggggaccacctcg
Q V R A N A R R L P A R P G T A G T T S 237
721 acccgggcgcgctgcgaagccgcgctcgtcgtggtgctgcgcacgctcagcgtgggtgctc
T R A R R K P R S L A L R T L S V V L 257
781 ctggcctctgtggcatgttggggccccctcttctcgtgctggttgctcagcgtggcgctgc
L A F V A C W G P L F L L L L L D V A C 277
841 ccggcgcgccacctgtcctgtactcctgcaggccgatcccttctcgtggactggccatggcc
P A R T C P V L L Q A D P F L G L A M A 297
901 aactcacttctgaaacccatcatctacacgctcaccacccgcgacctgcgccacgcgctc
N S L L N P I I Y T L T N R D L R H A L 317
961 ctgcgcctggtctgctgcggacgccactcctgcggcagagacccgagtggctcccagcag
L R L V C C G R H S C G R D P S G S Q Q 337
1021 tcggcgagcgcgggtgaggtctccggggcctgcgcgcgtgcctgccccgggccttgat
S A S A A E A S G G L R R C L P P G L D 357
1081 gggagcttcagcggctcggagcgtcctcgcgccagcgcgacgggtggacaccagcggc
G S F S G S E R S S P Q R D G L D T S G 377
1141 tccacaggcagcccggtgcacccacagcgcgcggactctggtatcagaacccggtgca
S T G S F G A P T A A R T L V S E P A A 397
1201 gactgacacccctcggcccaagactgtcttcccaagttttacagacttgttctttttacat
D * 398
1261 aaaggaattttgtacgaaatgcagccaaaggtgcagtcggaaaaagatgcaggggaaatgta
1321 tttatgcagcgcacaccccaaatgtgaacaaacagacaaaaaatctgtgcctcgtggaa
1381 ttgacgtctctgcttgggaacacagaaaagaactcgggtgatgaaataatggagatgattcc
1441 agtgacaaaacgacagagatggtgatggtggtcagggaagacctctctgcagaggttagtga
1501 cttgtgatgtgagctgagacctctgtcctgggaagacaaaagaaaagcatttcaggatg
1561 agggaaatggcatgcgcgaagccctgaggctgaaatgtgcccattgtgttctaagaaatgc
1621 agcgtatgctggtgtgcctggagcaggacgggaggggagaatggaggagacaggagct
1681 gaaggagtgttccgaaggaccttggtgggtgatataaggagacttcgcttttgctctgag
1741 tgagggtgggagccatagaagcttctaagcagaagagggaacttgccctaattcagggtgatc
1801 acagggtgtcttggtggcctccatgggaggttgaaaacacagaaggtgaaggggggctgca
1861 ctgagccacaggaacaatgatggagattccagctaagccagaccccgatggattctagat
1921 agatttttagaggcagcagacagaattactgaggaattgagtgtaagagtggaaataaagtt
1981 atcaaggacaatgccaaaggtggggcaccocccaaatttgactttggcagactcagccaaa
2041 tctatcttggaataaaaatttctttttttttttttttttttttttttttttttttttttt
2101 ttttttttttttttgagttgggatcttgctctgtcaccacaggctggagtgcaattggca
2161 caattatagctcactgcagcctggaactcctgggatcaagcctggagttcctgctcagc
2221 ctccctagtagctgggactacagcctgcaccacccatgccagtttaataaaatttcttca
2281 aatgcacaaaaaaaaaaaaaaaaaaaaa

FIG. 8

```

NRG-1 MESGLLRPAPVSEVIVLHNYTGKLRGARYQPGAGLRADAADVCLAVCAFIVLENLAVLLV
EDG-β MESGLLRPAPVSEVIVLHNYTGKLRGARYQPGAGLRADAADVCLAVCAFIVLENLAVLLV
SIP5 MESGLLRPAPVSEVIVLHNYTGKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV
*****

NRG-1 LGRHPRFHAPMFLLLGSLTSLDLAGAAYATNILLSGPLTLRLSPALWFAREGGVFVALA
EDG-β LGRHPRFHAPMFLLLGSLTSLDLAGAAYATNILLSGPLTLRLSPALWFAREGGVFVALA
SIP5 LGRHPRFHAPMFLLLGSLTSLDLAGAAYAANILLSGPLTLKLSPALWFAREGGVFVALT
*****:*****:*****:

NRG-1 ASVLSLLAIAIERHMTMARRGPAPAASRARTLAMAVAAGWGLLLTLGLLPALGWNCLGRLE
EDG-β ASVLSLLAIALERHMTMARRGPAPAASRARTLAMAVAAGWGLSLLTLGLLPALGWNCLGRLE
SIP5 ASVLSLLAIALERSLTMARRGPAPVSSRGRTLAMAAAAGVSLTLGLLPALGWNCLGRLE
*****: * *****: *****: * *****:

NRG-1 ACSTVLPVYAKAYVLCVLAFLGILAAICALYARIYCQVRANARRLRAGPGSRRATSSSR
EDG-β ACSTVLPVYAKAYVLCVLAFLGILAAICALYARIYCQVRANARRLRAGPGSRRATSSSR
SIP5 ACSTVLPVYAKAYVLCVLAFLGILAAICALYARIYCQVRANARRLRAGPGT-AGTTSTR
*****:*****:***** * *: .*:

NRG-1 SRHTPRSLALLRTLSTVLLAFVACWGPLEFLLLLLDVACPARACPVLLQADPFLGLAMANS
EDG-β SRHTPRSLALLRTLSTVLLAFVACWGPLEFLLLLLDVACPARACPVLLQADPFLGLAMANS
SIP5 ARKTPRSLALLRTLSTVLLAFVACWGPLEFLLLLLDVACPARTCPVLLQADPFLGLAMANS
*: *****:*****

NRG-1 LLNPITYTFTNRDLRHALLRLCCGRGPCNQDSSNSLQSPSAVGPGSGGLRRCLPPTLD
EDG-β LLNPITYTFTNRDLRHALLRLCCGRGPCNQDSSNSLQSPSAVGPGSGGLRRCLPPTLD
SIP5 LLNPITYTFTNRDLRHALLRLVCCGRHSCGRDPSGSQQ-SASAAEASGG-LRRCLPPGLD
*****:*****:*** .*:.* * *.** .*** ***** **

NRG-1 RSSSPSEHSCPQRDGMTSCSTGSPGAATANRTLVPDATD-
EDG-β RSSSPSEHSCPQRDGMTSCSTGSPGAATANRTLVPDATD-
SIP5 GSFGSERSSPQRDGLDTSGSTGSPGAPTAARTLVSEPAAD
* * *: *.*****:*** *****.* * *****:

```

SIP₅: Nrg - 85%

SIP₅: ECG - 86%

* - single, fully conserved residue
: - conservation of strong groups
. - conservation of weak groups
- no consensus

FIG. 9




FIG. 10




FIG. 11

SIP₅ MESGLLRPAPVSEVIVLHNYTGKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV
SIP₅-α MESGLLRPAPVSEVIVLHNYTGKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV

SIP₅ LGRHPRFHAPMFLLLGSLTSLDLAGAAYAANILLSGPLTLKLSPALWFAREGGVFVALT
SIP₅-α LGRHPRFHAPMFLLLGSLTSL-----

SIP₅ ASVLSLLAIALERSLTMARRGPAPVSSRGRTLAMAAAAWGVSLLLGLLPALGWNCLGRLD
SIP₅-α -----
SIP₅ ACSTVLPYAKAYVLCVLAFFVGILAAICALYARIYCQVRANARRLPARPGTAGTTSTRA
SIP₅-α -----VPARPGTAGTTSTRA
:*****
SIP₅ RRKPRSLALLRTL SVVLLAFVACWGPLFLLLLDVACPARTCPVLLQADPFLGLAMANSL
SIP₅-α RRKPRSLALLRTL SVVLLAFVACWGPLFLLLLDVACPARTCPVLLQADPFLGLAMANSL

SIP₅ LNPIIYTLTNRDLRHALLRLVCCGRHSCGRDPGSGSQQSASAAEASGGLRRCLPPGLDGSF
SIP₅-α LNPIIYTLTNRDLRHALLRLVCCGRHSCGRDPGSGSQQSASAAEASGGLRRCLPPGLDGSF

SIP₅ SGSESSPQRDGLDTSGTSGSPGAPTAARTLVSEPAAD
SIP₅-α SGSESSPQRDGLDTSGTSGSPGAPTAARTLVSEPAAD

FIG. 12A

SIP₅ MESGLLRPAPVSEVIVLHYNVTGKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV
SIP₅-β MESGLLRPAPVSEVIVLHYNVTGKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV

SIP₅ LGRHPRFHAPMFLLLGSLTILSDLLAGAAAYAANILLSGPLTLKLSPALWFAREGGVFVALT
SIP₅-β LGRHPRFHAPMFLLLGSLTILSDLLAGAAAYA-----

SIP₅ ASVLSLLAIALERSLTMARRGPAPVSSRGRTLAMAAAAWGVSLLLGLLPALGWNCLGRLD
SIP₅-β -----
SIP₅ ACSTVLPLYAKAYVLCVLAFFVGILAAICAIYARIYCQVRANARRLPARPGTAGTTSTRA
SIP₅-β -----
SIP₅ RRKPRSLALLRTL SVVLLAFVACWGPFLLLLLDVACPARTCPVLLQADPFLGLAMANSL
SIP₅-β -----
SIP₅ LNPITYTLNRLRHALRLVCCGRHSCGRDPSGSQQSASAAEASGGLRCLPPGLDGSF
SIP₅-β -----
SIP₅ SCSERSSPQRDGLDTSGSTGSPGAPTAARTLVSEPAAD
SIP₅-β -----AARTLVSEPAAD

FIG. 12B




FIG. 13


```
SIP  MESGLLRPAPVSEVIVLHYNITGKLRGARYQPGAGLRADAVVCLAVCAFIIVLENLAVLLV
SIP1 MESGLLRPAPVSEVIVLHYNITGKLRGARYQPGAGLRADAVVCLAVCAFIIVLENLAVLLV
*****
SIP  LGRHPRFHAPMFLLLGSLTSLDLAGAAYAANILLSGPLTLKLSPALWFAREGCVFVALT
SIP1 LGRHPRFHAPMFLLLGSLTSL-----
*****
SIP  ASVLSLLATALEPSLTMARRGAPAPVSSRGRTLAMAAAANGVSLLLGLLPALGWNCLGRLD
SIP1 -----
SIP  ACSTVLPDYAKAYVLFVLAFFVILAAICALYARIYQVVRANARLPARPGTAGTTSTRA
SIP1 -----VPARPGTAGTTSTRA
*****
SIP  RRKPRSLALLRTLSTVLLAFVACWGPFLLLLLDVACPARTCPVLLQADPFGLAMANSL
SIP1 RRKPRSLALLRTLSTVLLAFVACWGPFLLLLLDVACPARTCPVLLQADPFGLAMANSL
*****
SIP  LNPIIYTLTNRDLRHALLRLVCCGRHSCGRDPGSGQASASAAEASGGLRRCCLPPGLDGSF
SIP1 LNPIIYTLTNRDLRHALLRLVCCGRHSCGRDPGSGQASASAAEASGGLRRCCLPPGLDGSF
*****
SIP  SGSERSSPQRDGLDTSGSTGSPGAPTAARTLVSEPAAD
SIP1 SGSERSSPQRDGLDTSGSTGSPGAPTAARTLVSEPAAD
*****
```

SIP= Sphingosine-1-Phosphate receptor versus
SIP1=Sphingosine-1-Phosphate 1 receptor

FIG. 14

```

SIP MESGLLRPAVPVSEVIVLHYNITKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV
SIP2 MESGLLRPAVPVSEVIVLHYNITKLRGARYQPGAGLRADAVVCLAVCAFIVLENLAVLLV
*****

SIP LGRHPRFHAPMFLLGSLTSLDLLAGAAYAANILLSGPLTLKLSPALWFAREGGVFVALT
SIP2 LGRHPRFHAPMFLLGSLTSLDLLAGAAYAA-----
*****

SIP ASVLSLLAIALERSLTMARRGPAPVSSRGRTLAMAAAAAGVSLLLGLLPALGWNCLGRLD
SIP2 -----

SIP ACSTVLPYAKAYVLFVCLAFVUGILAACALYARIYCVQRANARRLPARPGTAGTTSTRA
SIP2 -----

SIP RRKPRSLALLRTL SVVLLAFVACWGPLFLLLLLDVACPARTCPVLLQADPFLGLAMANSI
SIP2 -----

SIP LNPFIYTLTNDRDLRHALLRLVCCGRHSCGRDPSGSQASAAEASGGLRRCLP PGLDGSF
SIP2 -----

SIP SGSEKSSPQRDGLDTSGSTGSPGAPTAARTLVSEPAAD
SIP2 -----AARTLVSEPAAD
*****

```

FIG. 15

SPHINGOSINE 1-PHOSPHATE RECEPTOR GENE, SPPR

CROSS REFERENCE TO RELATED APPLICATIONS

The present application is a continuation of U.S. application Ser. No. 12/193,428, filed Aug. 18, 2008, which is a continuation of U.S. application Ser. No. 11/653,811, filed Jan. 16, 2007, now abandoned, which is a continuation of U.S. application Ser. No. 10/024,019, filed Dec. 21, 2001, now U.S. Pat. No. 7,220,580, which claims benefit of U.S. Provisional Application Ser. No. 60/257,119, filed Dec. 22, 2000, each of which is hereby incorporated by reference herein in its entirety, including any figures, tables, nucleic acid sequences, amino acid sequences, and drawings.

FIELD OF THE INVENTION

The present invention relates to the genetics of autoimmune diseases, including lymphoproliferative diseases, such as large granular lymphocyte leukemia (LGL), and rheumatoid arthritis (RA). Specifically, the invention relates to a novel sphingosine 1-phosphate receptor gene, herein termed *sppr*, and its splice variants. *Sppr* is up-regulated in LGL and is useful, for example, in the diagnosis and treatment of certain lymphoproliferative, neurodegenerative and autoimmune diseases.

BACKGROUND OF THE INVENTION

Large granular lymphocyte leukemia (LGL) is a rare form of lymphoproliferative disorder often associated with autoimmune disease (Loughran T. P., Clonal diseases of large granular lymphocytes. Blood 82, 1-14, 1993).

The cause of LGL is still not fully understood. An increased count of large granular lymphocytes is characteristic of LGL leukemia. Patients with clonal CD3+LGL, as determined by cytogenetic or T-cell receptor (TCR) gene rearrangement studies, are classified as T-LGL. Some of these patients may resemble those with Felty's syndrome with clinical features of rheumatoid arthritis, neutropenia and splenomegaly (Ahem M. J., et al., P. Phenotypic and genotypic analysis of mononuclear cells from patients with Felty's syndrome. Ann. Rheum. 49, 103-108, 1990.) Morbidity and mortality in patients with LGL leukemia typically results from infections acquired during severe neutropenia.

The etiology of LGL leukemia is also not yet known. There is strong evidence that suggests that leukemic large granular lymphocytes are antigen activated cytotoxic T lymphocytes (CTL), but the nature of the antigen and of the initial stimulus leading to antigen driven expansion are not known.

LGL leukemic cells express FAS and FAS ligand, but they are not actively undergoing apoptosis (Perzova, R and Loughran, T. P., Jr. Constitutive expression of Fas ligand in large granular lymphocyte leukemia. British Jnl. Haematology, 1997). How they acquire resistance to apoptosis is not known.

Within the field of the diagnosis and treatment of LGL and other autoimmune diseases, there is a need for better tools for diagnosis and early detection of disease, specific therapeutic targets and treatments for the disease, and more specific reagents and tools with which to identify the pathogenic pathways of these diseases. The present invention provides a novel gene and splice variants that are linked to these diseases, and which address the aforementioned needs and more, as will become clear to one of skill in the art upon reading the following disclosure.

SUMMARY OF THE INVENTION

Large granular lymphocyte leukemia (LGL) is a lymphoproliferative disorder often associated with autoimmune disease. In order to identify differentially expressed genes in LGL leukemia, microarray analysis is performed from RNA isolated from PBMC of LGL leukemia patients and compared with normal healthy individual(s). By screening a human LGL leukemia library the full-length sequence of a human gene that showed 85% identity with rat sphingosine 1-phosphate receptor is obtained. Two different isoforms are also identified by RT-PCR, designated sphingosine 1-phosphate receptor 1, also referred to as S1p5- α and sphingosine 1-phosphate receptor 2, also referred to as S1P5- β . Sphingosine 1-phosphate receptor (*sppr*) is present in brain, spleen, PBMCs, liver and kidney. The present inventors found it is over-expressed in LGL leukemia patients when compare to normal individuals.

In a first embodiment, the invention provides a gene comprising *sppr* or a splice 5 variant, or *sppr* protein or modified proteins or fragments thereof.

In a further embodiment, the invention provides a nucleic acid capable of hybridizing to at least a portion of said *sppr* gene, including splice variants.

In a further embodiment, the invention provides methods for screening for autoimmune diseases, including LGL or rheumatoid arthritis, based on overexpression of *sppr*.

In a further embodiment, the invention provides for monoclonal antibodies to *sppr* and their use in detection, diagnosis and treatment of disease states.

In a further embodiment, the invention provides for screening of ligands, agonists, and antagonists of *sppr*.

In a further embodiment, the invention provides for inhibition or treatment of neurodegenerative disease.

In a preferred embodiment the present invention provides a sphingosine 1-phosphate receptor gene. The use of said gene makes it possible to produce the sphingosine 1-phosphate receptor protein with ease and in large quantities, and said protein, which has sphingosine 1-phosphate receptor activity, can be used in developing therapeutic agents for various diseases.

Throughout this document the nomenclature *sppr* and S1P5 are used interchangeably. The receptor was initially termed *sppr*. However, to be consistent with a new nomenclature system this receptor was renamed S1P5.

DESCRIPTION OF THE FIGURES

FIG. 1A-B illustrates a microarray of the differential expression of the selected EST. (EST (GenBank ID 1868427) is obtained Incyte Genomics.) FIG. 1A-B shows a microarray hybridized with the fluorescent labeled probes generated using mRNA isolated from PBMC of LGL leukemia patient and from mRNA isolated from normal healthy individual. FIG. 1A illustrates a microarray showing the expression of an LGL leukemia patient cDNAs. FIG. 1B illustrates a microarray showing the expression of a normal healthy individual. Arrows show the expression of EST in both patient and normal individual (GeneBank Id: N47089). Intensity bar shows the increased expression starting from left to right. A balanced differential expression of 3.0 is determined for this EST.

FIG. 2 shows Northern blot analysis performed with 10 ug of total RNA isolated from PBMC of LGL leukemia patients and normal healthy individuals. These results demonstrate

over-expression of EST in the PBMCs of LGL leukemia when compared to normal and normal activated PBMCs of healthy individuals.

FIG. 3 shows the complete nucleotide sequence, SEQ ID NO: 4, of human sphingosine 1-Phosphate receptor (SPPR) cDNA and amino acid sequence (SEQ ID NO: 3) as predicted by the nucleic acid sequence. The full-length (2.2 kb) nucleotide sequence of *sppr* is compiled from sequences of clones isolated from an LGL leukemia library and RT-PCR products obtained by using the gene specific primers designed using the corresponding sequence from chromosome 19.

FIG. 4 shows the alignment of the amino acid sequence of SPPR with other members of the sphingosine 1-phosphate receptor family. The deduced amino acid sequence of *sppr* is compared with rat *edg-1* and *nrg-1*. There is approximately 85% identity with these genes.

FIG. 5 shows the nucleotide sequence and deduced amino acid sequence of splice variant, sphingosine 1-phosphate receptor 1. 1.6 kb fragment is obtained by RT-PCR using total RNA isolated from PBMC of an LGL leukemia patient. The fragment is then cloned and sequenced.

FIG. 6 shows the nucleotide sequence and deduced amino acid sequence of splice variant, sphingosine 1-phosphate receptor 2. The nucleotide sequence of an alternative splice variant of *sppr* and deduced sequence. 1.2 kb fragment is obtained from RT-PCR using total RNA isolated from PBMC of LGL leukemia. The fragment is then cloned and sequenced.

FIG. 7 shows results of *sppr* Northern blot analysis with different tissues. Northern blot analysis is performed using a multiple tissue Northern blot (Clontech). Northern blots contain approximately 1 μ g of poly A+ per lane from twelve different human tissues. A 1.5 kb fragment containing the full-length open reading frame for *sppr* is used as a probe. Results show *sppr* is expressed in mainly brain, spleen, and peripheral blood leukocytes. Small amounts of *sppr* are also expressed in lung, placenta, liver and kidney.

FIG. 8 shows nucleotide and deduced amino acid sequence of human *S1P₅* cDNA. Full-length (2.2 kb) nucleotide sequence of *S1P₅* is compiled from the sequences of clones isolated from LGL leukemia library (clone 6) and RT-PCR products. GenBank Accession No. AF331840. The predicted amino acids of the coding region are shown underneath by a single letter abbreviation. The left side of the sequence shows nucleotide numbers and the right side shows amino acid numbers. Possible seven transmembrane helices are underlined. The putative polyadenylation sites are in bold.

FIG. 9 shows Alignment of the deduced amino acid sequence of *S1P₅* with other members: The deduced amino acid sequence of *S1P₅* is compared with predicted amino acid sequences of rat *edg-8* and *nrg-1*. There is approximately 86% identity with these genes. *—single, fully conserved residue, :—conservation of strong groups, .—conservation of weak groups, —no consensus.

FIGS. 10A-10C show activation of Erk2 by *S1P* in HEK293 cells transiently transfected with *S1P₅*. HEK 293 cells transfected with the HA-ERK2 plasmid (0.2 μ g) and either pcDNA *S1P₅* (0.5 μ g) or vector alone. Vector plasmid is added to each transfection reaction to equalize the amount of total DNA (2.1 μ g). After serum-starvation, the cells are treated with 1 μ M *S1P* or 1 μ M LPA for 5 min (BSA was added to the controls). HA-ERK2 is immunoprecipitated from one half of each whole cell lysate and used for measuring the kinase activity utilizing MBP as substrate, while HA-ERK2 immunoprecipitated from the other half is used for determining the amount of ERK2 protein in the immune complex. FIG. 10A illustrates a representative autoradiogram

of 32 P incorporation into MBP catalyzed by HA-ERK 2 immunoprecipitated from transiently transfected cells treated as indicated. FIG. 10B illustrates the corresponding Western blot demonstrating the amount of HA-ERK2 present in each of the immune complexes. FIG. 10C illustrates a plot of ERK 2 activity (fold) normalized to the amount of ERK2 protein (means \pm SD from three independent experiments).

FIG. 11 shows Northern blot analysis of *S1P₅* mRNA expression in PBMC of LGL leukemia patients and normal healthy individuals. Northern blot is performed with 10 μ g of total RNA isolated from PBMC of LGL leukemia patients and normal healthy individuals. LGL=LGL leukemia patients, N=Normal healthy individual, NA=Normal healthy individuals PBMCs activated by IL2 and PHA. These results demonstrate over-expression of *S1P₅* in the PBMC of LGL leukemia when compared to normal and normal activated. PBMC of healthy individuals.

FIG. 12A-B shows comparison of the predicted amino acid sequences of *S1P₅*, *S1P₅- α* and *S1P₅- β* . The predicted amino acid sequences are aligned using CLUSTAL program. FIG. 12A illustrates the nucleotide sequence of an alternative splice variant of *S1P₅- α* and deduced amino acid sequence. A 1.6 kb fragment is obtained from RT-PCR using total RNA isolated from PBMC of LGL leukemia patient. This fragment is cloned and sequenced. FIG. 12B illustrates the nucleotide sequence of an alternative splice variant of *S1P₅- β* and deduced sequence. A 1.2 kb fragment is obtained from RT-PCR using total RNA isolated from PBMC of LGL leukemia. This fragment is cloned and sequenced.

FIG. 13 shows tissue distribution of *S1P₅* message. Northern blot analysis is performed using the multiple tissue blot obtained from Clontech. The Northern Blot contains approximately 2 μ g of poly per lane from twelve different human tissues and a 1.5 kb fragment containing the full-length open reading frame of *S1P₅* is used as a probe. As shown above, *S1P₅* is expressed mainly in brain, spleen, and peripheral blood leukocytes. Trace amounts of *S1P₅* are also expressed in lung, placenta, liver and kidney. (Please note: Signals are significantly stronger in normal tissue on poly A+RNA Northern blot compared to total RNA Northern blot.)

DETAILED DESCRIPTION OF THE INVENTION

The abbreviations for amino acids, peptides, base sequences, nucleic acids and so forth as used herein in the present specification are those recommended by the International Union of Pure and Applied Chemistry (IUPAC) and the International Union of Biochemistry (IUB) and in the "Guidelines for drafting patent specifications relative to base sequences and/or amino acid sequences" edited by the Japanese Patent Office or those commonly used in the relevant field of art.

Although the genes of the present invention is represented by a single-stranded DNA sequence, as shown under, for example, SEQ ID NO:4, the present invention also includes the DNA sequence complementary to such a single-stranded DNA sequence as well as a component comprising both of these. The DNA sequence representing the gene of the present invention shown in the above-mentioned SEQ ID NO: 4 is an example of the codon combination coding for the respective amino acid residues according to the amino acid sequence shown in SEQ ID NO:7. The gene of the present invention is not limited to the above-mentioned one but may, of course, have any other DNA base sequence comprising a combination of codons arbitrarily selected for the respective amino acid residues without altering the above-mentioned amino acid sequence. Selection of said codons can be carried out by

5

the conventional method in which the codon usage or codon choice in the host to be used for gene recombination is taken into consideration [Nucl. Acids Res., 9, 43-74 (1981)], and these codons can be produced, for example by chemical synthesis, etc.

The gene of the present invention further includes DNA sequences coding for those equivalents to the above-mentioned amino acid sequence that are derived from the latter by deletion, addition or like modification of one or more amino acid residues or part of the amino acid sequence and have similar sphingosine 1-phosphate receptor activity to that of the sphingosine 1-phosphate receptor protein. While production, alteration (mutation) or the like of these polypeptides may occur spontaneously, they can also be produced by post-translational modification. Furthermore, any desired gene can be produced by gene engineering techniques such as the site-specific mutagenesis technique in which the natural gene (gene of the present invention) is altered, by a chemical synthesis technique such as the phosphite triester method in which mutant DNAs are synthesized or by combining both procedures. By utilizing the gene of the present invention, namely by incorporating the same into a vector for use with a microorganism, for instance, and cultivating the transformant microorganism, the sphingosine 1-phosphate receptor protein can be expressed readily and in large quantities, and said protein can be isolated and provided. Since said protein has sphingosine 1-phosphate receptor activity, it is effective for various pharmacological purposes, and it is also useful, among others, in elucidating the pathogenesis, the pathologies or the like of various diseases. More specifically, the recombinant sphingosine 1-phosphate receptor protein obtained by utilizing the gene of the present invention can effectively be used, for example, in elucidating the mechanism of immunosuppression in living bodies, developing or screening out therapeutic agents for autoimmune diseases (e.g. rheumatism, SLE (systemic lupus erythematoses), LGL, etc.), searching for endogenous ligands and substrates to the novel protein and developing therapeutic agents therefor.

Similarly, the gene of the present invention can effectively be used, for example, in elucidating the mechanism of neurodegeneration in living bodies, developing or screening out therapeutic agents for neurodegenerative diseases (e.g. alzheimers, parkinson's and the like), searching for endogenous ligands and substrates to the novel protein and developing therapeutic agents therefor.

In the following, the gene of the present invention will be described in more detail. The gene of the present invention can be isolated by general genetic engineering techniques, for example, by selecting an appropriate clone from among a human fetal brain cDNA library (cDNA synthesized in the conventional manner from mRNA isolated and purified from total RNA obtained in turn from appropriate origin cells containing a gene coding for the sphingosine 1-phosphate receptor protein) using appropriate probes, such as for example those of SEQ ID 1 and SEQ ID2, purifying said clone, and determining the base sequence thereof. In the above procedure, the origin cells may be any animal cells or tissues where the occurrence of sphingosine 1-phosphate receptor protein is known (see for example, the experiment producing the results shown in FIG. 6), or soluble fractions of cultured cells derived therefrom. This can be isolated and purified for the culture supernatant by various chromatographic processes.

In the practice of the present invention, it is also possible to use a part of the DNA fragment sequenced in the above manner as a probe, label this using a random prime DNA labeling kit (available from Takara Shuzo, Amersham, etc.) in

6

accordance with the random prime DNA labeling method (Feinberg, A. P., et al., Anal. Biochem., 137, 266-267 (1984)), for instance, and use the thus-obtained labeled probe in screening out the desired sphingosine 1-phosphate receptor protein gene.

Using the above-mentioned labeled probe, for instance, the desired DNA can be screened out by the plaque hybridization technique developed by Benton and Davis (Benton, W. and Davis, R., Science, 196, 383-394 (1977)).

The gene of the present invention as obtained in the above manner can be cloned in various plasmids in the conventional manner. For instance, after cleavage with an appropriate restriction enzyme and purification, the gene of the present invention can be inserted into a cloning vector (e.g. plasmid) cleaved with the same restriction enzyme and purified, at the cleavage site thereof, whereby a recombinant plasmid can be obtained. By introducing said recombinant into an appropriate host (e.g. *Escherichia coli*) for transformation, a restriction enzyme map of the clone containing said gene can be drawn using the transformant by a conventional known method, for example the method as described in Sambrook, J. Fritsch, E. F., and Maniatis. Molecular cloning. A laboratory Manual 2nd edition. Cold Spring Harbor laboratory Press. Cold Spring Harbor, N.Y. After digestion of the above clone with an appropriate restriction enzyme, the base sequence of said clone can be determined by the above-mentioned dideoxy method or the Maxam-Gilbert method, for instance. The base sequence determination mentioned above may also be readily performed using a commercially available kit or the like.

The thus-determined DNA base sequence of the sphingosine 1-phosphate receptor protein gene of the present invention and the corresponding amino acid sequence encoded thereby are as shown in the sequence listing under SEQ ID NO: 3 and SEQ ID NO:4.

Using the above-mentioned gene (DNA) of the present invention, the recombinant sphingosine 1-phosphate receptor protein can be obtained by various known gene recombination techniques [cf. for example Science, 224, 1431 (1984); Biochem. Biophys. Res. Comm., 130, 692 (1985); Proc. Natl. Acad. Sci. USA, 80, 5990 (1983)]. Said sphingosine 1-phosphate receptor protein is produced, in more detail, by constructing a recombinant DNA allowing expression of the gene of the present invention in host cells, introducing this into host cells for transformation thereof, and cultivating the transformant strain. The host cells may be either eukaryotic or prokaryotic. As an expression vector for use with vertebrate cells, it is possible to use one containing a promoter generally located upstream of the gene to be expressed, an RNA splicing site, a polyadenylation site and a transcription termination sequence and so on. This may further have a replication origin, as necessary. Yeasts are often and generally used as eukaryotic microorganisms and, among them, yeasts belonging to the genus *Saccharomyces* are advantageously used. Usable as expression vectors for use with said yeasts and other eukaryotic microorganisms are pAM82 (A. Miyano-hara et al., Proc. Natl. Acad. Sci. USA, 80, 1-5 (1983)) containing a promoter for the acid phosphatase gene, and like vectors. *Escherichia coli* and *Bacillus subtilis* are generally and very often used as prokaryotic host cells. When these are used as hosts in the practice of the present invention, an expression plasmid is preferably used which is derived, for instance, from a plasmid vector capable of replication in said host microorganisms and provided with a promoter, the SD (Shine and Dalgarno) base sequence and further an initiation codon (e.g. ATG) necessary for the initiation of protein synthesis, upstream from the gene of the present invention so that

said gene can be expressed. As the host *Escherichia coli* mentioned above, the strain *Escherichia coli* K12 and the like are often used and, as the vector, pBR322 is generally and often used. However, the host and vector are not limited thereto, but other various known microbial strains and vectors can also be used. As regards the promoter, the tryptophan (trp) promoter, 1 pp promoter, lac promoter and P.sub.L promoter, for instance, can be used.

The thus-obtained desired recombinant DNA can be introduced into host cells for transformation thereof by various conventional methods. The transformant obtained can be cultivated in the conventional manner, leading to production and accumulation of the desired sphingosine 1-phosphate receptor protein encoded by the gene of the present invention. The medium to be used in said cultivation can adequately be selected, according to the host cells employed, from among various media in common use. When *Escherichia coli* or like cells are used as host cells, for instance, transformant cultivation can be conducted using LB medium, E medium, M9 medium, M63 medium or the like. To these media, there may be added, as necessary, generally known various carbon sources, nitrogen sources, inorganic salts, vitamins, nature-derived extracts, physiologically active substances, etc. The above-mentioned transformant cultivation can be carried out under conditions suited for the growth of the host cells. In the case of *Escherichia coli*, such conditions can be employed, for instance, as a pH of about 5 to 8, preferably 7 or thereabout, and a temperature of about 20 to 43.degree. C., preferably 37.degree. C. or thereabout. In the above manner, the transformant cells produce and accumulate intracellularly or secrete extracellularly the desired recombinant FK506 binding protein.

Said desired protein can be isolated and purified by various separation techniques utilizing its physical, chemical and other properties [cf. for example "Seikagaku (Biochemistry) Data Book II", pages 1175-1259, 1st edition, 1st printing, published Jun. 23, 1980 by Kabushiki Kaisha Tokyo Kagaku Dojin; Biochemistry, vol. 25, No. 25, 8274-8277 (1986); Eur. J. Biochem., 163, 313-321 (1987)]. As specific examples of said techniques, there may be mentioned conventional reconstitution treatment, treatment with a protein precipitating agent (salting out), centrifugation, osmotic pressure shock treatment, ultrasonic disruption, ultrafiltration, various liquid chromatographic processes such as molecular sieve chromatography (gel filtration), adsorption chromatography, ion exchange chromatography, affinity chromatography and high performance liquid chromatography (HPLC), dialysis, and combinations of these. In the above manner, the desired recombinant protein can be produced on an industrial scale with ease and with high efficiency.

In order to provide diagnostics for LGL leukemia, and provide therapeutic targets for drugs directed to mitigate the pathogenesis of LGL leukemia, microarray analysis is performed to identify differentially expressed genes. A large number of genes are identified that are differentially expressed in LGL leukemia compared to normal controls. One of the ESTs of approximately 300 base pairs is fully characterized herein. Initial Blast analysis shows 100% homology with Homo-sapiens full-length insert cDNA clone YY 85D04 (gb/AF 088014). No open reading frame within the full-length insert cDNA. Therefore, in order get the complete sequence of the gene, the LGL leukemia library is screened and also RT-PCR is performed using the total RNA isolated from different LGL leukemia patients. 15 positive clones are selected from library screening. All of them give partial sequences with the longest one being approximately 340 base pairs shorter (clone 6). BLAST search with htgs,

shows that clone 6 shows 100% homology with genomic sequence present in the chromosome 19. Primers are designed based on the genomic sequence information to obtain full-length sequence of the gene. By using these primers in the PCR with genomic DNA and RT-PCR with total RNA, the full-length gene, SEQ ID:4 is obtained. This gene belongs to the G-protein-coupled receptor super-family of integral membrane proteins. BLAST analysis of the complete gene reveals 85% homology with rat sphingosine 1-phosphate receptor edg-8 and nrg-1 (Im, D., et al., Characterization of a Novel sphingosine 1-Phosphate receptor, Edg-8. J. Biol. Chem. 275. 14281-14286 (2000); Glickman, M., et al., Molecular cloning, tissue-specific expression and chromosomal localization of a novel nerve growth factor regulated G-protein-coupled receptor, nrg-1. Mol. Cell. Neurosci. 14, 141-152 (1999)), shown in FIG. 4. It is interesting to note that this gene is present mainly in brain, spleen and PBMCs (FIG. 7), and it is over expressed in PBMC of LGL leukemia patients and is involved in LGL leukemia cell survival or proliferation.

Material and Methods:

Isolation of Peripheral blood mononuclear cells (PBMC and RNA). PBMC are isolated from normal healthy individuals and from LGL leukemia patients. Trizole is obtained from GTBCO-BRL. EST (GenBank ID 1868427) is obtained Incyte Genomics. Oligotex mRNA mini-kit, plasmid isolation kits, gel extraction kits, and PCR reagents are purchased from Qiagen; RNA loading dye is from Sigma Chemical Co. The Prime-a-Gene labeling kit is from Promega Corp. (Madison, Wis.). Deoxycytidine 5' triphosphate dCTP a-32P (3,000 Ci/mmol) is from Dupont NEN (Boston, Mass.). Nytran membrane is obtained from Schleicher & Schuell, Inc., 10 optical Avenue, Keene, N. H. Nick translation columns are obtained from Pharmacia Chemical Co. The Topo-TA cloning kit is from Invitrogen.

PBMC are isolated from whole blood using Ficoll-Hypaque density gradient centrifugation. The PBMC cells are suspended in Trizole reagent (GIBCO-BRL, Rockville, Md.) and total RNA is immediately isolated according to the Oligotex mRNA mini-kit manufacturer's instructions and stored at -70° C. Poly A+ RNA is isolated from total RNA by using Oligo-Tex mini mRNA kit according to the manufacturer's recommendations. PBMCs are cultured in vitro and activated by Interleukin 2 and phytohemagglutinin (PHA) for 2 to 3 days. In a preferred embodiment, PBMC is cultured in vitro and activated by PHA, (Sigma Chemical Co. St. Louis, Mo.) (1 µg/ml, 2 days) and Interleukin-2 (IL-2) (100 U/ml, 10 days), Next, total RNA is isolated as described above.

Microarray probing and analysis is done by Incyte Genomics, (St. Louis, Mo.). Approximately 1 µg of Poly (A)+ RNA isolated from PBMCs of LGL leukemia and healthy individual is reverse transcribed to generate Cys3 and Cys 5 fluorescently labeled cDNA probes. In a preferred embodiment, more than 90% of PBMC from the LGL leukemia patient are leukemic LGL as indicated by CD 8+ staining. cDNA probes are competitively hybridized to a human UniGEM V cDNA microarray containing approximately 7075 immobilized cDNA fragments (4107 known genes and 2968 ESTs). Scanning and quantitation is performed by Incyte Genomics and balanced differential differentiation is given for all the genes. The balanced differential expression is calculated using the ratio between the P1 signal (intensity reading for probe 1) and the balanced P2 signal (intensity reading for probe 2 adjusted using the balanced coefficient). A balanced differential expression of 2.0 is considered indicative of up-regulation of a given gene.

Verification of clones: GEM cDNA clones are purchased from Incyte Genomics as individual bacterial stabs and streaked on LB/agar plates containing appropriate antibiotic(s). Individual colonies are picked and grown in LB medium. Plasmid DNA is isolated and sequenced in order to verify the correct identity of each clone.

Northern Blot analysis: Northern Blotting is done as described previously (Sambrook et al, 1998). Essentially, 10 ug of total RNA from each sample is denatured at 65° C. in a RNA loading buffer, electrophoresed in 1% agarose containing 2.2 M formaldehyde gel, and blotted onto a Nytran membrane. (Nytran membrane obtained from Schleicher & Schuell, Inc, Keene, N.H). The RNA is fixed to the membrane by UV cross-linking. cDNA is labeled with [³²P] (Prime-a-Gene labeling kit from Promega Corp. Madison, Wis., deoxycytidine 5' triphosphate (dCTP α -³²P, 3,000 Ci/mmol, Dupont NEN, Boston, Mass.) and purified by Nick columns (Amersham Pharmacia Biotech AB, Piscataway, N.J.). Hybridization and washings of the blots are performed as described by Engler-Blum, G., Meier, M., Frank, J., and Muller, G. A. Reduction of background in problems in non-radioactive Northern blot analysis enables higher sensitivity than 32P-based hybridizations. Anal. Biochem. 210, 235-244 (1993).

Library construction and screening. cDNA is synthesized from poly(A)⁺ RNA isolated from pooled PBMCs of multiple LGL leukemia patients using oligo dT primer. The cDNA is unidirectionally inserted the EcoRI/XhoI sites of Lambda ZAPII (Stratagene). cDNA library is screened using EST according to standard protocol (Sambrook et al., 1989). In a preferred embodiment, DNA libraries are plated at a density of 50,000 plaque-forming units per 150 mm plate. Following incubation for 8 h at 37° C., the plated phage are overlaid with nitrocellulose filters. After 1 min the filters are removed and the membranes are crossed linked by Autocross linker. A [³²P] labeled cDNA fragment derived from an EST (GenBank accession No. N 47089) of interest is used to probe the filters. Hybridizations, washings, exposure of the membranes to films and then picking up the colony of interest are performed as outlined in the standard methodology (Sambrook et al., 1989). Secondary and tertiary screenings were also performed as outlined in standard methodology (Sambrook et al., 1989). After isolation of pure phage containing the gene of interest, mini-preparations or macro-preparation are made to isolate plasmid cDNA containing the gene of interest.

RT-PCR: To obtain the full-length sequence, 5' and 3' primers are designed based on the sequence information available in GenBank:

5' GCGCGGCCCAT GGAGTC 3' (SEQ.ID # 1)

is used as forward primer and

5' CTTTCTGTGTTCCCAAGC AGAAC GTCAAT 3' (SEQ.ID # 2)

is used as reverse primer. Total RNA from PBMC isolated from LGL leukemia patients and normal healthy individuals is used as a template for reverse transcriptase for making cDNA using either oligo(dT) primer or random hexamer primers. The PCR reaction mixture is heated to 95° C. for 2 min and then cycled 40 times at 95° C. for 30 sec, 60° C. for 45 sec, and 72° C. for 1.5 min. Finally, the reaction mixture is heated at 72° C. for 7 min and stored at 4° C. The reaction product is electrophoresed in 1% agarose gels. For direct PCR, all the conditions are the same as above except that

genomic DNA, isolated from PBMC, is used as a DNA template. PCR products are analyzed in 1% agarose gel and the bands are excised and cloned into a TOPO-TA cloning vector (Invitrogen) and sequenced. The insert is subcloned into EcoRI sites of mammalian expression vector pcDNA3.1 to produce pcDNA3S1P₅.

Cell culture and transfection. HEK293 cells are grown in Dulbecco's modified eagle's medium supplemented with 10% fetal bovine serum. The cells are transiently transfected with a plasmid encoding HA-tagged Erk2 (HA-Erk2) and either pcDNA 3 S1P₅ or pcDNA 3.1. Transfection is achieved by incubating the cells in 60 mm plates with plasmid/Lipofectamine complexes (2.1 μ g total DNA/12 μ l Lipofectamine) in serum-free medium for 5 hours. The DNA complexes are removed from the medium and the cells are starved overnight in serum-free medium and then used for experimentation.

Erk2 Kinase Assay. The serum-starved transiently transfected HEK293 cells are treated for 5 min preferably with either 1 μ M sphingosine-1-phosphate (S1P) or with 1 μ M lysophosphatidic acid (LPA). The cells are lysed in buffer containing 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM DTT, 1% Triton X-100, 25 mM NaF, 5 mM sodium pyrophosphate, 20 mM ρ -nitrophenyl phosphate, 2 μ g/mL leupeptin, and 100 μ g/mL phenylmethylsulfonyl fluoride. HA-Erk2 is immunoprecipitated with the monoclonal antibody HA.11 (Convance, Richmond, Calif.). Half of the immunoprecipitate is used to determine Erk 2 activity and the other half is used for measuring Erk2 protein expression. For the kinase assay, immune complexes are incubated for 10 min at 30° C. in 40 μ l of buffer containing 20 mM Hepes, pH 7.5, 10 mM MgCl₂, 1 mM dithiothreitol, 10 mM p-nitrophenyl phosphate, 40 μ M ATP and 0.375 mg/mL myelin basic protein and 10 μ Ci of [γ -³²P] ATP (3000 Ci/mmol). The reaction is terminated with SDS-containing gel-loading buffer and the reaction mixtures are analyzed on 11% SDS-polyacrylamide gels. The gels are processed by autoradiography. The bands on the gels are quantitated with a Phosphorimager. Erk2 protein in the immunoprecipitate is determined by immunoblotting with a polyclonal antibody to Erk2.

EXAMPLES

Referring now to FIG. 1, approximately 50 genes are up-regulated in LGL leukemia, with balanced differential expression of between about 7.8 and about 2.0. In addition, one EST is particularly noteworthy that is up-regulated in LGL leukemia with balanced differential expression of 3.0 (GenBank Accession number N47089). A clone containing this EST is sequenced. The total length of the EST is approximately 300 base pairs. A search using Blast shows 100% homology with another EST (GenBank Accession No. AF088014) named as *homo sapiens* full length insert cDNA clone YY85D04. No other information regarding this EST is found in the literature. No open reading frame is found within this sequence. Northern blot analysis confirms that a gene related to EST (GenBank ID No. N47098) is upregulated in majority of LGL leukemia patients.

Using the microarray screening method, one LGL leukemia patient is compared with one normal healthy individual. To show the same pattern in a larger sample of patients, Northern blot analysis is performed. Total RNAs, isolated from the PBMC of normal healthy individuals and LGL leukemia patients, are used in Northern blots. Initially, a 300 base pair cDNA fragment is used as a probe in initial experiments. Up-regulation of EST is observed in all the LGL leukemia patients when compared to the normal healthy individuals.

11

This confirms the microarray results regarding EST expression. The probe hybridizes to a 2.2 kb transcript in the Northern Blots. (FIG.-2).

An LGL leukemia library is constructed from the mRNA isolated from the pooled PBMCs of the seven LGL leukemia patients. This library is screened to obtain full-length sequence of the gene. Approximately 15 positive clones are selected and the larger clones are sequenced. The largest clone is 1500 bp in length. Analysis using Blast indicates that this gene has 85% homology with Rat edg-8 (Im et al, 2000). All of the clones are missing 5' end of the gene. Blast search with htgs show 99% homology with the sequence present in chromosome 19. Based on the sequence information, primers are designed from the 5' end and from 3' end of the open reading frame of the gene. Three different products (1.5, 1.6, and 1.2 bp in length) are obtained using RT-PCR. These products are subjected to gel electrophoresis and bands are excised, cloned into TOPO-TA cloning vectors and sequenced. The largest PCR product contains the entire open reading frame (FIG. 3). The deduced amino acid sequence shows 85% homology with complete sequence of rat sphingosine 1-phosphate receptor edg-8 and nrg-1. (FIG. 4). Shorter bands are also identified. The shorter bands are excised, cloned, and sequenced. These clones are splice variants of sphingosine 1-phosphate receptor with deletions. They are herein termed "sphingosine 1-phosphate receptor-1" and "sphingosine 1-phosphate receptor-2" (FIGS. 5 & 6).

Expression of sphingosine 1-phosphate receptor is examined in different normal tissues by Northern blot analysis. It is found that *sppr* is expressed in several tissues such as brain, spleen and PBMCs. (FIG. 7). Only trace amounts are detected in Jukat and CEM cell lines (data not shown).

To obtain a full-length sequence of the gene, an LGL leukemic cDNA library is constructed and screened using the EST probe. Approximately 15 positive clones are selected and larger clones are sequenced. The BLAST search of the largest clone (1500 bp) indicates that this gene has strong homology with Rat edg-8/Nrg-1. However, all of the clones are missing the 5' end of the gene when compared to the rat gene. A BLAST search with the human genome shows 99% homology with a sequence present on chromosome 19. Based on this sequence information, primers are designed from the 5' and 3' ends of the open reading frame of the gene.

Three different RT-PCR products (1.5, 1.6, and 1.2 bp) are obtained. These products are subjected to gel electrophoresis. The resulting bands are excised and cloned into TOPO-TA cloning vectors and then sequenced. The largest PCR product contains a complete open reading frame. The nucleotide sequence and the deduced amino acids are shown in FIG. 8. The gene is designated as SIP₅ (see below). The nucleotide sequence shows very strong homology with G-protein coupled receptors, especially with the endothelial differentiation genes (EDGs). When the deduced amino acid sequence of the full-length sequence is aligned with other members of the family using the CLUSTALW (multi sequence alignment) program, it is approximately 26 to 44% identical and 58 to 72% similar with EDGs at amino acid level (Table 1). In addition, it shows 86% identity and 96% similarity with rat edg-8 or rat nrg-1 at amino acid level. (FIG. 9, Table I). Transient transfection of HEK293 cells with this gene results in activation of Erk2 activity in response to sphingosine-1-phosphate but not LPA, confirming that it is a sphingosine-1-phosphate receptor (FIG. 10). Therefore, this gene is named SIP₅.

Samples from 30 LGL leukemia patients are tested for the presence of SIP₅ transcript by Northern blot analysis using full-length gene as a probe. Constitutive expression of SIP₅

12

transcripts is found in 24 samples (FIG. 11). In comparison SIP₅ transcripts are expressed at only trace levels in normal PBMC (N=12). After activation of normal PBMC the expression of SIP₅ is reduced to undetectable levels (FIG. 12). Additionally, expression of two smaller bands is detected in samples from leukemic LGL by RT-PCR. Human SIP₅ transcripts are expressed mainly in normal brain, spleen, and PBMC and in trace amounts in lung, kidney and liver (FIG. 13). Whereas expression of Edg-8 is observed only in brain and spleen of rat when Northern Blots are probed. Several cell lines are examined for the presence of SIP₅ transcript. Trace amounts of SIP₅ transcripts are identified in CEM and Jurkat cells (data not shown). All other cell lines tested are negative for SIP₅ transcript including MT2 (HTLV-I infected cell line) and MO-T (HTLV-II infected cell line), Moe7 (megakaryoblastic leukemic cell line) and U293 (human embryonic kidney cells).

TABLE 1

Identity and similarity between SIP₅ and other members of the Edgs. The deduced amino acid sequence of SIP₅ is aligned with the amino acid sequences of various members of Edgs. using the CLUSTALW program. Except for Edg 8 and nrg-1, all other sequences are from human. All the sequence information is obtained from GenBank.

Name of the gene	% Identity	% Similarity
hSiP5	100	100
rEdg-8*	87	96
hNrg-1	86	98
h Edg-1*	44	72
hEdg-5*	41	66
h Edg-3*	40	70
h Edg-6*	39	67
h Edg-2*	35	67
h Edg-4*	30	60
hEdg-7*	26	58

* = Sphingosine 1-phosphate receptors

* = Lysophosphatidic acid receptors

Discussion

Leukemic LGL are resistant to Fas-induced apoptosis, in spite of over-expression of Fas and Fas-ligand (FasL) implying that the accumulation of circulating LGL can be due to dysregulation of apoptosis. The accumulation of circulating LGL in leukemic patients can also be due to clonal proliferation of LGL. In order to understand the molecular mechanisms involved in pathogenesis of LGL leukemia, microarray techniques are used to identify differentially expressed genes. Approximately 50 genes are identified that are up-regulated and 10 genes that are down regulated. Several ESTs are also identified which show differential expression. As a systematic study, one of the ESTs that is up-regulated in LGL Leukemia is characterized. The full-length gene is obtained by screening the LGL leukemia library and performing RT-PCR, which is 85% identical to the rat Sphingosine-1 Phosphate receptor. This gene belongs to G-protein coupled receptor super family and can act as a sphingosine-1-phosphate receptor. Several splice variants in LGL leukemia patients are also identified, and are named Sphingosine 1-phosphate receptor 1 and Sphingosine 1-Phosphate receptor 2. The deduced amino acid sequence of Sphingosine 1-Phosphate receptor with rat edg-8 or nrg shows 85% homology. It has seven transmembrane domains, which is a characteristic of GTP-coupled receptors. Thus, the Sphingosine-1 Phosphate is involved in the signal transduction from the sphingosine 1-Phosphate in human.

Although the gene has lot of homology with other members of edg family, it is preferably named sphingosine-1-

13

phosphate receptor (S1P₅) because it is mainly present in lymphocytes, brain and spleen, but not in endothelial cells.

Lysophosphatidic acid (LPA) and sphingosine 1-phosphate (S1P) mediate T cell function. Both LPA and S1P signaling pathways are implicated in cell proliferation, suppression of apoptosis, enhancement of cellular survival and T-lymphoma cell invasion. Although it has been suggested that S1P can act as an intracellular mediator, it has also been suggested that S1P acts as an extracellular ligand for cell surface receptors. Indeed several such receptors have been identified in a wide variety of tissues. For example, receptors Edg-1, -3, -5, -6 and -8, are specific for S1P, whereas Edg-2, -4, and -7 are LPA specific. In normal lymphocytes, there is differential constitutive expression of receptors for LPA and S1P. CD4⁺ cells express predominantly Edg-4, while CD8⁺ cells appeared to lack receptors for LPA and S1P as only traces of Edg-2 and Edg-5 are detected. Human T cell tumors express many Edgs for both LPA and S1P.

Rat edg-8/nrg-1 is shown to be a sphingosine-1-phosphate receptor based on specific binding of radio-labeled S1P to cell membranes, inhibition of forskolin-induced cAMP accumulation, increased GTP binding ability and calcium mobilization studies. Even though these properties are adequate to classify edg-8/nrg-1 as a sphingosine-1-phosphate receptor, it seems surprising that this acene is different from other members of the human sphingosine-1-phosphate receptor family. For example, activation of EDG-1, -3, -5 and -6 by S1P leads to activation of Erk1/2 and induction of cell proliferation. In contrast S1P inhibited serum-induced activation of Erk1/2 and also inhibits the cell proliferation in CHO cells expressing EDG-8. The reasons for these differences are not known and might be due to species variation. As shown herein, S1P

14

activates Erk2 in transiently transfected HEK293 cells while lysophosphatidic acid does not, suggesting that S1P₅ is a sphingosine-1-phosphate receptor and participates in sphingosine 1-phosphate mediated signal transduction. A computational model of the Edg-1 receptor predicts that Glu¹²¹ is essential for interaction with S1P [21]. The S1P receptors Edg-1, -3, -5 and -8 as well as S1P₅ share such an anionic residue.

Leukemic LGL are antigen driven CTL that survive in vivo, at least in part, because of defective apoptosis. For example, leukemic LGL express both Fas and Fas-ligand, but are resistant to Fas mediated death. It is noteworthy that S1P₅ gene transcripts are down regulated after activation of normal T cells. Leukemic cells are activated T cells. Based upon the results disclosed herein, constitutive expression of S1P₅ transcripts represents dysregulated expression. This dysregulated expression of S1P₅ may participate in protection of leukemic LGL from apoptosis.

Note: The full-length sequence was deposited in GenBank (Accession No. AF331840) on Dec. 22, 2000.

Throughout this application, various publications, including United States patents, have been referred to. The disclosures of these publications and patents in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

While the invention has been described in terms of various preferred embodiments, those skilled in the art will recognize that various modifications, substitutions, omissions, and changes may be made without departing from the spirit of the present invention. Accordingly, it is intended that the scope of the present invention be limited solely by the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 19

<210> SEQ ID NO 1
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Probe (see p. 8 of specification)

<400> SEQUENCE: 1

gcgcggccca tggagtc

17

<210> SEQ ID NO 2
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Probe (see p. 8 of specification)

<400> SEQUENCE: 2

cttttctgtg ttccaagca gaacgtcaat

30

<210> SEQ ID NO 3
 <211> LENGTH: 398
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <223> OTHER INFORMATION: Human sphingosine 1-Phosphate receptor (SPPR)
 amino acid sequence (Figure 3)

-continued

<400> SEQUENCE: 3

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
 1 5 10 15
 Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
 20 25 30
 Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
 35 40 45
 Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
 50 55 60
 Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
 65 70 75 80
 Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Asn Ile Leu Leu Ser
 85 90 95
 Gly Pro Leu Thr Leu Lys Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
 100 105 110
 Gly Gly Val Phe Val Ala Leu Thr Ala Ser Val Leu Ser Leu Leu Ala
 115 120 125
 Ile Ala Leu Glu Arg Ser Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
 130 135 140
 Val Ser Ser Arg Gly Arg Thr Leu Ala Met Ala Ala Ala Ala Trp Gly
 145 150 155 160
 Val Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
 165 170 175
 Gly Arg Leu Asp Ala Cys Ser Thr Val Leu Pro Leu Tyr Ala Lys Ala
 180 185 190
 Tyr Val Leu Phe Cys Val Leu Ala Phe Val Gly Ile Leu Ala Ala Ile
 195 200 205
 Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg
 210 215 220
 Arg Leu Pro Ala Arg Pro Gly Thr Ala Gly Thr Thr Ser Thr Arg Ala
 225 230 235 240
 Arg Arg Lys Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val Val
 245 250 255
 Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu Leu
 260 265 270
 Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala
 275 280 285
 Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
 290 295 300
 Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
 305 310 315 320
 Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
 325 330 335
 Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
 340 345 350
 Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
 355 360 365
 Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
 370 375 380
 Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
 385 390 395

<210> SEQ ID NO 4

<211> LENGTH: 2336

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: Human sphingosine 1-Phosphate receptor (SPPR)
        cDNA sequence (Figure 3)

<400> SEQUENCE: 4

gcgcgGCCCA tggagtcggg gctgctgcgg ccggcgccgg tgagcgaggt catcgctctg      60
cattacaact acaccgcaa gctccgcggt gcgcgctacc agccgggtgc cggcctgcgc      120
gccgacgcgc tgggtgtgct ggcggtgtgc gccttcacgc tgctagagaa tctagccgtg      180
ttgttggtgc tcggacgcca ccgcgccttc cacgcctcca tgttcctgct cctgggcagc      240
ctcacgttgt cggatctgct ggcaggcgcc gctacgcgc ccaacatcct actgtcgggg      300
ccgctcacgc tgaactgtc ccccgcgctc tggttcgcac gggaggagg cgtcttcgtg      360
gcactcactg cgtccgtgct gagcctcctg gccatcgcgc tggagcgcag cctcaccatg      420
gcgcgcaggg ggcgcgcgcc cgtctccagt cgggggcgca cgtcggcgat ggcagccgcg      480
gctcggggcg tgcgctgct cctcgggctc ctgccagcgc tgggctggaa ttgctgggt      540
cgcttgagcg cttgctccac tgtcttgccg ctctacgcca aggcctacgt gctctctctg      600
gtgctcgctt tcgtgggcat cctggccgcg atctgtgcac tctacgcgcg catctactgc      660
caggctacgc ccaacgcgcg gcgcctgcgc gcacggccgc ggactgcggg gaccacctcg      720
acccgggcgc gtcgcaagcc gcgctcgctg gccttgctgc gcacgctcag cgtggtgctc      780
ctggcctttg tggcatgtt gggccccctc ttcctgctgc tgttgcctga cgtggcgctg      840
ccggcgcgca cctgtctgt actcctgcag gccgatccct tcctgggact ggccatggcc      900
aactcacttc tgaaccccat catctacacg ctcaccaacc gcgacctgcg ccacgcgctc      960
ctgcgcctgg tctgctcggg acgccactcc tgcggcagag acccgagtgg ctcccagcag     1020
tcggcgagcg cggctgaggc ttccgggggc ctgcgcgcgt gcctgcccc gggccttgat     1080
gggagcttca gcgctcgga gcgctcatcg cccagcgcg acgggctgga caccagcggc     1140
tccacaggca gccccggtgc acccacagcc gcccggaetc tggatcaga accggtgca     1200
gactgacacc ctcggcccac gactgtcttc ccaagtttta cagacttgtt ctttttacat     1260
aaaggaattt gtaggaaatg cagccaaagg tgcagtcgga aaagatgcag gggaaatgta     1320
tttatgcagc gacacccac aatgtgaaca aacagacaaa aaatctgtgc cctcgtggaa     1380
ttgacgtttt ctttgggaac acagaaaaga actcgggtgat gaaataatgg agatgattcc     1440
agtgacaaa cagagagatg gtgatggtgg tcagggaaga cctctctgca gaggtagtga     1500
cttgtgatgt gagctgagac ctctgtcctg ggaagaccaa aagaaaagca tttcaggatg     1560
agggaatggc atgcgcaaag gcctgaggc tgaaatgtgc ccatgtgttc taagaaatgc     1620
agcgatgctg gtgtgcctgg agcaggagcg gagggggaga atgggaggag acaaggagct     1680
gaaggagtag ttcccgaagg accttgtggg tgatatagag gacttcgctt ttgctctgag     1740
tgaggtgagg gccatagaag cttctaagca gaagagggac tgcacctaat tcagggtgat     1800
acagtggtct tgtggcctcc atgggaggtt gaaaaccaca gaaggatgaag gggggctgca     1860
ctgagccaca ggaacaatga tggagattcc agctaagccc agaccccgctg gattctagat     1920
agattttaga ggcagcagac agaattactg aggaattgag tgtaagagtg gaataaagt     1980
atcaaggaca atgccaaggg tggggcacc ccaaatttga ctttgggaga ctgagccaaa     2040
tcctatctgg taataaaatt tcttttttat tttcttttct tttctttctt tctttctttc     2100
tttttttttt tttaggttgg gatcttgcgc tctgtcacc aggctggagt gcaatgggca     2160

```

-continued

```

caattatagc tcaactgcagc ctggaactcc tgggatcaag cctggagttc ctgettcagc 2220
ctccctagta gctgggacta caggcatgca ccaccatgcc cagttaataa aattttcttca 2280
aatgcaaaaa aaaaaaaaaa aaaaaactcg aggggggggcc cggtacccaa ttcgcc 2336

```

```

<210> SEQ ID NO 5
<211> LENGTH: 400
<212> TYPE: PRT
<213> ORGANISM: Rattus norvegicus
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: Nrg-1 rat genes (Figure 4)

```

```

<400> SEQUENCE: 5

```

```

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
1          5          10         15
Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
20        25        30
Gly Ala Gly Leu Arg Ala Asp Ala Ala Val Cys Leu Ala Val Cys Ala
35        40        45
Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
50        55        60
Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
65        70        75        80
Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Thr Asn Ile Leu Leu Ser
85        90        95
Gly Pro Leu Thr Leu Arg Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
100       105       110
Gly Gly Val Phe Val Ala Leu Ala Ala Ser Val Leu Ser Leu Leu Ala
115       120       125
Ile Ala Ile Glu Arg His Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
130       135       140
Ala Ala Ser Arg Ala Arg Thr Leu Ala Met Ala Val Ala Ala Trp Gly
145       150       155       160
Leu Leu Leu Thr Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
165       170       175
Gly Arg Leu Glu Ala Cys Ser Thr Val Leu Pro Val Tyr Ala Lys Ala
180       185       190
Tyr Val Leu Phe Cys Val Leu Ala Phe Leu Gly Ile Leu Ala Ala Ile
195       200       205
Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg
210       215       220
Arg Leu Arg Ala Gly Pro Gly Ser Arg Arg Ala Thr Ser Ser Ser Arg
225       230       235       240
Ser Arg His Thr Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val
245       250       255
Val Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu
260       265       270
Leu Leu Asp Val Ala Cys Pro Ala Arg Ala Cys Pro Val Leu Leu Gln
275       280       285
Ala Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro
290       295       300
Ile Ile Tyr Thr Phe Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg
305       310       315       320
Leu Leu Cys Cys Gly Arg Gly Pro Cys Asn Gln Asp Ser Ser Asn Ser
325       330       335

```

-continued

Leu Gln Arg Ser Pro Ser Ala Val Gly Pro Ser Gly Gly Gly Leu Arg
 340 345 350

Arg Cys Leu Pro Pro Thr Leu Asp Arg Ser Ser Ser Pro Ser Glu His
 355 360 365

Ser Cys Pro Gln Arg Asp Gly Met Asp Thr Ser Cys Ser Thr Gly Ser
 370 375 380

Pro Gly Ala Ala Thr Ala Asn Arg Thr Leu Val Pro Asp Ala Thr Asp
 385 390 395 400

<210> SEQ ID NO 6
 <211> LENGTH: 400
 <212> TYPE: PRT
 <213> ORGANISM: Rattus norvegicus
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <223> OTHER INFORMATION: EDG-8 rat genes (Figure 4)

<400> SEQUENCE: 6

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
 1 5 10 15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
 20 25 30

Gly Ala Gly Leu Arg Ala Asp Ala Ala Val Cys Leu Ala Val Cys Ala
 35 40 45

Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
 50 55 60

Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
 65 70 75 80

Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Thr Asn Ile Leu Leu Ser
 85 90 95

Gly Pro Leu Thr Leu Arg Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
 100 105 110

Gly Gly Val Phe Val Ala Leu Ala Ala Ser Val Leu Ser Leu Leu Ala
 115 120 125

Ile Ala Leu Glu Arg His Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
 130 135 140

Ala Ala Ser Arg Ala Arg Thr Leu Ala Met Ala Val Ala Ala Trp Gly
 145 150 155 160

Leu Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
 165 170 175

Gly Arg Leu Glu Ala Cys Ser Thr Val Leu Pro Leu Tyr Ala Lys Ala
 180 185 190

Tyr Val Leu Phe Cys Val Leu Ala Phe Leu Gly Ile Leu Ala Ala Ile
 195 200 205

Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg
 210 215 220

Arg Leu Arg Ala Gly Pro Gly Ser Arg Arg Ala Thr Ser Ser Ser Arg
 225 230 235 240

Ser Arg His Thr Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val
 245 250 255

Val Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu
 260 265 270

Leu Leu Asp Val Ala Cys Pro Ala Arg Ala Cys Pro Val Leu Leu Gln
 275 280 285

Ala Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro
 290 295 300

-continued

```

Ile Ile Tyr Thr Phe Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg
305                310                315                320

Leu Leu Cys Cys Gly Arg Gly Pro Cys Asn Gln Asp Ser Ser Asn Ser
                325                330                335

Leu Gln Arg Ser Pro Ser Ala Val Gly Pro Ser Gly Gly Gly Leu Arg
                340                345                350

Arg Cys Leu Pro Pro Thr Leu Asp Arg Ser Ser Ser Pro Ser Glu His
                355                360                365

Ser Cys Pro Gln Arg Asp Gly Met Asp Thr Ser Cys Ser Thr Gly Ser
                370                375                380

Pro Gly Ala Ala Thr Ala Asn Arg Thr Leu Val Pro Asp Ala Thr Asp
385                390                395                400

```

```

<210> SEQ ID NO 7
<211> LENGTH: 398
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: SPPR (Figure 4)

```

```

<400> SEQUENCE: 7

```

```

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
1                5                10                15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
                20                25                30

Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
                35                40                45

Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
50                55                60

Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
65                70                75                80

Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Asn Ile Leu Leu Ser
                85                90                95

Gly Pro Leu Thr Leu Lys Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
                100                105                110

Gly Gly Val Phe Val Ala Leu Thr Ala Ser Val Leu Ser Leu Leu Ala
115                120                125

Ile Ala Leu Glu Arg Ser Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
130                135                140

Val Ser Ser Arg Gly Arg Thr Leu Ala Met Ala Ala Ala Ala Trp Gly
145                150                155                160

Val Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
                165                170                175

Gly Arg Leu Asp Ala Cys Ser Thr Val Leu Pro Leu Tyr Ala Lys Ala
                180                185                190

Tyr Val Leu Phe Cys Val Leu Ala Phe Val Gly Ile Leu Ala Ala Ile
195                200                205

Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg
210                215                220

Arg Leu Pro Ala Arg Pro Gly Thr Ala Gly Thr Thr Ser Thr Arg Ala
225                230                235                240

Arg Arg Lys Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val Val
                245                250                255

Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu Leu
260                265                270

```

-continued

Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala
 275 280 285
 Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
 290 295 300
 Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
 305 310 315 320
 Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
 325 330 335
 Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
 340 345 350
 Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
 355 360 365
 Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
 370 375 380
 Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
 385 390 395

<210> SEQ ID NO 8
 <211> LENGTH: 254
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <223> OTHER INFORMATION: Sphingosine-1-phosphate receptor.1 (Figure 5)

<400> SEQUENCE: 8

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
 1 5 10 15
 Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
 20 25 30
 Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
 35 40 45
 Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
 50 55 60
 Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
 65 70 75 80
 Ser Val Pro Ala Arg Pro Gly Thr Ala Gly Thr Thr Ser Thr Arg Ala
 85 90 95
 Arg Arg Lys Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val Val
 100 105 110
 Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu Leu
 115 120 125
 Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala
 130 135 140
 Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
 145 150 155 160
 Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
 165 170 175
 Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
 180 185 190
 Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
 195 200 205
 Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
 210 215 220
 Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
 225 230 235 240

-continued

Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
 245 250

<210> SEQ ID NO 9
 <211> LENGTH: 1698
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <223> OTHER INFORMATION: Sphingosine-1-phosphate receptor.1 (Figure 5)

<400> SEQUENCE: 9

```
cgcgcgcccc atggagtcgg ggctgctgcg gccggcgccg gtgagcgagg tcacgtcct 60
gcattacaac tacaccggca agctccgagg tgcgcgtac cagccgggtg ccggcctgcg 120
cgccgacgcc gtggtgtgcc tggcgggtgtg cgccttcac gtgctagaga atctagccgt 180
gttgtttggtg ctcggaagcc acccgcgctt ccacgctccc atgttcctgc tcctgggcag 240
cctcacgttg tcggtgcccg caccgcccgg gactgcgggg accacctcga cccgggcgcg 300
tcgcaagcgg cgctcgctgg ccttgctgcg cagctcagc gtggtgctcc tggcctttgt 360
ggcatgttgg ggccccctct tcctgctgct gttgctcgac gtggcgtgcc cggcgcgcac 420
ctgtcctgta ctctgcagg ccgatccctt cctgggactg gccatggcca actcaattct 480
gaaccccatc atctacacgc tcaccaaccg cgacctgcgc cagcgctcc tgcgcctggt 540
ctgctgcgga cgccactcct gcggcagaga cccgagtggc tcccagcagt cggcgagcgc 600
ggctgaggct tccggggggc tgcgcgctg cctgcccccg ggcttgatg ggagcttcag 660
cggctcggag cgctcatcgc ccagcgcga cgggctggac accagcggct ccacaggcag 720
ccccggtgca cccacagcgg cccggactct ggtatcagaa ccggctgcag actgacaccc 780
tcggcccaag actgtcttcc caagtcttac agacttgctt tttttacata aaggaatttg 840
taggaaatgc agccaaaggt gcagtcggaa aagatgcagg ggaaatgtat ttatgcagcg 900
acacccca atgtgaacaa acagacaaaa aatctgtgcc ctctggaat tgacgttctg 960
cttggaaca cagaaaagaa ctccgtgatg aaataatgga gatgattcca gtgacaaacg 1020
acagagatgg tgatgggtgg caggggaagac ctctctgcag aggtagtgc ttgtgatgtg 1080
agctgagacc tctgtcctgg gaagacccaa agaaaagcat ttcaggatga gggaatggca 1140
tgcgcaaagg ccctgaggct gaaatgtgcc catgtgttct aagaaatgca gcgatgctgg 1200
tgtgcctgga gcagggacgg agggggagaa tgggaggaga caaggagctg aaggagtagt 1260
tcccgaagga ccttgtgggt gatatagagg acttcgcttt tgcctgagt gaggtgggag 1320
ccatagaagc ttctaagcag aagagggact tgccctaatt caggtgatca cagggtgtct 1380
gtggcctcca tgggaggttg aaaaccagag aaggtgaagg ggggctgcac tgagccacag 1440
gaacaatgat ggagattcca gctaagccca gaccccgagg attctagata gatttttagag 1500
gcagcagaca gaattactga ggaattgagt gtaagagtgg aataaagtta tcaaggacaa 1560
tgccaagggt ggggcacccc caaatttgac tctgggagac tcagccaaat cctatctggt 1620
aataaaatct cttttttatt tttctttctt ttctttcttt cttttttttt tttttgagtt 1680
gggatcttgt gctctgtc 1698
```

<210> SEQ ID NO 10
 <211> LENGTH: 398
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE

-continued

<223> OTHER INFORMATION: Sphingosine-1-phosphate receptor (SIP)
(Figures 14 and 15)

<400> SEQUENCE: 10

```

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
1      5      10      15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
20     25     30

Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
35     40     45

Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
50     55     60

Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
65     70     75     80

Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Asn Ile Leu Leu Ser
85     90     95

Gly Pro Leu Thr Leu Lys Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
100    105    110

Gly Gly Val Phe Val Ala Leu Thr Ala Ser Val Leu Ser Leu Leu Ala
115    120    125

Ile Ala Leu Glu Arg Ser Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
130    135    140

Val Ser Ser Arg Gly Arg Thr Leu Ala Met Ala Ala Ala Ala Trp Gly
145    150    155    160

Val Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
165    170    175

Gly Arg Leu Asp Ala Cys Ser Thr Val Leu Pro Leu Tyr Ala Lys Ala
180    185    190

Tyr Val Leu Phe Cys Val Leu Ala Phe Val Gly Ile Leu Ala Ala Ile
195    200    205

Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg
210    215    220

Arg Leu Pro Ala Arg Pro Gly Thr Ala Gly Thr Thr Ser Thr Arg Ala
225    230    235    240

Arg Arg Lys Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val Val
245    250    255

Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu Leu
260    265    270

Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala
275    280    285

Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
290    295    300

Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
305    310    315    320

Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
325    330    335

Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
340    345    350

Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
355    360    365

Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
370    375    380

Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
385    390    395

```

-continued

```

<210> SEQ ID NO 11
<211> LENGTH: 254
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: Sphingosine-1-phosphate receptor 1 (SIPI)
      (Figure 14)

```

```

<400> SEQUENCE: 11

```

```

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
 1             5             10             15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
      20             25             30

Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
      35             40             45

Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
      50             55             60

Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
      65             70             75             80

Ser Val Pro Ala Arg Pro Gly Thr Ala Gly Thr Thr Ser Thr Arg Ala
      85             90             95

Arg Arg Lys Pro Arg Ser Leu Ala Leu Leu Arg Thr Leu Ser Val Val
      100            105            110

Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu Leu
      115            120            125

Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala
      130            135            140

Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
      145            150            155            160

Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
      165            170            175

Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
      180            185            190

Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
      195            200            205

Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
      210            215            220

Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
      225            230            235            240

Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
      245            250

```

```

<210> SEQ ID NO 12
<211> LENGTH: 103
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: Sphingosine-1-Phosphate receptor 2 (Figure 6)

```

```

<400> SEQUENCE: 12

```

```

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
 1             5             10             15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
      20             25             30

Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
      35             40             45

```

-continued

Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
 50 55 60

Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
 65 70 75 80

Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Ala Arg Thr Leu
 85 90 95

Val Ser Glu Pro Ala Ala Asp
 100

<210> SEQ ID NO 13
 <211> LENGTH: 1245
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <223> OTHER INFORMATION: Sphingosine-1-Phosphate receptor 2 (Figure 6)

<400> SEQUENCE: 13

```

cgcgcgcccc atggagtcgg ggctgctgcg gccggcgccg gtgagcgagg tcatcgctct      60
gcattacaac tacaccggca agctccgagg tgccgcgtac cagccgggtg ccggcctgcg      120
cgccgacgcc gtggtgtgccc tggcgggtgtg cgccttcctc gtgctagaga atctagccgt      180
gttgttggtg ctcggacgcc acccgcgctt ccacgctccc atgttctctg tctggggcag      240
cctcacgttg tcggatctgc tggcaggcgc cgcctacgcc gcccgccccc ggactctggt      300
atcagaaccg gctgcagact gacaccctcg gccacgact gtcttcccaa gttttacaga      360
cttgttcttt ttacataaag gaattttagt gaaatgcagc caaagggtgca gtcggaaaaag      420
atgcaggggg aatgtattta tgcagcgaca ccccaaatg tgaacaaaca gacaaaaaat      480
ctgtgccctc gtggaattga cgttctgctt gggaacacag aaaagaactc ggtgatgaaa      540
taatggagat gattccagtg acaaacgaca gagatggtga tgggtggtcag ggaagacctc      600
tctgcagagg tagtgacttg tgatgtgagc tgagacctct gtcctgggaa gacaaaaaga      660
aaagcatttc aggatgaggg aatggcatgc gcaaaggccc tgagggtgaa atgtgccccat      720
gtgttctaa gaaatgcagc atgctgggtg gcctggagca gggacggagg gggagaaatgg      780
gaggagacaa ggagctgaag gagtagttcc cgaaggacct tgtgggtgat atagaggact      840
tcgcttttgc tctgagttag gtgggagcca tagaagcttc taagcagaag agggacttgc      900
cctaattcag gtgatcacag gtgtcttctg gcctccatgg gaggttgaaa accagagaag      960
gtgaaggggg gctgcactga gccacaggaa caatgatgga gattccagct aagcccagac     1020
cccgtggatt ctagatagat tttagaggca gcagacagaa ttactgagga attgagtgtg     1080
agagtggaat aaagtatatc aggacaatgc caagggtggg gcacccccaa atttgactct     1140
gggagactca gccaaatcct atctggtaat aaaatttctt ttttattttt cttttctttc     1200
tttctttctt tttttttttt ttgagttggg atcttggtct ctgtc                      1245

```

<210> SEQ ID NO 14
 <211> LENGTH: 103
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <223> OTHER INFORMATION: Sphingosine-1-Phosphate receptor 2 (SIP2)
 (Figure 15)

<400> SEQUENCE: 14

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
 1 5 10 15

-continued

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
 20 25 30
 Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
 35 40 45
 Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
 50 55 60
 Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
 65 70 75 80
 Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Ala Ala Arg Thr Leu
 85 90 95
 Val Ser Glu Pro Ala Ala Asp
 100

<210> SEQ ID NO 15
 <211> LENGTH: 2306
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <223> OTHER INFORMATION: Full-length (2.2 kb) nucleotide sequence of
 human S1P5 cDNA (Figure 8)
 <400> SEQUENCE: 15

gcgcggccca tggagtcggg gctgctgcgg ccggcgccgg tgagcggagt catcgtcctg 60
 cattacaact acaccggcaa gctccgcggg gcgcgctacc agccgggtgc cggcctgcgc 120
 gccgacgcgc tgggtgtgct ggcgggtgtgc gccttcacgc tgctagagaa tctagccgtg 180
 ttgttggtgc tcggacgcca ccgcgccttc cacgctccca tgttcctgct cctgggcagc 240
 ctcacgttgt cggatctgct ggcaggcgcc gcctacgcgc ccaacatcct actgtcgggg 300
 ccgctcacgc tgaactgtc ccccgcgctc tggttcgac gggaggagg cgtcttcgtg 360
 gcactcactg cgtccgtgct gaggcctcct gccatcgcgc tggagcgag cctcaccatg 420
 gcgcgcaggg ggcgcgcgc cgtctccagt cgggggcgca cgctggcgat ggcagccgcg 480
 gcctggggcg tgcgctgct cctcgggctc ctgccagcgc tgggctggaa ttgctgggt 540
 cgcttggaag cttgctccac tgtcttgccg ctctacgcca aggcctacgt gctctctcgc 600
 gtgctcgctc tcgtgggcat cctggccgcg atctgtgac tctacgcgcg catctactgc 660
 caggtacgcg ccaacgcgcg gcgcctgccc gcacggcccc ggactgcggg gaccacctcg 720
 acccgggcgc gtcgcaagcc gcgctcgctg gccttgctgc gcacgctcag cgtgggtgctc 780
 ctggcctttg tggcatgttg gggccccctc ttcctgctgc tgttgctcga cgtggcgtgc 840
 ccggcgcgca cctgtcctgt actcctgcag gccgatccct tcctgggact ggccatggcc 900
 aactcacttc tgaaccccat catctacacg ctcaccaacc gcgacctgcg ccacgcgctc 960
 ctgcgcctgg tctgctgcgg acgccactcc tgcggcagag acccgagtgg ctcccagcag 1020
 tcggcgagcg cggtgagggc ttccgggggc ctgcgccgct gcctgcccc gggccttgat 1080
 gggagcttca gcggtcggga gcgctcatcg cccagcgcg acgggctgga caccagcggc 1140
 tccacaggca gccccggtgc acccacagcc gcccggaactc tggatcaga accggctgca 1200
 gactgacacc ctcggcccaac gactgtcttc ccaagtttta cagacttgtt ctttttacat 1260
 aaaggaattt gtaggaaatg cagccaaagg tgcagtcgga aaagatgcag gggaaatgta 1320
 tttatgcagc gacacccac aatgtgaaca aacagacaaa aaatctgtgc cctcgtggaa 1380
 ttgacgttct gcttgggaac acagaaaaga actcgggtgat gaaataatgg agatgattcc 1440
 agtgacaaac gacagagatg gtgatggtgg tcagggaaga cctctctgca gaggtagtga 1500

-continued

```

cttgtgatgt gagctgagac ctctgtcctg ggaagaccaa aagaaaagca tttcaggatg 1560
agggaaatggc atgcgcaaag gccttgaggc tgaaatgtgc ccatgtgttc taagaaatgc 1620
agcgatgctg gtgtgcctgg agcagggacg gagggggaga atgggaggag acaaggagct 1680
gaaggagtag tccccgaagg accttggtgg tgatatagag gacttcgctt ttgctctgag 1740
tgaggtggga gccatagaag cttctaagca gaagagggac ttgccctaat tcaggtgatc 1800
acaggtgtct tgtggcctcc atgggaggtt gaaaaccaca gaaggtgaag gggggctgca 1860
ctgagccaca ggaacaatga tggagattcc agctaagccc agaccccgctg gattctagat 1920
agattttaga ggcagcagac agaattactg aggaattgag tgtaagagtg gaataaagt 1980
atcaaggaca atgccaaggg tggggcacc ccaaatttga ctttgggaga ctcagccaaa 2040
tcctatctgg taataaaatt tcttttttat ttttctttt tttctttctt tctttctttc 2100
tttttttttt tttgagttgg gatcttgtgc tctgtcacc aggtggagt gcaatgggca 2160
caattatagc tcaactgcagc ctggaactcc tgggatcaag cctggagtgc ctgcttcagc 2220
ctccctagta gctgggacta caggcatgca ccaccatgcc cagttaataa aattttcttca 2280
aatgcaaaaa aaaaaaaaaa aaaaaa 2306

```

```

<210> SEQ ID NO 16
<211> LENGTH: 398
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: Deduced amino acid sequence of human S1P5 cDNA
coding region (Figure 8)

```

```

<400> SEQUENCE: 16

```

```

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
1      5      10     15
Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
20     25     30
Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
35     40     45
Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
50     55     60
Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
65     70     75     80
Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Asn Ile Leu Leu Ser
85     90     95
Gly Pro Leu Thr Leu Lys Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu
100    105    110
Gly Gly Val Phe Val Ala Leu Thr Ala Ser Val Leu Ser Leu Leu Ala
115    120    125
Ile Ala Leu Glu Arg Ser Leu Thr Met Ala Arg Arg Gly Pro Ala Pro
130    135    140
Val Ser Ser Arg Gly Arg Thr Leu Ala Met Ala Ala Ala Ala Trp Gly
145    150    155    160
Val Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu
165    170    175
Gly Arg Leu Asp Ala Cys Ser Thr Val Leu Pro Leu Tyr Ala Lys Ala
180    185    190
Tyr Val Leu Phe Cys Val Leu Ala Phe Val Gly Ile Leu Ala Ala Ile
195    200    205
Cys Ala Leu Tyr Ala Arg Ile Tyr Cys Gln Val Arg Ala Asn Ala Arg

```

-continued

210	215	220
Arg Leu Pro Ala Arg	Pro Gly Thr Ala Gly Thr	Thr Ser Thr Arg Ala
225	230	235 240
Arg Arg Lys Pro Arg	Ser Leu Ala Leu Leu Arg Thr	Leu Ser Val Val
	245	250 255
Leu Leu Ala Phe Val Ala Cys Trp Gly Pro Leu Phe Leu Leu Leu		
	260	265 270
Leu Asp Val Ala Cys Pro Ala Arg Thr Cys Pro Val Leu Leu Gln Ala		
	275	280 285
Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile		
	290	295 300
Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu		
	305	310 315 320
Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln		
	325	330 335
Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu		
	340	345 350
Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro		
	355	360 365
Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala		
	370	375 380
Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp		
	385	390 395

<210> SEQ ID NO 17

<211> LENGTH: 398

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<223> OTHER INFORMATION: Predicted amino acid sequence of S1P5 (Figures 12A and 12B)

<400> SEQUENCE: 17

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val		
1	5	10 15
Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro		
	20	25 30
Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala		
	35	40 45
Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His		
	50	55 60
Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu		
	65	70 75 80
Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Asn Ile Leu Leu Ser		
	85	90 95
Gly Pro Leu Thr Leu Lys Leu Ser Pro Ala Leu Trp Phe Ala Arg Glu		
	100	105 110
Gly Gly Val Phe Val Ala Leu Thr Ala Ser Val Leu Ser Leu Leu Ala		
	115	120 125
Ile Ala Leu Glu Arg Ser Leu Thr Met Ala Arg Arg Gly Pro Ala Pro		
	130	135 140
Val Ser Ser Arg Gly Arg Thr Leu Ala Met Ala Ala Ala Ala Trp Gly		
	145	150 155 160
Val Ser Leu Leu Leu Gly Leu Leu Pro Ala Leu Gly Trp Asn Cys Leu		
	165	170 175

-continued

Gly	Arg	Leu	Asp	Ala	Cys	Ser	Thr	Val	Leu	Pro	Leu	Tyr	Ala	Lys	Ala
			180					185					190		
Tyr	Val	Leu	Phe	Cys	Val	Leu	Ala	Phe	Val	Gly	Ile	Leu	Ala	Ala	Ile
		195					200					205			
Cys	Ala	Leu	Tyr	Ala	Arg	Ile	Tyr	Cys	Gln	Val	Arg	Ala	Asn	Ala	Arg
	210					215					220				
Arg	Leu	Pro	Ala	Arg	Pro	Gly	Thr	Ala	Gly	Thr	Thr	Ser	Thr	Arg	Ala
225					230					235					240
Arg	Arg	Lys	Pro	Arg	Ser	Leu	Ala	Leu	Leu	Arg	Thr	Leu	Ser	Val	Val
				245					250					255	
Leu	Leu	Ala	Phe	Val	Ala	Cys	Trp	Gly	Pro	Leu	Phe	Leu	Leu	Leu	Leu
		260						265					270		
Leu	Asp	Val	Ala	Cys	Pro	Ala	Arg	Thr	Cys	Pro	Val	Leu	Leu	Gln	Ala
	275						280					285			
Asp	Pro	Phe	Leu	Gly	Leu	Ala	Met	Ala	Asn	Ser	Leu	Leu	Asn	Pro	Ile
	290					295					300				
Ile	Tyr	Thr	Leu	Thr	Asn	Arg	Asp	Leu	Arg	His	Ala	Leu	Leu	Arg	Leu
305					310					315					320
Val	Cys	Cys	Gly	Arg	His	Ser	Cys	Gly	Arg	Asp	Pro	Ser	Gly	Ser	Gln
				325					330					335	
Gln	Ser	Ala	Ser	Ala	Ala	Glu	Ala	Ser	Gly	Gly	Leu	Arg	Arg	Cys	Leu
		340						345					350		
Pro	Pro	Gly	Leu	Asp	Gly	Ser	Phe	Ser	Gly	Ser	Glu	Arg	Ser	Ser	Pro
		355					360					365			
Gln	Arg	Asp	Gly	Leu	Asp	Thr	Ser	Gly	Ser	Thr	Gly	Ser	Pro	Gly	Ala
	370					375					380				
Pro	Thr	Ala	Ala	Arg	Thr	Leu	Val	Ser	Glu	Pro	Ala	Ala	Asp		
385					390					395					

<210> SEQ ID NO 18
 <211> LENGTH: 254
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <223> OTHER INFORMATION: Predicted amino acid sequence of S1P5-alpha
 (Figure 12A)

<400> SEQUENCE: 18

Met	Glu	Ser	Gly	Leu	Leu	Arg	Pro	Ala	Pro	Val	Ser	Glu	Val	Ile	Val
1				5					10					15	
Leu	His	Tyr	Asn	Tyr	Thr	Gly	Lys	Leu	Arg	Gly	Ala	Arg	Tyr	Gln	Pro
		20					25						30		
Gly	Ala	Gly	Leu	Arg	Ala	Asp	Ala	Val	Val	Cys	Leu	Ala	Val	Cys	Ala
		35				40					45				
Phe	Ile	Val	Leu	Glu	Asn	Leu	Ala	Val	Leu	Leu	Val	Leu	Gly	Arg	His
	50				55						60				
Pro	Arg	Phe	His	Ala	Pro	Met	Phe	Leu	Leu	Leu	Gly	Ser	Leu	Thr	Leu
65				70						75				80	
Ser	Val	Pro	Ala	Arg	Pro	Gly	Thr	Ala	Gly	Thr	Thr	Ser	Thr	Arg	Ala
			85					90						95	
Arg	Arg	Lys	Pro	Arg	Ser	Leu	Ala	Leu	Leu	Arg	Thr	Leu	Ser	Val	Val
		100					105					110			
Leu	Leu	Ala	Phe	Val	Ala	Cys	Trp	Gly	Pro	Leu	Phe	Leu	Leu	Leu	Leu
	115						120					125			
Leu	Asp	Val	Ala	Cys	Pro	Ala	Arg	Thr	Cys	Pro	Val	Leu	Leu	Gln	Ala
	130					135						140			

-continued

```

Asp Pro Phe Leu Gly Leu Ala Met Ala Asn Ser Leu Leu Asn Pro Ile
145                150                155                160

Ile Tyr Thr Leu Thr Asn Arg Asp Leu Arg His Ala Leu Leu Arg Leu
                165                170                175

Val Cys Cys Gly Arg His Ser Cys Gly Arg Asp Pro Ser Gly Ser Gln
                180                185                190

Gln Ser Ala Ser Ala Ala Glu Ala Ser Gly Gly Leu Arg Arg Cys Leu
                195                200                205

Pro Pro Gly Leu Asp Gly Ser Phe Ser Gly Ser Glu Arg Ser Ser Pro
                210                215                220

Gln Arg Asp Gly Leu Asp Thr Ser Gly Ser Thr Gly Ser Pro Gly Ala
225                230                235                240

Pro Thr Ala Ala Arg Thr Leu Val Ser Glu Pro Ala Ala Asp
                245                250

<210> SEQ ID NO 19
<211> LENGTH: 103
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<223> OTHER INFORMATION: Predicted amino acid sequence of S1P5-beta
(Figure 12B)

<400> SEQUENCE: 19

Met Glu Ser Gly Leu Leu Arg Pro Ala Pro Val Ser Glu Val Ile Val
1          5          10          15

Leu His Tyr Asn Tyr Thr Gly Lys Leu Arg Gly Ala Arg Tyr Gln Pro
20         25         30

Gly Ala Gly Leu Arg Ala Asp Ala Val Val Cys Leu Ala Val Cys Ala
35         40         45

Phe Ile Val Leu Glu Asn Leu Ala Val Leu Leu Val Leu Gly Arg His
50         55         60

Pro Arg Phe His Ala Pro Met Phe Leu Leu Leu Gly Ser Leu Thr Leu
65         70         75         80

Ser Asp Leu Leu Ala Gly Ala Ala Tyr Ala Ala Ala Ala Arg Thr Leu
85         90         95

Val Ser Glu Pro Ala Ala Asp
100

```

We claim:

1. A method of screening for rheumatoid arthritis, comprising screening a sample from a patient for over-expression of a nucleic acid molecule encoding the sphingosine 1-phosphate receptor (SPPR) protein, wherein said protein comprises the amino acid sequence of SEQ ID NO:3, and wherein over-expression of said nucleic acid molecule is indicative of rheumatoid arthritis.
2. The method of claim 1, wherein said screening comprises measuring the amount of SPPR protein in the sample.
3. The method of claim 1, wherein said screening comprises measuring the amount of mRNA of SEQ ID NO:4 in the sample.
4. A method of screening for large granular lymphocyte (LGL) leukemia, comprising screening a sample from a patient for over-expression of a nucleic acid molecule encoding the sphingosine 1-phosphate receptor (SPPR) protein,

wherein said protein comprises the amino acid sequence of SEQ ID NO:3, and wherein over-expression of the nucleic acid molecule is indicative of LGL leukemia.

5. The method of claim 4, wherein said screening comprises measuring the amount of SPPR protein in the sample.

6. The method of claim 4, wherein said sample comprises blood.

7. A method of screening for large granular lymphocyte (LGL) leukemia in peripheral blood mononuclear cells (PBMC), comprising screening a sample of PBMC from a patient for over-expression of a nucleic acid molecule encoding the SPPR protein, wherein said protein comprises the amino acid sequence of SEQ ID NO:3, and wherein over-expression of the nucleic acid molecule is indicative of LGL leukemia.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,964,358 B2
APPLICATION NO. : 12/879918
DATED : June 21, 2011
INVENTOR(S) : Thomas P. Loughran and Ravi Kothapalli

Page 1 of 3

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 1.

Line 36, "not hilly" should read --not fully--.

Column 2.

Line 17, "when compare to" should read --when compared to--.

Line 47, "renamed S1P5.

DESCRIPTION OF THE FIGURES"

should read

--renamed S1P5.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is a forward primer used according to the subject invention.

SEQ ID NO:2 is a reverse primer used according to the subject invention.

SEQ ID NO:3 is the predicted amino acid sequence of the human sphingosine 1-Phosphate receptor (SPPR) cDNA of **SEQ ID NO:4**.

SEQ ID NO:4 is the complete nucleotide sequence of human sphingosine 1-Phosphate receptor (SPPR) cDNA.

SEQ ID NO:5 is the amino acid sequence of rat nrg-1.

SEQ ID NO:6 is the amino acid sequence of rat cdg-8.

SEQ ID NO:7 is the amino acid sequence of SPPR.


SEQ ID NO:8 is the deduced amino acid sequence of splice variant, sphingosine 1-phosphate receptor 1 of **SEQ ID NO:9**.

SEQ ID NO:9 is the nucleotide sequence of splice variant, sphingosine 1-phosphate receptor 1.

SEQ ID NO:10 is the amino acid sequence of the sphingosine 1-phosphate receptor (S1P).

SEQ ID NO:11 is the amino acid sequence of the sphingosine 1-phosphate receptor 1 (S1P₁).

Signed and Sealed this
Eleventh Day of October, 2011



David J. Kappos
Director of the United States Patent and Trademark Office

SEQ ID NO:12 is the deduced amino acid sequence of splice variant, sphingosine 1-phosphate receptor 2 of **SEQ ID NO:13**.

SEQ ID NO:13 is the nucleotide sequence of splice variant, sphingosine 1-phosphate receptor 2.

SEQ ID NO:14 is the amino acid sequence of sphingosine 1-phosphate receptor 2 (S1P₂).

SEQ ID NO:15 is the full-length (2.2 kb) nucleotide sequence of human S1P5 cDNA (Figure 8).

SEQ ID NO:16 is the deduced amino acid sequence of human S1P5 cDNA coding region (Figure 8).

SEQ ID NO:17 is the predicted amino acid sequence of S1P5 (Figures 12A and 12B).

SEQ ID NO:18 is the predicted amino acid sequence of S1P5-alpha (Figure 12A).

SEQ ID NO:19 is the predicted amino acid sequence of S1P5-beta (Figure 12B).

DESCRIPTION OF THE FIGURES--.

Column 3.

Line 14, "sppr is" should read --sppr (SEQ ID NO:7) is--.

Line 15, "rat edg-1 and nrg-1. There" should read

--rat edg-1 (SEQ ID NO:6) and nrg-1 (SEQ ID NO:5). There--.

Line 17, "sequence and" should read --sequence (SEQ ID NO:9) and--.

Line 18, "sequence of" should read --sequence (SEQ ID NO:8) of--.

Line 22, "sequence and" should read --sequence (SEQ ID NO:13) and--.

Line 23, "sequence of" should read --sequence (SEQ ID NO:12) of--.

Line 40, "S1P₅ is" should read --S1P₅ (SEQ ID NO:15) is--.

Line 44, "abbreviation. The" should read --abbreviation (SEQ ID NO:16). The--.

Line 50, "S1P₅ is" should read --S1P₅ (SEQ ID NO:7) is--.

Line 51, "rat edg-8 and nrg-1. There" should read

--rat edg-8 (SEQ ID NO:6) and nrg-1 (SEQ ID NO:5). There--.

Column 4.

Line 18, "shows comparison of" should read --shows comparisons of--.

Line 19, "S1P₅-β. The" should read

--S1P₅-β (SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, respectively). The--.

Line 22, "sequence. A" should read --sequence (SEQ ID NO:18). A--.

Line 27, "sequence. A" should read --sequence (SEQ ID NO:19). A--.

Line 33, "poly per" should read --poly A⁺ per--.

Line 40, "Northern blot.)

Column 4,

Line 40, “Northern blot.)

DETAILED DESCRIPTION OF THE INVENTION”

should read

--Northern blot.)

FIGURE 14 shows a comparison of the amino acid sequences of the sphingosine-1-phosphate receptor (S1P) (SEQ ID NO:10) and the sphingosine-1-phosphate 1 receptor (S1P1) (SEQ ID NO:11).

FIGURE 15 shows a comparison of the amino acid sequences of S1P (SEQ ID NO:10) and S1P2 (SEQ ID NO:14).

DETAILED DESCRIPTION OF THE INVENTION--.

Line 53, “is represented” should read --are represented--.

Column 10,

Line 38, “Phosphorimager” should read --PhosphorImager--.

Line 61, “sample of patents” should read --sample of patients--.

Column 13,

Line 25, “this acne is” should read --this gene is--.

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,964,358 B2
APPLICATION NO. : 12/879918
DATED : June 21, 2011
INVENTOR(S) : Thomas P. Loughran and Ravi Kothapalli

Page 1 of 3

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 1,

Line 36, "not hilly" should read --not fully--.

Column 2,

Line 17, "when compare to" should read --when compared to--.

Line 47, "renamed S1P5.

DESCRIPTION OF THE FIGURES"

should read

--renamed S1P5.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is a forward primer used according to the subject invention.

SEQ ID NO:2 is a reverse primer used according to the subject invention.

SEQ ID NO:3 is the predicted amino acid sequence of the human sphingosine 1-Phosphate receptor (SPPR) cDNA of SEQ ID NO:4.

SEQ ID NO:4 is the complete nucleotide sequence of human sphingosine 1-Phosphate receptor (SPPR) cDNA.

SEQ ID NO:5 is the amino acid sequence of rat nrg-1.

SEQ ID NO:6 is the amino acid sequence of rat edg-8.

SEQ ID NO:7 is the amino acid sequence of SPPR.

SEQ ID NO:8 is the deduced amino acid sequence of splice variant, sphingosine 1-phosphate receptor 1 of SEQ ID NO:9.

SEQ ID NO:9 is the nucleotide sequence of splice variant, sphingosine 1-phosphate receptor 1.

This certificate supersedes the Certificate of Correction issued October 11, 2011.

Signed and Sealed this
Twenty-seventh Day of March, 2012



David J. Kappos
Director of the United States Patent and Trademark Office

SEQ ID NO:10 is the amino acid sequence of the sphingosine 1-phosphate receptor (S1P).

SEQ ID NO:11 is the amino acid sequence of the sphingosine 1-phosphate receptor 1 (S1P₁).

SEQ ID NO:12 is the deduced amino acid sequence of splice variant, sphingosine 1-phosphate receptor 2 of SEQ ID NO:13.

SEQ ID NO:13 is the nucleotide sequence of splice variant, sphingosine 1-phosphate receptor 2.

SEQ ID NO:14 is the amino acid sequence of sphingosine 1-phosphate receptor 2 (S1P₂).

SEQ ID NO:15 is the full-length (2.2 kb) nucleotide sequence of human S1P5 cDNA (Figure 8).

SEQ ID NO:16 is the deduced amino acid sequence of human S1P5 cDNA coding region (Figure 8).

SEQ ID NO:17 is the predicted amino acid sequence of S1P5 (Figures 12A and 12B).

SEQ ID NO:18 is the predicted amino acid sequence of S1P5-alpha (Figure 12A).

SEQ ID NO:19 is the predicted amino acid sequence of S1P5-beta (Figure 12B).

DESCRIPTION OF THE FIGURES--.

Column 3,

Line 14, "sppr is" should read --sppr (SEQ ID NO:7) is--.

Line 15, "rat edg-1 and nrg-1. There" should read

--rat edg-1 (SEQ ID NO:6) and nrg-1 (SEQ ID NO:5). There--.

Line 17, "sequence and" should read --sequence (SEQ ID NO:9) and--.

Line 18, "sequence of" should read --sequence (SEQ ID NO:8) of--.

Line 22, "sequence and" should read --sequence (SEQ ID NO:13) and--.

Line 23, "sequence of" should read --sequence (SEQ ID NO:12) of--.

Line 40, "S1P₅ is" should read --S1P₅ (SEQ ID NO:15) is--.

Line 44, "abbreviation. The" should read --abbreviation (SEQ ID NO:16). The--.

Line 50, "S1P₅ is" should read --S1P₅ (SEQ ID NO:7) is--.

Line 51, "rat edg-8 and nrg-1. There" should read

--rat edg-8 (SEQ ID NO:6) and nrg-1 (SEQ ID NO:5). There--.

Column 4,

Line 18, "shows comparison of" should read --shows comparisons of--.

Line 19, "S1P₅-β. The" should read

--S1P₅-β (SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, respectively). The--.

Line 22, "sequence. A" should read --sequence (SEQ ID NO:18). A--.

Line 27, "sequence. A" should read --sequence (SEQ ID NO:19). A--.

Line 33, "poly per" should read --poly A+ per--.

Column 4,

Line 40, “Northern blot.)

DETAILED DESCRIPTION OF THE INVENTION”

should read

--Northern blot.)

FIGURE 14 shows a comparison of the amino acid sequences of the sphingosine-1-phosphate receptor (S1P) (SEQ ID NO:10) and the sphingosine-1-phosphate 1 receptor (S1P1) (SEQ ID NO:11).

FIGURE 15 shows a comparison of the amino acid sequences of S1P (SEQ ID NO:10) and S1P2 (SEQ ID NO:14).

DETAILED DESCRIPTION OF THE INVENTION--.

Line 53, “is represented” should read --are represented--.

Column 10,

Line 38, “Phosphorimager” should read --PhosphorImager--.

Line 61, “sample of patents” should read --sample of patients--.

Column 13,

Line 25, “this acne is” should read --this gene is--.