

---

USF Patents

---

August 2011

## SHIP-deficiency to increase megakaryocyte progenitor production

William G. Kerr

Caroline Despons

Lia Elena Perez

Follow this and additional works at: [https://digitalcommons.usf.edu/usf\\_patents](https://digitalcommons.usf.edu/usf_patents)

---

### Recommended Citation

Kerr, William G.; Despons, Caroline; and Perez, Lia Elena, "SHIP-deficiency to increase megakaryocyte progenitor production" (2011). *USF Patents*. 426.  
[https://digitalcommons.usf.edu/usf\\_patents/426](https://digitalcommons.usf.edu/usf_patents/426)

This Patent is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Patents by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact [digitalcommons@usf.edu](mailto:digitalcommons@usf.edu).



US008008273B2

(12) **United States Patent**  
**Kerr et al.**

(10) **Patent No.:** **US 8,008,273 B2**  
(45) **Date of Patent:** **\*Aug. 30, 2011**

(54) **SHIP-DEFICIENCY TO INCREASE  
MEGAKARYOCYTE PROGENITOR  
PRODUCTION**

(75) Inventors: **William G. Kerr**, Syracuse, NY (US);  
**Caroline Desponte**, La Jolla, CA (US);  
**Lia Elena Perez**, Tampa, FL (US)

(73) Assignee: **University of South Florida**, Tampa, FL  
(US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.  
  
This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **12/819,711**

(22) Filed: **Jun. 21, 2010**

(65) **Prior Publication Data**

US 2010/0260730 A1 Oct. 14, 2010

#### **Related U.S. Application Data**

(60) Division of application No. 11/451,004, filed on Jun.  
12, 2006, now Pat. No. 7,763,592, which is a  
continuation-in-part of application No. 10/904,667,  
filed on Nov. 22, 2004, now Pat. No. 7,807,646.

(60) Provisional application No. 60/481,677, filed on Nov.  
20, 2003.

(51) **Int. Cl.**  
**A61K 48/00** (2006.01)

(52) **U.S. Cl.** ..... **514/44**; 536/24.5; 536/24.1

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

#### **U.S. PATENT DOCUMENTS**

4,603,112 A 7/1986 Paoletti et al.  
4,769,330 A 9/1988 Paoletti et al.  
4,777,127 A 10/1988 Suni et al.  
5,017,487 A 5/1991 Stunnenberg et al.  
5,166,057 A 11/1992 Palese et al.  
5,804,412 A 9/1998 Gill et al.  
6,025,198 A 2/2000 Bennett et al.  
6,090,621 A 7/2000 Kavanaugh et al.  
6,117,850 A 9/2000 Patchen et al.  
6,200,807 B1 3/2001 Bennett et al.  
6,506,559 B1 1/2003 Fire et al.  
6,703,215 B2 3/2004 Erneux et al.  
7,691,821 B2 4/2010 Desponte et al.  
7,713,945 B2 5/2010 Kerr  
7,763,592 B1 7/2010 Kerr et al.  
7,807,646 B1 10/2010 Kerr et al.  
2002/0137711 A1 9/2002 Kerr  
2002/0165192 A1 11/2002 Kerr et al.  
2003/0114401 A1 6/2003 Bennett et al.  
2003/0143732 A1 7/2003 Fosnaugh et al.  
2003/0166282 A1 9/2003 Brown et al.  
2004/0072298 A1 4/2004 Sauvageau et al.  
2004/0235765 A1 11/2004 Kerr et al.  
2004/0259247 A1 12/2004 Tuschl et al.  
2005/0054103 A1 3/2005 Peled et al.

2006/0223749 A1 10/2006 Desponte et al.  
2007/0224124 A1 9/2007 Kerr et al.  
2008/0076731 A1 3/2008 Kerr  
2010/0136026 A1 6/2010 Kerr et al.  
2010/0144841 A1 6/2010 Kerr

#### **FOREIGN PATENT DOCUMENTS**

EP 0 345 242 A2 12/1989  
EP 0 440 219 A1 8/1991  
GB 2 200 651 A 8/1988  
WO WO 89/01973 A2 3/1989  
WO WO 91/02805 A2 3/1991  
WO WO 92/06693 A1 4/1992  
WO WO 97/10252 A1 3/1997  
WO WO 97/12039 A2 4/1997  
WO WO 99/32619 A1 7/1999  
WO WO 02/24233 A2 3/2002  
WO WO 02/44321 A2 6/2002  
WO WO 02/078614 A3 10/2002  
WO WO 2008/022468 A1 2/2008  
WO WO 2009/042910 A2 4/2009

#### **OTHER PUBLICATIONS**

Crooke, S.T. "Progress in Antisense Technology" *Ann Rev Medicine*,  
2004, 55:61-95.  
Jang, J.H. et al. "Gene delivery from polymer scaffolds for tissue  
engineering" *Expert Rev. Medical Devices*, 2004, 1(1):127-138.  
Office Action mailed Dec. 29, 2005 in U.S. Appl. No. 09/955,174,  
filed Sep. 19, 2001.  
Office Action mailed Aug. 9, 2005 in U.S. Appl. No. 10/605,452, filed  
Sep. 30, 2003.  
Office Action mailed Apr. 12, 2006 in U.S. Appl. No. 10/605,542,  
filed Sep. 30, 2003.  
Office Action mailed Sep. 28, 2009 in U.S. Appl. No. 11/451,004,  
filed Jun. 12, 2006.  
Office Action mailed Apr. 7, 2009 in U.S. Appl. No. 11/451,004, filed  
Jun. 12, 2006.  
Office Action mailed Aug. 5, 2008 in U.S. Appl. No. 11/451,004, filed  
Jun. 12, 2006.  
Notice of Allowance mailed Oct. 7, 2009 in U.S. Appl. No.  
11/787,064, filed Apr. 13, 2007.  
U.S. Appl. No. 10/605,452, Kerr et al.  
U.S. Appl. No. 10/709,801, Desponte et al.  
U.S. Appl. No. 10/904,667, Kerr et al. U.S. Appl. No. 12/670,360,  
Kerr et al.  
U.S. Appl. No. 12/689,167, Desponte et al.  
U.S. Appl. No. 12/651,809, Kerr.  
Adams, A. "RNA Therapeutics Enter Clinical Trials" *The Scientist*,  
Jan. 17, 2005, 19(1):1-4.  
Agrawal, S. "Antisense oligonucleotides: towards clinical trials"  
*Tibtech*, Oct. 1996, 14:376-387.  
Agrawal, S. et al. "Antisense therapeutics: is it as simple as comple-  
mentary base recognition" *Molecular Medicine Today*, Feb. 2000,  
6:72-81.

(Continued)

*Primary Examiner* — Kimberly Chong

(74) *Attorney, Agent, or Firm* — Saliwanchik, Lloyd &  
Eisenschenk

(57) **ABSTRACT**

The invention concerns a method for increasing megakaryo-  
cyte and megakaryocyte progenitor numbers in vitro or in  
vivo by suppressing SH2-containing inositol-5-phosphatase  
(SHIP) function in megakaryocytes or megakaryocyte pro-  
genitors expressing the SHIP gene. SHIP function can be  
suppressed by administering an interfering RNA, or other  
SHIP inhibitor, to the megakaryocytes or megakaryocyte pro-  
genitors in vitro or in vivo.

**13 Claims, 13 Drawing Sheets**

## OTHER PUBLICATIONS

- Agrawal, N. et al. "RNA Interference: Biology, Mechanism, and Applications" *Microbiology and Molecular Biology*, Dec. 2003, 67(4):657-685.
- Ahmed, N. et al. "Cytokine-Induced Expansion of Human CD34<sup>+</sup> Stem/Progenitor and CD34<sup>+</sup>CD41<sup>+</sup> Early Megakaryocytic Marrow Cells Cultured on Normal Osteoblasts" *Stem Cells*, 1999, 17:92-99.
- Akagi, K. et al. "Cre-mediated somatic site-specific recombination in mice" *Nucleic Acids Research*, 1997, 25(9):1766-1773.
- Bender, M.A. et al. "Description and Targeted Deletion of 5' Hypersensitive Site 5 and 6 of the Mouse  $\beta$ -Globin Locus Control Region" *Blood*, Dec. 1998, 92(11):4394-4403.
- Bjorklund, L.M. et al. "Embryonic stem cells develop into functional dopaminergic neurons after transplantation in a Parkinson rat model" *PNAS*, Feb. 19, 2003, 99(4):2344-2349.
- Bolland, S. et al. "SHIP Modulates Immune Receptor Responses by Regulating Membrane Association of Btk" *Immunity*, Apr. 1998, 8:509-516.
- Bonetta, L. "RNAi: Silencing never sounded better" *Nature Methods*, Oct. 2004, 1(1):79-86.
- Braasch, D.A. et al. "Novel Antisense and Peptide Nucleic Acid Strategies for Controlling Gene Expression" *Biochemistry*, Apr. 9, 2002 41(14):4503-4510.
- Branch, A.D. "A good antisense molecule is hard to find" *TIBS*, Feb. 1998, 23:45-50.
- Cantley, L.C. et al. "Oncogenes and Signal Transduction" *Cell*, Jan. 25, 1991, 64:281-302.
- Caplen, N.J. et al. "Specific inhibition of gene expression by small double-stranded RNAs in invertebrate and vertebrate systems" *PNAS*, Aug. 14, 2001, 98(17):9742-9747.
- Caplen, N.J. "RNAi as a gene therapy approach" *Expert Opinion Biol Ther.*, 2003, 3(4):575-586.
- Chirila, T.V. et al. "The use of synthetic polymers for delivery of therapeutic antisense oligodeoxynucleotides" *Biomaterials*, 2002, 23:321-342.
- Crooke, S.T. "Basic principles of antisense therapeutics" in *Antisense Research and Application*, Chapter 1, pp. 1-50, S. Crooke, Editor, Springer-Verlag, 1999.
- Damen, J.E. et al. "The 145-kDa protein induced to associate with Shc by multiple cytokines is an inositol tetrakisphosphate and phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase" *Proc Natl Acad Sci USA*, Feb. 1996, 93:1689-1693.
- Despons, C. et al. "MHC Class I inhibitory receptors on natural killer cells recruit SHIP in an attempt to control cell survival" *FASEB Journal*, Mar. 20, 2002, 16(4):A707, abstract.
- De Souza, A.T. et al. "Transcriptional and phenotypic comparisons of *Ppara* knockout and siRNA knockdown mice" *Nucleic Acids Research*, 2006, 34(16):4486-4494.
- Elbashir, S.M. et al. "RNA interference is mediated by 21- and 22-nucleotide RNAs" *Genes & Development*, 2001, 15:188-200.
- Elbashir, S.M. et al. "Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells" *Nature*, May 2001, 411:494-498.
- Evans, D.J. et al. "An engineered poliovirus chimera elicits broadly reactive HIV-1 neutralizing antibodies" *Nature*, Jun. 1989, 339:385-388.
- Examination Report in European Application No. 01973144.7, dated Nov. 11, 2006.
- Fire, A. et al. "Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*" *Nature*, Feb. 1998, 391:806-811.
- Fisher-Hoch, S.P. et al. "Protection of rhesus monkeys from fatal Lassa fever by vaccination with a recombinant vaccinia virus containing the Lassa virus glycoprotein gene" *Proc Natl Acad Sci USA*, Jan. 1989, 86:317-321.
- Geier, S.J. et al. "The Human SHIP Gene Is Differentially Expressed in Cell Lineages of the Bone Marrow and Blood" *Blood*, 1997, 89:1876-1885.
- Gewirtz, A.M. et al. "Facilitating oligonucleotide delivery: Helping antisense deliver on its promise" *Proc Natl Acad Sci USA*, Apr. 1996, 93:3161-3163.
- Ghansah, T. et al. "A role for the SH2-containing Inositol phosphatase in the biology of natural killer cells and stem cells" in *Activating and inhibitory immunoglobulin-like Receptors*, 2001, pp. 129-140.
- Ghansah, T. et al. "Target disruption of Src homology 2-containing 5' inositol phosphatase murine natural killer cells" *FASEB Journal*, Mar. 20, 2002, 16(4):A706, abstract.
- Ghansah, T. et al. "The Src homology 2 containing inositol phosphatase is vital for the function and homeostasis of Natural Killer cells" *FASEB Journal*, Mar. 7, 2001, 15(4):A655, abstract.
- Ghansah, T. et al. "Expansion of Myeloid Suppressor Cells in SHIP-Deficient Mice Represses Allogeneic T Cell Responses" *The Journal of Immunology*, 2004, 173:7324-7330.
- Guzman, R.J. et al. "Efficient and Selective Adenovirus-Mediated Gene Transfer Into Vascular Neointima" *Circulation*, Dec. 1993, 88(6):2838-2848.
- Hannon, G.J. et al. "Unlocking the potential of the human genome with RNA interference" *Nature*, Sep. 16, 2004, 431:371-378.
- Harborth, J. et al. "Identification of essential genes in cultured mammalian cells using small interfering RNAs" *Journal of Cell Science*, 114(24):4557-4565, 2001.
- Hawkins, P.T. et al. "Platelet-derived growth factor stimulates synthesis of PtdIns(3,4,5)P<sub>3</sub> by activating a PtdIns(4,5)P<sub>2</sub> 3-OH kinase" *Nature*, Jul. 1992, 358(6382):157-159.
- Held, W. et al. "Transgenic Expression of the Ly49A Natural Killer Cell Receptor Confers Class I Major Histocompatibility Complex (MHC)-specific Inhibition and Prevents Bone Marrow Allograft Rejection" *J Exp Med*, Nov. 1996, 184:2037-2041.
- Helgason, C.D. et al. "Targeted disruption of SHIP leads to homopoietic perturbations, lung pathology, and a shortened life span" *Genes & Development*, 1998, 12:1610-1620.
- Helgason, C.D. et al. "Homeostasis and regeneration of the hematopoietic stem cell pool are altered in SHIP-deficient mice" *Blood*, 2003, 102(10):3541-3547.
- Hemann, M.T. et al. "An epi-allelic series of p53 hypomorphs created by stable RNAi produces distinct tumor phenotypes in vivo" *Nature Genetics*, Mar. 2003, 33:396-400.
- Hemmati-Brivanlou, A. et al. "Vertebrate Embryonic Cells Will Become Nerve Cells Unless Told Otherwise" *Cell*, Jan. 10, 1997, 88:13-17.
- Huber, M. et al. "The src homology 2-containing inositol phosphatase (SHIP0) is the gatekeeper of mast cell degranulation" *Proc Natl Acad Sci USA*, Sep. 1998, 95:11330-11335.
- Huber, M. et al. "The role of SHIP in growth factor induced signaling" *Progress in Biophysics & Molecular Biology*, 1999, 71:423-434.
- Jefferson, A.B. et al. "Properties of Type II Inositol Polyphosphate 5-Phosphatase" *The Journal of Biological Chemistry*, Apr. 1995, 270(16):9370-9377.
- Jen, K.Y. et al. "Suppression of Gene Expression by Targeted Disruption of Messenger RNA: Available Options and Current Strategies" *Stem Cells*, 2000, 18:307-319.
- Jolly, D. "Viral vector systems for gene therapy" *Cancer Gene Therapy*, 1994, 1(1):51-64.
- Kass-Eisler, A. et al. "Quantitative determination of adenovirus-mediated gene delivery to rat cardiac myocytes in vitro and in vivo" *Proc Natl Acad Sci USA*, Dec. 1993, 90:11498-11502.
- Kawasaki, H. et al. "Induction of Midbrain Dopaminergic Neurotechnique Neurons from ES Cells by Stromal Cell-Derived Inducing Activity" *Neuron*, Oct. 2000, 28:31-40.
- Kerr, W.G., "The SH2 Containing Inositol Phosphatase (SHIP) is a Crucial Regulator of NK Cell Repertoire and Function" Abstract #34, presented at Core Research for Evolutional Science and Technology (CREST) International Symposium on Immunoglobulin-like Receptors, held Sep. 19-20, 2000, at the Sendai International Center, Sendai City, Japan.
- Kerr, W.G. et al. "Critical role for SHIP in engraftment of histoincompatible stem cells" *Oncology Research*, 2001, 12:285.
- Kim, J.H. et al. "Dopamine neurons derived from embryonic stem cells function in an animal model of Parkinson's disease" *Nature*, Jun. 20, 2002, pp. 1-7, doi.1038/nature00900, advance online publication.

- Klippel, A. et al. "Membrane Localization of Phosphatidylinositol 3-Kinase Is Sufficient to Activate Multiple Signal-Transducing Kinase Pathways" *Molecular and Cellular Biology*, Aug. 1996, 16(8):4117-4127.
- Koh, C.Y. et al. "Augmentation of antitumor effects by NK cell inhibitory receptor blockade in vitro and in vivo" *Blood*, 2001, 97(10):3132-3137.
- Kolls, J. et al. "Prolonged and effective blockade of tumor necrosis factor activity through adenovirus-mediated gene transfer" *Proc Natl Acad Sci USA*, Jan. 1994, 91:215-219.
- Krystal, G. et al. "Molecules in focus: SHIPs ahoy" *The International Journal of Biochemistry & Cell Biology*, 1999, 31:1007-1010.
- Lanier, L.L. "NK Cell Receptors" *Annu Rev Immunol*, 1998, 16:359-393.
- Lewis, D.L. et al. "Efficient delivery of siRNA for inhibition of gene expression in postnatal mice" *Nature Genetics*, Sep. 2002, 32:107-108.
- Liu, L. et al. "The Src Homology 2 (SH2) Domain of SH2-containing Inositol Phosphatase (SHIP) Is Essential for Tyrosine Phosphorylation of SHIP, Its Association with Shc, and Its Induction of Apoptosis" *The Journal of Biological Chemistry*, Apr. 4, 1997, 272(14):8983-8988.
- Liu, Q. et al. "Molecular Cloning and Chromosomal Localization in Human and Mouse of the SH2-Containing Inositol Phosphatase, INPP5D (SHIP)" *Genomics*, 1997, 39:109-112.
- Liu, Q. et al. "The Inositol Polyphosphate 5-Phosphatase Ship Is a Crucial Negative Regulator of B Cell Antigen Receptor Signaling" *J Exp Med*, Oct. 5, 1998, 188(7):1333-1342.
- Liu, Q. et al. "SHIP is a negative regulator of growth factor receptor-mediated PKB/Akt activation and myeloid cell survival" *Genes & Development*, 1999, 13(7):786-791.
- Lotzova, E. et al. "Prevention of Rejection of Allogeneic Bone Marrow Transplants by NK-1.1 Anti Serum" *Transplantation*, 1983, 35(5):490-494, abstract.
- Lucas, D.M. et al. "A Novel Spliced Form of SH2-Containing Inositol Phosphatase is Expressed During Myeloid Development" *Blood*, Mar. 15, 1999, 93(6):1922-1933.
- Muraille, E. et al. "Distribution of the Src-homology-2-domain-containing inositol 5-phosphatase SHIP-2 in both non-haemopoietic and haemopoietic cells and possible involvement of SHIP-2 in negative signaling of B-cells" *Biochem J*, 1999, 342:697-705.
- Montgomery, M.K. et al. "RNA as a target of double-stranded RNA-mediated genetic interference in *Caenorhabditis elegans*" *Proc Natl Acad Sci USA*, Dec. 1998, 95:15502-15507.
- Moody, J.L. et al. "Anemia, thrombocytopenia, leukocytosis, extramedullary hematopoiesis, and impaired progenitor function in *Pten<sup>+/+</sup>SHIP<sup>-/-</sup>* mice: a novel model of myelodysplasia" *Blood*, 2004, 103:4503-4510.
- Novina, C.D. et al. "The RNAi revolution" *Nature*, Jul. 8, 2004, 430:161-164.
- Okada, H. et al. "Cutting Edge: Role of the Inositol Phosphatase SHIP in B Cell Receptor-Induced Ca<sup>2+</sup> Oscillatory Response" *The Journal of Immunology*, 1998, 161:5129-5132.
- Opalinska, J.B. et al. "Nucleic-Acid Therapeutics: Basic Principles and Recent Applications" *Nature Reviews: Drug Discovery*, Jul. 2002, 1:503-514.
- Odorico, J.S. et al. "Multilineage Differentiation from Human Embryonic Stem Cell Lines" *Stem Cells*, 2001, 19:193-204.
- Overbaugh, J. et al. "Molecular Cloning of a Feline Leukemia Virus That Induces Fatal Immunodeficiency Disease in Cats" *Science*, 1988, 239:906-910.
- Palu, G. et al. "In pursuit of new developments for gene therapy of human diseases" *Journal of Biotechnology*, 1999, 68:1-13.
- Paroo, Z. et al. "Challenges for RNAi in vivo" *TRENDS in Biotechnology*, Aug. 2004, 22(8):390-394.
- Pasquet, J.M. et al. "Phosphatidylinositol 3,4,5-trisphosphate regulates Ca<sup>2+</sup> entry via Btk in platelets and megakaryocytes without increasing phospholipase C activity" *EMBO Journal*, Jun. 15, 2000, 19(12):2793-2802.
- Pera, M.F. et al. "Commentary: Human embryonic stem cells" *Journal of Cell Science*, 2000, 113:5-10.
- Peracchi, A. "Prospects for antiviral ribozymes and deoxyribozymes" *Reviews in Medical Virology*, 2004, 14:47-64.
- Perez, L.E. et al. "SH<sub>2</sub>-Inositol Phosphatase 1 Negatively Influences Early Megakaryocyte Progenitors" *PLOS One*, Oct. 2008, 3(10):e3565, pp. 1-8.
- Pesesse, X. et al. "The SH2 domain containing inositol 5-phosphatase SHIP2 displays phosphatidylinositol 3,4,5-trisphosphate and inositol 1,3,4,5-tetrakisphosphate 5-phosphatase activity" *FEBS Letters*, 1998, 437:301-303.
- Pihl-Carey, K. "Disease Drug fails in Phase III" *BioWorld Today*, Dec. 16, 1999, 10:1-2.
- Poznansky, M. et al. "Gene Transfer into Human Lymphocytes by a Defective Human Immunodeficiency Virus Type 1 Vector" *Journal of Virology*, Jan. 1991, 65(1):532-536.
- Program and Abstracts for CREST International Symposium on Immunoglobulin-like Receptors, held Sep. 19-20, 2000.
- Puente, X.S. et al. "Human and Mouse Proteases: A Comparative Genomic Approach" *Nature Reviews: Genetics*, Jul. 2003, 4:544-558.
- Rauh, M.J. et al. "The role of SHIP1 in macrophage programming and activation" *Biochemical Society Transactions*, 2004, 32(5):785-788.
- Rehli, M. et al. "The Membrane-Bound Ecto-peptidase CPM as a Marker of Macrophage Maturation In Vitro and In Vivo" *Cellular Peptidases in Immune Functions and Diseases 2*, 2000, pp. 205-216, eds. Lagner and Ansorge, Kluwer Academic/Plenum Publishers.
- Rohrschneider, L.R. et al. "Structure, function, and biology of SHIP proteins" *Genes & Development*, 2000, 14:505-520.
- Ruggeri, L. et al. "Role of Natural Killer Cell Alloreactivity in HLA-Mismatched Hematopoietic Stem Cell Transplantation" *Blood*, Jul. 1, 1999, 94(1):333-339.
- Sabin, A.B. et al. "History of Sabin attenuated poliovirus oral live vaccine strains" *Journal of Biological Standardization*, 1973, 1:115-118.
- Samulski, R.J. et al. "Helper-Free Stocks of Recombinant Adeno-Associated Viruses: Normal Integration Does Not Require Viral Gene Expression" *Journal of Virology*, Sep. 1989, 63(9):3822-3828.
- Statement of Dr. Toshiyuki Takai dated Apr. 10, 2007, an organizer of the CREST International Symposium on Immunoglobulin-like Receptors, held Sep. 19-20, 2000, 1 page.
- Sauer, B. "Inducible Gene Targeting in Mice Using the Cre/lox System" *Methods: A Companion to Methods in Enzymology*, 1998, 14:381-392.
- Sawyers, C.L. "Chronic Myeloid Leukemia" *The New England Journal of Medicine*, Apr. 29, 1999, 340(17):1330-1340.
- Sly, L.M. et al. "SHIP, SHIP2, and PTEN activities are regulated in vivo by modulation of their protein levels: SHIP is up-regulated in macrophages and mast cells by lipopolysaccharide" *Experimental Hematology*, 2003, 31:1170-1181.
- Stephens, L.R. et al. "Agonist-stimulated synthesis of phosphatidylinositol(3,4,5)-trisphosphate: a new intracellular signaling system?" *Biochimica et Biophysica Acta*, 1993, 1179:27-75.
- Svoboda, P. et al. "Selective reduction of dormant maternal mRNAs in mouse oocytes by RNA interference" *Development*, 2000, 127:4147-4156.
- Tamm, I. et al. "Antisense therapy in oncology: new hope for an old idea?" *Lancet*, 2001, 358:489-497.
- Tu, Z. et al. "Embryonic and hematopoietic stem cells express a novel SH2-containing inositol 5'-phosphatase isoform that partners with the Grb2 adapter protein" *Blood*, 2001, 98:2028-2038.
- Tuschl, T. et al. "Targeted mRNA degradation by double-stranded RNA in vitro" *Genes & Development*, 13:3191-3197, 1999.
- Tuschl, T. "RNA Interference and Small Interfering RNAs" *ChemBioChem*, 2001, 2:239-245.
- Verfaillie, C.M. et al. "Hematopoietic stem cells for transplantation" *Nature Immunology*, 2002, 3(4):314-317.
- Wang, C.Y. et al. "pH-sensitive immunoliposomes mediate target-cell-specific delivery and controlled expression of a foreign gene in mouse" *Proc Natl Acad Sci USA*, Nov. 1987, 84:7851-7855.
- Wang, J.W. et al. "Influence of SHIP on the NK Repertoire and Allogeneic Bone Marrow Transplantation" *Science*, Mar. 15, 2002, 295:2094-2097.

- Wang, J.W. et al. Identification of a Novel Lipopolysaccharide-Inducible Gene with Key Features of Both a Kinase Anchor Proteins and chsl/beige Proteins, *The Journal of Immunology*, 2001, 166:4586-4595.
- Ware, M.D. et al. "Cloning and Characterization of Human SHIP, the 145-kD Inositol 5-Phosphatase That Associates With SHC After Cytokine Stimulation" *Blood*, Oct. 15, 1996, 88(8):2833-2840.
- Wisniewski, D. et al. "Neoplasia: A Novel SH2-Containing Phosphatidylinositol 3,4,5-Trisphosphate 5-Phosphatase (SHIP2) Is Constitutively Tyrosine Phosphorylated and Associated With src Homologous and Collagen Gene (SHC) in Chronic Myelogenous Leukemia Progenitor Cells" *Blood*, Apr. 1999, 93(8):2707-2720.
- Wolf, I. et al. "Cloning of the Genomic Locus of Mouse SH2 Containing Inositol 5-Phosphatase (SHIP) and a Novel 110-kDa Splice Isoform, SHIP8" *Genomics*, 2000, 69:104-112.
- Yokoyama, W.M. et al. "Natural killer cell receptors" *Current Opinion in Immunology*, 1998, 10:298-305.
- Zamore, P.D. et al. "RNAi: Double-Stranded RNA Directs the ATP-Dependent Cleavage of mRNA at 21 to 23 Nucleotide Intervals" *Cell*, Mar. 31, 2000, 101:25-33.
- Zandstra, P.W. et al. "Leukemia Inhibitory Factor (LIF) Concentration Modulates Embryonic Stem Cell Self-Renewal and Differentiation Independently of Proliferation" *Biotechnol Bioeng*, 2000, 69:607-617.
- Zwaka, T.P. et al. "Homologous recombination in human embryonic stem cells" *Nature Biotechnology*, Feb. 10, 2003, pp. 1-3, doi: 10.1038/nbt788, advance online publication.
- Office Action mailed Jan. 14, 2009 in U.S. Appl. No. 09/955,174, filed Sep. 19, 2001.
- Office Action mailed Feb. 10, 2006 in U.S. Appl. No. 10/097,101, filed Feb. 14, 2002.
- Office Action mailed Oct. 17, 2006 in U.S. Appl. No. 10/605,452, filed Sep. 30, 2003.
- Office Action mailed Apr. 28, 2009 in U.S. Appl. No. 10/709,801, filed May 28, 2004.
- Office Action mailed Sep. 22, 2008 in U.S. Appl. No. 10/709,801, filed May 28, 2004.
- Office Action mailed Jan. 7, 2009 in U.S. Appl. No. 10/904,667, filed Nov. 22, 2004.
- Office Action mailed Jul. 22, 2009 in U.S. Appl. No. 10/904,667, filed Nov. 22, 2004.
- Office Action mailed Jun. 5, 2009 in U.S. Appl. No. 11/787,064, filed Apr. 13, 2007.
- Office Action mailed Dec. 16, 2008 in U.S. Appl. No. 11/787,064, filed Apr. 13, 2007.




FIG. 1A

FIG. 1B-1




FIG. 1B-2




FIG. 1B-3




FIG. 1C-1




FIG. 1C-2





FIG. 1C-3








FIG. 1D




FIG. 1E

FIG. 2

1 aaacaggaag tcagtcagtt aagctggtgg cagcagccga ggccaccaag aggcaacggg  
61 cggcaggttg cagtggagg gctcgcgtc ccctcgggtg tgtgtgggtc ctgggggtgc  
121 ctgcccggccc ggccgaggag gcccacgccc accatggtcc cctgctggaa ccatggcaac  
181 atcacccgct ccaaggcggg ggagctgctt tccaggacag gcaaggacgg gagcttcctc  
241 gtgcgtgcc aagagtcctat ctcccgggca tacgcgtctt gcgtgctgta tcggaattgc  
301 gtttacactt acagaattct gcccaatgaa gatgataaat tcactgttca ggcatccgaa  
361 ggcgtctcca tgaggttctt caccaagctg gaccagctca tcgagtttta caagaaggaa  
421 aacatggggc tggtgaccca tctgcaatac cctgtgccgc tggaggaa gaacacaggc  
481 gacgaacctg aggaggacac agaaagtgtc gtgtctccac ccgagctgcc cccaagaaac  
541 atcccgcgtg ctgccagctc ctgtgaggcc aaggaggttc ctttttcaaa cgagaatccc  
601 cgagcgaccg agaccagccg gccgagcctc tccgagacat tgttccagcg actgcaaagc  
661 atggacacca gtgggcttcc agaagagcat cttaggcca tocaagatta ttttaagcact  
721 cagctcgccc aggactctga atttgtgaag acagggtcca gcagtottcc tcacctgaag  
781 aaactgacca cactgctctg caaggagctc tatggagaag tcatccggac cctccatcc  
841 ctggagtctc tgagagggtt atttgaccag cagctctccc cgggcctccg tccacgtcct  
901 caggttctct gtgaggccaa tcccatcaac atggtgtcca agctcagcca actgacaagc  
961 ctggtgtcat ccattgaaga caaggccaag gccttgctgc acgagggtcc tagtctccg  
1021 caccggccct cccttatccc tccagtcacc tttgagggtg aggcagagtc tctggggatt  
1081 cctcagaaaa tgcagctcaa agtcgacgtt gagtctggga aactgatcat taagaagtc  
1141 aaggatggtt ctgaggacaa gttctacagc cacaagaaaa tctgcagct cattaagtca  
1201 cagaaatttc tgaataagtt ggtgatcttg gtgaaacag agaaggagaa gatcctgcgg  
1261 aaggaatatg tttttgctga ctccaaaaag agagaaggct tctgccagct cctgcagcag  
1321 atgaagaaca agcactcaga gcagccggag cccgacatga tcacctctt catcggcacc  
1381 tggaacatgg gtaacgcccc ccctcccaag aagatcacgt cctggtttct ctccaagggg  
1441 cagggaagaa cgcgggacga ctctgcccag tacatccccc atgacattta cgtgatccgc  
1501 acccaagagg acccctgag tgagaaggag tggctggaga tccctcaaca ctccctgcaa  
1561 gaaatcacca gtgtgacttt taaaacagtc gccatccaca cgtctggaa catccgcac  
1621 gtggtgctgg ccaagcctga gcacgagaac cggatcagcc acatctgtac tgacaacgtg  
1681 aagacaggca ttgcaaacac actggggaac aaggagccg tgggggtgtc gttcatgttc  
1741 aatggaacct ccttagggtt cgtcaacagc cacttgactt caggaagtga aaagaaactc  
1801 aggcgaaacc aaaactatat caacattctc cgttctctgg ccctgggcga caagaagctg  
1861 agtcccttta acatcactca ccgcttcacg cacctcttct ggtttgggga tcttaactac  
1921 cgtgtggatc tgccctacct ggaggcagaa accatcatcc agaaaatcaa gcagcagcag  
1981 tacgcagacc tctgtccca cgaccagctg ctcacagaga ggaggagca gaaggtcttc  
2041 ctacacttcg aggaggaaga aatcacgttt gcccaacct accgttttga gagactgact  
2101 cgggacaaat acgcctacaa caagcagaaa cgcacagggg tgaaagtaca cttgccttcc  
2161 tgggtgtgacc gagtccctctg gaagtcttat cccctggtgc acgtggtgtg tcagtcttat  
2221 ggcagtacca gcgacatcat gacgagtgtc cacagccctg tctttgcac atttgaggca  
2281 ggagtcaact ccagtttgt ctccaagaac ggtcccggga ctgttgacag ccaaggacag  
2341 attgagtttc tcaggtgcta tgccacattg aagaccaagt cccagaccaa attctacctg  
2401 gagttccact cgagctgctt ggagagtttt gtcaagagtc aggaaggaga aaatgaagaa  
2461 ggaagtgagg gggagctggt ggtgaagttt ggtgagactc ttccaaagct gaagcccatt  
2521 atctctgacc ctgagtacct gctagaccag cacatcctca toagcatcaa gtccctctgac  
2581 agcgaagaat cctatggcga gggctgcatt gcccttcggt tagaggccac agaaacgcag  
2641 ctgcccactc acacgcctct caccacatt ggggagttga caggccactt ccagggggag  
2701 atcaagctgc agacctctca gggcaagacg agggagaagc totatgactt tbtgaagacg  
2761 gagegtgatg aatccagtgg gccaaagacc ctgaagagcc tcaccagcca cgaccccatg  
2821 aagcagtggg aagtcactag cagggccctt ccgtgcagtg gotccagcat cactgaaatc  
2881 atcaacccca actacatggg agtggggccc tttgggccac caatgcccct gcacgtgaag  
2941 cagaccttgt cccctgacca gcagcccaca gcctggagct acgaccagcc gcccaaggac  
3001 tccccgctgg gccctgcag gggagaaagt cctccgacac ctcccggcca gccgccata  
3061 tcaccaaga agtttttaac ctcaacagca aaccgggggtc tccctcccag gaacacaggag  
3121 tcaaggccca gtgacctggg gaagaacgca ggggacagc tgcctcagga ggacctgccc  
3181 ctgacgaagc ccgagatggt tgagaacccc ctgtatgggt ccttgagttc cttccctaag  
3241 cctgctccca ggaaggacca ggaatcccc aaaatgcgcg ggaaggaaac ccgcocctgc  
3301 ccggaacccc gcatcttgtc gccagcctc gtgctacca aagcccagga ggtgatcgc

## FIG. 2-continued

```
3361 ggcgaggggc ccggcaagca ggtgcccgcg ccccggtgc gctccttcac gtgctcatcc
3421 tctgccgagg gcagggcggc cggcggggac aagagccaag ggaagcccaa gaccccggtc
3481 agtcccagg ccccggtgcc ggccaagagg cccatcaagc cttccagatc ggaaatcaac
3541 cagcagaccc cgcaccaccc gacgcgcggc cgcgcgctgc cagtcaagag cccggcggtg
3601 ctgcacctcc agcactccaa gggcgcgcgc taccgcgcgc acaccgagct cccgcatacc
3661 ggcaagcacc ggccggagga ggggccacca gggcctctag gcaggactgc catgcagtga
3721 agccctcagt gagctgccac tgagtcggga gccagagga acggcgtgaa gccactggac
3781 cctctcccgg gacctcctgc tggctcctcc tgcccagctt cctatgcaag gctttgtgtt
3841 ttcaggaaaag ggccctagctt ctgtgtggcc cacagagtcc actgcctgtg agacttagca
3901 ccaagtgtctg aggttggaag aaaaacgcac accagacggg caacaaacag tctgggtccc
3961 cagctcgtct ttggtacttg ggaccccagt gcctcgttga gggcgccatt ctgaagaaag
4021 gaactgcagc gccgatttga ggggtggagat atagataata ataataataa taataataat
4081 ggccacatgg atcgaacact catgatgtgc caagtgtgtg gctaagtgtt ttacgaacat
4141 tcgtcatatc aggatgacct cgagagctga ggctctagcc acctaaaacc acgtgcccaa
4201 acccaccagt ttaaaacggg gtgtgttcgg aggggtgaaa gcattaagaa gcccagtgcc
4261 ctcttgaggt gagacaaggc ctgcgcctta aggagctgaa gagtctgggt agcttgttta
4321 ggggtacaaga agcctgttct gtccagcttc agtgacacaa gctgctttag ctaaagtccc
4381 gcgggttccg gcatggctag gctgagagca gggatctacc tggcttctca gttctttggt
4441 tggaaggagc aggaaatcag ctccatttct ccagtggaga gatctggcct cagcttgggc
4501 tagagatgcc aaggcctgtg ccaggttccc tgtgccctcc tcgaggtggg cagccatcac
4561 cagccacagt taagccaagc cccccaacat gtattccatc gtgctggtag aagagtcttt
4621 gctgtttgct ccgaaagccg tgctctccag cctggctgcc agggaggggt ggccctcttg
4681 ttccaggctc ttgaaatagt gcagcctttt ctccctatct ctgtggcttt cagctctgct
4741 tccttggtta ttaggagaat agatgggtga tgtctttcct tatgttgctt tttcaacata
4801 gcagaattaa tgtagggagc taaatccagt ggtgtgtgtg aatgcagaag ggaatgcacc
4861 ccacattccc atgatggaag tctgcgtaac caataaattg tgcccttctc actcaaaaaa
4921 aaaaa
```

## FIG. 3A

MPAMVPGWNHGNITRSKAEELLSRAGKDGSLVRASESIPRAYA  
LCVLFRNCVYTYRILPNEDDKFTVQASEGVPMRFFTKLDQLIDFYKKENMGLVTHLQY  
PVPLEEEDAIDEAEEDTVESVMSPPPELPPRNIPMSAGPSEAKDLPLATENPRAPEVTR  
LSLSETLFQRLQSMDTSGLP EEHLKAIQDYLSTQLLLDSDFLKTGSSNLP HLKKLMSL  
LCKELHGEVIRTLPSLES LQRLFDQQLSPGLRPRPQVPGEASPITMVAKLSQLTSLLS  
SIEDKVKSLLHEGSESTNRRSLIPPVTFEVKSESLGIPQKMHLKVDVESGKLIVKKSK  
DGSEDKFYSHKKILQLIKSQKFLNKLVLVETEKEKILRKEYVFADSKKREGFCQLLQ  
QMKNKHSEQPEPDMITIFIGTWNMGNA PPKKITSWFLSKGQGKTRDDSADYIPHDIY  
VIGTQEDPLGEKEWLELLRHSLQEVTSMTFKTVAIHTLWNIRIVVLAKPEHENRISHI  
CTDNVKTGIAN TLGNKGAAGVSFMFIGTSLGFVNSHLTSGSEKKLRNQNYMNILRFL  
ALGDKKLSPFNITHRFTHLFWLGDLNYRVELPTWEAEAI IQKIKQQQYSDLLAHDQLL  
LERKDQKVFLHFEEEEITFAPTYRFERLTRDKYAYTKQKATGMKYNLPSWCDRVLWKS  
YPLVHVVCQSYGSTSDIMTSDHSPVFATFEAGVTSQFVSKNGPGTVDSQGQIEFLACY  
ATLKTKSQTKFYLEFHSSCLESFVKSQEGENE EGSEGELVVRFGETLPKLKPIISDPE  
YLLDQHILISIKSSDSDESYGEGCIALRLETTEAQHP IYTPLTHHGEMTG HFRGEIKL  
QTSQGKMREKLYDFVKTERDESSGMKCLKNLTSHDPMRQWEPSGRVPACGVSSLNEMI  
NPNYIGMPFGQPLHGKSTLSPDQQLTAWSYDQLPKDSSLGPGRGEGPPTPPSQPPLS  
PKKFSSSTANRGPCPRVQEARPGDLGKVEALLQEDLLLTKPEMFENPLYGSVSSF PKL  
VPRKEQESPKMLRKEPPPCPDPGISSPSIVLPKAQEVE SVKGTSKQAPVPVLGPTPRI  
RSFTCSSSAEGRMTSGDKSQGKPKASASSQAPVPVKRPVKPSRSEMSQQTTPIPAPRP  
PLPVKSPAVLQLQHSGKGRDYRDNTELP HHGKHRQEEGLLGRTAMQ

## FIG. 3B

ggcaattttct gagaggcaac aggcggcagg tctcagccta gagagggccc tgaactactt  
61 tgctggagtg tccgtccctgg gagggtgctgc tgacccagtc caggagaccc atgcctgcca  
121 tgggtccctgg gtggaacat ggcaacatca cccgctccaa ggagaggag ctactttcca  
181 gagccggcaa ggacgggagc ttccttgctgc gtgccagcga gtccatcccc cgggectacg  
241 cactctgctg gctgttccgg aattgtgttt acacttacag gattctgccc aatgaggacg  
301 ataaattcac tgttcaggca tccgaagggtg tccccatgag gttcttcacg aagctggacc  
361 agctcatcga cttttacaag aaggaaaaca tggggctggg gaccacctg cagtaccccg  
421 tgccccctgga ggaggaggat gctattgatg aggctgagga ggacactgta gaaagtgtca  
481 tgtcaccacc tgagctgcct ccagaaaaca ttcctatgtc tgccgggccc agcgaggcca  
541 aggaccttcc tcttgcaaca gagaaccccc gagccccctga ggtcaccggg ctgagtctct  
601 ccgagacact gtttcagcgt ctacagagca tggataccag tgggcttccc gaggagcacc  
661 tgaaagccat ccaggattat ctgagcactc agctcctcct ggattccgac tttttgaaga  
721 cgggctccag caacctccct cacctgaaga agctgatgtc actgctctgc aaggagctcc  
781 atggggaagt catcaggact ctgccatccc tggagtctct gcagagggtg tttgaccaac  
841 agctctcccc aggccttcgc ccacgacctc aggtgcccgg agaggccagt cccatcacca  
901 tggttgocaa actcagccaa ttgacaagtc tgctgtcttc cattgaagat aaggtcaagt  
961 ccttgctgca cgagggtca gaatctacca acaggcgctc ccttatccct ccggtcacct  
1021 ttgaggtgaa gtcagagtcc ctgggcattc ctcagaaaat gcatctcaaa gtggacgttg  
1081 agtctgggaa actgatcgtt aagaagtcca aggatggctc tgaggacaag ttctacagcc  
1141 acaaaaaaat cctgcagctc attaagtccc agaagtttct aaacaagttg gtgattttgg  
1201 tggagacgga gaaggagaaa atcctgagga aggaatatgt ttttgctgac tctaagaaaa  
1261 gagaaggctt ctgtcaactc ctgcagcaga tgaagaacaa gcattcggag cagccagagc  
1321 ctgacatgat caccatcttc attggcactt ggaacatggg taatgcacc cctcccaaga  
1381 agatcacgtc ctggtttctc tccaaggggc agggaaagac acgggacgac tctgtgact  
1441 acatccccc tgacatctat gtgattggca cccaggagga tccccttgga gagaaggagt  
1501 ggctggagct actcaggcac tccctgcaag aagtcaccag catgacattt aaaacagttg  
1561 ccatccacac cctctggaac attcgcatag tgggtgcttg caagccagag catgagaatc  
1621 ggatcagcca tatctgact gacaacgtga agacaggcat cgccaacacc ctgggaaaca  
1681 agggagcagc gggagtgtcc ttcattgttc ttggaacctc cttgggggtc gtcaacagcc  
1741 acttgacttc tggaagtga aaaaagctca ggagaaatca aaactatatg aacatcctgc  
1801 ggttcctggc cctgggagac aagaagctaa gccatttaa catcacccac cgcttcaccc  
1861 acctcttctg gcttggggat ctcaactacc gcgtggagct gccacttg gaggcagagg  
1921 ccatcatcca gaagatcaag caacagcagt attcagacct tctggcccac gaccaactgc  
1981 tcctggagag gaaggaccag aaggtcttcc tgcactttga ggaggaagag atcaccttcg  
2041 cccccacctc tcgatttgaa agactgacct gggacaagta tgcatacacg aagcagaaag  
2101 caacagggat gaagtacaac ttgcctcct ggtgcgacc agtctctg aagtcttacc  
2161 cgctggtgca tgtggtctgt cagtctatg gcagtaccag tgacatcatg acgagtgacc  
2221 acagccctgt ctttgccacg tttgaagcag gagtacatc tcaattcgtc tccaagaatg  
2281 gtccctggcac tgtagatagc caagggcaga tcgagtttct tgcattgctac gccacactga  
2341 agaccaagtc ccagactaag ttctacttg agttccactc aagctgttta gagagttttg  
2401 tcaagagtca ggaaggagag aatgaagagg gaagtgaagg agagctggtg gtacggtttg  
2461 gagagactct tcccaagcta aagcccatta tctctgacct cgagtactta ctggaccagc  
2521 atatcctgat cagcattaaa tctctgaca gtgacgagtc ctatggtgaa ggctgcattg  
2581 cccttcgctt ggagaccaca gaggctcagc atcctatcta cacgctctc acccaccatg  
2641 gggagatgac tggccacttc aggggagaga ttaagctgca gacctccag ggcaagatga  
2701 gggagaagct ctatgacttt gtgaagacag agcgggatga atccagtggga atgaaatgct  
2761 tgaagaacct caccagccat gaccctatga ggcaatggga gccttctggc aggtccctg  
2821 catgtggtgt ctccagcctc aatgagatga tcaatccaaa ctacattggt atggggcctt  
2881 ttggacagcc cctgcatggg aatcaacc gactcctccc tggggcctgg gaggggggag ggtcctccaa  
2941 gttatgacca gctacccaaa gactcctccc tggggcctgg gaggggggag ggtcctccaa  
3001 cccctccctc ccaaccacct ctgtcgccaa agaagtttct atcttcaca gccaaaccgag  
3061 gtccttgccc cagggtgcaa gaggcaagac ctggggatct gggaaaggtg gaagctctgc  
3121 tccaggagga cctgctgctg acgaagccc agatgtttga gaaccactg tatggatccg  
3181 tgagttcctt ccctaagctg gtgccaggga aagagcagga gtctcccaag atgctgogga

## FIG. 3B-continued

3241 aggagcctccc gccctgtcca gaccacaggaa tctcatcacc cagcatcgtg ctccccaaag  
3301 cccaagaggt ggagagtgtc aaggggacaa gcaaacaggc ccctgtgcct gtccttgcc  
3361 ccacaccccg gatccgctcc ttacctgtt cttcttctgc tgagggcaga atgaccagt  
3421 gggacaagag ccaaggaag cccaaggcct cagccagttc ccaagcccca gtgccagtca  
3481 agaggcctgt caagccttcc aggtcagaaa tgagccagca gacaacaccc atcccagctc  
3541 cacggccacc cctgccagtc aagagtcctg ctgtcctgca gctgcaacat tccaaaggca  
3601 gagactaccg tgacaacaca gaactccccc accatggcaa gcaccgccaa gaggaggggc  
3661 tgcttggcag gactgccatg cagtgcagctg ctgggtgatcg gagcctggag gaacagcaca  
3721 aagcagacct gcgcctctct caggatgcct ctctcaggat gcctcttgga ggacctcctg  
3781 ctactctctt ttgcctagct tcaagtccca ggctgtgtat tttttttcag gaaacggcct  
3841 cacttctctg tggccaaga agtgtgctgc tggctgccac actgtgcggc agatgctaaa  
3901 gctggatgac aaacgcacgc catacagaca gcagacagcg gcaactgggtc tcagaacttg  
3961 gattcctggg ccttcttcca gtcgcggttt taaagaaagg aactaacgga gctgctcatc  
4021 cgagggtgaa gatataaata ataattattat taataataat aacagtcagg tgccatgtgc  
4081 tgtgttaagt gctttatgaa catttgctcg gctggcctcc agtgctgagg tgccagtcag  
4141 cctgaacct atgccaggc ccactaatcc caaatgggtg gtccctgagat gtttttaaaa  
4201 agcattaaag aaaaccatcg gtctcttaga gctaaccggc cgggctctac tgcagggacc  
4261 cgaacagtct gcatggctaa gtggcacaag gagcctggcc ctgtccagct tcagagatcc  
4321 aagctgcttt ttgctggggt tctgtcacag gcctgatcct cttgggtttt atgggggttc  
4381 aagtctgcca gagtcagaaa tcagctctaa ctgcagctg aagagatctg gccttaactt  
4441 aagccagcca cgtcaggccc ctgctgagcc tatggaccaa taaatactcc ccgtgccact  
4501 ggaggtgggc agctatcacc ataccctgag ttgggccaag cccacccac ccctaccctg  
4561 caacatttct gatgtwctga ggaagagtct ccacatagt cccaagggc tgagttctcc  
4621 agcctgctat cagggaaggt gagcattggt ccaggctct caaaatagtg cagcctcttc  
4681 ttcccaagct ctgggggtgca ccctgtgtcc ttggttacca ggagactagg gttgtgatat  
4741 cttttcttct cttgcttttt gatatatcag gattaatgta ggaaaccaga cctagattat  
4801 tcaggagagt aggtatatcc cctgtgtttc ccagtctgag tgaccaataa aattgtgcct  
4861 ttcta

1 aaacaggaag tcaatcagtt aagctgggtg cagcagccga ggccaccaag agggcaacggg  
61 cggcagggtg cagtggaggg gctcccgctc cctcgggtg tgtgtgggtc ctgggggtg  
121 tgcggcgccc ggcgaggagg gccacgccc accatgggtc cctgctggaa ccatggcaac  
181 atcacccgct ccaagcgga ggagctgctt tccaggacag gcaaggacgg gagcttctc  
241 gtgggtgcca gcgagtcacat ccccgggcca taccgctctt gctgtgctga tgcggaatga  
301 gctgctcaca cgaatctt gcccaatgaa gatgataaat tcaactgttca ggcatccgaa  
361 ggcggtctcca tgagggtctt caccagctg gaccagctca tccagtttta caagaaggaa  
421 aacatggggc tggtagccca tctgcaatac cctgtgcggc tggaggaa gaacacaggc  
481 gacgacctg aggaggacac agaaagtgtc gtgtctccac ccgagctgco ccaagaaac  
541 atcccgctga ctggcagctc ctgtgaggcc aaggaggttc ctttttcaaa cgagaatccc  
601 cgagcgagcc agaccagccg gccgagcttc tccgagacat tgttccagcg actgcaaac  
661 atggacacca gtcgggtctc ggaagacac cttgaaggcca tccaagatta ttttaagcact  
721 cagctcgccc aggaactctga atttgtgaag acagggtcca gcagttcttc tcaactgaag  
781 aaactgacca cactgctctg caaggagctc tatggagaag tcatccggac cctcccatcc  
841 ctggagtgct tgcagaggtt atttgaccac cagctctccc cgggctctcc tccactgct  
901 caggttctcg gtgaggccaa tccc atcaac atgtgttcca agctcagcca actgacaagc  
961 ctgttgcctc cacttgcctc caaggtcgaag gcttgcctgc acagggttcc tgaagtctcc  
1021 caccggccct cctttatccc tccagtcacc tttgagtgta aggcagagtc totggggatt  
1081 cctcagaaaa tgcagctcaa agtcgagctt gactctggga aactgatcat taagaagtc  
1141 aaggatggtt ctgaggaaca gtltctacag cacaagaaaa tctgtcagct catbaagtc  
1201 cagaatattc tgataaagt ggtgatcttg gtggaacagc agaaggagaa gatcctgagg  
1261 aaggaaatag tttttgctga ctcacaaaa agagaaggct tctgccagct cctgcagcag  
1321 atgaagaaca agcactcaga gcagccggag cccgacatga tccactctt catcgggacc  
1381 tggaaacttg gtaacgcccc cctcccaaac aagatcacgt cctgtgttct ctcacagggg  
1441 cagggaaga cgcgggagca cctcgggac taccatcccc atgacattta cgtgatcggc  
1501 acccaaggag accccctgag tgagaaggag tgggtggaga tctcacaaca cctcctgcaa  
1561 gaaatcacc a gtgtgacttt taaaacagtc gccatccaca cgtctctggaa catccgcatc  
1621 gtgtgtgctg ccaagcctga gccagagaa cggatcacgc acatctgtac tgacaacttg  
1681 aagacaggca ttgcaaacac actggggaac aaggagagcc tgggggtgtc gttcatg ttc  
1741 aatggaaact ccttaggggt cgtcaacagc cacttgactt caggaaagtg aagaagactc  
1801 aggcgaagac aaaaactatat gaacattctc cgttctctgg ccttggggca caagaagctg  
1861 agtcccttta acatcactca ccgcttcacg cactctctct ggtttgggga tcttaactac  
1921 cgtgtggatc tgcctacctg ggaggcagaa accatcctcc agaaaatcaa gcagcagcag  
1981 taccgagacc tctgttccca cagccagctg ctccacagaa ggaggagca gaaggtcttc  
2041 ctacacttcc agggagaaag aatcacgttt gccccaacct accgttttga gagactgact  
2101 cgggacaaat acgctcaac caagcagaaa cgcagagggc tgaagtacaa cttgctctcc  
2161 tgggtgtgac gactctctct gaagtcttat cctcgtgtgc acgtgtgtgt tcaactctat  
2221 ggcagtagca gcgacatcat gacgagtgac cagcgcctgt tctttgccc atttgaggca  
2281 ggagtcactt cccagtttgt ctcacagaac ggtcccgga ctgttgacag ccaaggagac  
2341 attgagtttc tcaagtgtct tgccacattg aagaccaaag cccagaccaa attctactct  
2401 gacttccact cagactgctt ggagagtttt gtcaagagtc aggaaggaga aatgaagaa  
2461 ggaagtgaag gggagctggt ggtgaagttt ggtgagactc ttcacaaagc gaagccatt  
2521 atctctgacc ctgagtaact gctagaccag cactctctca tcaagatcaa gtctctgac  
2581 agcagcgaat cctatggcga gggctgcat tgcctctggt tagaggccac agaaaagcag  
2641 ctgcccactt acacgctct caccacacat ggggagttga caggccactt ccagggggag  
2701 atcaagctgc agacctctca gggc aagacg agggagaaag tctatgactt tgtgaagacg  
2761 gagcgtgatg aatccagtg gccaaaacac ctagaagagc tcaccagcca cgaacccatg  
2821 aagcagtgag aagtcactag caggggccct cgtgagctg gctccagcat cactgaaac  
2881 atcaacccca actacatgg agtggggccc tttgggccc caatgcccct gcaagtgaag  
2941 cagaccttgt cccctgacca gcagcccaca gcctggagct acgaccagcc gcccaaggac  
3001 tcccgcgtg ggcctgca gggagaaagt cctccgacac ctcggggcca gcccgcata  
3061 tcccccagga agtttttacc ctcacacaga aaccggggtc tccctccagc gacacaggag  
3121 tcaaggccca gtgacctggg gaagaacgca ggggacagc tgcctcagga ggacctgccc  
3181 ctgacgagc cagagatgtt tgagaacccc ctgtatgggt ccttgagttc cttccctaa  
3241 cctgctccca ggaaggacca ggaatccccc aaaaatgcgc ggaaggaaac cccgcccctg  
3301 ccggaacccc gcatctgtc gccacagc g tgcctacca aagcccagga ggctgactgc  
3361 ggcgaggggg ccggcaagca ggtgcccgcg ccccggtgct gctccttcac gtgctcatcc  
3421 tctgcccagg ccaaggcggc cggcggggac aagagccaa ggaagcccaa gaccocggtc  
3481 agtcaccagg ccccggtgcc ggccaagag cccatcaagc cttccagatc ggaaatc aac  
3541 cagcagaccc cgcacacccc gacgcccggc cgcgcgctgc cagtcagag cccggcggtg  
3601 ctgacactcc agcactccaa gggccgcgac taccgcgaca acacagagct cccgcatcac  
3661 ggcaagcacc ggcgggagga ggggcccaca gggcctctag gcagagctgc catgcaagtga  
3721 agccctcagt gactgcccac tgaagtggga gccacagga acggcggtga gccactggac  
3781 cctctcccg gactcctgc tggctctctc tgcacagctt cctatgcaag gctttgtgtt  
3841 ttcaggaaag ggcctagctt ctgtgtgggc cagcagttc actgctgtg agacttagca  
3901 ccaagtgtgt aggtgtgaa aaaaaagcac accagacggg caacaaacag tctgggtccc  
3961 cagctcgtc tttgtacttg ggaccccagt gctcgttga gggcgccatt ctgaagaaag  
4021 gaactgcagc ggcgatttga ggtgtggagt atagataata ataataatga taataataat  
4081 ggccacatgg atcgaaact catgatgtgc caagtgtctg gc taagtgtc ttacgaacat  
4141 tgcctcatatc aggatgaact cgagagctga ggcctctagcc acctaaaaac acgtgcccac  
4201 accacacagt taaaaacggt gtgtgttctg aggggtgaaa gcattaaaga gccacgtgcc  
4261 cctctgagat gagacaaggg ctcggccctta aggaagctga gactcgggt agcttgttta  
4321 gggtagaaga agcctgttct gtcacagctt agtgacacaa gctgcttag ctaaaagccc  
4381 ggcgggtccc gcatggctag gctgagagca gggatctacc tggctctcca gttctttggt  
4441 tggaaaggagc aggaatcag cctctattct ccagttggga gatctggcct cagcttgggc  
4501 tagagatgcc aaggcctgtg ccag gttccc tgtgcccctc tcaaggtggg cagccatcac  
4561 cagccacagt taagccaaag ccccaaacat gtattccatc gtgctggtag aagagtcttt  
4621 gctgttgtct ccgaaagccg tgcctctcag cctgctgccc agggaggggt ggccctctgg  
4681 tctcaggtct ttgaatatgt gcagcctttt cttcctatct ctgtggcttt cagctctgct  
4741 tcttgggtta ttgagaaat agatgggtga tgttcttctc tatgtgtgct tttcaacata  
4801 gcagaattaa ttagggagc taatccaggt ggtgtgtgtg aatgcagaag ggaatgcacc  
4861 ccactctccc atgatggaag tctgcgtaac caataaattg tgccttctc actcaaaaa  
4921 aaaaa

SHIP-1 shRNA 1  
SHIP-1 shRNA 2  
SHIP-1 siRNA H1  
SHIP-1 siRNA H2

FIG. 4




FIG. 5A



FIG. 5B-1




FIG. 5B-2





FIG. 5B-3






FIG. 6

### siRNA Knockdown in NKL Cells




FIG. 7

# SHIP-DEFICIENCY TO INCREASE MEGAKARYOCYTE PROGENITOR PRODUCTION

## CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a divisional of U.S. application Ser. No. 11/451,004, filed Jun. 12, 2006, now U.S. Pat. No. 7,763,592, which is a continuation-in-part of U.S. application Ser. No. 10/904,667, filed Nov. 22, 2004, now U.S. Pat. No. 7,807,646, which claims the benefit of U.S. Provisional Application Ser. No. 60/481,677, filed Nov. 20, 2003, each of which is hereby incorporated by reference herein in its entirety, including any figures, tables, nucleic acid sequences, amino acid sequences, and drawings.

## GOVERNMENT SUPPORT

This invention was made with government support under grant numbers HL072523 and CA087989 awarded by the Leukemia and Lymphoma Society of America and the National Institutes of Health. The government has certain rights in the invention.

## BACKGROUND OF THE INVENTION

Platelets are critical for blood clotting. However, in various human anemias, and in bone marrow transplant patients, platelets and the megakaryocytes they are derived from can drop below a critical threshold that is required to maintain normal clotting. This can require platelet transfusions that are very expensive and which place the patient at risk for infection by blood-borne pathogens (e.g. HIV, HepB and C).

SH2-containing-5' inositol phosphatase-1 (SHIP) can catalyze the removal of the 5' phosphate group from  $PI_{(3,4,5)}P_3$  (PIP3) (Damen, J. E. et al. *Proc Natl Acad Sci USA*, 1996, 93:1689-1693). In this manner, SHIP regulates survival and proliferation of various hematopoietic cell types. The numbers of myeloid cells and osteoclasts are increased in SHIP-deficient mice due to enhanced activity of the phosphatidyl inositol 3-kinase (PI3K)/Akt signaling pathway that promotes their survival (Takeshita, S. et al. *Nat Med.*, 2002, 8:943-949; Helgason, C. D. et al. *Genes Dev.*, 1998, 12:1610-1620; Liu, Q. et al. *Genes Dev.*, 1999, 13:786-791).

Furthermore, the present inventors have shown that the number of natural killer cells are increased in SHIP-deficient mice resulting in an enhancement of engraftment of allogeneic hematopoietic stem cell grafts (Wang, J. W. et al. *Science*, 2002, 295:2094-2097). SHIP is also known to influence signaling pathways downstream of receptors for chemokines and cytokines involved in megakaryocytopoiesis and thrombopoiesis, such as Stromal-cell-derived-Factor 1 (SDF-1/CXCL-12) (Wang, J. F. et al. *Blood*, 1998, 92:756-764; Hamada, T. et al. *J Exp Med.*, 1998, 188:539-548; Hattori, K. et al. *Blood*, 2001, 97:3354-3360; Avecilla, S. T. et al., *Nat Med.*, 2004, 10:64-71; Chernock, R. D. et al. *Blood*, 2001, 97:608-615), interleukin-3 (Liu, L. et al. *Mol Cell Biol.*, 1994, 14:6926-6935), and thrombopoietin (TPO) (Lok, S. et al. *Nature*, 1994, 369:565-568; Drachman, J. G. et al. *Proc Natl Acad Sci USA*, 1997, 94:2350-2355). SHIP is phosphorylated after TPO binding to its receptor, c-mpl, leading to activation of PI3K that promotes cycling of megakaryocytes (MK) (Drachman, J. G. et al. *Proc Natl Acad Sci USA*, 1997, 94:2350-2355; Drachman, J. G. et al. *Blood*, 1997, 89:483-492; Geddis, A. E. et al. *J Biol Chem.*, 2001, 276:34473-34479). TPO influences MK development by controlling

their proliferation, differentiation, survival and endoduplication (Kaushansky, K. et al. *Nature*, 1994, 369:568-571). Circulating platelets sequester free TPO, and thereby limit megakaryocytopoiesis during steady-state hematopoiesis (Kaushansky, K. *N Engl J Med.*, 1998, 339:746-754). Furthermore, SDF-1/CXCL12 induces transendothelial MK migration and platelet production in vitro (Wang, J. F. et al. *Blood*, 1998, 92:756-764; Hamada, T. et al. *J Exp Med.*, 1998, 188:539-548) and in vivo (Hattori, K. et al. *Blood*, 2001, 97:3354-3360). The present inventors have also shown that it enhances human thrombocytopoiesis in xenotransplanted NOD/SCID mice (Perez, L. E. et al. *Exp Hematol.*, 2004, 32:300-307). SHIP-deficient myeloid progenitors exhibit enhanced chemotaxis towards SDF-1/CXCL-12, indicating SHIP influences signaling downstream of CXCR-4 (Kim, C. H. et al. *J Clin Invest.*, 1999, 104:1751-1759). In addition, SHIP has been shown to regulate PIP3 levels after thrombin or collagen activation of platelets (Giuriato, S. et al. *J Biol Chem.*, 1997, 272:26857-26863; Giuriato, S. et al. *Biochem J.*, 2003, 376:199-207).

Thus, the present inventors hypothesized that SHIP may also be involved in the regulation of megakaryocytopoiesis and platelet production in vivo. It has been reported that colonyforming-unit megakaryocyte (CFU-Mk) are decreased in SHIP<sup>-/-</sup> bone marrow (BM) (Moody, J. L. et al. *Blood*, 2004, 103:4503-10).

A naturally occurring gene-silencing mechanism triggered by double-stranded RNA (dsRNA), designated as small interfering RNA (siRNA), has emerged as a very important tool to suppress or knock down gene expression in many systems. RNA interference is triggered by dsRNA that is cleaved by an RNase-III-like enzyme, Dicer, into 21-25 nucleotide fragments with characteristic 5' and 3' termini (Provost, P. D. et al. *Embo J*, 2002, 21:5864). These siRNAs act as guides for a multi-protein complex, including a PAZ/PIWI domain containing the protein Argonaute2, that cleaves the target mRNA (Hammond, S. M. et al. *Science*, 2001, 293:1146-1150). These gene-silencing mechanisms are highly specific and potent and can potentially induce inhibition of gene expression throughout an organism. The short interference RNA (siRNA) approach has proven effective in silencing a number of genes of different viruses (Fire, A. *Trends Genet.*, 1999, 15:358-363).

RNA interference (RNAi) is a polynucleotide sequence-specific, post-transcriptional gene silencing mechanism effected by double-stranded RNA that results in degradation of a specific messenger RNA (mRNA), thereby reducing the expression of a desired target polypeptide encoded by the mRNA (see, e.g., WO 99/32619; WO 01/75164; U.S. Pat. No. 6,506,559; Fire et al., *Nature* 391:806-11 (1998); Sharp, *Genes Dev.* 13:139-41 (1999); Elbashir et al. *Nature* 411:494-98 (2001); Harborth et al., *J. Cell Sci.* 114:4557-65 (2001)). RNAi is mediated by double-stranded polynucleotides, such as double-stranded RNA (dsRNA), having sequences that correspond to exonic sequences encoding portions of the polypeptides for which expression is compromised. RNAi reportedly is not effected by double-stranded RNA polynucleotides that share sequence identity with intronic or promoter sequences (Elbashir et al., 2001). RNAi pathways have been best characterized in *Drosophila* and *Caenorhabditis elegans*, but "small interfering RNA" (siRNA) polynucleotides that interfere with expression of specific polynucleotides in higher eukaryotes such as mammals (including humans) have also been considered (e.g., Tuschl, 2001 *Chem-biochem.* 2:239-245; Sharp, 2001 *Genes Dev.* 15:485; Bernstein et al., 2001 *RNA* 7:1509; Zamore, 2002 *Science* 296:1265; Plasterk, 2002 *Science* 296:1263; Zamore 2001 *Nat.*

*Struct. Biol.* 8:746; Matzke et al., 2001 *Science* 293:1080; Scadden et al., 2001 *EMBO Rep.* 2:1107).

According to a current non-limiting model, the RNAi pathway is initiated by ATP-dependent cleavage of long dsRNA into double-stranded fragments of about 18-27 (e.g., 19, 20, 21, 22, 23, 24, 25, 26, etc.) nucleotide base pairs in length, called small interfering RNAs (siRNAs) (see review by Hutvagner et al., *Curr. Opin. Gen. Dev.* 12:225-32 (2002); Elbashir et al., 2001; Nyknen et al., *Cell* 107:309-21 (2001); Zamore et al., *Cell* 101:25-33 (2000)). In *Drosophila*, an enzyme known as "Dicer" cleaves the longer double-stranded RNA into siRNAs; Dicer belongs to the RNase III family of dsRNA-specific endonucleases (WO 01/68836; Bernstein et al., *Nature* 409:363-66 (2001)). Further, according to this non-limiting model, the siRNA duplexes are incorporated into a protein complex, followed by ATP-dependent unwinding of the siRNA, which then generates an active RNA-induced silencing complex (RISC) (WO 01/68836). The complex recognizes and cleaves a target RNA that is complementary to the guide strand of the siRNA, thus interfering with expression of a specific protein (Hutvagner et al., supra).

In *C. elegans* and *Drosophila*, RNAi may be mediated by long double-stranded RNA polynucleotides (WO 99/32619; WO 01/75164; Fire et al., 1998; Clemens et al., *Proc. Natl. Acad. Sci. USA* 97:6499-6503 (2000); Kisielow et al., *Biochem. J.* 363:1-5 (2002); see also WO 01/92513 (RNAi-mediated silencing in yeast)). In mammalian cells, however, transfection with long dsRNA polynucleotides (i.e., greater than 30 base pairs) leads to activation of a non-specific sequence response that globally blocks the initiation of protein synthesis and causes mRNA degradation (Bass, *Nature* 411:428-29 (2001)). Transfection of human and other mammalian cells with double-stranded RNAs of about 18-27 nucleotide base pairs in length interferes in a sequence-specific manner with expression of particular polypeptides encoded by messenger RNAs (mRNA) containing corresponding nucleotide sequences (WO 01/75164; Elbashir et al., 2001; Elbashir et al., *Genes Dev.* 15:188-200 (2001)); Harborth et al., *J. Cell Sci.* 114:4557-65 (2001); Carthew et al., *Curr. Opin. Cell Biol.* 13:244-48 (2001); Mailand et al., *Nature Cell Biol.* Advance Online Publication (Mar. 18, 2002); Mailand et al. 2002 *Nature Cell Biol.* 4:317).

siRNA may offer certain advantages over other polynucleotides known to the art for use in sequence-specific alteration or modulation of gene expression to yield altered levels of an encoded polypeptide product. These advantages include lower effective siRNA polynucleotide concentrations, enhanced siRNA stability, and shorter siRNA oligonucleotide lengths relative to such other polynucleotides (e.g., antisense, ribozyme or triplex polynucleotides). By way of a brief background, "antisense" polynucleotides bind in a sequence-specific manner to target nucleic acids, such as mRNA or DNA, to prevent transcription of DNA or translation of the mRNA (see, e.g., U.S. Pat. Nos. 5,168,053; 5,190,931; 5,135,917; 5,087,617; see also, e.g., Clusel et al., 1993 *Nucl. Acids Res.* 21:3405-11, describing "dumbbell" antisense oligonucleotides). "Ribozyme" polynucleotides can be targeted to an RNA transcript and are capable of catalytically cleaving such transcripts, thus impairing translation of mRNA (see, e.g., U.S. Pat. Nos. 5,272,262; 5,144,019; and 5,168,053, 5,180,818, 5,116,742 and 5,093,246; U.S. Ser. No. 2002/193579). "Triplex" DNA molecules refers to single DNA strands that bind duplex DNA to form a colinear triplex molecule, thereby preventing transcription (see, e.g., U.S. Pat. No. 5,176,996, describing methods for making synthetic oligonucleotides that bind to target sites on duplex DNA). Such triple-stranded structures are unstable and form only

transiently under physiological conditions. Because single-stranded polynucleotides do not readily diffuse into cells and are therefore susceptible to nuclease digestion, development of single-stranded DNA for antisense or triplex technologies often requires chemically modified nucleotides to improve stability and absorption by cells. siRNAs, by contrast, are readily taken up by intact cells, are effective at interfering with the expression of specific polynucleotides at concentrations that are several orders of magnitude lower than those required for either antisense or ribozyme polynucleotides, and do not require the use of chemically modified nucleotides.

Due to its advantages, RNAi has been applied as a target validation tool in research and as a potential strategy for in vivo target validation and therapeutic product development (Novina, C. D. and Sharp, P. A., *Nature*, 2004, 430:161-164). In vivo gene silencing with RNAi has been reported using viral vector delivery and high-pressure, high-volume intravenous (i.v.) injection of synthetic iRNAs (Scherr, M. et al. *Oligonucleotides*, 2003, 13:353-363; Song, E. et al. *Nature Med.*, 2003, 347-351). In vivo gene silencing has been reported after local direct administration (intravitreal, intranasal, and intrathecal) of siRNAs to sequestered anatomical sites in various models of disease or injury, demonstrating the potential for delivery to organs such as the eye, lungs, and central nervous system (Reich, S. J. et al. *Mol. Vis.*, 2003, 9:210-216; Zhang, X. et al. *J. Biol. Chem.*, 2004, 279:10677-10684; Dorn, G. et al. *Nucleic Acids Res.*, 2004, 32, e49). Silencing of endogenous genes by systemic administration of siRNAs has also been demonstrated (Soutschek, J. et al. *Nature*, 2004, 432:173-178). It has been shown that siRNAs delivered systemically in a liposomal formulation can silence the disease target apolipoprotein B (ApoB) in non-human primates (Zimmermann T. S. et al., *Nature*, 2006, 441:111-114).

#### BRIEF SUMMARY OF THE INVENTION

The unexpected observations below provide the basis for this invention, which is directed to a method of increasing the yield of megakaryocytes and megakaryocyte progenitors through the inhibition of SHIP gene function.

As indicated above, SHIP influences signals downstream of cytokine and chemokine receptors that play a role in megakaryocytopoiesis, including the receptors for thrombopoietin and Stromal-cell-derived-Factor 1/CXCL-12. Herein, the present inventors show that two separate strains of mice with different SHIP mutations exhibit profound increases in megakaryocyte progenitors (MKP; Lin<sup>-</sup>cKit<sup>+</sup>CD41<sup>+</sup>) and megakaryocytes (MK; Lin<sup>-</sup>cKit<sup>+</sup>CD41<sup>+</sup>) in the hematopoietic compartment. However, despite increased MKP and MK numbers, platelet production is significantly lower in SHIP-deficient mice. These findings demonstrate that SHIP is a key regulator of signaling pathways that control megakaryocytopoiesis in vivo.

Mice that lack expression of a SHIP gene exhibit increased levels of both megakaryocyte progenitors and megakaryocytes in the bone marrow and spleen. In fact, megakaryocytes, the immediate precursor of platelets, are increased in the periphery of SHIP-deficient mice approximately 10- to 100-fold. Therefore, methods that inhibit SHIP expression, its enzymatic activity, or its signaling functions could be used in human patients in vivo to temporarily increase megakaryocytes during periods when their platelets drop below numbers sufficient to promote normal blood clotting. In a similar way, SHIP expression or activity could be used to increase the

yield of megakaryocytes and megakaryocyte progenitors in ex vivo expansion regimens that use human growth factors.

One embodiment of the invention is a method for increasing the yield of megakaryocytes or megakaryocyte progenitors in a patient, in vivo, comprising the steps of administering a therapeutically effective amount of a substance that inhibits SHIP function (a SHIP inhibitor) to the patient. The SHIP inhibiting substance can be, for example, one or more of the following: interfering RNA, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, non-standard nucleic acids, aptamers, decoys, oligonucleotide based gene regulation, or dominant/negative mutants.

Another embodiment relates to a method for increasing the yield of megakaryocytes or megakaryocyte progenitors in a sample in vitro (e.g., a sample obtained from a patient ex vivo), comprising providing a sample containing target cells and contacting the target cells in vitro with an efficacious amount of a substance that inhibits SHIP function. The SHIP inhibiting substance can be, for example, one or more of the following: interfering RNA, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, non-standard nucleic acids, aptamers, decoys, oligonucleotide-based gene regulation, and dominant/negative mutants. Target cells include megakaryocytes and/or megakaryocyte progenitors. The sample can be any composition containing megakaryocytes and/or megakaryocyte progenitors. For example, the sample can be a biological sample obtained from a patient (such as peripheral blood, bone marrow, spleen, or other tissue or fluid containing megakaryocytes and/or megakaryocyte progenitors), or a cell culture. Thus, the step of providing a sample containing target cells can comprise harvesting a sample from a patient.

Another embodiment of the invention relates to a method for improving haematopoietic recovery in a patient in need thereof, comprising administering a therapeutically effective amount of a substance that inhibits SHIP function to the patient. The SHIP inhibiting substance can be, for example, one or more of the following: interfering RNA, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, non-standard nucleic acids, aptamers, decoys, oligonucleotide-based gene regulation, and dominant/negative mutants. Target cells include megakaryocytes and/or megakaryocyte progenitors.

Another embodiment of the invention relates to a method for improving haematopoietic recovery in a patient in need thereof, ex vivo, comprising harvesting target cells from a patient, contacting the target cells with an efficacious amount of a substance that inhibits SHIP function, and delivering (e.g., re-infusing) the expanded target cells back into the patient. The target cells are chosen from among megakaryocytes and/or megakaryocyte progenitors.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A, 1B-1, 1B-2, 1B-3, 1C-1, 1C-2, 1C-3, 1D, and 1E show increased numbers of MKP and MK in SHIP-deficient mice. FIG. 1A shows results of flow cytometric analysis of bone marrow (BM), peripheral blood (PB) and spleen of SHIP<sup>-/-</sup> (-/-) and SHIP<sup>ΔIP/ΔIP</sup> (ΔIP/ΔIP) and their respective WT littermates. Shown is a representative flow cytometry plot of c-Kit vs. CD41 after gating on live cells and Lin<sup>-</sup> cells. MKP are in the upper right quadrant and MK in the lower right quadrant. Percentages for each population are indicated on the plot. FIGS. 1B-1, 1B-2, and 1B-3 show absolute numbers of MKP (Lin<sup>-</sup>cKit<sup>+</sup>CD41<sup>+</sup>) in BM (sum for 2 femurs+2 tibias), intact spleen, and PB. FIGS. 1C-1, 1C-2, and 1C-3 show absolute numbers of MK (Lin<sup>-</sup>cKit<sup>+</sup>CD41<sup>+</sup>) in BM

(sum for 2 femurs+2 tibias), intact spleen, and PB. The bar graphs show the different SHIP-deficient models in black and their respective WT littermates in gray. (n=3 mice/genotype). Statistical significance was assessed using the two-tailed Student t-test, \*\*p=0.0001, \*p<0.0005, ++p<0.005, +p<0.05. FIGS. 1D and 1E show hematoxylin-eosin (H & E) staining of SHIP<sup>ΔIP/ΔIP</sup> and WT spleen section, and SHIP<sup>-/-</sup> and WT BM section, respectively. The images were photographed at 40× (spleen) and 63×(BM) magnification.

FIG. 2 shows the human SHIP cDNA sequence (GenBank accession no. NM\_005541).

FIGS. 3A and 3B show mouse SHIP amino acid and cDNA sequences (GenBank accession no. NM\_10566).

FIG. 4 shows the human SHIP cDNA sequence with the targets for siRNA sequence H1 (SEQ ID NO:16), siRNA sequence H2 (SEQ ID NO:17), shRNA 63332 (SEQ ID NO:18), and shRNA 63331 (SEQ ID NO:19) in boxes.

FIGS. 5A, 5B-1, 5B-2, and 5B-3 demonstrate a significant increase in the percentage of MKP cells in SHIP-ablated BM and spleen. MxCre<sup>+</sup> and MxCre<sup>-</sup> mice with floxed SHIP alleles were treated with polyIC 3 times prior to being analyzed. FIG. 5A shows representative FACS plots showing detection of and MK and MKP in the BM and spleen of MxCre<sup>+</sup> and MxCre<sup>-</sup> mice after treatment. FIG. 5B-1 shows the percentage of MKP in BM (top), spleen (Spl) (bottom) and of SHIP-ablated (black) and WT (grey) mice. FIG. 5B-2 shows the absolute number of MKP cells in BM (per femur and tibia pair). FIG. 5B-3 shows the percentage of MK found in the spleen. Data was acquired on a FACS Calibur with CellQuest software (BD Biosciences, San Jose, Calif.), and analyzed with FlowJo. Significance was established using the unpaired student t test (Prism 4). +++p<0.0005, ++p<0.005, and +p<0.05. (mean±SEM, n=3).

FIG. 6 is a general vector map representing shRNA 63332 and shRNA 63331.

FIG. 7 is a Western blot demonstrating SHIP knockdown in NK cell lines using siRNA sequence H1 (SEQ ID NO:16) and siRNA sequence H2 (SEQ ID NO:17). Actin was used as the control. As shown in FIG. 7, H1 was more effective at silencing SHIP.

#### BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is the human SHIP cDNA sequence (GenBank accession nos. NM\_005541 and NM\_001017915) (FIG. 2).

SEQ ID NO:2 is the mouse SHIP cDNA sequence (GenBank accession no. NM\_10566) (FIG. 3).

SEQ ID NOS:3-15 are examples of target regions within the human SHIP open reading frame.

SEQ ID NOS:16-17 are siRNA sequences (designated H1 and H2, respectively) that have been identified as effective at SHIP knockdown in human NK cell lines.

SEQ ID NOS:18-19 are shRNA sequences (designated 63332 and 63331, respectively) targeting human SHIP, including the loop and termini portions of the shRNA.

SEQ ID NOS:20-21 are the target sequences within human SHIP for shRNA 6332 (SEQ ID NO:18) and shRNA 63331 (SEQ ID NO:19), respectively.

#### DETAILED DESCRIPTION OF THE INVENTION

As described in detail in the Examples, BM of SHIP-deficient mice exhibit increased numbers of megakaryocyte progenitors (MKP) (Lin<sup>-</sup>, c-Kit<sup>+</sup>, CD41<sup>+</sup>), but decreased numbers of MK (Lin<sup>-</sup>, c-Kit<sup>+</sup>, CD41<sup>+</sup>). In addition, the present inventors observed expanded numbers of both MKP and MK

in SHIP-deficient spleens. Peripheral blood (PB) of SHIP-deficient mice also contains increased numbers of MK relative to WT controls. However, platelet levels in the PB of SHIP-deficient mice were not significantly altered relative to WT controls.

The invention is based on a method of modulating megakaryocytes and their progenitors, as a significant leap forward in the treatment of various human anemias. The present inventors reasoned that the identification of a novel gene involved in megakaryocyte production would lead to increased efficacy of current treatments. The clinical potential of such an approach is significant, as it allows for modulation of a gene-specific determinate of megakaryocyte production.

One embodiment of the invention is a method of increasing the yield of megakaryocytes in a patient, in vivo, comprising the steps of administering a therapeutically effective amount of a substance that inhibits SHIP function to the patient. The SHIP inhibiting substance can be, for example, one or more of the following: interfering RNA, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, non-standard nucleic acids, aptamers, decoys, oligonucleotide based gene regulation, or dominant/negative mutants.

Another embodiment relates to a method of increasing the yield of megakaryocytes in a sample in vitro (e.g., a sample obtained from a patient ex vivo), comprising providing a sample containing target cells and contacting the target cells in vitro with an efficacious amount of a substance that inhibits SHIP function. The SHIP inhibiting substance can be, for example, one or more of the following: interfering RNA, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, non-standard nucleic acids, aptamers, decoys, oligonucleotide-based gene regulation, and dominant/negative mutants. Target cells include megakaryocytes and/or megakaryocyte progenitors. The sample can be any composition containing megakaryocytes and/or megakaryocyte progenitors. For example, the sample can be a biological sample obtained from a patient (such as peripheral blood, bone marrow, spleen, or other tissue or fluid containing megakaryocytes and/or megakaryocyte progenitors), or a cell culture. Thus, the step of providing a sample containing target cells can comprise harvesting a sample from a patient. Optionally, the method of the invention further comprises detecting the presence of megakaryocytes or their progenitors in the sample before, during, and/or after administration of the SHIP inhibitor. Optionally, the method of the invention further comprises isolating the megakaryocytes or their progenitors from the sample. Isolation of the cells may be carried out before, during, and/or after detection of the megakaryocytes or megakaryocyte progenitors within the sample.

Methods for detecting megakaryocytes and their progenitors are known in the art (see, for example, Nakorn T. N. et al., *Proc. Natl. Acad. Sci. USA*, 2003, 100:205-210; Bruno E. and R. Hoffman., *Semin. Hematol.*, 1998, 35(3):183-191; Ivanyi J. L. et al., *Act Histochem.*, 1993, 95(1):79-88; Vannucchi A. S. et al., *Blood*, 2000, 95:2559-2568; Boque C. et al., *J. Clinical Pathology*, 1989, 42:982-984; Long M. W. and C. H. Heffner, *Exp. Hematol.*, 1998, 16(1):62-70; Bruno E. et al., *Exp. Hematol.*, 1996, 24(4):552-558; Long M. W. et al., *J. Clin. Invest.*, 1998, 82(5):1779-1786; U.S. patent publication US 2005/0003471 A1, Wang et al., each of which are incorporated herein by reference in their entirety). For example, megakaryocytes may be detected based on cell surface markers (such as CD9, CD41, CD61, actin, FVIIIIRAg) and optionally separated using flow cytometry.

Another embodiment of the invention relates to a method for improving haematopoietic recovery in a patient in need thereof, comprising administering a therapeutically effective

amount of a substance that inhibits SHIP function to the patient. The SHIP inhibiting substance can be, for example, one or more of the following: interfering RNA, antisense oligonucleotides, ribozymes, DNazymes, nucleic acid modifiers, PNAs, non-standard nucleic acids, aptamers, decoys, oligonucleotide-based gene regulation, and dominant/negative mutants. Target cells include megakaryocytes and/or megakaryocyte progenitors.

Another embodiment of the invention relates to a method for improving haematopoietic recovery in a patient in need thereof, ex vivo, comprising harvesting target cells from a patient, contacting the target cells with an efficacious amount of a substance that inhibits SHIP function, and delivering (e.g., re-infusing) the expanded target cells back into the patient. The target cells include megakaryocytes and/or megakaryocyte progenitors. Optionally, the method of the invention further comprises isolating the megakaryocytes or their progenitors from the sample before, during, and/or after administration of the SHIP inhibitor. Isolation of the cells may be carried out before, during and/or after detection of the megakaryocytes or megakaryocyte progenitors within the sample.

As used herein, the term "SHIP" refers to hematopoietic-specific SH2-containing inositol-5-phosphatase-1, which catalyzes the removal of the 5' phosphate group from PI<sub>(3,4,5)</sub>P<sub>3</sub> (PIP<sub>3</sub>) and inositol 1,3,4,5-tetrakisphosphate (IP<sub>4</sub>). SHIP, which is also known in the scientific literature as SHIP-1, SHIP1, SHIP1, and SHIP-I was also the subject of Helgason, et al., *Genes Dev.*, 1998, 12(11):1610-1620; Huber et al., *Proc. Natl. Aca. Sci. USA*, 1998, 95(19):11330-11335; Liu et al., *Genes Dev.*, 1999, 13(7):789-791; Liu et al.; *J. Exp. Med.*, 1998, 188(7):1333-1342; Rohrschneider et al., *Genes & Development*, 2000, 14:505-520, U.S. Pat. No. 6,090,621 (Kavanaugh et al.), PCT publication WO 9710252A1 (Rohrschneider, L. R.), and PCT publication WO 9712039A2 (Krystal, G.). The nucleotide sequences of mouse SHIP and human SHIP, for example, have been publicly available for several years (GenBank Accession Numbers NM\_10566 and NM\_005541, respectively, on the National Center for Biotechnology Information (NCBI) database).

The terms "inhibitor of SHIP", "SHIP inhibitor", and "SHIP inhibiting substance" are used herein interchangeably to refer to any molecule that decreases the activity of SHIP (inositol phosphatase activity) or decreases the protein level of SHIP. Thus, a SHIP inhibitor can be a small molecule that decreases activity of SHIP, e.g., by interfering with interaction of the inositol phosphatase with another molecule, e.g., its substrate. The SHIP inhibitor can also be a small molecule that decreases expression of the gene encoding the inositol phosphatase. An inhibitor can also be an interfering RNA molecule, antisense oligonucleotide, a ribozyme, an antibody, or a dominant negative mutant of SHIP. Dominant negative mutants of SHIP have been developed (Gupta N. et al., *J. Exp. Med.*, 1997, 186(3):473-478; and Tridandapani S. et al., *J. Immunol.*, 2002, 169(8):4370-4378, which are each incorporated herein by reference in their entirety). A "direct inhibitor" of SHIP is an inhibitor that interacts with the SHIP enzyme, or substrate thereof, or with a nucleic acid encoding SHIP (e.g., the SHIP gene or its mRNA) or its regulatory sequences. An "indirect inhibitor" of SHIP is an inhibitor that interacts upstream or downstream of the SHIP enzyme in the regulatory pathway, and which does not interact with the enzyme or substrate thereof or with a nucleic acid encoding SHIP or its regulatory sequences.

The methods of the invention can further include a step of determining the amount or concentration of megakaryocytes or megakaryocyte progenitors in vitro (e.g., in a sample) or in

vivo (e.g., in a mammal) before introduction of the SHIP inhibitor, after introduction of the SHIP inhibitor, or both. The terms “detecting”, “detection”, “analyzing”, “analysis”, and other grammatical variations thereof (e.g., “detecting megakaryocytes,” etc.) refer to any quantitative, semi-quantitative, or qualitative method for determining an analyte in general, and a megakaryocyte or megakaryocyte progenitor in particular. For example, a method that merely detects the presence or absence of a megakaryocyte or its progenitor in a sample lies within the scope of the term, as do methods that provide data as to the amount or concentration of the cells in the sample.

As used herein, the term “polypeptide” refers to any polymer comprising any number of amino acids, and is interchangeable with “protein”, “gene product”, and “peptide”.

As used herein, the term “nucleoside” refers to a molecule having a purine or pyrimidine base covalently linked to a ribose or deoxyribose sugar. Exemplary nucleosides include adenosine, guanosine, cytidine, uridine and thymidine.

The term “nucleotide” refers to a nucleoside having one or more phosphate groups joined in ester linkages to the sugar moiety. Exemplary nucleotides include nucleoside monophosphates, diphosphates and triphosphates. The terms “polynucleotide” and “nucleic acid molecule” are used interchangeably herein and refer to a polymer of nucleotides joined together by a phosphodiester linkage between 5' and 3' carbon atoms. The terms “nucleic acid” or “nucleic acid sequence” encompass an oligonucleotide, nucleotide, polynucleotide, or a fragment of any of these, DNA or RNA of genomic or synthetic origin, which may be single-stranded or double-stranded and may represent a sense or antisense strand, peptide nucleic acid (PNA), or any DNA-like or RNA-like material, natural or synthetic in origin. As will be understood by those of skill in the art, when the nucleic acid is RNA, the deoxynucleotides A, G, C, and T are replaced by ribonucleotides A, G, C, and U, respectively.

As used herein, the term “RNA” or “RNA molecule” or “ribonucleic acid molecule” refers generally to a polymer of ribonucleotides. The term “DNA” or “DNA molecule” or “deoxyribonucleic acid molecule” refers generally to a polymer of deoxyribonucleotides. DNA and RNA molecules can be synthesized naturally (e.g., by DNA replication or transcription of DNA, respectively). RNA molecules can be post-transcriptionally modified. DNA and RNA molecules can also be chemically synthesized. DNA and RNA molecules can be single-stranded (i.e., ssRNA and ssDNA, respectively) or multi-stranded (e.g., double stranded, i.e., dsRNA and dsDNA, respectively). Based on the nature of the invention, however, the term “RNA” or “RNA molecule” or “ribonucleic acid molecule” can also refer to a polymer comprising primarily (i.e., greater than 80% or, preferably greater than 90%) ribonucleotides but optionally including at least one non-ribonucleotide molecule, for example, at least one deoxyribonucleotide and/or at least one nucleotide analog.

As used herein, the term “nucleotide analog”, also referred to herein as an “altered nucleotide” or “modified nucleotide” refers to a non-standard nucleotide, including non-naturally occurring ribonucleotides or deoxyribonucleotides. Preferred nucleotide analogs are modified at any position so as to alter certain chemical properties of the nucleotide yet retain the ability of the nucleotide analog to perform its intended function.

As used herein, the term “RNA analog” refers to a polynucleotide (e.g., a chemically synthesized polynucleotide) having at least one altered or modified nucleotide as compared to a corresponding unaltered or unmodified RNA but retaining the same or similar nature or function as the corresponding unaltered or unmodified RNA. As discussed above,

the oligonucleotides may be linked with linkages which result in a lower rate of hydrolysis of the RNA analog as compared to an RNA molecule with phosphodiester linkages. Exemplary RNA analogues include sugar- and/or backbone-modified ribonucleotides and/or deoxyribonucleotides. Such alterations or modifications can further include addition of non-nucleotide material, such as to the end(s) of the RNA or internally (at one or more nucleotides of the RNA). An RNA analog need only be sufficiently similar to natural RNA that it has the ability to mediate (mediates) RNA interference or otherwise reduce target gene expression.

The terms “operably-linked” or “operatively-linked” are used herein interchangeably to refer to an arrangement of flanking sequences wherein the flanking sequences so described are configured or assembled so as to perform their usual function. Thus, a flanking sequence operably-linked to a coding sequence may be capable of effecting the replication, transcription and/or translation of the coding sequence. For example, a coding sequence is operably-linked to a promoter when the promoter is capable of directing transcription of that coding sequence. A flanking sequence need not be contiguous with the coding sequence, so long as it functions correctly. Thus, for example, intervening untranslated yet transcribed sequences can be present between a promoter sequence and the coding sequence, and the promoter sequence can still be considered “operably-linked” to the coding sequence. Each nucleotide sequence coding for a siRNA will typically have its own operably-linked promoter sequence.

The term “vector” is used to refer to any molecule (e.g., nucleic acid, plasmid, or virus) used to transfer coding information (e.g., a polynucleotide of the invention) to a host cell. The terms “expression vector” and “transcription vector” are used interchangeably to refer to a vector that is suitable for use in a host cell (e.g., a subject’s cell) and contains nucleic acid sequences that direct and/or control the expression of exogenous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

As used herein, the term “RNA interference” (“RNAi”) refers to a selective intracellular degradation of RNA. RNAi occurs in cells naturally to remove foreign RNAs (e.g., viral RNAs). Natural RNAi proceeds via fragments cleaved from free dsRNA which direct the degradative mechanism to other similar RNA sequences. Alternatively, RNAi can be initiated by the hand of man, for example, to silence the expression of endogenous target genes, such as SHIP.

As used herein, the term “small interfering RNA” (“siRNA”) (also referred to in the art as “short interfering RNAs”) refers to an RNA (or RNA analog) comprising between about 10-50 nucleotides (or nucleotide analogs) which is capable of directing or mediating RNA interference.

As used herein, a siRNA having a “sequence sufficiently complementary to a target mRNA sequence to direct target-specific RNA interference (RNAi)” means that the siRNA has a sequence sufficient to trigger the destruction of the target mRNA (e.g., SHIP mRNA) by the RNAi machinery or process. “mRNA” or “messenger RNA” or “transcript” is single-stranded RNA that specifies the amino acid sequence of one or more polypeptides. This information is translated during protein synthesis when ribosomes bind to the mRNA.

As used herein, the term “cleavage site” refers to the residues, e.g., nucleotides, at which RISC\* cleaves the target RNA, e.g., near the center of the complementary portion of the target RNA, e.g., about 8-12 nucleotides from the 5' end of the complementary portion of the target RNA.

11

The term “dominant negative mutant” is art-recognized and refers to the mutant form of a wild-type protein that interferes with the function of the wild-type protein (e.g., by interacting with the wild-type protein). Thus, overexpression of the dominant negative mutant can be expected to interfere with the function of the wild-type version of the protein.

As used herein, the term “mismatch” refers to a basepair consisting of noncomplementary bases, e.g., not normal complementary G:C, A:T or A:U base pairs.

As used herein, the term “isolated” molecule (e.g., isolated nucleic acid molecule) refers to molecules which are substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

As used herein, the term “in vitro” has its art recognized meaning, e.g., involving purified reagents or extracts, e.g., cell extracts. The term “in vivo” also has its art recognized meaning, e.g., involving living cells in an organism, e.g., immortalized cells, primary cells, and/or cell lines, in an organism.

A gene “involved in” or “associated with” a disorder includes a gene, the normal or aberrant expression or function of which affects or causes a disease or disorder or at least one symptom of the disease or disorder. The methods of the invention are useful in decreasing expression of SHIP in megakaryocytes and their progenitors in vitro or in vivo, consequently increasing their proliferation. Thus, the methods of the invention are useful in the treatment of human or non-human animal subjects suffering from, or at risk of developing, disorders associated with impaired megakaryocyte production.

The methods of the invention may include further steps. In some embodiments, a subject with the relevant condition or disease (e.g., disorders associated with impaired megakaryocyte production) is identified or a patient at risk for the condition or disease is identified prior to administration of the SHIP inhibitor. A patient may be someone who has not been diagnosed with the disease or condition (diagnosis, prognosis, and/or staging) or someone diagnosed with the disease or condition (diagnosis, prognosis, monitoring, and/or staging), including someone treated for the disease or condition (prognosis, staging, and/or monitoring). Alternatively, the person may not have been diagnosed with the disease or condition but suspected of having the disease or condition based either on patient history or family history, or the exhibition or observation of characteristic symptoms.

As used herein, an “effective amount” of a SHIP inhibitor (such as an interfering RNA, an antisense oligonucleotide, or a ribozyme, which selectively interferes with expression of SHIP) is that amount effective to bring about the physiological changes desired in the cells to which the SHIP inhibitor is administered in vitro (e.g., ex vivo) or in vivo. The term “therapeutically effective amount” as used herein, means that amount of SHIP inhibitor alone or in combination with another agent according to the particular aspect of the invention, that elicits the biological or medicinal response in cells (e.g., tissue(s)) that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation and/or prevention of the symptoms of the disease or disorder being treated. Preferably, suppression of SHIP function (e.g., by reduction of SHIP expression) results in increased megakaryocytopoiesis.

Various methods of the present invention can include a step that involves comparing a value, level, feature, characteristic, property, etc. to a “suitable control”, referred to interchangeably herein as an “appropriate control”. A “suitable control”

12

or “appropriate control” is any control or standard familiar to one of ordinary skill in the art useful for comparison purposes. In one embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc. determined prior to performing an RNAi methodology, as described herein. For example, a transcription rate, mRNA level, translation rate, protein level, biological activity, cellular characteristic or property, genotype, phenotype, etc. can be determined prior to introducing a siRNA of the invention into a cell or organism. In another embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc. determined in a cell or organism, e.g., a control or normal cell or organism, exhibiting, for example, normal traits. In yet another embodiment, a “suitable control” or “appropriate control” is a predefined value, level, feature, characteristic, property, etc.

Reduction (suppression) of expression results in a decrease of SHIP mRNA and/or protein. For example, in a given cell, the suppression of SHIP mRNA by administration of a SHIP inhibitor that reduces SHIP function by reducing SHIP expression (such as interfering RNA, antisense oligonucleotide, or ribozyme) results in a decrease in the quantity of SHIP mRNA relative to an untreated cell. Suppression may be partial. Preferred degrees of suppression are at least 50%, more preferably one of at least 60%, 70%, 80%, 85%, or 90%. A level of suppression between 90% and 100% is generally considered a “silencing” of expression. Where an increase in megakaryocytopoiesis is desired, the level of suppression is sufficient to increase megakaryocyte or megakaryocyte progenitor production.

SHIP gene expression can be determined before and/or after introduction of the SHIP inhibitor in vitro or in vivo. Reduction in SHIP gene expression can be detected at either the protein or mRNA level. Protein expression analysis can be performed by Western blotting, immunofluorescence, or flow cytometry and cell sorting (FACS). Reduction in SHIP gene expression can be detected at the mRNA level by real-time RT-PCR, microarray analysis, or Northern blotting, for example. Preferably, all expression data is compared with levels of a “house keeping” gene to normalize for variable amounts of RNA in different samples.

#### RNA Interference

RNAi is an efficient process whereby double-stranded RNA (dsRNA, also referred to herein as siRNAs or ds siRNAs, for double-stranded small interfering RNAs) induces the sequence-specific degradation of targeted mRNA in animal and plant cells (Hutvagner and Zamore, *Curr. Opin. Genet. Dev.*: 12, 225-232 (2002); Sharp, *Genes Dev.*, 15:485-490 (2001)). In mammalian cells, RNAi can be triggered by 21-nucleotide (nt) duplexes of small interfering RNA (siRNA) (Chiu et al., *Mol. Cell.* 10:549-561 (2002); Elbashir et al., *Nature* 411:494-498 (2001)), or by micro-RNAs (miRNA), functional small-hairpin RNA (shRNA), or other dsRNAs which can be expressed in vivo using DNA templates with RNA polymerase III promoters (Zeng et al., *Mol. Cell* 9:1327-1333 (2002); Paddison et al., *Genes Dev.* 16:948-958 (2002); Lee et al., *Nature Biotechnol.* 20:500-505 (2002); Paul et al., *Nature Biotechnol.* 20:505-508 (2002); Tuschl, T., *Nature Biotechnol.* 20:440-448 (2002); Yu et al., *Proc. Natl. Acad. Sci. USA* 99(9):6047-6052 (2002); McManus et al., *RNA* 8:842-850 (2002); Sui et al., *Proc. Natl. Acad. Sci. USA* 99(6):5515-5520 (2002)), each of which are incorporated herein by reference in their entirety.

The scientific literature contains many reports of endogenous and exogenous gene expression silencing using siRNA, highlighting their therapeutic potential (Gupta, S. et al. *PNAS*, 2004, 101:1927-1932; Takaku, H. *Antivir Chem.*



*Chemother.*, 2004, 15:57-65; Pardridge, W. M. *Expert Opin. Biol. Ther.*, 2004, 4:1103-1113; Zheng, B. J. *Antivir. Ther.*, 2004, 9:365-374; Shen, W. G. *Chin. Med. J. (Engl)*, 2004, 117:1084-1091; Fuchs, U. et al. *Curr. Mol. Med.*, 2004, 4:507-517; Wadhwa, R. et al. *Mutat. Res.*, 2004, 567:71-84; Ichim, T. E. et al. *Am. J. Transplant*, 2004, 4:1227-1236; Jana, S. et al. *Appl. Microbiol. Biotechnol.*, 2004, 65:649-657; Ryther, R. C. et al. *Gene Ther.*, 2005, 12:5-11; Chae, S-S. et al., *J. Clin. Invest.*, 2004, 114:1082-1089; Fougerolles, A. et al., *Methods Enzymol.*, 2005, 392:278-296), each of which is incorporated herein by reference in its entirety. Therapeutic silencing of endogenous genes by systemic administration of siRNAs has been described in the literature (Kim B. et al., *American Journal of Pathology*, 2004, 165:2177-2185; Soutschek J. et al., *Nature*, 2004, 432:173-178; Pardridge W. M., *Expert Opin. Biol. Ther.*, 2004, July, 4(7):1103-1113), each of which is incorporated herein by reference in its entirety.

Accordingly, the invention includes such interfering RNA molecules that are targeted to SHIP mRNA. The interfering RNA molecules are capable, when suitably introduced into or expressed within a cell that otherwise expresses SHIP mRNA, of suppressing expression of the SHIP gene by RNAi. The interfering RNA may be a double-stranded siRNA. As the skilled person will appreciate, and as explained further herein, an siRNA molecule may include a short 3' DNA sequence also. Alternatively, the nucleic acid may be a DNA (usually double-stranded DNA) which, when transcribed in a cell, yields an RNA having two complementary portions joined via a spacer, such that the RNA takes the form of a hairpin when the complementary portions hybridize with each other. In a mammalian cell, the hairpin structure may be cleaved from the molecule by the enzyme Dicer, to yield two distinct, but hybridized, RNA molecules.

In one embodiment, the invention provides an interfering RNA that is capable, when suitably introduced or expressed within a cell that normally expresses SHIP mRNA, suppresses its expression by RNAi, wherein the interfering RNA is generally targeted to the SHIP enzymatic domain (inositol 5'-phosphatase domain), within the human SHIP cDNA (SEQ ID NO:1). Examples of SHIP target sequences include GCCTGTTGTCATCCATTGA (SEQ ID NO:3), ATAAGT-TGGTGATCTTGGT (SEQ ID NO:4), GCCACATCTG-TACTGACAA (SEQ ID NO:5), AGACAGGCATTGCAAA-CAC (SEQ ID NO:6), ACATCACTCACCCTTCAC (SEQ ID NO:7), TCTTAACCTACCGTGTGGAT (SEQ ID NO:8), AATACGCCTACACCAAGCA (SEQ ID NO:9), GTAC-CAGCGACATCATGAC (SEQ ID NO:10), GCGACATCAT-GACGAGTGA (SEQ ID NO:11), AGGACAGAT-TGAGTTTCTC (SEQ ID NO:12), GGTGCTATGCCACATTGAA (SEQ ID NO:13), GTTTG-TGAGACTCTTCCA (SEQ ID NO:14), AGACG-GAGCGTGTGAATC (SEQ ID NO:15), GCTTCCAGAA-GAGCATCTTAT (SEQ ID NO:20), and GCCCATATCACCCAAGAAGTTT (SEQ ID NO:21). In a specific embodiment, the interfering RNA comprises a sequence selected from the group consisting of SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, and SEQ ID NO:19. Preferably, the interfering RNA sequence is within the range of about 19 to 23 nucleotides. For example, in those embodiments in which an shRNA is utilized, that portion of the shRNA targeting SHIP is preferably within the range of about 19 to 23 nucleotides.

It is expected that perfect identity/complementarity between the interfering RNA used in the method of the invention and the target sequence, although preferred, is not essential. Accordingly, the interfering RNA may include a single mismatch compared to the target sequence within the SHIP

mRNA. It is expected, however, that the presence of even a single mismatch is likely to lead to reduced efficiency, so the absence of mismatches is preferred. When present, 3' overhangs may be excluded from the consideration of the number of mismatches.

The term "complementarity" is not limited to conventional base pairing between nucleic acid consisting of naturally occurring ribo- and/or deoxyribonucleotides, but also includes base pairing between mRNA and nucleic acids of the invention that include non-natural nucleotides.

#### siRNA Molecules

Short interfering RNAs (siRNAs) induce the sequence-specific suppression or silencing (i.e., reducing expression which may be to the extent of partial or complete inhibition) genes by the process of RNAi. Thus, siRNA is the intermediate effector molecule of the RNAi process. The interfering RNA that function as SHIP inhibitors include dsRNA molecules comprising 16-30, e.g., 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in each strand, wherein one of the strands is substantially identical, e.g., at least 80% (or more, e.g., 85%, 90%, 95%, or 100%) identical, e.g., having 3, 2, 1, or 0 mismatched nucleotide(s), to a target region in the SHIP mRNA, and the other strand is identical or substantially identical to the first strand. The dsRNA molecules that function as SHIP inhibitors can be chemically synthesized, or can be transcribed in vitro from a DNA template, or in vivo from, e.g., shRNA. The dsRNA molecules can be designed using any method known in the art, for instance, by using the following protocol:

1. Using any method known in the art, compare the potential targets to the appropriate genome database (human, mouse, rat, etc.) and eliminate from consideration any target sequences with significant homology to other coding sequences. One such method for sequence homology searches is known as BLAST, which is available at the National Center for Biotechnology Information (NCBI) web site of the National Institutes of Health. Also available on the NCBI web site is the HomoloGene database, which is a publicly available system for automated detection of homologs among the annotated genes of several completely sequenced eukaryotic genomes and is readily utilized by those of ordinary skill in the art.

2. Select one or more sequences that meet your criteria for evaluation. Further general information regarding the design and use of siRNA can be found in "The siRNA User Guide," available at the web site of the laboratory of Dr. Thomas Tuschl at Rockefeller University (Elbashir et al., *EMBO J.*, 2001, 20:6877-6888).

3. Negative control siRNAs preferably have the same nucleotide composition as the selected siRNA, but without significant sequence complementarity to the appropriate genome. Such negative controls can be designed by randomly scrambling the nucleotide sequence of the selected siRNA; a homology search can be performed to ensure that the negative control lacks homology to any other gene in the appropriate genome. In addition, negative control siRNAs can be designed by introducing one or more base mismatches into the sequence.

Initially, basic criteria were defined for identification of efficient siRNA, such as GC content and position of the targeted sequence in the context of the mRNA (Elbashir S M. et al., *Methods*, 2002, 26:199-213). Further progress was achieved more recently, when the assembly of the RNAi enzyme complex was described as being dependent on thermodynamic characteristics of the siRNA (Khvorovova A. et al., *Cell*, 2003, 115:209-216; Schwarz D. S. et al., *Cell*, 2003, 115:199-208). The relative stability of both ends of the duplex

was determined to have effects on the extent to which the individual strands enter the RNAi pathway. In addition, certain sequence motifs at defined positions of the siRNA were reported to influence its potency (Amarzguoui M. and H. Prydz, *Biochem. Biophys. Res. Commun.*, 2004, 316:1050-1058; Reynolds A. et al., *Nature Biotechnol.*, 2004, 22:326-330). On this basis, sophisticated algorithms have been developed to increase the success rate of siRNA design and are available to those skilled in the art (Amarzguoui M. and H. Prydz, 2004; Reynolds A. et al., 2004; and Ui-Tei K. et al., *Nucl. Acids Res.*, 2004, 32:936-948, each of which is incorporated herein in its entirety).

Other computational tools that may be used to select siRNAs of the present invention include the Whitehead siRNA selection Web Server from the bioinformatics group at the Whitehead Institute for Biomedical Research in Cambridge, Mass., and other disclosed in Yuan, B. et al. ("siRNA Selection Server: an automated siRNA oligonucleotide prediction server", *Nucleic Acids Research*, 2004, Vol. 32, W130-W134, Web Server issue) and Bonetta L. ("RNAi: Silencing never sounded better", *Nature Methods*, October, 2004, 1(1):79-86), each of which are incorporated by reference herein in their entirety.

The efficiencies of different siRNAs may differ significantly. However, strategies for rational design of effective interfering RNA exist (Gong D. and J. E. Ferrell Jr., *TRENDS in Biotechnology*, 2004, 22(9):451; Schubert S. et al., *J. Mol. Biol.*, 2005, 348:883-893; Pancoska P. et al., *Nucleic Acids Research*, 2004, 32(4):1469-1479; Mittal V., *Nat. Rev. Genet.*, 2004, 5(5):355-365, each of which is incorporated herein by reference in its entirety).

Screening for the most efficient siRNAs using cell cultures may be carried out. Several in vitro screening methods based on the use of siRNA mixtures, which may contain a particular efficient siRNA (or several), have been developed. These include the preparation of siRNA mixtures using RNase III or Dicer enzymes to digest longer double-stranded RNAs, such as BLOCK-IT products (INVITROGEN, Carlsbad Calif.) (Yang D. et al., *Proc. Natl. Acad. Sci. USA*, 2002, 99:9942-9947; Myers J. W. et al., *Nat. Biotechnol.*, 2003, 21:324-328). The short RNAs produced as a result of these digestions have been found to be efficient in RNAi. Oligonucleotide arrays can also be used for the efficient preparation of defined mixtures of siRNAs for reducing the expression of exogenous and endogenous genes such as SHIP (Oleinikov A. V. et al., *Nucleic Acids Research*, 2005, 33(10):e92).

The SHIP inhibitors of the invention can include both unmodified siRNAs and modified siRNAs as known in the art. Thus, the invention includes siRNA derivatives that include siRNA having two complementary strands of nucleic acid, such that the two strands are crosslinked. For example, a 3' OH terminus of one of the strands can be modified, or the two strands can be crosslinked and modified at the 3' OH terminus. The siRNA derivative can contain a single crosslink (e.g., a psoralen crosslink). In some embodiments, the siRNA derivative has at its 3' terminus a biotin molecule (e.g., a photocleavable biotin), a peptide (e.g., a Tat peptide), a nanoparticle, a peptidomimetic, organic compounds (e.g., a dye such as a fluorescent dye), or dendrimer. Modifying siRNA derivatives in this way can improve cellular uptake or enhance cellular targeting activities of the resulting siRNA derivative as compared to the corresponding siRNA, are useful for tracing the siRNA derivative in the cell, or improve the stability of the siRNA derivative compared to the corresponding siRNA.

The SHIP inhibitors of the invention can be unconjugated or can be conjugated to another moiety, such as a nanoparticle, to enhance a property of the compositions, e.g., a phar-

macokinetic parameter such as absorption, efficacy, bioavailability, and/or half-life. The conjugation can be accomplished by methods known in the art, e.g., using the methods of Lambert et al., *Drug Deliv. Rev.* 47(1): 99-112 (2001) (describes nucleic acids loaded to polyalkylcyanoacrylate (PACA) nanoparticles); Fattal et al., *J. Control Release* 53(1-3):137-43 (1998) (describes nucleic acids bound to nanoparticles); Schwab et al., *Ann. Oncol.* 5 Suppl. 4:55-8 (1994) (describes nucleic acids linked to intercalating agents, hydrophobic groups, polycations or PACA nanoparticles); and Godard et al., *Eur. J. Biochem.* 232(2):404-10 (1995) (describes nucleic acids linked to nanoparticles).

The SHIP inhibitors of the present invention can also be labeled using any method known in the art; for instance, nucleic acids can be labeled with a fluorophore, e.g., Cy3, fluorescein, or rhodamine. The labeling can be carried out using a kit, e.g., the SILENCER siRNA labeling kit (AM-BION). Additionally, the siRNA can be radiolabeled, e.g., using  $^3\text{H}$ ,  $^{32}\text{P}$ , or other appropriate isotope.

Because RNAi is believed to progress via at least one single stranded RNA intermediate, the skilled artisan will appreciate that ss-siRNAs (e.g., the antisense strand of a ds-siRNA) can also be designed as described herein and utilized according to the claimed methodologies.

There are a number of companies that will generate interfering RNAs for a specific gene. Thermo Electron Corporation (Waltham, Mass.) has launched a custom synthesis service for synthetic short interfering RNA (siRNA). Each strand is composed of 18-20 RNA bases and two DNA bases overhang on the 3' terminus. Dharmacon, Inc. (Lafayette, Colo.) provides siRNA duplexes using the 2'-ACE RNA synthesis technology. Qiagen (Valencia, Calif.) uses TOM-chemistry to offer siRNA with high individual coupling yields (Li, B. et al., *Nat. Med.*, 2005, 11(9), 944-951).

siRNA Delivery for Longer-Term Expression

Synthetic siRNAs can be delivered into cells by methods known in the art, including cationic liposome transfection (LIPOFECTAMINE 2000 reagent, for example) and electroporation, for example. However, these exogenous siRNA generally show short term persistence of the silencing effect (4 to 5 days in cultured cells), which may be beneficial in certain embodiments. To obtain longer suppression of SHIP expression and to facilitate delivery under certain circumstances, one or more siRNA duplexes, e.g., SHIP ds siRNA, can be expressed within cells from recombinant DNA constructs (McIntyre G. J. and G. C. Fanning, *BMC Biotechnology*, 2006, 6:1-8). Such methods for expressing siRNA duplexes within cells from recombinant DNA constructs to allow longer-term target gene suppression in cells are known in the art, including mammalian Pol III promoter systems (e.g., H1 or U6/snRNA promoter systems (Tuschl (2002), supra) capable of expressing functional double-stranded siRNAs; Bagella et al., *J. Cell. Physiol.* 177:206-213 (1998); Lee et al. (2002), supra; Miyagishi et al. (2002), supra; Paul et al. (2002), supra; Yu et al. (2002), supra; Sui et al. (2002), supra). Transcriptional termination by RNA Pol III occurs at runs of four consecutive T residues in the DNA template, providing a mechanism to end the siRNA transcript at a specific sequence. The siRNA is complementary to the sequence of the target gene in 5'-3' and 3'-5' orientations, and the two strands of the siRNA can be expressed in the same construct or in separate constructs. Hairpin siRNAs, driven by an H1 or U6 snRNA promoter can be expressed in cells, and can inhibit target gene expression (Bagella et al. (1998), supra; Lee et al. (2002), supra; Miyagishi et al. (2002), supra; Paul et al. (2002), supra; Yu et al. (2002), supra; Sui et al. (2002) supra). Constructs containing siRNA sequence(s)

under the control of a T7 promoter also make functional siRNAs when co-transfected into the cells with a vector expressing T7 RNA polymerase (Jacque (2002), supra). A single construct may contain multiple sequences coding for siRNAs, such as multiple regions of SHIP mRNA, and can be driven, for example, by separate PolIII promoter sites.

Animal cells express a range of noncoding RNAs of approximately 22 nucleotides termed micro RNA (miRNAs) which can regulate gene expression at the post transcriptional or translational level during animal development. One common feature of miRNAs is that they are all excised from an approximately 70 nucleotide precursor RNA stem-loop, probably by Dicer, an RNase III-type enzyme, or a homolog thereof. By substituting the stem sequences of the miRNA precursor with miRNA sequence complementary to the target mRNA, a vector construct that expresses the novel miRNA can be used to produce siRNAs to initiate RNAi against specific mRNA targets in mammalian cells (Zeng (2002), supra). When expressed by DNA vectors containing polymerase III promoters, micro-RNA designed hairpins can silence gene expression (McManus (2002), supra). Viral-mediated delivery mechanisms can also be used to induce specific silencing of targeted genes through expression of siRNA, for example, by generating recombinant adenoviruses harboring siRNA under RNA Pol II promoter transcription control (Xia et al. (2002), supra). Infection of HeLa cells by these recombinant adenoviruses allows for diminished endogenous target gene expression. Injection of the recombinant adenovirus vectors into transgenic mice expressing the target genes of the siRNA results in in vivo reduction of target gene expression. In an animal model, whole-embryo electroporation can efficiently deliver synthetic siRNA into post-implantation mouse embryos (Calegari et al., *Proc. Natl. Acad. Sci. USA* 99(22):14236-40 (2002)). In adult mice, efficient delivery of siRNA can be accomplished by the "high-pressure" delivery technique, a rapid injection (within 5 seconds) of a large volume of siRNA containing solution into animal via the tail vein (Liu (1999), supra; McCaffrey (2002), supra; Lewis, *Nature Genetics* 32:107-108 (2002)). Nanoparticles, liposomes and other cationic lipid molecules can also be used to deliver siRNA into animals. It has been shown that siRNAs delivered systemically in a liposomal formulation can silence the disease target apolipoprotein B (ApoB) in non-human primates (Zimmermann T. S. et al., *Nature*, 2006, 441: 111-114). A gel-based agarose/liposome/siRNA formulation is also available (Jiang M. et al., *Oligonucleotides*, 2004, Winter, 14(4):239-48).

#### Uses of Engineered RNA Precursors to Induce RNAi

Engineered RNA precursors, introduced into cells or whole organisms as described herein, will lead to the production of a desired siRNA molecule. Such an siRNA molecule will then associate with endogenous protein components of the RNAi pathway to bind to and target a specific mRNA sequence for cleavage and destruction. In this fashion, the SHIP mRNA to be targeted by the siRNA generated from the engineered RNA precursor will be depleted from the cell or organism, leading to a decrease in the concentration of any translational product encoded by that mRNA in the cell or organism. The RNA precursors are typically nucleic acid molecules that individually encode either one strand of a dsRNA or encode the entire nucleotide sequence of an RNA hairpin loop structure.

#### Antisense

An "antisense" nucleic acid sequence (antisense oligonucleotide) can include a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded

cDNA molecule or complementary to the SHIP mRNA. Antisense nucleic acid sequences and delivery methods are well known in the art (Goodchild J., *Curr. Opin. Mol. Ther.*, 2004, April, 6(2):120-128; Clawson G. A. et al., *Gene Ther.*, 2004, September, 11(17):1331-1341), which are incorporated herein by reference in their entirety. The antisense nucleic acid can be complementary to an entire coding strand of a target sequence, or to only a portion thereof. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence within the SHIP mRNA. An antisense oligonucleotide can be, for example, about 7, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, or more nucleotides in length.

An antisense nucleic acid sequence can be designed such that it is complementary to the entire SHIP mRNA sequence, but can also be an oligonucleotide that is antisense to only a portion of the SHIP mRNA. For example, the antisense oligonucleotide can be complementary to a portion of the SHIP enzymatic domain (inositol 5'-phosphatase domain) or a portion of the amino-terminal src-homology domain (SH2).

An antisense nucleic acid of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. The antisense nucleic acid also can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject (e.g., systemically or locally by direct injection at a tissue site), or generated in situ such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding the SHIP to thereby inhibit expression of the SHIP gene. Alternatively, antisense nucleic acid molecules can be modified to target selected cells (such as megakaryocytes and/or megakaryocyte progenitors) and then administered systemically. For systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens (such as CD9, CD41, CD61, actin, or FVIIIRAg) expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter can be used.

In yet another embodiment, the antisense oligonucleotide of the invention is an alpha-anomeric nucleic acid molecule. An alpha-anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual beta-units, the strands run parallel to each other (Gaultier et al., *Nucleic Acids. Res.* 15:6625-6641 (1987)). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue et al. *Nucleic Acids Res.* 15:6131-6148 (1987)) or a chimeric RNA-DNA analogue (Inoue et al. *FEBS Lett.*, 215:327-330 (1987)).

SHIP expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the SHIP gene to form triple helical structures that prevent expression of SHIP in target cells. See generally, Helene, C. *Anticancer Drug Des.* 6:569-84 (1991); Helene, C. *Ann. N.Y. Acad. Sci.* 660:27-36 (1992); and Maher, *Bioassays* 14:807-15 (1992). The potential sequences that can be targeted for triple helix formation can be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3',3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex. Ribozymes

Ribozymes are a type of RNA that can be engineered to enzymatically cleave and inactivate other RNA targets in a specific, sequence-dependent fashion. Ribozymes and methods for their delivery are well known in the art (Hendry P. et al., *BMC Chem. Biol.*, 2004, Dec., 4(1):1; Grassi G. et al., *Curr. Pharm. Biotechnol.*, 2004, Aug., 5(4):369-386; Bagheri S. et al., *Curr. Mol. Med.*, 2004, Aug., 4(5):489-506; Kashani-Sabet M., *Expert Opin. Biol. Ther.*, 2004, Nov., 4(11):1749-1755), each of which are incorporated herein by reference in its entirety. By cleaving the target RNA, ribozymes inhibit translation, thus preventing the expression of the target gene. Ribozymes can be chemically synthesized in the laboratory and structurally modified to increase their stability and catalytic activity using methods known in the art. Alternatively, ribozyme genes can be introduced into cells through gene-delivery mechanisms known in the art. A ribozyme having specificity for SHIP mRNA can include one or more sequences complementary to a nucleotide sequence within the SHIP mRNA, and a sequence having a known catalytic sequence responsible for mRNA cleavage (see U.S. Pat. No. 5,093,246 or Haselhoff and Gerlach *Nature* 334:585-591 (1988)). For example, a derivative of a Tetrahymena L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in the mRNA encoded by a uORF of an extended, overlapping 5'-UTR AS mRNA species (see, e.g., Cech et al. U.S. Pat. No. 4,987,071; and Cech et al. U.S. Pat. No. 5,116,742). Alternatively, SHIP mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules (see, e.g., Bartel, D. and Szostak, J. W. *Science* 261:1411-1418 (1993)). Nucleic Acid Targets

The nucleic acid targets of SHIP inhibitors that are polynucleotides (referred to herein as "polynucleotide SHIP inhibitors" or "nucleic acid SHIP inhibitors") such as the antisense, interfering RNA molecules, and ribozymes described herein, may be any portion of the SHIP gene or SHIP mRNA. In some embodiments, the nucleic acid target is the SHIP enzymatic domain (inositol 5'-phosphatase) or the amino-terminal src-homology domain (SH2). In other embodiments, the nucleic acid target is a translation initiation site, 3' untranslated region, or 5' untranslated region

As indicated above, the nucleotide sequences of mouse SHIP and human SHIP have been publicly available for several years (GenBank Accession Numbers NM\_10566 and NM\_005541, respectively, on the NCBI database). Pair-wise alignment scoring of orthologues shows high levels of homology among SHIP sequences of humans, mice, and rats. Each sequence has the SHIP enzymatic domain (inositol 5'-phosphatase domain), and the degree of nucleotide homology between human, mouse, and rat is over 85%. Furthermore, mice and humans are believed to have the same five SHIP protein isoforms. In a preferred embodiment, the polynucle-

otide SHIP inhibitor (e.g., interfering RNA, antisense oligonucleotide, ribozyme) targets an mRNA sequence shared by all known hematopoietic SHIP isoforms in humans. Such target sequence can be readily determined by those skilled in the art due to the extensive amount of sequence overlap between the isoforms.

The target SHIP sequence can be within any orthologue of SHIP, such as sequences substantially identical to human, mouse, rat, or bovine, or a portion of any of the foregoing, including but not limited to GenBank Accession Numbers NM\_005541 and NM\_001017915 (human), NM\_10566 (mouse), and U55192 (rat).

Table 1 lists thirteen target sequences predicted to have good specificity and knockdown potential against the human SHIP cDNA sequence.

TABLE 1

Seq. Identifier	Target Sequence	Region (nt)	GC Content
SEQ ID NO: 3	GCCTGTTGTCATCCATTGA	ORF	890 47.37%
SEQ ID NO: 4	ATAAGTTGGTGATCTTGGT	ORF	1145 36.84%
SEQ ID NO: 5	GCCACATCTGTACTGACAA	ORF	1589 47.37%
SEQ ID NO: 6	AGACAGGCATTGCAACAC	ORF	1613 47.37%
SEQ ID NO: 7	ACATCACTCACCGCTTCAC	ORF	1802 52.63%
SEQ ID NO: 8	TCTTAACACCGTGTGGAT	ORF	1842 42.11%
SEQ ID NO: 9	AATACGCCTACACCAAGCA	ORF	2039 47.37%
SEQ ID NO: 10	GTACCAGCGACATCATGAC	ORF	2156 52.63%
SEQ ID NO: 11	GCGACATCATGACGAGTGA	ORF	2162 52.63%
SEQ ID NO: 12	AGGACAGATTGAGTTTCTC	ORF	2265 42.11%
SEQ ID NO: 13	GGTGCTATGCCACATTGAA	ORF	2285 47.37%
SEQ ID NO: 14	GTTTGGTGAGACTCTTCCA	ORF	2418 47.37%
SEQ ID NO: 15	AGACGGAGCGTGATGAATC	ORF	2687 52.63%

The term "orthologue" as used herein refers to a sequence which is substantially identical to a reference sequence. The term "substantially identical" is used herein to refer to a first amino acid or nucleotide sequence that contains a sufficient or minimum number of identical or equivalent (e.g., with a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences have a common structural domain or common functional activity. For example, amino acid or nucleotide sequences that contain a common structural domain having at least about 60%, or 65% identity, likely 75% identity, more likely 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity are defined herein as substantially identical.

Calculations of homology or sequence identity between sequences (the terms are used interchangeably herein) are performed as follows.

To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In one embodiment, the length of a reference

sequence aligned for comparison purposes is at least 50%, at least 60%, at least 70%, 80%, 90%, or 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In one embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* 48:444-453 (1970)) algorithm, which has been incorporated into the GAP program in the GCG software package (available at the official Accelrys web site), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at the official Accelrys web site), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. One set of parameters (and the one that can be used if the practitioner is uncertain about what parameters should be applied to determine if a molecule is within a sequence identity or homology limitation of the invention) are a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

The nucleic acid and protein sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other orthologs, e.g., family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. *J. Mol. Biol.* 215:403-10 (1990). BLAST nucleotide searches can be performed with the NBLAST program, score=100, word length=12, to obtain nucleotide sequences homologous to known SHIP nucleic acid sequences. BLAST protein searches can be performed with the XBLAST program, score=50, word length=3, to obtain amino acid sequences homologous to known polypeptide products of the SHIP gene. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used (see the National Center for Biotechnology Information web site of the National Institutes of Health).

Orthologs can also be identified using any other routine method known in the art, such as screening a cDNA library, e.g., a human cDNA library, using a probe designed to identify sequences which are substantially identical to a reference sequence.

Pharmaceutical Compositions and Methods of Administration

The SHIP inhibitors of the subject invention (such as interfering RNA molecules, antisense molecules, and ribozymes) can be incorporated into pharmaceutical compositions. Such compositions typically include the SHIP inhibitor and a pharmaceutically acceptable carrier. As used herein, the term "pharmaceutically acceptable carrier" includes saline, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions. Formulations (compositions) are described in a number of sources that are well known and readily available to those skilled in the art. For example, *Remington's Pharmaceutical Sciences* (Martin E. W., Easton Pa., Mack Publishing Company, 19<sup>th</sup> ed., 1995) describes formulations which can be used in connection with the subject invention.

A pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), topical, transdermal, transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, CREMOPHOR EL (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. Isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride can also be included in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polynucleotide of the invention) in the required amount in an appropriate solvent with

one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, suitable methods of preparation include vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, e.g., gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, PRIMOGEL, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the SHIP inhibitors can be delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer. Such inhalation methods and inhalant formulations include those described in U.S. Pat. No. 6,468,798.

Systemic administration of SHIP inhibitors can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compound (e.g., polynucleotides of the invention) are formulated into ointments, salves, gels, or creams, as generally known in the art.

The pharmaceutical compositions can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

The SHIP inhibitors can also be administered by transfection or infection using methods known in the art, including but not limited to the methods described in McCaffrey et al., *Nature* 418(6893):38-39 (2002) (hydrodynamic transfection); Xia et al., *Nature Biotechnol.* 20(10):1006-10 (2002) (viral-mediated delivery); or Putnam, *Am. J. Health Syst. Pharm.* 53(2):151-160 (1996), erratum at *Am. J. Health Syst. Pharm.* 53(3):325 (1996).

SHIP inhibitors that are polynucleotides can also be administered by any method suitable for administration of nucleic acid agents, such as a DNA vaccine. These methods include gene guns, bio injectors, and skin patches as well as needle-free methods such as the micro-particle DNA vaccine technology disclosed in U.S. Pat. No. 6,194,389, and the mammalian transdermal needle-free vaccination with powder-form vaccine as disclosed in U.S. Pat. No. 6,168,587. Additionally, intranasal delivery is possible, as described in Hamajima et al., *Clin. Immunol. Immunopathol.* 88(2):205-

10 (1998). Liposomes (e.g., as described in U.S. Pat. No. 6,472,375) and microencapsulation can also be used. Biodegradable targetable microparticle delivery systems can also be used (e.g., as described in U.S. Pat. No. 6,471,996).

In one embodiment, the polynucleotide SHIP inhibitors are prepared with carriers that will protect the polynucleotide against rapid elimination from, or degradation in, the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Such formulations can be prepared using standard techniques. Liposomal suspensions (including liposomes targeted to antigen-presenting cells with monoclonal antibodies) can also be used as pharmaceutically acceptable carriers. Examples of such antigens presented by megakaryocytes and their progenitors include CD9, CD41, CD61, actin, and FVIIIIRAg. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811. Strategies that inhibit members of the RNase A family of enzymes or can otherwise protect polynucleotide SHIP inhibitors from these enzymes may be utilized. For example, U.S. Pat. No. 6,096,720 (Love et al.) describes oligonucleotides targeted to human raf mRNA, which are entrapped in sterically stabilized liposomes. In one embodiment, the oligonucleotide in Love et al. is a chimeric oligonucleotide containing a first region to enhance target affinity and a second region that is a substrate for RNase. siSHIELD RNase inhibitor is designed to prevent degradation of siRNA by RNase (MP BIOMEDICALS, Irvine, Calif.). A strategy for the compaction of short oligonucleotides into well-defined condensates may also be used to deliver the polynucleotides of the subject invention (Sarkar T. et al., *Nucleic Acids Research*, 2005, 33(1):143-151), which is incorporated herein by reference in its entirety.

In particular, suitable techniques for cellular administration of the polynucleotide SHIP inhibitors, such as interfering RNA, in vitro or in vivo are disclosed in the following articles: General Reviews:

Borkhardt, A. *Cancer Cell*, 2002, 2:167-8; Hannon, G. J. *Nature*, 2002, 418:244-51; McManus, M. T. and Sharp, P. A. *Nat Rev Genet.*, 2002, 3:737-47; Scherr, M. et al. *Curr Med. Chem.*, 2003, 10:245-56; Shuey, D. J. et al. *Drug Discov Today*, 2002, 7:1040-6; Gilmore, I. R. et al., *J. Drug Target.*, 2004, 12(6):315-340; Dykxhoorn, D. M. and Lieberman J., *Annu. Rev. Med.*, 2005, 56:401-423.

Systemic Delivery Using Liposomes:

Lewis, D. L. et al. *Nat Genet.*, 2002, 32:107-8; Paul, C. P. et al. *Nat Biotechnol.*, 2002, 20:505-8; Song, E. et al. *Nat Med.*, 2003, 9:347-51; Sorensen, D. R. et al. *J Mol Biol.*, 2003, 327:761-6.

Virus Mediated Transfer:

Abbas-Terki, T. et al. *Hum Gene Ther.*, 2002, 13:2197-201; Barton, G. M. and Medzhitov, R. *Proc Natl Acad Sci USA*, 2002, 99:14943-5; Devroe, E. and Silver, P. A. *BMC Biotechnol.*, 2002, 2:15; Lori, F. et al. *Am J Pharmacogenomics*, 2002, 2:245-52; Matta, H. et al. *Cancer Biol Ther.*, 2003, 2:206-10; Qin, X. F. et al. *Proc Natl Acad Sci USA*, 2003, 100:183-8; Scherr, M. et al. *Cell Cycle*, 2003, 2:251-7; Shen, C. et al. *FEBS Lett.*, 2003, 539:111-4; Lee S. K. et al. *Blood*, 2005, 106(3):818-826, epub Apr. 14, 2005.

Peptide Delivery:

Morris, M. C. et al. *Curr Opin Biotechnol.*, 2000, 11:461-6; Simeoni, F. et al. *Nucleic Acids Res.*, 2003, 31:2717-24.

Song E. et al. describe antibody mediated in vivo delivery of siRNAs via cell-surface receptors (Song E. et al., *Nat. Biotechnol.*, 2005, 23(6):709-717, epub May 22, 2005). This

cell-specific delivery technique can be used to target interfering RNA molecules to the cell-surface receptors on megakaryocytes and megakaryocyte progenitors.

Other technologies that may be suitable for delivery of polynucleotide SHIP inhibitors, such as interfering RNA, to the target cells are based on nanoparticles or nanocapsules such as those described in U.S. Pat. Nos. 6,649,192B and 5,843,509B. Recent technologies that may be employed for selecting, delivering, and monitoring interfering RNA molecules include Raab, R. M. and Stephanopoulos, G. *Biotechnol. Bioeng.*, 2004, 88:121-132; Huppi, K. et al. *Mol. Cell*, 2005, 17:1-10; Spagnou, S. et al. *Biochemistry*, 2004, 43:13348-13356; Muratovska, A. and Eccles, M. R. *FEBS Lett.*, 2004, 558:63-68; Kumar, R. et al. *Genome Res.*, 2003, 13:2333-2340; Chen, A. A. et al. *Nucleic Acids Res.*, 2005, 33:e190; Dykxhoorn, D. M. et al. *Gene Ther.*, 2006, epub ahead of print; Rodriguez-Lebron, E. and Paulson, H. L. *Gene Ther.*, 2005, epub ahead of print; Pai, S. I. et al. *Gene Ther.*, 2005, epub ahead of print; Raoul, C. et al. *Gene Ther.*, 2005, epub ahead of print; Manfredsson, F. P. et al. *Gene Ther.*, 2005, epub ahead of print; Downward, J. *BMJ*, 2004, 328:1245-1248.

A mixture of SHIP inhibitors, of the same type or different types, may be introduced into cells in vitro or in vivo. For example, a mixture or pool of polynucleotide SHIP inhibitors such as interfering RNA molecules (e.g., 2-4 interfering molecules or more) can be introduced into cells (Oleinikov A. V. et al., *Nucleic Acids Research*, 2005, 33(10):e92). Preferably, the interfering RNA molecules target different regions of the SHIP mRNA. Preferably, the interfering RNA molecules have been previously validated as individually functioning to reduce SHIP expression. The individual interfering RNAs of the mixture can be chemically synthesized (Elbashir S. M. et al., *Genes Dev.*, 2001, 15:188-200) or introduced as short DNA templates containing RNA polymerase promoter, which are transcribed within the cells in vitro or in vivo (Yu J. Y. et al., *Proc. Natl. Acad. Sci. USA*, 99:6047-6052).

Toxicity and therapeutic efficacy of compositions can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compositions which exhibit high therapeutic indices can be used. While compositions that exhibit toxic side effects can be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

Data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compositions generally lies within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any composition used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the test composition which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

The SHIP inhibitor can be administered on any appropriate schedule, e.g., from one or more times per day to one or more times per week; including once every other day, for any number of days or weeks, e.g., 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 1 week, 10 days, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 2 months, 3 months, 6 months, or more, or any variation thereon. The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a SHIP inhibitor can include a single treatment or can include a series of treatments.

The polynucleotide SHIP inhibitors (e.g., interfering RNA, antisense oligonucleotide, or ribozyme) can be introduced (administered) into cells (such as mammalian cells) in vitro or in vivo using known techniques, as those described herein, to suppress gene expression. Similarly, genetic constructs (e.g., transcription vectors) containing DNA of the invention may be introduced into cells in vitro or in vivo using known techniques, as described herein, for transient or stable expression of RNA, to suppress gene expression. When administered to the cells in vivo, the polynucleotide SHIP inhibitors can be administered to a subject systemically (e.g., intravenously), for example, or administered locally at the site of the cells (such as the peripheral blood, bone marrow, or spleen).

The cells in which the polynucleotide SHIP inhibitors are introduced may be any cell, such as a megakaryocyte or megakaryocyte progenitor, containing SHIP mRNA. The cells can be primary cells, cultured cells, cells of cell lines, etc. In one embodiment, the cells are from bone marrow. In another embodiment, the cells are from the spleen. In another embodiment, the cells are from peripheral blood.

Mammalian species which benefit from the disclosed methods include, but are not limited to, primates, such as humans, apes, chimpanzees, orangutans, monkeys; domesticated animals (e.g., pets) such as dogs, cats, guinea pigs, hamsters, Vietnamese pot-bellied pigs, rabbits, and ferrets; domesticated farm animals such as cows, buffalo, bison, horses, donkey, swine, sheep, and goats; exotic animals typically found in zoos, such as bear, lions, tigers, panthers, elephants, hippopotamus, rhinoceros, giraffes, antelopes, sloth, gazelles, zebras, wildebeests, prairie dogs, koala bears, kangaroo, opossums, raccoons, pandas, hyena, seals, sea lions, elephant seals, otters, porpoises, dolphins, and whales. As used herein, the terms "subject", "patient", and "individual" are used interchangeably and intended to include such human and non-human mammalian species. Likewise, in vitro methods of the present invention can be carried out on cells of such mammalian species. Host cells comprising exogenous polynucleotides of the invention may be administered to the subject, and may, for example, be autogenic (use of one's own cells), allogenic (from one person to another), or transgenic or xenogenic (from one mammalian species to another mammalian species), relative to the subject.

The polynucleotide SHIP inhibitors of the invention can be inserted into genetic constructs, e.g., viral vectors, retroviral vectors, expression cassettes, or plasmid viral vectors, e.g., using methods known in the art, including but not limited to those described in Xia et al., (2002), supra. Genetic constructs can be delivered to a subject by, for example, inhalation, orally, intravenous injection, local administration (see U.S. Pat. No. 5,328,470) or by stereotactic injection (see, e.g., Chen et al., *Proc. Natl. Acad. Sci. USA* 91:3054-3057 (1994)). The pharmaceutical preparation of the delivery vector can



include the vector in an acceptable diluent, or can comprise a slow release matrix in which the delivery vehicle is imbedded. Alternatively, where the complete delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the polynucleotide delivery system.

The polynucleotide SHIP inhibitors can be small hairpin RNAs (shRNAs), and expression constructs engineered to express shRNAs. Transcription of shRNAs is initiated at a polymerase III (pol III) promoter, and is thought to be terminated at position 2 of a 4-5-thymine transcription termination site. Upon expression, shRNAs are thought to fold into a stem-loop structure with 3' UU-overhangs; subsequently, the ends of these shRNAs are processed, converting the shRNAs into siRNA-like molecules of about 21 nucleotides (Brummelkamp et al., *Science* 296:550-553 (2002); Lee et al., (2002), supra; Miyagishi and Taira, *Nature Biotechnol.* 20:497-500 (2002); Paddison et al. (2002), supra; Paul (2002), supra; Sui (2002) supra; Yu et al. (2002), supra.

siRNAs targeting SHIP mRNA may be fused to other nucleotide molecules, or to polypeptides, in order to direct their delivery or to accomplish other functions. Thus, for example, fusion proteins comprising a siRNA oligonucleotide that is capable of specifically interfering with expression of SHIP may comprise affinity tag polypeptide sequences, which refers to polypeptides or peptides that facilitate detection and isolation of the polypeptide via a specific affinity interaction with a ligand. The ligand may be any molecule, receptor, counter-receptor, antibody or the like with which the affinity tag may interact through a specific binding interaction as provided herein. Such peptides include, for example, poly-His or "FLAG" or the like, e.g., the antigenic identification peptides described in U.S. Pat. No. 5,011,912 and in Hopp et al., (*Bio/Technology* 6:1204, 1988), or the XPRESS epitope tag (INVITROGEN, Carlsbad, Calif.). The affinity sequence may be a hexa-histidine tag as supplied, for example, by a pBAD/His (INVITROGEN) or a pQE-9 vector to provide for purification of the mature polypeptide fused to the marker in the case of a bacterial host, or, for example, the affinity sequence may be a hemagglutinin (HA) tag when a mammalian host, e.g., COS-7 cells, is used. The HA tag corresponds to an antibody defined epitope derived from the influenza hemagglutinin protein (Wilson et al., 1984 *Cell* 37:767).

The present invention also relates to vectors and to constructs that include or encode polynucleotide SHIP inhibitors (e.g., siRNA), and in particular to "recombinant nucleic acid constructs" that include any nucleic acid such as a DNA polynucleotide segment that may be transcribed to yield SHIP-specific siRNA polynucleotides according to the invention as provided above; to host cells which are genetically engineered with vectors and/or constructs of the invention and to the production of siRNA polynucleotides, polypeptides, and/or fusion proteins of the invention, or fragments or variants thereof, by recombinant techniques. siRNA sequences disclosed herein as RNA polynucleotides may be engineered to produce corresponding DNA sequences using well-established methodologies such as those described herein. Thus, for example, a DNA polynucleotide may be generated from any siRNA sequence described herein, such that the present siRNA sequences will be recognized as also providing corresponding DNA polynucleotides (and their complements). These DNA polynucleotides are therefore encompassed within the contemplated invention, for example, to be incorporated into the subject invention recombinant nucleic acid constructs from which siRNA may be transcribed.

According to the present invention, a vector may comprise a recombinant nucleic acid construct containing one or more promoters for transcription of an RNA molecule, for example, the human U6 snRNA promoter (see, e.g., Miyagishi et al., *Nat. Biotechnol.* 20:497-500 (2002); Lee et al., *Nat. Biotechnol.* 20:500-505 (2002); Paul et al., *Nat. Biotechnol.* 20:505-508 (2002); Grabarek et al., *BioTechniques* 34:73544 (2003); see also Sui et al., *Proc. Natl. Acad. Sci. USA* 99:5515-20 (2002)). Each strand of a siRNA polynucleotide may be transcribed separately each under the direction of a separate promoter and then may hybridize within the cell to form the siRNA polynucleotide duplex. Each strand may also be transcribed from separate vectors (see Lee et al., supra). Alternatively, the sense and antisense sequences specific for a SHIP mRNA sequence may be transcribed under the control of a single promoter such that the siRNA polynucleotide forms a hairpin molecule (Paul et al., supra). In such instance, the complementary strands of the siRNA specific sequences are separated by a spacer that comprises at least four nucleotides, but may comprise at least 5, 6, 7, 8, 9, 10, 11, 12, 14, 16, or 18 or more nucleotides as described herein. In addition, siRNAs transcribed under the control of a U6 promoter that form a hairpin may have a stretch of about four uridines at the 3' end that act as the transcription termination signal (Miyagishi et al., supra; Paul et al., supra). By way of illustration, if the target sequence is 19 nucleotides, the siRNA hairpin polynucleotide (beginning at the 5' end) has a 19-nucleotide sense sequence followed by a spacer (which as two uridine nucleotides adjacent to the 3' end of the 19-nucleotide sense sequence), and the spacer is linked to a 19 nucleotide antisense sequence followed by a 4-uridine terminator sequence, which results in an overhang. siRNA polynucleotides with such overhangs effectively interfere with expression of the target polypeptide. A recombinant construct may also be prepared using another RNA polymerase III promoter, the H1 RNA promoter, that may be operatively linked to siRNA polynucleotide specific sequences, which may be used for transcription of hairpin structures comprising the siRNA specific sequences or separate transcription of each strand of a siRNA duplex polynucleotide (see, e.g., Brummelkamp et al., *Science* 296:550-53 (2002); Paddison et al., supra). DNA vectors useful for insertion of sequences for transcription of an siRNA polynucleotide include pSUPER vector (see, e.g., Brummelkamp et al., supra); pAV vectors derived from pCWRSVN (see, e.g., Paul et al., supra); and pIND (see, e.g., Lee et al., supra), or the like.

Polynucleotide SHIP inhibitors can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters, providing ready systems for evaluation of siRNA polynucleotides that are capable of interfering with SHIP expression as provided herein. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described, for example, by Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, Third Edition, Cold Spring Harbor, N.Y., (2001).

The appropriate DNA sequence(s) may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Standard techniques for cloning, DNA isolation, amplification and purification, for enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like, and various separation techniques are those known and commonly employed by those skilled in the art. A number of standard techniques are described, for example, in Ausubel et al. (1993 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., Boston, Mass.); Sambrook et



al. (2001 Molecular Cloning, Third Ed., Cold Spring Harbor Laboratory, Plainview, N.Y.); Maniatis et al. (1982 Molecular Cloning, Cold Spring Harbor Laboratory, Plainview, N.Y.); and elsewhere.

The DNA sequence in the expression vector is operatively linked to at least one appropriate expression control sequence (e.g., a promoter or a regulated promoter) to direct mRNA synthesis. Representative examples of such expression control sequences include LTR or SV40 promoter, the *E. coli* lac or trp, the phage lambda  $P_L$  promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Examples of Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of the appropriate vector and promoter is well within the level of ordinary skill in the art, and preparation of certain particularly preferred recombinant expression constructs comprising at least one promoter or regulated promoter operably linked to a polynucleotide of the invention is described herein.

As noted above, in certain embodiments the vector may be a viral vector such as a mammalian viral vector (e.g., retrovirus, adenovirus, adeno-associated virus, lentivirus). For example, retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The viral vector includes one or more promoters. Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CMV) promoter described in Miller, et al., *Biotechniques* 7:980-990 (1989), or any other promoter (e.g., cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and beta-actin promoters). Other viral promoters that may be employed include, but are not limited to, adenovirus promoters, adeno-associated virus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein, and may be from among either regulated promoters (e.g., tissue-specific or inducible promoters) or promoters as described above. A tissue-specific promoter allows preferential expression of the polynucleotide SHIP inhibitor in a given target tissue, thereby avoiding expression in other tissues. For example, to target genes specifically in the heart, a number of cardiac-specific regulatory elements can be used. An example of a cardiac-specific promoter is the ventricular form of MLC-2v promoter (see, Zhu et al., *Mol. Cell. Biol.* 13:4432-4444, 1993; Navankasattusas et al., *Mol. Cell Biol.* 12:1469-1479, 1992) or a variant thereof such as a 281 bp fragment of the native MLC-2v promoter (nucleotides -264 to +17, GenBank Accession No. U26708). Examples of other cardiac-specific promoters include alpha myosin heavy chain (Minamino et al., *Circ. Res.* 88:587-592, 2001) and myosin light chain-2 (Franz et al., *Circ. Res.* 73:629-638, 1993). Endothelial cell gene promoters include endoglin and ICAM-2. See Velasco et al., *Gene Ther.* 8:897-904, 2001. Liver-specific promoters include the human phenylalanine hydroxylase (PAH) gene promoters (Bristeau et al., *Gene* 274:283-291, 2001), hB1F (Zhang et al., *Gene* 273:239-249, 2001), and the human C-reactive protein (CRP) gene promoter (Ruther et al., *Oncogene*

8:87-93, 1993). Promoters that are kidney-specific include CLCN5 (Tanaka et al., *Genomics* 58:281-292, 1999), renin (Sinn et al., *Physical Genomics* 3:25-31, 2000), androgen-regulated protein, sodium-phosphate cotransporter, renal cytochrome P-450, parathyroid hormone receptor and kidney-specific cadherin. See *Am. J. Physiol. Renal Physiol.* 279:F383-392, 2000. An example of a pancreas-specific promoter is the pancreas duodenum homeobox 1 (PDX-1) promoter (Samara et al., *Mol. Cell Biol.* 22:4702-4713, 2002). A number of brain-specific promoters may be useful in the invention and include the thy-1 antigen and gamma-enolase promoters (Vibert et al., *Eur. J. Biochem.* 181:33-39, 1989), the glial-specific glial fibrillary acidic protein (GFAP) gene promoter (Cortez et al., *J. Neurosci. Res.* 59:39-46, 2000), and the human FGF1 gene promoter (Chiu et al., *Oncogene* 19:6229-6239, 2000). The GATA family of transcription factors have promoters directing neuronal and thymocyte-specific expression (see Asnagli et al., *J. Immunol.* 168:4268-4271, 2002).

In another aspect, the present invention relates to host cells containing the above described recombinant constructs. Host cells are genetically engineered/modified (transduced, transformed or transfected) with the vectors and/or expression constructs of this invention that may be, for example, a cloning vector, a shuttle vector, or an expression construct. The vector or construct may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying particular genes such as genes encoding siRNA polynucleotides or fusion proteins thereof. The culture conditions for particular host cells selected for expression, such as temperature, pH and the like, will be readily apparent to the ordinarily skilled artisan.

The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Representative examples of appropriate host cells according to the present invention include, but need not be limited to, bacterial cells, such as *E. coli*, *Streptomyces*, *Salmonella typhimurium*; fungal cells, such as yeast; insect cells, such as *Drosophila* S2 and *Spodoptera* Sf9; animal cells, such as CHO, COS or 293 cells; adenoviruses; plant cells, or any suitable cell already adapted to in vitro propagation or so established de novo.

Various mammalian cell culture systems can also be employed to produce polynucleotide SHIP inhibitors from recombinant nucleic acid constructs of the present invention. The invention is therefore directed in part to a method of producing a polynucleotide, such as an siRNA, by culturing a host cell comprising a recombinant nucleic acid construct that comprises at least one promoter operably linked to a polynucleotide SHIP inhibitor. In certain embodiments, the promoter may be a regulated promoter as provided herein, for example a tetracycline-repressible promoter. In certain embodiments the recombinant expression construct is a recombinant viral expression construct as provided herein. Examples of mammalian expression systems include the COS-7 lines of monkey kidney fibroblasts, described by Gluzman, *Cell* 23:175 (1981), and other cell lines capable of expressing a compatible vector, for example, the C127, 3T3, CHO, HeLa, HEK, and BHK cell lines. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and enhancer, and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences, for example as described herein

regarding the preparation of recombinant polynucleotide constructs. DNA sequences derived from the SV40 splice, and polyadenylation sites may be used to provide the required nontranscribed genetic elements. Introduction of the construct into the host cell can be effected by a variety of methods with which those skilled in the art will be familiar, including but not limited to, for example, liposomes including cationic liposomes, calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis et al., 1986 Basic Methods in Molecular Biology), or other suitable technique.

The expressed polynucleotides may be useful in intact host cells; in intact organelles such as cell membranes, intracellular vesicles or other cellular organelles; or in disrupted cell preparations including but not limited to cell homogenates or lysates, microsomes, uni- and multilamellar membrane vesicles or other preparations. Alternatively, expressed polynucleotides can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Finally, high performance liquid chromatography (HPLC) can be employed for final purification steps.

As used herein, the terms “administer”, “introduce”, “apply”, “treat”, “transplant”, “implant”, “deliver”, and grammatical variations thereof, are used interchangeably to provide SHIP inhibitors to target cells in vitro (e.g., ex vivo) or in vivo, or provide genetically modified (engineered) cells of the subject invention to a subject.

As used herein, the term “co-administration” and variations thereof refers to the administration of two or more agents simultaneously (in one or more preparations), or consecutively. For example, one or more types of genetically modified cells of the invention can be co-administered with other agents.

As used in this specification, including the appended claims, the singular “a”, “an”, and “the” include plural reference unless the context dictates otherwise. Thus, for example, a reference to “a polynucleotide” includes more than one such polynucleotide. A reference to “a nucleic acid sequence” includes more than one such sequence. A reference to “a cell” includes more than one such cell.

As used herein, the term “or” is inclusive of “both” (i.e., and/or). For example, as used herein, reference to megakaryocytes “or” megakaryocyte progenitors includes “either” or “and” (i.e., and/or). When the amount or concentration of megakaryocytes or megakaryocyte progenitors is to be determined, the amount or concentration of either or both cell types is intended.

The terms “comprising”, “consisting of” and “consisting essentially of” are defined according to their standard meaning. The terms may be substituted for one another throughout the instant application in order to attach the specific meaning associated with each term.

In general, the target nucleic acid is DNA or RNA. However, inventive methods may employ, for example, samples that contain DNA, or DNA and RNA, including messenger RNA, wherein DNA or RNA may be single stranded or double stranded, or a DNA-RNA hybrid may be included in the sample. A mixture of nucleic acids may also be employed. The specific nucleic acid sequence to be detected may be a fraction of a larger molecule or can be present initially as a discrete molecule, so that the specific sequence constitutes the entire nucleic acid. It is not necessary that the sequence to

be studied be present initially in a pure form; the nucleic acid may be a minor fraction of a complex mixture, such as contained in whole human DNA. The nucleic acid-containing sample used for determination of the sensitivity of the target cells to radiation therapy may be extracted by a variety of techniques such as that described by Sambrook, et al. (Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, N.Y., 1989; incorporated in its entirety herein by reference).

Cells expressing the target nucleic acid isolated from a subject can be obtained in a biological specimen from the subject. The cells, or nucleic acid, can be isolated from tumor tissue, blood, plasma, serum, lymph, lymph nodes, spleen, bone marrow, or any other biological specimen containing the target nucleic acid. Tumor tissue, blood, plasma, serum, lymph, spleen, and bone marrow are obtained by various medical procedures known to those of skill in the art.

The inventive methods are useful for producing a clinical response to treatment of various human anemias, bone marrow transplants, or cell proliferative disorders. A cell proliferative disorder as described herein may be a neoplasm. Such neoplasms are either benign or malignant. The term “neoplasm” refers to a new, abnormal growth of cells or a growth of abnormal cells that reproduce faster than normal. A neoplasm creates an unstructured mass (a tumor) which can be either benign or malignant. The term “benign” refers to a tumor that is noncancerous, e.g., its cells do not invade surrounding tissues or metastasize to distant sites. The term “malignant” refers to a tumor that is metastatic, invades contiguous tissue or no longer under normal cellular growth control.

As used herein, “a clinical response” is the response of a subject to modulation of the gene of interest. Criteria for determining a response to therapy are widely accepted and enable comparisons of the efficacy alternative treatments (see Slapak and Kufe, Principles of Cancer Therapy, in Harrison’s Principles of Internal Medicine, 13<sup>th</sup> edition, eds. Isselbacher et al., McGraw-Hill, Inc. 1994). A complete response (or complete remission) is the disappearance of all detectable malignant disease. A partial response is an approximately 50 percent decrease in the product of the greatest perpendicular diameters of one or more lesions. There can be no increase in size of any lesion or the appearance of new lesions. Progressive disease means at least an approximately 25 percent increase in the product of the greatest perpendicular diameter of one lesion or the appearance of new lesions. The response to treatment is evaluated after the subjects had completed therapy.

SHIP inhibitors (also referred to herein as “active compounds”) can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the SHIP-inhibiting nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal

administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL (BASF; Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a

similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring. For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to target cells with monoclonal antibodies) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

**Screening Assays.** The invention provides a method (also referred to herein as a "screening assay") for identifying SHIP inhibitors, i.e., candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other drugs) that bind to the SHIP protein or have a stimulatory or inhibitory effect on, for example, the SHIP gene expression or SHIP gene activity. Such identified compounds may be useful for the modulation of drug resistance. In one embodiment, the invention provides assays for screening candidate or test

compounds which bind to or modulate the activity of the target gene protein or polypeptide or biologically active portion thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; natural products libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam (1997) *Anti-cancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. *Proc. Natl. Acad. Sci. USA*, 1993, 90:6909; Erb et al. *Proc. Natl. Acad. Sci. USA*, 1994, 91:11422; Zuckermann et al. *J. Med. Chem.*, 1994, 37:2678; Cho et al. *Science*, 1993, 261:1303; Carrell et al. *Angew. Chem. Int. Ed. Engl.*, 1994, 33:2059; Carrell et al. *Angew. Chem. Int. Ed. Engl.*, 1994, 33:2061; and Gallop et al. *J. Med. Chem.*, 1994, 37:1233.

Libraries of compounds may be presented in solution (e.g., Houghten *Bio/Techniques*, 1992, 13:412-421), or on beads (Lam *Nature*, 1991, 354:82-84), chips (Fodor *Nature*, 1993, 364:555-556), bacteria (U.S. Pat. No. 5,223,409), spores (U.S. Pat. Nos. 5,571,698; 5,403,484; and 5,223,409), plasmids (Cull et al. *Proc. Natl. Acad. Sci. USA*, 1992, 89:1865-1869) or on phage (Scott and Smith *Science*, 1990, 249:386-390; Devlin *Science*, 1990, 249:404-406; Cwirla et al. *Proc. Natl. Acad. Sci.*, 1990, 87:6378-6382; and Felici *J. Mol. Biol.*, 1991, 222:301-310).

In one embodiment, an assay is a cell-based assay in which a cell which expresses the target gene protein, or a biologically active portion thereof, is contacted with a test compound and the ability of the test compound to bind to the target gene protein determined. The cell, for example, can be a yeast cell or a cell of mammalian origin. Determining the ability of the test compound to bind to the target gene protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the target gene protein or biologically active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^3\text{H}$ , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In a preferred embodiment, the assay comprises contacting a cell which expresses the target gene protein, or a biologically active portion thereof, with a known compound which binds the target gene to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the target gene protein, wherein determining the ability of the test compound to interact with the target gene protein comprises determining the ability of the test compound to preferentially bind to the target gene or a biologically active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing the target gene protein, or a biologically active portion thereof, with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the target

gene protein or biologically active portion thereof. Determining the ability of the test compound to modulate the activity of the target gene or a biologically active portion thereof can be accomplished, for example, by determining the ability of the target gene protein to bind to or interact with the target gene target molecule. As used herein, a "target molecule" is a molecule with which the target gene protein binds or interacts in nature, for example, a molecule in the nucleus or cytoplasm of a cell which expresses the target gene protein. The target gene target molecule can be a non-target gene molecule or the target gene protein or polypeptide. The target, for example, can be a second intracellular protein which has catalytic activity, a protein which naturally binds to the target gene, or a protein which facilitates the association of DNA with the target gene.

Determining the ability of the target gene protein to bind to or interact with the target gene target molecule can be accomplished by one of the methods described above for determining direct binding. In a preferred embodiment, determining the ability of the target gene protein to bind to or interact with the target gene target molecule can be accomplished by determining the activity of the target molecule or detecting a cellular response, for example, cell survival or cell proliferation in the presence of a chemotherapeutic drug.

In yet another embodiment, an assay of the present invention is a cell-free assay comprising contacting the target gene protein or biologically active portion thereof with a test compound and determining the ability of the test compound to bind to the target gene protein or biologically active portion thereof. Binding of the test compound to the target gene protein can be determined either directly or indirectly as described above. In a preferred embodiment, the assay includes contacting the target gene protein or biologically active portion thereof with a known compound which binds the target gene to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the target gene protein, wherein determining the ability of the test compound to interact with the target gene protein comprises determining the ability of the test compound to preferentially bind to the target gene or biologically active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-free assay comprising contacting the target gene protein or biologically active portion thereof with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the target gene protein or biologically active portion thereof. Determining the ability of the test compound to modulate the activity of the target gene can be accomplished, for example, by determining the ability of the target gene protein to bind to the target gene target molecule by one of the methods described above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of the target gene can be accomplished by determining the ability of the target gene protein further modulate the target gene target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as previously described.

In yet another embodiment, the cell-free assay comprises contacting the target gene protein or biologically active portion thereof with a known compound which binds the target gene to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the target gene protein, wherein determining the ability of the test compound to interact with the target gene protein comprises determining the ability of

the target gene protein to preferentially bind to or modulate the activity of the target gene target molecule.

The cell-free assays of the present invention are amenable to use of both native and variant forms (e.g., peptide fragments and fusion proteins) of the target gene. In the case of cell-free assays comprising a hydrophobic form of the target gene, it may be desirable to utilize a solubilizing agent such that the hydrophobic form of the target gene is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl-N,N-dimethyl-3-ammonio-1-propane sulfonate.

In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either the target gene or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to the target gene, or interaction of the target gene with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase/target gene fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein or the target gene protein, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of the target gene binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the target gene or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated target gene or target molecules can be prepared from biotin-NHS(N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with the target gene or target molecules but which do not interfere with binding of the target gene protein to its target molecule can be derivatized to the wells of the plate, and unbound target or the target gene trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the target gene or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target gene or target molecule.

In another embodiment, modulators of the target gene expression are identified in a method in which a cell is contacted with a candidate compound and the expression of the target gene (mRNA or protein, or the copy number of the target gene) in the cell is determined. The level of expression of the target gene in the presence of the candidate compound is compared to the level of expression of the target gene in the absence of the candidate compound. The candidate compound can then be identified as a modulator of the target gene expression based on this comparison. For example, when expression of the target gene mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of the target gene mRNA or protein expression. Alternatively, when expression of the target gene mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of the target gene mRNA or protein expression. The level of the target gene mRNA or protein expression in the cells, or the number of the target gene copies per cell can be determined by methods described herein for detecting the target gene genomic DNA, mRNA, or protein.

Target gene proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (see, e.g., U.S. Pat. No. 5,283,317; Zervos et al. *Cell*, 1993, 72:223-232; Madura et al. *J. Biol. Chem.*, 1993, 268:12046-12054; Bartel et al. *Bio/Techniques*, 1993, 14:920-924; Iwabuchi et al. *Oncogene*, 1993, 8:1693-1696; and W094/10300), to identify other proteins, which bind to or interact with the target gene ("target gene-binding proteins" or "target gene-bp") and modulate the target gene activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for the target gene is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, in vivo, forming the target gene-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the target gene.

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

Predictive Medicine. The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the present invention relates to diagnostic assays for determining the target gene protein and/or nucleic acid expression as well as the target gene activity, in the context of a biological sample (e.g., blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant target gene expression or activity

(e.g., altered drug resistance). The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with the target gene protein, nucleic acid expression or activity (e.g., altered drug resistance). For example, mutations in the target gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with the target gene protein, nucleic acid expression or activity. For example, because megakaryocyte production is inhibited where the target gene is expressed at a higher level in cells than normal, expression of the target gene can be used as an indicator of diminished megakaryocyte production.

Another aspect of the invention provides methods for determining the target gene protein, nucleic acid expression or the target gene activity in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (e.g., drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (e.g., the genotype of the individual examined to determine the ability of the individual to respond to a particular agent).

Diagnostic Assays. The invention provides a method of assessing expression, especially undesirable expression, of a cellular target gene. Undesirable (e.g., excessive) expression may indicate the presence, persistence or reappearance of reduced megakaryocyte production in an individual's tissue (e.g. spleen or bone marrow). More generally, aberrant expression may indicate the occurrence of a deleterious or disease-associated phenotype contributed to by the target gene.

An exemplary method for detecting the presence or absence of the target gene in a biological sample involves obtaining a biological sample (preferably from a body site implicated in a possible diagnosis of diseased or malignant tissue) from a test subject and contacting the biological sample with a compound or an agent capable of detecting the target gene protein or nucleic acid (e.g., mRNA, genomic DNA) that encodes the target gene protein such that the presence of the target gene is detected in the biological sample. The presence and/or relative abundance of the target gene indicates aberrant or undesirable expression of a cellular target gene, and correlates with the occurrence in situ of reduced megakaryocytes in the periphery.

A preferred agent for detecting the target gene mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to the target gene mRNA or genomic DNA. The nucleic acid probe can be, for example, a full-length target gene nucleic acid, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to the target gene mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

A preferred agent for detecting the target gene protein is an antibody capable of binding to the target gene protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')<sub>2</sub>) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include

detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect the target gene mRNA, protein, or genomic DNA in a biological sample in vitro as well as in vivo. For example, in vitro techniques for detection of the target gene mRNA include Northern hybridizations and in situ hybridizations. In vitro techniques for detection of the target gene protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. In vitro techniques for detection of the target gene genomic DNA include Southern hybridizations.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting the target gene protein, mRNA, or genomic DNA, such that the presence of the target gene protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of the target gene protein, mRNA or genomic DNA in the control sample with the presence of the target gene protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of the target gene in a biological sample (a test sample). Such kits can be used to determine if a subject is suffering from or is at increased risk of developing a disorder associated with aberrant expression of the target gene (e.g., the presence of a drug resistance cancer). For example, the kit can comprise a labeled compound or agent capable of detecting the target gene protein or mRNA in a biological sample and means for determining the amount of the target gene in the sample (e.g., an anti-target gene antibody or an oligonucleotide probe which binds to DNA encoding the target gene). Kits may also include instruction for observing that the tested subject is suffering from or is at risk of developing a disorder associated with aberrant expression of the target gene if the amount of the target gene protein or mRNA is above or below a normal level.

For antibody-based kits, the kit may comprise, for example: (1) a first antibody (e.g., attached to a solid support) which binds to the target gene protein; and, optionally, (2) a second, different antibody which binds to the target gene protein or the first antibody and is conjugated to a detectable agent.

For oligonucleotide-based kits, the kit may comprise, for example: (1) an oligonucleotide, e.g., a detectably labeled oligonucleotide, which hybridizes to the target gene nucleic acid sequence or (2) a pair of primers useful for amplifying the target gene nucleic acid molecule;

The kit may also comprise, e.g., a buffering agent, a preservative, or a protein stabilizing agent. The kit may also comprise components necessary for detecting the detectable agent (e.g., an enzyme or a substrate). The kit may also contain a control sample or a series of control samples which can be assayed and compared to the test sample contained. Each component of the kit is usually enclosed within an individual container and all of the various containers are

within a single package along with instructions for observing whether the tested subject is suffering from or is at risk of developing a disorder associated with aberrant expression of the target gene.

Pharmacogenomics. Agents, or modulators which have a stimulatory or inhibitory effect on the target gene activity (e.g., SHIP) as identified by a screening assay can be administered to individuals to treat (prophylactically or therapeutically) disorders (e.g., anemias) associated with aberrant target gene activity. In conjunction with such treatment, the pharmacogenomics (i.e., the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of the individual may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (e.g., drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to determine appropriate dosages and therapeutic regimens. Accordingly, the activity of the target gene protein, expression of the target gene nucleic acid, or mutation content of the target genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See, e.g., Linder, *Clin. Chem.*, 1997, 43(2): 254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

Thus, the activity of the target gene product (SHIP), expression of the target gene nucleic acid, or mutation content of the target genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition, pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with the target gene modulator, such as a modulator identified by one of the exemplary screening assays described herein.

Monitoring of Effects During Clinical Trials. Monitoring the influence of agents (e.g., drugs, compounds) on the expression or activity of the target gene (e.g., the ability to modulate the SHIP phenotype of a cell) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay to decrease the target gene expression, protein levels, or downregulate the target gene activity, can be

monitored in clinical trials of subjects exhibiting increased target gene expression, protein levels, or upregulated target gene activity.

Alternatively, the effectiveness of an agent determined by a screening assay to increase the target gene expression, protein levels, or upregulate target gene activity (e.g., to decrease megakaryocyte production), can be monitored in clinical trials of compounds designed to increase the target gene expression, protein levels, or upregulate target gene activity. In such clinical trials, the expression or activity of the target gene and, preferably, other genes that have been implicated in, for example, a cellular proliferation disorder, can be used as a "read out" or markers of the drug resistance of a particular cell.

For example, and not by way of limitation, genes, including the target gene, that are modulated in cells by treatment with an agent (e.g., compound, drug or small molecule) which modulates the target gene activity (e.g., identified in a screening assay) can be identified. Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of the target gene and other genes implicated in the disorder. The levels of gene expression (i.e., a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, or as is otherwise known in the art, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of the target gene or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during, treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (e.g., an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a preadministration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of the target gene protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the target gene protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the target gene protein, mRNA, or genomic DNA in the pre-administration sample with the target gene protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to decrease the expression or activity of the target gene to lower levels than detected, i.e., to increase the effectiveness of the agent.

Methods of Treatment. The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant target gene expression or activity. Alternatively, the target gene expression or activity may be normal (non-aberrant) but a decrease in target gene expression or activity would nonetheless have a beneficial effect on the subject. Such disorders include various human anemias and those in need of bone marrow transplants.

Prophylactic Methods. In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant target gene expression or



activity (e.g., the development of drug resistance), by administering to the subject an agent which modulates the target gene expression. Subjects at risk for a condition which is caused or contributed to by aberrant target gene expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as is known in the art. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of target gene aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. For example, administration of a prophylactic agent to a patient in need of a bone marrow transplant may prevent or delay the development of platelet production dropping below a critical threshold. Depending on the type of the target gene aberrancy, for example, the target gene agonist or the target gene antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein.

Therapeutic Methods. Another aspect of the invention pertains to methods of modulating the target gene expression or activity for therapeutic purposes. For example, the effectiveness of a bone marrow transplant is "potentiated" (enhanced) by increasing megakaryocyte production. The modulation of expression of the target gene disclosed in the method of the invention involves contacting a cell with an agent that modulates one or more of the activities of the target gene protein activity associated with the cell. An agent that modulates the target gene protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the target gene protein, a peptide, the target gene peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more of the biological activities of the target gene protein. Examples of such stimulatory agents include active the target gene protein and a nucleic acid molecule encoding the target gene that has been introduced into the cell. In another embodiment, the agent inhibits one or more of the biological activities of the target gene protein. Examples of such inhibitory agents include antisense target gene nucleic acid molecules and anti-target gene antibodies. These modulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of the target gene protein or nucleic acid molecule. In one embodiment, the method involves administering an agent, or combination of agents that modulates (e.g., upregulates or downregulates) the target gene expression or activity. In another embodiment, the method involves administering the target gene protein or nucleic acid molecule as therapy to compensate for reduced or aberrant target gene expression or activity.

For example, in one embodiment, the method involves administering a desired drug to an individual with a cell population expressing relatively high target gene levels, and coadministering an inhibitor of the target gene expression or activity. The administration and coadministration steps can be carried out concurrently or in any order, and can be separated by a time interval sufficient to allow uptake of either compound by the cells to be eradicated. For example, an antisense pharmaceutical composition (or a cocktail composition comprising an the target gene antisense oligonucleotide in combination with one or more other antisense oligonucleotides) can be administered to the individual sufficiently in advance of administration of the drug to allow the antisense composition to permeate the individual's tissues, especially tissue comprising the transformed cells to be eradicated; to be

internalized by transformed cells; and to disrupt the target gene expression and/or protein production.

Inhibition of the target gene activity is desirable in situations in which the target gene is abnormally upregulated and/or in which decreased target gene activity is likely to have a beneficial effect, e.g., increasing megakaryocyte production in the tissue of patient. Conversely, stimulation of the target gene activity is desirable in situations in which the target gene is abnormally downregulated and/or in which increased the target gene activity is likely to have a beneficial effect, e.g., in decreasing megakaryocyte and megakaryocyte progenitor production.

#### Materials And Methods

Mice strains. SHIP<sup>-/-</sup> mice (F9 or F10 X C57BL/6/J) produced in the inventors' laboratory have a deletion of the SHIP promoter and first exon (Wang, J. W. et al. *Science*, 2002, 295:2094-2097). A second SHIP-deficient mouse model, SHIP<sup>ΔIP/ΔIP</sup> (129SvJ) (Karlsson, M. C. et al. *J Exp Med.*, 2003, 198:333-340) in which the inositol phosphatase domain is deleted, was also analyzed (kindly provided by Dr. Jeffrey Ravetch, Rockefeller University, NY, USA). All studies described herein were conducted on six to eight week-old adult mice. Experiments were performed in compliance with institutional guidelines of the University of South Florida.

Cell isolation. Isolation of BM cells and splenocytes was as described (Wang, J. W. et al. *Science*, 2002, 295:2094-2097). Following red blood cell (RBC) lysis, the cells were re-suspended in staining medium (Wang, J. W. et al. *Science*, 2002, 295:2094-2097). PB was obtained from the retro-orbital sinus. For MKP analysis of PB, RBC were lysed in 1×RBC lysis buffer (EBIOSCIENCE, San Diego, Calif.) twice. Cells were then re-suspended for antibody staining.

Flow cytometry analysis and antibodies. Staining of MKP and MK was performed as per Hodohara et al. (Hodohara, K. et al. *Blood*, 2000, 95:769-775). All antibodies were from BD PHARMINGEN (San Diego, Calif.) except when mentioned otherwise. The cells were treated with anti-CD16/CD32 (2.4G2) to block Fc receptors and then stained with a lineage panel (PE), CD41-FITC(MWReg30), and cKit-APC(2B8). The Lineage panel was CD3e(17A2), CD4(GK1.5), CD8a (53-6.7), B220(RA3-6B2), Gr-1(RB6-8C5), Mac-1 (M1/70) (CALTAG, Burlingame, Calif.) and Ter119(TER-119). Dead cells were excluded using 7-AAD (BD PHARMINGEN, San Diego, Calif.). Analysis was done on a FACS Calibur and display of FACS data was performed using FlowJo 4.5.

Platelet analysis. Platelets were quantified using the Cell-Dyn 3700 hematology analyzer (ABBOTT DIAGNOSTIC, Dallas, Tex. USA).

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

#### EXAMPLE 1

##### SHIP-deficient Mice have Increased Numbers of MKP and MK

BM, spleen, and PB from SHIP<sup>-/-</sup>, SHIP<sup>ΔIP/ΔIP</sup>, and WT mice were analyzed by flow cytometry to determine the size of the megakaryocyte progenitor (MKP) and megakaryocyte (MK) compartment in vivo (FIG. 1A), using an immunophenotype defined by Hodohara and colleagues (Hodohara, K. et al. *Blood*, 2000, 95:769-775), Lin<sup>-</sup>cKit<sup>+</sup>CD41<sup>+</sup>, which con-



tains the majority of CFU-Mk activity. In BM, an expansion of the MKP compartment in SHIP<sup>-/-</sup> and SHIP<sup>ΔIP/ΔIP</sup> mice was observed (FIG. 1B). SHIP<sup>-/-</sup> and SHIP<sup>ΔIP/ΔIP</sup> spleens also show higher percentages of MKP compared to WT littermates (data not shown). Thus, SHIP<sup>-/-</sup> and SHIP<sup>ΔIP/ΔIP</sup> BM show a mean 18.1-fold and 50-fold increase, respectively, in the absolute number of MKP relative to WT controls (FIG. 1B). PB contained very few MKP, 3 to 5 MKP/μl, and their numbers were not significantly increased in SHIP<sup>-/-</sup> mice when compared to WT. However, MKP numbers were slightly, but significantly higher, in SHIP<sup>ΔIP/ΔIP</sup> PB as compared to WT littermates (FIG. 1B).

Assessing the MK compartment, it was observed that SHIP<sup>-/-</sup> and SHIP<sup>ΔIP/ΔIP</sup> BM contains significantly less MK compared to WT controls (FIG. 1C). However, in the spleen, there was a greater number of MK in SHIP<sup>-/-</sup> and SHIP<sup>ΔIP/ΔIP</sup> mice compared to their respective WT littermates (FIG. 1C). In SHIP<sup>-/-</sup> spleen there is a mean 10.9-fold increase in the absolute number of MK (FIG. 1C). Furthermore, in the PB of SHIP<sup>-/-</sup> mice there is a mean 7.7-fold increase in the absolute number of MK and the same trend was observed for SHIP<sup>ΔIP/ΔIP</sup> mice (FIG. 1C).

The increase in MK numbers found by flow cytometry in the SHIP-deficient spleens was corroborated by morphology, where the spleen of SHIP-deficient mice contained an increased number of MK (FIG. 1D). Furthermore, the BM histopathology revealed that MK in SHIP<sup>-/-</sup> BM have a hypolobulated micromegakaryocytic morphology when compared to WT BM, which contains mature hyperlobulated MK (FIG. 1E).

EXAMPLE 2

Platelet Levels Remain Unchanged in Ship-deficient Mice

Despite the profound expansion of the MKP and the MK compartment in SHIP<sup>-/-</sup> and SHIP<sup>ΔIP/ΔIP</sup> mice, they do not exhibit increased platelet levels relative to WT controls, when measured by hematology or flow cytometry. Platelet levels limit megakaryocytopoiesis by sequestering TPO (Kaushansky, K. *N Engl J Med.*, 1998, 339:746-754). Since platelet levels are not significantly increased in SHIP-deficient mice, it is possible that SHIP-deficiency increases the sensitivity of MK and MKP to steady-state TPO levels. Thus, proliferation and/or survival of MKP and MK is increased in SHIP mutant mice, leading to an expansion of the megakaryocytic compartment. Consistent with the present inventors' hypothesis, TPO stimulation of primary MK and c-mpl transfected Ba/F3 cells leads to SHIP phosphorylation and activation of downstream effectors of the cell cycle and survival (Geddis, A. E. et al. *J Biol Chem.*, 2001, 276:34473-34479).

Table 2 shows the different SHIP-deficient models and their WT littermates (n>6 mice/genotype). Data was analyzed using an unpaired, two-tailed. Students' T-test.

TABLE 2

Platelet counts in SHIP-deficient mice.	
Mice genotype	Platelet levels (# × 10 <sup>3</sup> /μl)
SHIP <sup>-/-</sup>	672.8 ± 43.4 <sup>++</sup>
SHIP <sup>+/-</sup>	848.9 ± 35.7
SHIP <sup>+/+</sup> (C57Bl6)	803.7 ± 31.5
SHIP <sup>ΔIP/ΔIP</sup>	455.4 ± 81.1 <sup>++</sup>

TABLE 2-continued

Platelet counts in SHIP-deficient mice.	
Mice genotype	Platelet levels (# × 10 <sup>3</sup> /μl)
SHIP <sup>+/ΔIP</sup>	652.3 ± 28.0
SHIP <sup>+/+</sup> (129SvJ)	647.9 ± 28.0

<sup>++</sup>p < 0.05 compared to their respective WT and SHIP heterozygous littermates

A decrease in the number of MK in the BM of SHIP-deficient mice and an increase in MK numbers in the PB and spleen were observed. This may be due to increased responsiveness of MK to SDF-1, which would cause trans-endothelial migration of MK from the BM to the circulation (Wang, J. F. et al. *Blood*, 1998, 92:756-764; Hamada, T. et al. *J Exp Med.*, 1998, 188:539-548; Avecilla, S. T. et al., *Nat Med.*, 2004, 10:64-71). Thus, SHIP may also control pathways that mediate MK migration in response to SDF-1, as it does in myeloid progenitors (Kim, C. H. et al. *J Clin Invest.*, 1999, 104:1751-1759).

Despite an increase in the level of MKP and MK, the platelet counts were reduced in SHIP-deficient mice. It is well established that SHIP<sup>-/-</sup> mice suffer from splenomegaly, resulting from extramedullary hematopoiesis, where SHIP-deficient spleen are enlarged by 5-fold (Helgason et al., *Genes Dev.*, 1998, 12:1610-20). Since splenomegaly has been associated with splenic platelet sequestration previously (Aster R H Br. J., *Haematol.*, 1972, 22:259-63) (Naouri et al. *Acta Haematol.*, 1993, 89:200-3), the present inventors hypothesize that SHIP-deficient spleens sequester platelets, leading to a reduction in circulating platelet numbers in the peripheral blood.

Thus, overall, these findings suggest that SHIP can be targeted in vivo to increase the pool of MKP and MK, and thus enable this compartment to replenish platelets more rapidly following myeloablative chemotherapy and radiation treatment, for example.

EXAMPLE 3

MKP and MK are Increased in BM and Spleen of SHIP-ablated Mice

To observe if inhibition of SHIP during adulthood could also result in an increase in MKP production, the present inventors used the MxCre model (Wang et al. *Science*, 2002, 295:2094-2097). Briefly, the treatment of MxCre<sup>+/fl</sup> mice with polyIC will lead to Cre recombinase expression through Type interferon-inducible Mx1 promoter, and deletion of the gene section between two loxP sites (Kuhn et al. *Science*, 1995, 259:1427-1429; Oberdoerffer et al. *Nucleic Acids Res.*, 2003, 31:e140). In this case, the promoter and the first exon of the SHIP gene will be deleted resulting in the ablation of SHIP expression. As a control, MxCre<sup>-</sup>/SHIP<sup>fl/fl</sup> are treated with polyIC in the same manner than the MxCre<sup>+/fl</sup> mice. Twenty-one days after the last polyIC treatment, mice were euthanized and the level of MKP was evaluated by flow cytometry (FIG. 5A). As observed in FIG. 5B-1, there was an increase in the percentage of MKP in the BM and spleen of SHIP-ablated mice as compared to MxCre<sup>-</sup> mice. Furthermore, it was observed that SHIP-ablated BM contains approximately 4 times more MKP than their MxCre<sup>-</sup> counterpart (FIG. 5B-2). As for the germline SHIP<sup>-/-</sup>, an increase in the percentage of MK present in the spleen was also observed (shown in FIG. 5B-3). This result suggests that mice that undergo normal development can also exhibit increased MKP numbers once SHIP is deleted during adulthood. Thus,

47

methods and agents that inhibit SHIP function could be used as a therapy in vitro (e.g., ex vivo) or in vivo to increase megakaryocytopoiesis in adult patients.

## EXAMPLE 4

## SHIP Knockdown Using Interfering RNA

Four potential interfering RNA sequences targeting human SHIP were obtained from Open Biosystems:

siRNA sequence H1: (SEQ ID NO: 16)  
AAGGAAUUGCGUUUACACUUA

siRNA sequence H2: (SEQ ID NO: 17)  
AAAAUUGCGUUUACACUUA

shRNA sequence 63332: (SEQ ID NO: 18)  
TGCTGTTGACAGTGAGCGAGGCTTCCAGAAGAGCATCTTATAGTGAAGC  
CACAGATGTATAAGATGCTCTTCTGGAAGCCCTGCCTACTGCCTCGGA

shRNA sequence 63331: (SEQ ID NO: 19)  
TGCTGTTGACAGTGAGCGAGCCCATATCACCCAAGAAGTTTATAGTGAAGC  
CACAGATGTAACTTCTTGGGTGATATGGGCGTGCCTACTGCCTCGGA

The underlined portion of shRNA 63332 (SEQ ID NO:18): GCTTCCAGAAGAGCATCTTAT (SEQ ID NO:20), and shRNA 63331 (SEQ ID NO:19): GCCCATATCACCCAAGAGTTT (SEQ ID NO:21), represent the target sequences

48

in human SHIP. The other non-underlined portions are the loop and termini of the shRNA.

FIG. 4 shows the human SHIP cDNA sequence with the targets for siRNA sequence H1 (SEQ ID NO:16), siRNA sequence H2 (SEQ ID NO:17), shRNA 63332 (SEQ ID NO:18), and shRNA 63331 (SEQ ID NO:19) in boxes. FIG. 6 is a general vector map representing shRNA 63332 and shRNA 63331.

H1 was predicted to have the best knockdown effectiveness by the siRNA design tool available on the DHARMACON website, using the default parameters. H1 and H2 siRNA were delivered to human NK cells in vitro using the nucleofection system (AMAXA, Maryland, USA). Twenty-four hours later, cells were lysed and a Western Blot was performed to determine the level of SHIP protein in the test cells following administration of the siRNAs. The  $\beta$ -actin protein level was also observed to ensure equal loading of the protein lysate. Results are shown in FIG. 7. Both H1 and H2 effectively reduced SHIP expression; however, H1 was more effective, which was consistent with the siRNA design tool.

It will be seen that the objects set forth above, and those made apparent from the foregoing description, are efficiently attained and since certain changes may be made in the above construction without departing from the scope of the invention, it is intended that all matters contained in the foregoing description or shown in the accompanying drawings shall be interpreted as illustrative and not in a limiting sense.

All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety, including all figures and tables, to the extent they are not inconsistent with the explicit teachings of this specification.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 21

<210> SEQ ID NO 1

<211> LENGTH: 4925

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

aaacaggaag tcagtcagtt aagctggttg cagcagccga ggccaccaag aggcaacggg      60
cggcaggttg cagtgagggg gcctccgctc ccctcggttg tgtgtgggtc ctgggggtgc      120
ctgccggccc ggccgaggag gcccacgccc accatggtcc cctgctggaa ccatggcaac      180
atcacccgct ccaaggcgga ggagctgctt tccaggacag gcaaggacgg gagcttcctc      240
gtgcgtgcc a gtagtccat ctccgggca tacgcgtctt gcgtgctgta tcggaattgc      300
gtttacactt acagaattct gcccaatgaa gatgataaat tcaactgttca ggcatccgaa      360
ggcgtctcca tgaggttctt caccaagctg gaccagctca tcgagtttta caagaaggaa      420
aacatggggc tggtgaccca tctgcaatac cctgtgccgc tggaggaaga ggacacaggc      480
gacgacctg aggaggacac agaaagtgtc gtgtctccac ccgagctgcc cccaagaaac      540
atcccgtga ctgccagctc ctgtgaggcc aaggaggttc ctttttcaaa cgagaatccc      600
cgagcgaccg agaccagccg gccgagcctc tccgagacat tgttccagcg actgcaaagc      660
atggacacca gtgggtcttc agaagagcat cttaaggcca tccaagatta tttaagcact      720
cagctcgccc aggactctga atttgtgaag acagggtcca gcagtcttcc tcacctgaag      780
aaactgacca cactgctctg caaggagctc tatggagaag tcatccggac cctcccatcc      840

```

-continued

---

ctggagtctc	tgcagaggtt	atttgaccag	cagctctccc	cgggcctccg	tccacgtcct	900
caggttcctg	gtgaggccaa	tccatcaac	atggtgtcca	agctcagcca	actgacaagc	960
ctgttgctcat	ccattgaaga	caaggccaag	gccttgctgc	acgaggggtc	tgagtctccg	1020
caccggccct	cccttatccc	tccagtcacc	tttgagggtg	aggcagagtc	tctggggatt	1080
cctcagaaaa	tgcagctcaa	agtcgacgtt	gagtcctggg	aactgatcat	taagaagtcc	1140
aaggatggtt	ctgaggacaa	gttctacagc	cacaagaaaa	tcctgcagct	cattaagtca	1200
cagaaatttc	tgaataagtt	ggtgatcttg	gtggaacag	agaaggagaa	gacctgcggy	1260
aaggaatatg	tttttgctga	ctccaaaaag	agagaaggct	tctgccagct	cctgcagcag	1320
atgaagaaca	agcactcaga	gcagccggag	cccacatga	tcaccatctt	catcggcacc	1380
tggaaacatg	gtaacgcccc	ccctcccaag	aagatcacgt	cctggtttct	ctccaagggg	1440
cagggaaga	cgcgggacga	ctctgcggac	tacatcccc	atgacattta	cgtgatcggc	1500
acccaagagg	acccctgtag	tgagaaggag	tggtgggaga	tcctcaaaac	ctccctgcaa	1560
gaaatcacca	gtgtgacttt	taaaacagtc	gccatccaca	cgctctggaa	catccgcac	1620
gtggtgctgg	ccaagcctga	gcacgagaa	cggatcacgc	acatctgtac	tgacaacgtg	1680
aagacaggca	ttgcaaacac	actgggggaa	aaggagcg	tggtgggtgc	gttcatgttc	1740
aatggaacct	ccttaggggt	cgtaacagc	cacttgactt	caggaagtga	aaagaaactc	1800
aggcgaaacc	aaaactatat	gaacattctc	cggttcctgg	ccctgggcca	caagaagctg	1860
agtcccttta	acatcactca	ccgcttcacg	cacctcttct	ggtttgggga	tcttaactac	1920
cgtgtggatc	tgctacctg	ggaggcagaa	accatcatcc	agaaaatcaa	gcagcagcag	1980
tacgcagacc	tcctgtccca	cgaccagctg	ctcacagaga	ggaggggagca	gaaggctctc	2040
ctacacttcg	aggaggaaga	aatcacgttt	gccccaacct	accgttttga	gagactgact	2100
cgggacaaat	acgcctacac	caagcagaaa	gcgacaggga	tgaagtacaa	cttgcccttc	2160
tggtgtgacc	gagtcctctg	gaagtcttat	cccctgggtc	acgtgggtgtg	tcagtcttat	2220
ggcagtacca	gcgacatcat	gacgagtgc	cacagccctg	tccttgccac	atttgaggca	2280
ggagtcaatt	cccagtttgt	ctccaagaa	ggtcccgagg	ctgttgacag	ccaaggacag	2340
attgagtttc	tcagggtgta	tgccacattg	aagaccaagt	cccagaccaa	attctacctg	2400
gagttccact	cgagctgctt	ggagagtgtt	gtcaagagtc	aggaaggaga	aatgaagaa	2460
ggaagtgagg	gggagctggt	ggtgaagtgt	ggtgagactc	ttccaaagct	gaagcccatt	2520
atctctgacc	ctgagtacct	gctagaccag	cacatcctca	tcagcatcaa	gtcctctgac	2580
agcgacgaat	cctatggcga	gggctgcatt	gcccttcggt	tagaggccac	agaaacgcag	2640
ctgcccatct	acacgcctct	cacccaccat	ggggagttag	caggccactt	ccagggggag	2700
atcaagctgc	agacctctca	gggcaagacg	agggagaagc	tctatgactt	tgtgaagacg	2760
gagcgtgatg	aatccagtgg	gcccagagcc	ctgaagagcc	tcaccagcca	cgaccccatg	2820
aagcagtggg	aagtcactag	cagggccctt	ccgtgcagtg	gctccagcat	caactgaaatc	2880
atcaacccca	actacatggg	agtggggccc	tttggggccc	caatgcccct	gcacgtgaag	2940
cagaccttgt	cccctgacca	gcagcccaca	gcctggagct	acgaccagcc	gcccagggac	3000
tccccgctgg	ggcctgcag	gggagaaaag	cctccgacac	ctcccgccca	gcccgcata	3060
tcacccaaga	agtttttacc	ctcaacagca	aaccgggggtc	tcctccag	gacacaggag	3120
tcaaggccca	gtgacctggg	gaagaacgca	ggggacacgc	tgctcagga	ggacctgccg	3180
ctgacgaagc	ccgagatggt	tgagaacccc	ctgtatgggt	ccctgagttc	cttccctaag	3240

-continued

---

```

cctgctccca ggaaggacca ggaatcccc aaaatgccgc ggaaggaaacc cccgccctgc 3300
ccggaacccg gcattctgtc gccacgcatc gtgctcacca aagcccagga ggctgatcgc 3360
ggcgaggggc ccggcaagca ggtgcccgcg ccccggtgc gctccttcac gtgctcatcc 3420
tctgccgagg gcaggggcgc cggcggggac aagagccaag ggaagcccaa gaccccggtc 3480
agctcccagg ccccggtgcc ggccaagagg cccatcaagc cttccagatc ggaaatcaac 3540
cagcagaccc cgcccacccc gacgcccgcg ccgcgctgc cagtcaagag cccggcggtg 3600
ctgcacctcc agcactccaa gggcccgac taccgcgaca acaccgagct cccgcatcac 3660
ggcaagcacc ggccggagga ggggccacca gggcctctag gcaggactgc catgcagtga 3720
agccctcagt gagctgccac tgagtcggga gccagagga acggcgtaa gccactggac 3780
cctctcccgg gacctctgc ttgctctcc tgcacagctt cctatgcaag gctttgtgtt 3840
ttcaggaaag ggcctagctt ctgtgtggcc cacagagttc actgcctgtg agacttagca 3900
ccaagtgtg aggtggaag aaaaacgcac accagacggg caacaaacag tctgggtccc 3960
cagctcgctc ttggtacttg ggaccccagt gcctcgttga gggcgccatt ctgaagaaag 4020
gaactgcagc gccgatttga ggttgagat atagataata ataataata taataataat 4080
ggccacatgg atcgaacact catgatgtgc caagtgtgt gctaagtgtt ttacgaacat 4140
tcgtcatatc aggatgacct cgagagctga ggctctagcc acctaaaacc acgtgcccaa 4200
acccaccagt ttaaaacggt gtgtgttcgg aggggtgaaa gcattaagaa gccagtgcc 4260
ctcctggagt gagacaaggg ctgcgcctta aggagctgaa gagtctgggt agcttgttta 4320
gggtacaaga agcctgttct gtccagcttc agtgacacaa gctgctttag ctaaagtccc 4380
gcgggttcgc gcattggctag gctgagagca gggatctacc tggcttctca gttctttggt 4440
tggaaggagc aggaatcag ctccattct ccagtggaga gatctggcct cagcttgggc 4500
tagagatgcc aaggcctgtg ccaggttccc tgtgccctcc tcgaggtggg cagccatcac 4560
cagccacagt taagccaagc ccccaacat gtattccatc gtgctggtag aagagctttt 4620
gctgttgcgc ccgaaagccg tgctctccag cctggctgcc agggaggggt ggctcttgg 4680
ttccaggctc ttgaaatagt gcagcctttt ctccctatct ctgtggcttt cagctctgct 4740
tccttggtta ttaggagaat agatgggtga tgtctttcct tatgttgctt tttcaacata 4800
gcagaattaa tgtaggagc taaatccagt ggtgtgtgtg aatgcagaag ggaatgcacc 4860
ccacattccc atgatggaag tctgcgtaac caataaattg tgcctttctc actcaaaaaa 4920
aaaaa 4925

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 4865

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Mus musculus

&lt;400&gt; SEQUENCE: 2

```

Gly Gly Cys Ala Ala Thr Thr Thr Cys Thr Gly Ala Gly Ala Gly Gly
1          5          10         15
Cys Ala Ala Cys Ala Gly Gly Cys Gly Gly Cys Ala Gly Gly Thr Cys
20         25         30
Thr Cys Ala Gly Cys Cys Thr Ala Gly Ala Gly Ala Gly Gly Gly Cys
35         40         45
Cys Cys Thr Gly Ala Ala Cys Thr Ala Cys Thr Thr Thr Gly Cys Thr
50         55         60
Gly Gly Ala Gly Thr Gly Thr Cys Cys Gly Thr Cys Cys Thr Gly Gly

```

-continued

65	70	75	80
Gly Ala Gly Thr Gly 85	Gly Cys Thr Gly Cys 90	Thr Gly Ala Cys Cys 95	Cys
Ala Gly Thr Cys Cys 100	Ala Gly Gly Ala Gly 105	Ala Cys Cys Cys 110	Ala Thr
Gly Cys Cys Thr Gly 115	Cys Cys Ala Thr Gly 120	Gly Gly Thr Cys Cys 125	Cys Thr
Gly Gly Gly Thr Gly 130	Gly Ala Ala Cys Cys 135	Ala Thr Gly Gly Cys 140	Ala
Ala Cys Ala Thr Cys 145	Ala Cys Cys Cys Gly 150	Cys Thr Cys Cys Ala 155	Ala
Gly Gly Cys Ala Gly 165	Ala Gly Gly Ala Gly 170	Cys Thr Ala Cys Thr 175	Thr
Thr Cys Cys Ala Gly 180	Ala Gly Cys Cys Gly 185	Gly Gly Cys Ala Ala 190	Gly Gly
Ala Cys Gly Gly Gly 195	Ala Gly Cys Thr Thr 200	Cys Cys Thr Thr Gly 205	Thr
Gly Cys Gly Thr Gly 210	Cys Cys Ala Gly Cys 215	Gly Ala Gly Thr Cys 220	Cys
Ala Thr Cys Cys Cys 225	Cys Cys Gly Gly Gly 230	Cys Cys Thr Ala Cys 235	Gly
Cys Ala Cys Thr Cys 245	Thr Gly Cys Gly Thr 250	Gly Cys Thr Gly Thr 255	Thr
Cys Cys Gly Gly Ala 260	Ala Thr Thr Gly Thr 265	Thr Thr Thr Ala Cys 270	
Ala Cys Thr Thr Ala 275	Cys Ala Gly Gly Ala 280	Thr Thr Cys Thr Gly 285	Cys
Cys Cys Ala Ala Thr 290	Gly Ala Gly Gly Ala 295	Cys Gly Ala Thr Ala 300	Ala
Ala Thr Thr Cys Ala 305	Cys Thr Gly Thr Thr 310	Cys Ala Gly Gly Cys 315	Ala
Thr Cys Cys Gly Ala 325	Ala Gly Gly Thr Gly 330	Cys Cys Cys Cys Ala 335	
Thr Gly Ala Gly Gly 340	Thr Thr Cys Thr Thr 345	Cys Ala Cys Gly Ala 350	Ala
Gly Cys Thr Gly Gly 355	Ala Cys Cys Ala Gly 360	Cys Thr Cys Ala Thr 365	Cys
Gly Ala Cys Thr Thr 370	Thr Thr Ala Cys Ala 375	Gly Ala Ala Gly Gly 380	
Ala Ala Ala Ala Cys 385	Ala Thr Gly Gly Gly 390	Gly Cys Thr Gly Gly 395	Thr
Gly Ala Cys Cys Cys 405	Ala Cys Cys Thr Gly 410	Cys Ala Gly Thr Ala 415	Cys
Cys Cys Cys Gly Thr 420	Gly Cys Cys Cys Cys 425	Thr Gly Gly Ala Gly 430	Gly
Ala Gly Gly Ala Gly 435	Gly Ala Thr Gly Cys 440	Thr Ala Thr Thr Gly 445	Ala
Thr Gly Ala Gly Gly 450	Cys Thr Gly Ala Gly 455	Gly Ala Gly Gly Ala 460	Cys
Ala Cys Thr Gly Thr 465	Ala Gly Ala Ala Ala 470	Gly Thr Gly Thr Cys 475	Ala
Thr Gly Thr Cys Ala 485	Cys Cys Ala Cys Cys 490	Thr Gly Ala Gly Cys 495	Thr

-continued

Gly	Cys	Cys	Thr	Cys	Cys	Cys	Ala	Gly	Ala	Ala	Ala	Cys	Ala	Thr	Thr
			500					505					510		
Cys	Cys	Thr	Ala	Thr	Gly	Thr	Cys	Thr	Gly	Cys	Cys	Gly	Gly	Gly	Cys
		515					520					525			
Cys	Cys	Ala	Gly	Cys	Gly	Ala	Gly	Gly	Cys	Cys	Ala	Ala	Gly	Gly	Ala
		530				535					540				
Cys	Cys	Thr	Thr	Cys	Cys	Thr	Cys	Thr	Thr	Gly	Cys	Ala	Ala	Cys	Ala
545					550					555					560
Gly	Ala	Gly	Ala	Ala	Cys	Cys	Cys	Cys	Cys	Gly	Ala	Gly	Cys	Cys	Cys
			565						570					575	
Cys	Thr	Gly	Ala	Gly	Gly	Thr	Cys	Ala	Cys	Cys	Cys	Gly	Gly	Cys	Thr
			580					585					590		
Gly	Ala	Gly	Thr	Cys	Thr	Cys	Thr	Cys	Cys	Gly	Ala	Gly	Ala	Cys	Ala
		595					600					605			
Cys	Thr	Gly	Thr	Thr	Thr	Cys	Ala	Gly	Cys	Gly	Thr	Cys	Thr	Ala	Cys
610						615					620				
Ala	Gly	Ala	Gly	Cys	Ala	Thr	Gly	Gly	Ala	Thr	Ala	Cys	Cys	Ala	Gly
625					630					635					640
Thr	Gly	Gly	Gly	Cys	Thr	Thr	Cys	Cys	Cys	Gly	Ala	Gly	Gly	Ala	Gly
				645					650					655	
Cys	Ala	Cys	Cys	Thr	Gly	Ala	Ala	Ala	Gly	Cys	Cys	Ala	Thr	Cys	Cys
			660					665					670		
Ala	Gly	Gly	Ala	Thr	Thr	Ala	Thr	Cys	Thr	Gly	Ala	Gly	Cys	Ala	Cys
		675					680					685			
Thr	Cys	Ala	Gly	Cys	Thr	Cys	Cys	Thr	Cys	Cys	Thr	Gly	Gly	Ala	Thr
690						695					700				
Thr	Cys	Cys	Gly	Ala	Cys	Thr	Thr	Thr	Thr	Thr	Gly	Ala	Ala	Gly	Ala
705					710					715					720
Cys	Gly	Gly	Gly	Cys	Thr	Cys	Cys	Ala	Gly	Cys	Ala	Ala	Cys	Cys	Thr
				725					730					735	
Cys	Cys	Cys	Thr	Cys	Ala	Cys	Cys	Thr	Gly	Ala	Ala	Gly	Ala	Ala	Gly
			740					745				750			
Cys	Thr	Gly	Ala	Thr	Gly	Thr	Cys	Ala	Cys	Thr	Gly	Cys	Thr	Cys	Thr
		755					760					765			
Gly	Cys	Ala	Ala	Gly	Gly	Ala	Gly	Cys	Thr	Cys	Cys	Ala	Thr	Gly	Gly
770					775						780				
Gly	Gly	Ala	Ala	Gly	Thr	Cys	Ala	Thr	Cys	Ala	Gly	Gly	Ala	Cys	Thr
785					790					795					800
Cys	Thr	Gly	Cys	Cys	Ala	Thr	Cys	Cys	Cys	Thr	Gly	Gly	Ala	Gly	Thr
			805						810					815	
Cys	Thr	Cys	Thr	Gly	Cys	Ala	Gly	Ala	Gly	Gly	Thr	Thr	Gly	Thr	Thr
			820					825					830		
Thr	Gly	Ala	Cys	Cys	Ala	Ala	Cys	Ala	Gly	Cys	Thr	Cys	Thr	Cys	Cys
		835					840					845			
Cys	Cys	Ala	Gly	Gly	Cys	Cys	Thr	Thr	Cys	Gly	Cys	Cys	Cys	Ala	Cys
850							855				860				
Gly	Ala	Cys	Cys	Thr	Cys	Ala	Gly	Gly	Thr	Gly	Cys	Cys	Cys	Gly	Gly
865					870					875					880
Ala	Gly	Ala	Gly	Gly	Cys	Cys	Ala	Gly	Thr	Cys	Cys	Cys	Ala	Thr	Cys
				885					890					895	
Ala	Cys	Cys	Ala	Thr	Gly	Gly	Thr	Thr	Gly	Cys	Cys	Ala	Ala	Ala	Cys
			900					905					910		
Thr	Cys	Ala	Gly	Cys	Cys	Ala	Ala	Thr	Thr	Gly	Ala	Cys	Ala	Ala	Gly
		915					920					925			

-continued

---

Thr	Cys	Thr	Gly	Cys	Thr	Gly	Thr	Cys	Thr	Thr	Cys	Cys	Ala	Thr	Thr
930						935					940				
Gly	Ala	Ala	Gly	Ala	Thr	Ala	Ala	Gly	Gly	Thr	Cys	Ala	Ala	Gly	Thr
945					950					955					960
Cys	Cys	Thr	Thr	Gly	Cys	Thr	Gly	Cys	Ala	Cys	Gly	Ala	Gly	Gly	Gly
				965					970					975	
Cys	Thr	Cys	Ala	Gly	Ala	Ala	Thr	Cys	Thr	Ala	Cys	Cys	Ala	Ala	Cys
			980					985					990		
Ala	Gly	Gly	Cys	Gly	Thr	Thr	Cys	Cys	Cys	Thr	Thr	Ala	Thr	Cys	Cys
	995						1000						1005		
Cys	Thr	Cys	Cys	Gly	Gly	Thr	Cys	Ala	Cys	Cys	Thr	Thr	Thr	Thr	Gly
1010						1015						1020			
Ala	Gly	Gly	Thr	Gly	Ala	Ala	Gly	Thr	Cys	Ala	Gly	Ala	Gly	Thr	
1025						1030					1035				
Cys	Cys	Cys	Thr	Gly	Gly	Gly	Cys	Ala	Thr	Thr	Cys	Cys	Thr	Cys	
1040						1045					1050				
Ala	Gly	Ala	Ala	Ala	Ala	Thr	Gly	Cys	Ala	Thr	Cys	Thr	Cys	Ala	
1055						1060					1065				
Ala	Ala	Gly	Thr	Gly	Gly	Ala	Cys	Gly	Thr	Thr	Gly	Ala	Gly	Thr	
1070						1075					1080				
Cys	Thr	Gly	Gly	Gly	Ala	Ala	Ala	Cys	Thr	Gly	Ala	Thr	Cys	Gly	
1085						1090					1095				
Thr	Thr	Ala	Ala	Gly	Ala	Ala	Gly	Thr	Cys	Cys	Ala	Ala	Gly	Gly	
1100						1105					1110				
Ala	Thr	Gly	Gly	Thr	Thr	Cys	Thr	Gly	Ala	Gly	Gly	Ala	Cys	Ala	
1115						1120					1125				
Ala	Gly	Thr	Thr	Cys	Thr	Ala	Cys	Ala	Gly	Cys	Cys	Ala	Cys	Ala	
1130						1135					1140				
Ala	Ala	Ala	Ala	Ala	Ala	Thr	Cys	Cys	Thr	Gly	Cys	Ala	Gly	Cys	
1145						1150					1155				
Thr	Cys	Ala	Thr	Thr	Ala	Ala	Gly	Thr	Cys	Cys	Cys	Ala	Gly	Ala	
1160						1165					1170				
Ala	Gly	Thr	Thr	Thr	Cys	Thr	Ala	Ala	Ala	Cys	Ala	Ala	Gly	Thr	
1175						1180					1185				
Thr	Gly	Gly	Thr	Gly	Ala	Thr	Thr	Thr	Thr	Gly	Gly	Thr	Gly	Gly	
1190						1195					1200				
Ala	Gly	Ala	Cys	Gly	Gly	Ala	Gly	Ala	Ala	Gly	Gly	Ala	Gly	Ala	
1205						1210					1215				
Ala	Ala	Ala	Thr	Cys	Cys	Thr	Gly	Ala	Gly	Gly	Ala	Ala	Gly	Gly	
1220						1225					1230				
Ala	Ala	Thr	Ala	Thr	Gly	Thr	Thr	Thr	Thr	Thr	Gly	Cys	Thr	Gly	
1235						1240					1245				
Ala	Cys	Thr	Cys	Thr	Ala	Ala	Gly	Ala	Ala	Ala	Ala	Gly	Ala	Gly	
1250						1255					1260				
Ala	Ala	Gly	Gly	Cys	Thr	Thr	Cys	Thr	Gly	Thr	Cys	Ala	Ala	Cys	
1265						1270					1275				
Thr	Cys	Cys	Thr	Gly	Cys	Ala	Gly	Cys	Ala	Gly	Ala	Thr	Gly	Ala	
1280						1285					1290				
Ala	Gly	Ala	Ala	Cys	Ala	Ala	Gly	Cys	Ala	Thr	Thr	Cys	Gly	Gly	
1295						1300					1305				
Ala	Gly	Cys	Ala	Gly	Cys	Cys	Ala	Gly	Ala	Gly	Cys	Cys	Thr	Gly	
1310						1315					1320				
Ala	Cys	Ala	Thr	Gly	Ala	Thr	Cys	Ala	Cys	Cys	Ala	Thr	Cys	Thr	

-continued

1325	1330	1335
Thr Cys Ala Thr Thr Gly Gly Cys Ala Cys Thr Thr Gly Gly Ala 1340 1345 1350		
Ala Cys Ala Thr Gly Gly Gly Thr Ala Ala Thr Gly Cys Ala Cys 1355 1360 1365		
Cys Cys Cys Cys Thr Cys Cys Cys Ala Ala Gly Ala Ala Gly Ala 1370 1375 1380		
Thr Cys Ala Cys Gly Thr Cys Cys Thr Gly Gly Thr Thr Thr Cys 1385 1390 1395		
Thr Cys Thr Cys Cys Ala Ala Gly Gly Gly Gly Cys Ala Gly Gly 1400 1405 1410		
Gly Ala Ala Ala Gly Ala Cys Ala Cys Gly Gly Gly Ala Cys Gly 1415 1420 1425		
Ala Cys Thr Cys Thr Gly Cys Thr Gly Ala Cys Thr Ala Cys Ala 1430 1435 1440		
Thr Cys Cys Cys Cys Cys Ala Thr Gly Ala Cys Ala Thr Cys Thr 1445 1450 1455		
Ala Thr Gly Thr Gly Ala Thr Thr Gly Gly Cys Ala Cys Cys Cys 1460 1465 1470		
Ala Gly Gly Ala Gly Gly Ala Thr Cys Cys Cys Cys Thr Thr Gly 1475 1480 1485		
Gly Ala Gly Ala Gly Ala Ala Gly Gly Ala Gly Thr Gly Gly Cys 1490 1495 1500		
Thr Gly Gly Ala Gly Cys Thr Ala Cys Thr Cys Ala Gly Gly Cys 1505 1510 1515		
Ala Cys Thr Cys Cys Cys Thr Gly Cys Ala Ala Gly Ala Ala Gly 1520 1525 1530		
Thr Cys Ala Cys Cys Ala Gly Cys Ala Thr Gly Ala Cys Ala Thr 1535 1540 1545		
Thr Thr Ala Ala Ala Ala Cys Ala Gly Thr Thr Gly Cys Cys Ala 1550 1555 1560		
Thr Cys Cys Ala Cys Ala Cys Cys Cys Thr Cys Thr Gly Gly Ala 1565 1570 1575		
Ala Cys Ala Thr Thr Cys Gly Cys Ala Thr Ala Gly Thr Gly Gly 1580 1585 1590		
Thr Gly Cys Thr Thr Gly Cys Cys Ala Ala Gly Cys Cys Ala Gly 1595 1600 1605		
Ala Gly Cys Ala Thr Gly Ala Gly Ala Ala Thr Cys Gly Gly Ala 1610 1615 1620		
Thr Cys Ala Gly Cys Cys Ala Thr Ala Thr Cys Thr Gly Cys Ala 1625 1630 1635		
Cys Thr Gly Ala Cys Ala Ala Cys Gly Thr Gly Ala Ala Gly Ala 1640 1645 1650		
Cys Ala Gly Gly Cys Ala Thr Cys Gly Cys Cys Ala Ala Cys Ala 1655 1660 1665		
Cys Cys Cys Thr Gly Gly Gly Ala Ala Ala Cys Ala Ala Gly Gly 1670 1675 1680		
Gly Ala Gly Cys Ala Gly Cys Gly Gly Gly Ala Gly Thr Gly Thr 1685 1690 1695		
Cys Cys Thr Thr Cys Ala Thr Gly Thr Thr Cys Ala Thr Thr Gly 1700 1705 1710		
Gly Ala Ala Cys Cys Thr Cys Cys Thr Thr Gly Gly Gly Gly Thr 1715 1720 1725		



-continued

---

Thr	Cys	Gly	Thr	Cys	Ala	Ala	Cys	Ala	Gly	Cys	Cys	Ala	Cys	Thr
1730						1735					1740			
Thr	Gly	Ala	Cys	Thr	Thr	Cys	Thr	Gly	Gly	Ala	Ala	Gly	Thr	Gly
1745						1750					1755			
Ala	Ala	Ala	Ala	Ala	Ala	Ala	Gly	Cys	Thr	Cys	Ala	Gly	Gly	Ala
1760						1765					1770			
Gly	Ala	Ala	Ala	Thr	Cys	Ala	Ala	Ala	Ala	Cys	Thr	Ala	Thr	Ala
1775						1780					1785			
Thr	Gly	Ala	Ala	Cys	Ala	Thr	Cys	Cys	Thr	Gly	Cys	Gly	Gly	Thr
1790						1795					1800			
Thr	Cys	Cys	Thr	Gly	Gly	Cys	Cys	Cys	Thr	Gly	Gly	Gly	Ala	Gly
1805						1810					1815			
Ala	Cys	Ala	Ala	Gly	Ala	Ala	Gly	Cys	Thr	Ala	Ala	Gly	Cys	Cys
1820						1825					1830			
Cys	Ala	Thr	Thr	Thr	Ala	Ala	Cys	Ala	Thr	Cys	Ala	Cys	Cys	Cys
1835						1840					1845			
Ala	Cys	Cys	Gly	Cys	Thr	Thr	Cys	Ala	Cys	Cys	Cys	Ala	Cys	Cys
1850						1855					1860			
Thr	Cys	Thr	Thr	Cys	Thr	Gly	Gly	Cys	Thr	Thr	Gly	Gly	Gly	Gly
1865						1870					1875			
Ala	Thr	Cys	Thr	Cys	Ala	Ala	Cys	Thr	Ala	Cys	Cys	Gly	Cys	Gly
1880						1885					1890			
Thr	Gly	Gly	Ala	Gly	Cys	Thr	Gly	Cys	Cys	Cys	Ala	Cys	Thr	Thr
1895						1900					1905			
Gly	Gly	Gly	Ala	Gly	Gly	Cys	Ala	Gly	Ala	Gly	Gly	Cys	Cys	Ala
1910						1915					1920			
Thr	Cys	Ala	Thr	Cys	Cys	Ala	Gly	Ala	Ala	Gly	Ala	Thr	Cys	Ala
1925						1930					1935			
Ala	Gly	Cys	Ala	Ala	Cys	Ala	Gly	Cys	Ala	Gly	Thr	Ala	Thr	Thr
1940						1945					1950			
Cys	Ala	Gly	Ala	Cys	Cys	Thr	Thr	Cys	Thr	Gly	Gly	Cys	Cys	Cys
1955						1960					1965			
Ala	Cys	Gly	Ala	Cys	Cys	Ala	Ala	Cys	Thr	Gly	Cys	Thr	Cys	Cys
1970						1975					1980			
Thr	Gly	Gly	Ala	Gly	Ala	Gly	Gly	Ala	Ala	Gly	Gly	Ala	Cys	Cys
1985						1990					1995			
Ala	Gly	Ala	Ala	Gly	Gly	Thr	Cys	Thr	Thr	Cys	Cys	Thr	Gly	Cys
2000						2005					2010			
Ala	Cys	Thr	Thr	Thr	Gly	Ala	Gly	Gly	Ala	Gly	Gly	Ala	Ala	Gly
2015						2020					2025			
Ala	Gly	Ala	Thr	Cys	Ala	Cys	Cys	Thr	Thr	Cys	Gly	Cys	Cys	Cys
2030						2035					2040			
Cys	Cys	Ala	Cys	Cys	Thr	Ala	Thr	Cys	Gly	Ala	Thr	Thr	Thr	Gly
2045						2050					2055			
Ala	Ala	Ala	Gly	Ala	Cys	Thr	Gly	Ala	Cys	Cys	Cys	Gly	Gly	Gly
2060						2065					2070			
Ala	Cys	Ala	Ala	Gly	Thr	Ala	Thr	Gly	Cys	Ala	Thr	Ala	Cys	Ala
2075						2080					2085			
Cys	Gly	Ala	Ala	Gly	Cys	Ala	Gly	Ala	Ala	Ala	Gly	Cys	Ala	Ala
2090						2095					2100			
Cys	Ala	Gly	Gly	Gly	Ala	Thr	Gly	Ala	Ala	Gly	Thr	Ala	Cys	Ala
2105						2110					2115			
Ala	Cys	Thr	Thr	Gly	Cys	Cys	Gly	Thr	Cys	Cys	Thr	Gly	Gly	Thr
2120						2125					2130			

-continued

---

Gly Cys 2135	Gly Ala Cys Cys 2140	Ala Gly Thr Cys Cys 2145	Thr Cys Thr
Gly Gly 2150	Ala Ala Gly Thr Cys 2155	Thr Thr Ala Cys Cys 2160	Cys Gly Cys
Thr Gly 2165	Gly Thr Gly Cys Ala 2170	Thr Gly Thr Gly Gly 2175	Thr Cys Thr
Gly Thr 2180	Cys Ala Gly Thr Cys 2185	Cys Thr Ala Thr Gly 2190	Gly Cys Ala
Gly Thr 2195	Ala Cys Cys Ala Gly 2200	Thr Gly Ala Cys Ala 2205	Thr Cys Ala
Thr Gly 2210	Ala Cys Gly Ala Gly 2215	Thr Gly Ala Cys Cys 2220	Ala Cys Ala
Gly Cys 2225	Cys Cys Thr Gly Thr 2230	Cys Thr Thr Thr Gly 2235	Cys Cys Ala
Cys Gly 2240	Thr Thr Thr Gly Ala 2245	Ala Gly Cys Ala Gly 2250	Gly Ala Gly
Thr Cys 2255	Ala Cys Ala Thr Cys 2260	Thr Cys Ala Ala Thr 2265	Thr Cys Gly
Thr Cys 2270	Thr Cys Cys Ala Ala 2275	Gly Ala Ala Thr Gly 2280	Gly Thr Cys
Cys Thr 2285	Gly Gly Cys Ala Cys 2290	Thr Gly Thr Ala Gly 2295	Ala Thr Ala
Gly Cys 2300	Cys Ala Ala Gly Gly 2305	Gly Cys Ala Gly Ala 2310	Thr Cys Gly
Ala Gly 2315	Thr Thr Thr Cys Thr 2320	Thr Gly Cys Ala Thr 2325	Gly Cys Thr
Ala Cys 2330	Gly Cys Cys Ala Cys 2335	Ala Cys Thr Gly Ala 2340	Ala Gly Ala
Cys Cys 2345	Ala Ala Gly Thr Cys 2350	Cys Cys Ala Gly Ala 2355	Cys Thr Ala
Ala Gly 2360	Thr Thr Cys Thr Ala 2365	Cys Thr Thr Gly Gly 2370	Ala Gly Thr
Thr Cys 2375	Cys Ala Cys Thr Cys 2380	Ala Ala Gly Cys Thr 2385	Gly Cys Thr
Thr Ala 2390	Gly Ala Gly Ala Gly 2395	Thr Thr Thr Thr Gly 2400	Thr Cys Ala
Ala Gly 2405	Ala Gly Thr Cys Ala 2410	Gly Gly Ala Ala Gly 2415	Gly Ala Gly
Ala Gly 2420	Ala Ala Thr Gly Ala 2425	Ala Gly Ala Gly Gly 2430	Gly Ala Ala
Gly Thr 2435	Gly Ala Ala Gly Gly 2440	Ala Gly Ala Gly Cys 2445	Thr Gly Gly
Thr Gly 2450	Gly Thr Ala Cys Gly 2455	Gly Thr Thr Thr Gly 2460	Gly Ala Gly
Ala Gly 2465	Ala Cys Thr Cys Thr 2470	Thr Cys Cys Cys Ala 2475	Ala Gly Cys
Thr Ala 2480	Ala Ala Gly Cys Cys 2485	Cys Ala Thr Thr Ala 2490	Thr Cys Thr
Cys Thr 2495	Gly Ala Cys Cys Cys 2500	Cys Gly Ala Gly Thr 2505	Ala Cys Thr
Thr Ala 2510	Cys Thr Gly Gly Ala 2515	Cys Cys Ala Gly Cys 2520	Ala Thr Ala
Thr Cys	Cys Thr Gly Ala Thr	Cys Ala Gly Cys Ala	Thr Thr Ala

-continued

2525	2530	2535
Ala Ala Thr Cys Cys Thr Cys Thr Gly Ala Cys Ala Gly Thr Gly	2545	2550
Ala Cys Gly Ala Gly Thr Cys Cys Thr Ala Thr Gly Gly Thr Gly	2560	2565
Ala Ala Gly Gly Cys Thr Gly Cys Ala Thr Thr Gly Cys Cys Cys	2575	2580
Thr Thr Cys Gly Cys Thr Thr Gly Gly Ala Gly Ala Cys Cys Ala	2590	2595
Cys Ala Gly Ala Gly Gly Cys Thr Cys Ala Gly Cys Ala Thr Cys	2605	2610
Cys Thr Ala Thr Cys Thr Ala Cys Ala Cys Gly Cys Cys Thr Cys	2620	2625
Thr Cys Ala Cys Cys Cys Ala Cys Cys Ala Thr Gly Gly Gly Gly	2635	2640
Ala Gly Ala Thr Gly Ala Cys Thr Gly Gly Cys Cys Ala Cys Thr	2650	2655
Thr Cys Ala Gly Gly Gly Gly Ala Gly Ala Gly Ala Thr Thr Ala	2665	2670
Ala Gly Cys Thr Gly Cys Ala Gly Ala Cys Cys Thr Cys Cys Cys	2680	2685
Ala Gly Gly Gly Cys Ala Ala Gly Ala Thr Gly Ala Gly Gly Gly	2695	2700
Ala Gly Ala Ala Gly Cys Thr Cys Thr Ala Thr Gly Ala Cys Thr	2710	2715
Thr Thr Gly Thr Gly Ala Ala Gly Ala Cys Ala Gly Ala Gly Cys	2725	2730
Gly Gly Gly Ala Thr Gly Ala Ala Thr Cys Cys Ala Gly Thr Gly	2740	2745
Gly Ala Ala Thr Gly Ala Ala Ala Thr Gly Cys Thr Thr Gly Ala	2755	2760
Ala Gly Ala Ala Cys Cys Thr Cys Ala Cys Cys Ala Gly Cys Cys	2770	2775
Ala Thr Gly Ala Cys Cys Cys Thr Ala Thr Gly Ala Gly Gly Cys	2785	2790
Ala Ala Thr Gly Gly Gly Ala Gly Cys Cys Thr Thr Cys Thr Gly	2800	2805
Gly Cys Ala Gly Gly Gly Thr Cys Cys Cys Thr Gly Cys Ala Thr	2815	2820
Gly Thr Gly Gly Thr Gly Thr Cys Thr Cys Cys Ala Gly Cys Cys	2830	2835
Thr Cys Ala Ala Thr Gly Ala Gly Ala Thr Gly Ala Thr Cys Ala	2845	2850
Ala Thr Cys Cys Ala Ala Ala Cys Thr Ala Cys Ala Thr Thr Gly	2860	2865
Gly Thr Ala Thr Gly Gly Gly Gly Cys Cys Thr Thr Thr Thr Gly	2875	2880
Gly Ala Cys Ala Gly Cys Cys Cys Cys Thr Gly Cys Ala Thr Gly	2890	2895
Gly Gly Ala Ala Ala Thr Cys Ala Ala Cys Cys Cys Thr Gly Thr	2905	2910
Cys Cys Cys Cys Ala Gly Ala Thr Cys Ala Gly Cys Ala Ala Cys	2920	2925

-continued

---

Thr	Cys	Ala	Cys	Ala	Gly	Cys	Thr	Thr	Gly	Gly	Ala	Gly	Thr	Thr
2930						2935					2940			
Ala	Thr	Gly	Ala	Cys	Cys	Ala	Gly	Cys	Thr	Ala	Cys	Cys	Cys	Ala
2945						2950					2955			
Ala	Ala	Gly	Ala	Cys	Thr	Cys	Cys	Thr	Cys	Cys	Cys	Thr	Gly	Gly
2960						2965					2970			
Gly	Gly	Cys	Cys	Thr	Gly	Gly	Gly	Ala	Gly	Gly	Gly	Gly	Gly	Gly
2975						2980					2985			
Ala	Gly	Gly	Gly	Thr	Cys	Cys	Thr	Cys	Cys	Ala	Ala	Cys	Cys	Cys
2990						2995					3000			
Cys	Thr	Cys	Cys	Cys	Thr	Cys	Cys	Cys	Ala	Ala	Cys	Cys	Ala	Cys
3005						3010					3015			
Cys	Thr	Cys	Thr	Gly	Thr	Cys	Gly	Cys	Cys	Ala	Ala	Ala	Gly	Ala
3020						3025					3030			
Ala	Gly	Thr	Thr	Thr	Thr	Cys	Ala	Thr	Cys	Thr	Thr	Cys	Cys	Ala
3035						3040					3045			
Cys	Ala	Gly	Cys	Cys	Ala	Ala	Cys	Cys	Gly	Ala	Gly	Gly	Thr	Cys
3050						3055					3060			
Cys	Cys	Thr	Gly	Cys	Cys	Cys	Cys	Ala	Gly	Gly	Gly	Thr	Gly	Cys
3065						3070					3075			
Ala	Ala	Gly	Ala	Gly	Gly	Cys	Ala	Ala	Gly	Ala	Cys	Cys	Thr	Gly
3080						3085					3090			
Gly	Gly	Gly	Ala	Thr	Cys	Thr	Gly	Gly	Gly	Ala	Ala	Ala	Gly	Gly
3095						3100					3105			
Thr	Gly	Gly	Ala	Ala	Gly	Cys	Thr	Cys	Thr	Gly	Cys	Thr	Cys	Cys
3110						3115					3120			
Ala	Gly	Gly	Ala	Gly	Gly	Ala	Cys	Cys	Thr	Gly	Cys	Thr	Gly	Cys
3125						3130					3135			
Thr	Gly	Ala	Cys	Gly	Ala	Ala	Gly	Cys	Cys	Cys	Gly	Ala	Gly	Ala
3140						3145					3150			
Thr	Gly	Thr	Thr	Thr	Gly	Ala	Gly	Ala	Ala	Cys	Cys	Cys	Ala	Cys
3155						3160					3165			
Thr	Gly	Thr	Ala	Thr	Gly	Gly	Ala	Thr	Cys	Cys	Gly	Thr	Gly	Ala
3170						3175					3180			
Gly	Thr	Thr	Cys	Cys	Thr	Thr	Cys	Cys	Cys	Thr	Ala	Ala	Gly	Cys
3185						3190					3195			
Thr	Gly	Gly	Thr	Gly	Cys	Cys	Cys	Ala	Gly	Gly	Ala	Ala	Ala	Gly
3200						3205					3210			
Ala	Gly	Cys	Ala	Gly	Gly	Ala	Gly	Thr	Cys	Thr	Cys	Cys	Cys	Ala
3215						3220					3225			
Ala	Gly	Ala	Thr	Gly	Cys	Thr	Gly	Cys	Gly	Gly	Ala	Ala	Gly	Gly
3230						3235					3240			
Ala	Gly	Cys	Cys	Cys	Cys	Cys	Gly	Cys	Cys	Cys	Thr	Gly	Thr	Cys
3245						3250					3255			
Cys	Ala	Gly	Ala	Cys	Cys	Cys	Ala	Gly	Gly	Ala	Ala	Thr	Cys	Thr
3260						3265					3270			
Cys	Ala	Thr	Cys	Ala	Cys	Cys	Cys	Ala	Gly	Cys	Ala	Thr	Cys	Gly
3275						3280					3285			
Thr	Gly	Cys	Thr	Cys	Cys	Cys	Cys	Ala	Ala	Ala	Gly	Cys	Cys	Cys
3290						3295					3300			
Ala	Ala	Gly	Ala	Gly	Gly	Thr	Gly	Gly	Ala	Gly	Ala	Gly	Thr	Gly
3305						3310					3315			
Thr	Cys	Ala	Ala	Gly	Gly	Gly	Gly	Ala	Cys	Ala	Ala	Gly	Cys	Ala
3320						3325					3330			

-continued

---

Ala	Ala	Cys	Ala	Gly	Gly	Cys	Cys	Cys	Cys	Thr	Gly	Thr	Gly	Cys
3335						3340					3345			
Cys	Thr	Gly	Thr	Cys	Cys	Thr	Thr	Gly	Gly	Cys	Cys	Cys	Cys	Ala
3350						3355					3360			
Cys	Ala	Cys	Cys	Cys	Cys	Gly	Gly	Ala	Thr	Cys	Cys	Gly	Cys	Thr
3365						3370					3375			
Cys	Cys	Thr	Thr	Thr	Ala	Cys	Cys	Thr	Gly	Thr	Thr	Cys	Thr	Thr
3380						3385					3390			
Cys	Thr	Thr	Cys	Thr	Gly	Cys	Thr	Gly	Ala	Gly	Gly	Gly	Cys	Ala
3395						3400					3405			
Gly	Ala	Ala	Thr	Gly	Ala	Cys	Cys	Ala	Gly	Thr	Gly	Gly	Gly	Gly
3410						3415					3420			
Ala	Cys	Ala	Ala	Gly	Ala	Gly	Cys	Cys	Ala	Ala	Gly	Gly	Gly	Ala
3425						3430					3435			
Ala	Gly	Cys	Cys	Cys	Ala	Ala	Gly	Gly	Cys	Cys	Thr	Cys	Ala	Gly
3440						3445					3450			
Cys	Cys	Ala	Gly	Thr	Thr	Cys	Cys	Cys	Ala	Ala	Gly	Cys	Cys	Cys
3455						3460					3465			
Cys	Ala	Gly	Thr	Gly	Cys	Cys	Ala	Gly	Thr	Cys	Ala	Ala	Gly	Ala
3470						3475					3480			
Gly	Gly	Cys	Cys	Thr	Gly	Thr	Cys	Ala	Ala	Gly	Cys	Cys	Thr	Thr
3485						3490					3495			
Cys	Cys	Ala	Gly	Gly	Thr	Cys	Ala	Gly	Ala	Ala	Ala	Thr	Gly	Ala
3500						3505					3510			
Gly	Cys	Cys	Ala	Gly	Cys	Ala	Gly	Ala	Cys	Ala	Ala	Cys	Ala	Cys
3515						3520					3525			
Cys	Cys	Ala	Thr	Cys	Cys	Cys	Ala	Gly	Cys	Thr	Cys	Cys	Ala	Cys
3530						3535					3540			
Gly	Gly	Cys	Cys	Ala	Cys	Cys	Cys	Cys	Thr	Gly	Cys	Cys	Ala	Gly
3545						3550					3555			
Thr	Cys	Ala	Ala	Gly	Ala	Gly	Thr	Cys	Cys	Thr	Gly	Cys	Thr	Gly
3560						3565					3570			
Thr	Cys	Cys	Thr	Gly	Cys	Ala	Gly	Cys	Thr	Gly	Cys	Ala	Ala	Cys
3575						3580					3585			
Ala	Thr	Thr	Cys	Cys	Ala	Ala	Ala	Gly	Gly	Cys	Ala	Gly	Ala	Gly
3590						3595					3600			
Ala	Cys	Thr	Ala	Cys	Cys	Gly	Thr	Gly	Ala	Cys	Ala	Ala	Cys	Ala
3605						3610					3615			
Cys	Ala	Gly	Ala	Ala	Cys	Thr	Cys	Cys	Cys	Cys	Cys	Ala	Cys	Cys
3620						3625					3630			
Ala	Thr	Gly	Gly	Cys	Ala	Ala	Gly	Cys	Ala	Cys	Cys	Gly	Cys	Cys
3635						3640					3645			
Ala	Ala	Gly	Ala	Gly	Gly	Ala	Gly	Gly	Gly	Gly	Cys	Thr	Gly	Cys
3650						3655					3660			
Thr	Thr	Gly	Gly	Cys	Ala	Gly	Gly	Ala	Cys	Thr	Gly	Cys	Cys	Ala
3665						3670					3675			
Thr	Gly	Cys	Ala	Gly	Thr	Gly	Ala	Gly	Cys	Thr	Gly	Cys	Thr	Gly
3680						3685					3690			
Gly	Thr	Gly	Ala	Thr	Cys	Gly	Gly	Ala	Gly	Cys	Cys	Thr	Gly	Gly
3695						3700					3705			
Ala	Gly	Gly	Ala	Ala	Cys	Ala	Gly	Cys	Ala	Cys	Ala	Ala	Ala	Gly
3710						3715					3720			
Cys	Ala	Gly	Ala	Cys	Cys	Thr	Gly	Cys	Gly	Cys	Cys	Thr	Cys	Thr

3725					3730					3735				
Cys	Thr	Cys	Ala	Gly	Gly	Ala	Thr	Gly	Cys	Cys	Thr	Cys	Thr	Cys
	3740					3745					3750			
Thr	Cys	Ala	Gly	Gly	Ala	Thr	Gly	Cys	Cys	Thr	Cys	Thr	Thr	Gly
	3755					3760					3765			
Gly	Ala	Gly	Gly	Ala	Cys	Cys	Thr	Cys	Cys	Thr	Gly	Cys	Thr	Ala
	3770					3775					3780			
Gly	Cys	Thr	Cys	Thr	Thr	Cys	Thr	Thr	Gly	Cys	Cys	Thr	Ala	Gly
	3785					3790					3795			
Cys	Thr	Thr	Cys	Ala	Ala	Gly	Thr	Cys	Cys	Cys	Ala	Gly	Gly	Cys
	3800					3805					3810			
Thr	Gly	Thr	Gly	Thr	Ala	Thr	Thr	Thr	Thr	Thr	Thr	Thr	Thr	Cys
	3815					3820					3825			
Ala	Gly	Gly	Ala	Ala	Ala	Cys	Gly	Gly	Cys	Cys	Thr	Cys	Ala	Cys
	3830					3835					3840			
Thr	Thr	Cys	Thr	Cys	Thr	Gly	Thr	Gly	Thr	Cys	Cys	Ala	Ala	
	3845					3850					3855			
Gly	Ala	Ala	Gly	Thr	Gly	Thr	Gly	Cys	Thr	Gly	Cys	Thr	Gly	Gly
	3860					3865					3870			
Cys	Thr	Gly	Cys	Cys	Ala	Cys	Ala	Cys	Thr	Gly	Thr	Gly	Cys	Gly
	3875					3880					3885			
Gly	Cys	Ala	Gly	Ala	Thr	Gly	Cys	Thr	Ala	Ala	Ala	Gly	Cys	Thr
	3890					3895					3900			
Gly	Gly	Ala	Thr	Gly	Ala	Cys	Ala	Ala	Cys	Gly	Cys	Ala	Cys	
	3905					3910					3915			
Gly	Cys	Cys	Ala	Thr	Ala	Cys	Ala	Ala	Cys	Ala	Gly	Cys	Ala	
	3920					3925					3930			
Gly	Ala	Cys	Ala	Gly	Cys	Gly	Cys	Ala	Cys	Thr	Gly	Gly	Gly	
	3935					3940					3945			
Thr	Cys	Thr	Cys	Ala	Gly	Ala	Cys	Thr	Thr	Gly	Gly	Ala	Thr	
	3950					3955					3960			
Thr	Cys	Cys	Thr	Gly	Gly	Gly	Cys	Cys	Thr	Thr	Cys	Thr	Thr	Cys
	3965					3970					3975			
Cys	Ala	Gly	Thr	Cys	Gly	Cys	Cys	Gly	Thr	Thr	Thr	Ala	Ala	
	3980					3985					3990			
Ala	Gly	Ala	Ala	Ala	Gly	Gly	Ala	Ala	Cys	Thr	Ala	Ala	Cys	Gly
	3995					4000					4005			
Gly	Ala	Gly	Cys	Thr	Gly	Cys	Thr	Cys	Ala	Thr	Cys	Cys	Gly	Ala
	4010					4015					4020			
Gly	Gly	Gly	Thr	Gly	Ala	Ala	Gly	Ala	Thr	Ala	Thr	Ala	Ala	
	4025					4030					4035			
Thr	Ala	Ala	Thr	Ala	Ala	Thr	Thr	Ala	Ala	Thr	Thr	Ala	Ala	
	4040					4045					4050			
Thr	Ala	Ala	Thr	Ala	Ala	Thr	Ala	Cys	Ala	Gly	Thr	Cys	Ala	

-continued

---

Gly	Gly	Thr	Gly	Cys	Cys	Ala	Gly	Thr	Cys	Ala	Gly	Cys	Cys	Thr
4130						4135					4140			
Gly	Ala	Ala	Cys	Cys	Cys	Thr	Ala	Thr	Gly	Cys	Cys	Cys	Ala	Gly
4145						4150					4155			
Gly	Cys	Cys	Cys	Ala	Cys	Thr	Ala	Ala	Thr	Cys	Cys	Cys	Ala	Ala
4160						4165					4170			
Ala	Thr	Gly	Gly	Thr	Gly	Gly	Gly	Thr	Cys	Cys	Thr	Gly	Ala	Gly
4175						4180					4185			
Ala	Thr	Gly	Thr	Thr	Thr	Thr	Thr	Ala	Ala	Ala	Ala	Ala	Gly	Cys
4190						4195					4200			
Ala	Thr	Thr	Ala	Ala	Ala	Gly	Ala	Ala	Ala	Ala	Cys	Cys	Ala	Thr
4205						4210					4215			
Cys	Gly	Gly	Thr	Cys	Thr	Cys	Thr	Thr	Ala	Gly	Ala	Gly	Cys	Thr
4220						4225					4230			
Ala	Ala	Cys	Cys	Gly	Gly	Cys	Cys	Gly	Gly	Gly	Cys	Thr	Cys	Thr
4235						4240					4245			
Ala	Cys	Thr	Gly	Cys	Ala	Gly	Gly	Gly	Ala	Cys	Cys	Cys	Gly	Ala
4250						4255					4260			
Ala	Cys	Ala	Gly	Thr	Cys	Thr	Gly	Cys	Ala	Thr	Gly	Gly	Cys	Thr
4265						4270					4275			
Ala	Ala	Gly	Thr	Gly	Gly	Cys	Ala	Cys	Ala	Ala	Gly	Gly	Ala	Gly
4280						4285					4290			
Cys	Cys	Thr	Gly	Gly	Cys	Cys	Cys	Thr	Gly	Thr	Cys	Cys	Ala	Gly
4295						4300					4305			
Cys	Thr	Thr	Cys	Ala	Gly	Ala	Gly	Ala	Thr	Cys	Cys	Ala	Ala	Gly
4310						4315					4320			
Cys	Thr	Gly	Cys	Thr	Thr	Thr	Thr	Thr	Gly	Cys	Thr	Gly	Gly	Gly
4325						4330					4335			
Gly	Thr	Thr	Cys	Thr	Gly	Thr	Cys	Ala	Cys	Ala	Gly	Gly	Cys	Cys
4340						4345					4350			
Thr	Gly	Ala	Thr	Cys	Cys	Thr	Cys	Thr	Thr	Gly	Gly	Thr	Thr	Thr
4355						4360					4365			
Thr	Thr	Ala	Thr	Gly	Gly	Gly	Gly	Thr	Thr	Thr	Cys	Ala	Ala	Gly
4370						4375					4380			
Thr	Cys	Thr	Gly	Cys	Cys	Ala	Gly	Ala	Gly	Thr	Cys	Ala	Gly	Ala
4385						4390					4395			
Ala	Ala	Thr	Cys	Ala	Gly	Cys	Thr	Cys	Thr	Ala	Ala	Cys	Thr	Cys
4400						4405					4410			
Gly	Cys	Cys	Ala	Gly	Thr	Gly	Ala	Ala	Gly	Ala	Gly	Ala	Thr	Cys
4415						4420					4425			
Thr	Gly	Gly	Cys	Cys	Thr	Thr	Ala	Ala	Cys	Thr	Thr	Ala	Ala	Gly
4430						4435					4440			
Cys	Cys	Ala	Gly	Cys	Cys	Ala	Cys	Gly	Thr	Cys	Ala	Gly	Gly	Cys
4445						4450					4455			
Cys	Cys	Cys	Thr	Gly	Cys	Thr	Gly	Ala	Gly	Cys	Cys	Thr	Ala	Thr
4460						4465					4470			
Gly	Gly	Ala	Cys	Cys	Ala	Ala	Thr	Ala	Ala	Ala	Thr	Ala	Cys	Thr
4475						4480					4485			
Cys	Cys	Cys	Cys	Gly	Thr	Gly	Cys	Cys	Ala	Cys	Thr	Gly	Gly	Ala
4490						4495					4500			
Gly	Gly	Thr	Gly	Gly	Gly	Cys	Ala	Gly	Cys	Thr	Ala	Thr	Cys	Ala
4505						4510					4515			
Cys	Cys	Ala	Thr	Ala	Cys	Cys	Cys	Thr	Gly	Ala	Gly	Thr	Thr	Gly
4520						4525					4530			

-continued

---

Gly	Gly	Cys	Cys	Ala	Ala	Gly	Cys	Cys	Cys	Ala	Cys	Cys	Cys	Cys
4535						4540					4545			
Ala	Cys	Cys	Cys	Cys	Thr	Ala	Cys	Cys	Cys	Thr	Gly	Cys	Ala	Ala
4550						4555					4560			
Cys	Ala	Thr	Thr	Thr	Cys	Thr	Gly	Ala	Thr	Gly	Thr	Trp	Cys	Thr
4565						4570					4575			
Gly	Ala	Gly	Gly	Ala	Ala	Gly	Ala	Gly	Thr	Cys	Thr	Cys	Cys	Ala
4580						4585					4590			
Cys	Cys	Ala	Thr	Ala	Gly	Thr	Cys	Cys	Cys	Cys	Ala	Ala	Gly	Gly
4595						4600					4605			
Gly	Cys	Thr	Gly	Ala	Gly	Thr	Thr	Cys	Thr	Cys	Cys	Ala	Gly	Cys
4610						4615					4620			
Cys	Thr	Gly	Cys	Thr	Ala	Thr	Cys	Ala	Gly	Gly	Gly	Ala	Ala	Gly
4625						4630					4635			
Gly	Thr	Gly	Ala	Gly	Cys	Ala	Thr	Thr	Gly	Gly	Thr	Cys	Cys	Cys
4640						4645					4650			
Ala	Gly	Gly	Cys	Thr	Cys	Thr	Cys	Ala	Ala	Ala	Ala	Thr	Ala	Gly
4655						4660					4665			
Thr	Gly	Cys	Ala	Gly	Cys	Cys	Thr	Cys	Thr	Thr	Cys	Thr	Thr	Cys
4670						4675					4680			
Cys	Cys	Ala	Ala	Gly	Cys	Thr	Cys	Thr	Gly	Gly	Gly	Gly	Thr	Gly
4685						4690					4695			
Cys	Ala	Cys	Cys	Cys	Thr	Gly	Thr	Gly	Thr	Cys	Cys	Thr	Thr	Gly
4700						4705					4710			
Gly	Thr	Thr	Ala	Cys	Cys	Ala	Gly	Gly	Ala	Gly	Ala	Cys	Thr	Ala
4715						4720					4725			
Gly	Gly	Gly	Thr	Thr	Gly	Thr	Gly	Ala	Thr	Ala	Thr	Cys	Thr	Thr
4730						4735					4740			
Thr	Thr	Cys	Thr	Thr	Gly	Thr	Cys	Thr	Thr	Gly	Cys	Thr	Thr	Thr
4745						4750					4755			
Thr	Thr	Gly	Ala	Thr	Ala	Thr	Ala	Thr	Cys	Ala	Gly	Gly	Ala	Thr
4760						4765					4770			
Thr	Ala	Ala	Thr	Gly	Thr	Ala	Gly	Gly	Ala	Ala	Ala	Cys	Cys	Ala
4775						4780					4785			
Gly	Ala	Cys	Cys	Thr	Ala	Gly	Ala	Thr	Thr	Ala	Thr	Thr	Cys	Ala
4790						4795					4800			
Gly	Gly	Ala	Gly	Ala	Gly	Thr	Ala	Gly	Gly	Thr	Ala	Thr	Ala	Thr
4805						4810					4815			
Cys	Cys	Cys	Cys	Thr	Gly	Thr	Gly	Thr	Thr	Thr	Cys	Cys	Cys	Ala
4820						4825					4830			
Gly	Thr	Cys	Thr	Gly	Ala	Gly	Thr	Gly	Ala	Cys	Cys	Ala	Ala	Thr
4835						4840					4845			
Ala	Ala	Ala	Ala	Thr	Thr	Gly	Thr	Gly	Cys	Cys	Thr	Thr	Thr	Cys
4850						4855					4860			
Thr	Ala													
4865														

<210> SEQ ID NO 3  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

gcctgttgtc atccattga



-continued

<210> SEQ ID NO 4  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

ataagttggt gatcttggt 19

<210> SEQ ID NO 5  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

gccacatctg tactgacaa 19

<210> SEQ ID NO 6  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

agacaggcat tgcaaacac 19

<210> SEQ ID NO 7  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

acatcactca ccgcttcac 19

<210> SEQ ID NO 8  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

tcttaactac cgtgtggat 19

<210> SEQ ID NO 9  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

aatacgccta caccaagca 19

<210> SEQ ID NO 10  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

gtaccagcga catcatgac 19

<210> SEQ ID NO 11  
<211> LENGTH: 19  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

gcgacatcat gacgagtga 19

-continued

---

<210> SEQ ID NO 12  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 12  
  
 aggacagatt gagtttctc 19

<210> SEQ ID NO 13  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 13  
  
 ggtgctatgc cacattgaa 19

<210> SEQ ID NO 14  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 14  
  
 gtttggtgag actcttcca 19

<210> SEQ ID NO 15  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 15  
  
 agacggagcg tgatgaatc 19

<210> SEQ ID NO 16  
 <211> LENGTH: 21  
 <212> TYPE: RNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 16  
  
 aaggaaugc guuuacacuu a 21

<210> SEQ ID NO 17  
 <211> LENGTH: 21  
 <212> TYPE: RNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 17  
  
 aaaaauugcg uuacacuuac a 21

<210> SEQ ID NO 18  
 <211> LENGTH: 97  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 18  
  
 tgctgttgac agtgagcgag gcttccagaa gagcatctta tagtgaagcc acagatgtat 60  
 aagatgctct tctggaagcc ctgcctactg cctcgga 97

<210> SEQ ID NO 19  
 <211> LENGTH: 97  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 19

-continued

---

 tgctgttgac agtgagcgag cccatatacac ccaagaagtt tagtgaagcc acagatgtaa 60

acttcttggg tgatatgggc gtgcctactg cctcgga 97

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 20

gcttcagaa gagcatctta t 21

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 22

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 21

gccatatca cccaagaagt tt 22

We claim:

1. A method for improving haematopoietic recovery in a patient in need thereof, comprising:

harvesting megakaryocytes or megakaryocyte progenitors from a patient;

contacting the megakaryocytes or megakaryocyte progenitors with an effective amount of an inhibitor of SH2-containing inositol-5-phosphatase (SHIP) function, thereby expanding the megakaryocytes or megakaryocyte progenitors, wherein the inhibitor of SHIP function is selected from the group consisting of interfering RNA, antisense oligonucleotide, ribozyme, DNase, nucleic acid modifier, PNA, nonstandard nucleic acid, aptamer, decoy, and dominant/negative mutant; and reinfusing the expanded megakaryocytes or megakaryocyte progenitors into the patient.

2. The method of claim 1, wherein said method further comprises determining the amount or concentration of megakaryocytes or megakaryocyte progenitors in the patient before said harvesting, after said reinfusing, or both.

3. The method of claim 1, wherein the inhibitor of SHIP function is an interfering RNA targeted to SHIP mRNA.

4. The method of claim 3, wherein the interfering RNA is targeted to the SHIP enzymatic domain (inositol 5'-phosphatase domain).

5. The method of claim 3, wherein the interfering RNA is targeted to the amino-terminal src-homology domain (SH2).

6. The method of claim 1, wherein the patient is human.

7. The method of claim 1, wherein the inhibitor of SHIP function is an interfering RNA or antisense oligonucleotide, and wherein the interfering RNA or antisense oligonucleotide is targeted to SHIP mRNA.

8. The method of claim 1, wherein the inhibitor of SHIP function is an antisense oligonucleotide targeted to SHIP mRNA.

9. The method of claim 3, wherein the interfering RNA is a small interfering RNA (shRNA).

10. The method of claim 3, wherein the interfering RNA is a small hairpin RNA (siRNA).

11. The method of claim 7, wherein the patient is human.

12. The method of claim 6, wherein the human patient has an anemia.

13. The method of claim 6, wherein the human patient has undergone a bone marrow transplant.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 8,008,273 B2  
APPLICATION NO. : 12/819711  
DATED : August 30, 2011  
INVENTOR(S) : William G. Kerr, Caroline Despons and Lia Elena Perez

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 9,

Line 9, "in a samples" should read --in a sample--.

Lines 38-39, "molecule" or "deoxyribonucleic acid molecule" refers" should read  
--molecule" or "deoxyribonucleic acid molecule" refers--.

Column 10,

Line 63, "the tens" should read --the term--.

Column 12,

Line 55, "polymerase III promoters" should read --polymerase III promoters--.

Column 17,

Line 13, "RNase III-type" should read --RNase III-type--.

Column 19,

Line 2, "complementary' to" should read --complementary to--.

Column 20,

Lines 45-46, "The tem" should read --The term--.

Column 25,

Line 6, "nanoparticles or nanoeapsules" should read --nanoparticles or nanocapsules--.


Column 27,

Line 20, "siRNAs targeting" should read --SiRNAs targeting--.

Column 29,

Line 56, "a 281 by fragment" should read --a 281 bp fragment--.

Signed and Sealed this  
Sixth Day of December, 2011

A handwritten signature in black ink that reads "David J. Kappos". The signature is written in a cursive, flowing style.

David J. Kappos  
*Director of the United States Patent and Trademark Office*

**CERTIFICATE OF CORRECTION (continued)**

Page 2 of 2

**U.S. Pat. No. 8,008,273 B2**

Column 35,

Line 45, “or<sup>3</sup>H” should read --or <sup>3</sup>H--.

Column 40,

Line 43, “which binds to DNA” should read --which binds to DNA--.

Line 56, “detectably labelled” should read --detectably labeled--.

Column 45,

Line 56, “WI littermates” should read --WT littermates--.

Column 82,

Line 37, “RNA (shRNA)” should read --RNA (siRNA)--.

Line 39, “RNA (siRNA)” should read --RNA (shRNA)--.