

1985

Noosing Adult Cormorants for Banding

Geoffrey G. Hogan

Follow this and additional works at: <https://digitalcommons.usf.edu/nabb>

Recommended Citation

Hogan, Geoffrey G. (1985) "Noosing Adult Cormorants for Banding," *North American Bird Bander*. Vol. 10 : Iss. 3 , Article 4.

Available at: <https://digitalcommons.usf.edu/nabb/vol10/iss3/4>

This Contents is brought to you for free and open access by the Searchable Ornithological Research Archive at Digital Commons @ University of South Florida. It has been accepted for inclusion in North American Bird Bander by an authorized editor of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

Noosing Adult Cormorants for Banding

Geoffrey G. Hogan
Department of Biological Sciences
Brock University
St. Catharine's, Ontario L2S 3A1

Noosing is a well known means of capturing birds for banding and marking, although noose designs and capture techniques vary widely (see McNicholl 1983). The noosing technique reported here was used to capture adult Great Cormorants (*Phalacrocorax carbo*) at a breeding colony on Prince Edward Island, during a study of breeding parameters of this species and Double-crested Cormorants (*P. auritus*) from 1976 to 1979 (Hogan 1979). According to Lewis (1929), Double-crested Cormorants were captured for food by Indians with nooses, and adult Pelagic Cormorants (*P. pelagicus*) have been captured with padded leghold traps (Tenaza 1966).

Great Cormorants in North America breed in colonies on seaciff ledges, on the level ground at the tops of cliffs, or on small, coastal islands. They rarely nest in trees (Godfrey 1966, pers. obs.), a habit common to other races of this widespread species elsewhere (Cramp and Simmons 1977). While they often nest apart from other species, they occasionally share nesting colonies with Double-crests.


Between 1976 and 1979 there were 6 Great Cormorant colonies on Prince Edward Island containing a maximum 738 nests (Hogan 1979). By 1983 there were 8 colonies containing 1324 nests (Hogan 1983). All colonies are on sandstone seaciffs ranging in height from 6 to 33 m.

Between 1974 and 1978 I banded 538 Great Cormorant nestlings at the breeding colony on Durell Point, on the east coast of the island. The majority of nests (approx. 95%) were built on the upper sloping shelves of the 12-27 m high cliff and were readily accessible. In addition, 51 adults were captured with a noosing pole modified from a similar technique used by Edgar (1968) for Australian Gannets (*Sula serrator*). A diagram and details of the device are shown in Figure 1. The noosing pole was constructed by a local tinsmith.

Present Address: Department of Biology, University of Prince Edward Island, Charlottetown, Prince Edward Island C1A 4P3

Figure 1. Diagram of noosing pole used to capture adult Great Cormorants.

- A. ½-in thinwall steel tubing, 120 in
- B. ¾-in thinwall steel tubing, 30 in
- C. ½-in tin or steel band
- D. ⅜-in steel cable, 96 in
- E. ⅜-in nipple
- F. ⅜-in copper tubing, 48 in
- G. ¾-in clamp



To capture a bird, the steel cable forming the noose was fully extended by sliding the outer steel tube (to which the cable is attached) forward. The noose was then moved slowly towards the bird and carefully slipped over the bird's head and down to cover the upper wings and breast. If lowered further, the bird simply stepped out of it. Once in position, the noose is tightened by sliding the outer steel tube back to pin the wings firmly to the body. The bird was then either lifted away from the nest or held in place. A helper was required to grab the bird. The entire procedure generally took 5 to 15 minutes. Some birds were captured easily, while others would not tolerate the approach of the noose. Once captured and processed, the birds were released, whereupon it invariably flew to the water, bathed, and returned to the nest, usually within 10 to 15 minutes. Neighboring birds exhibited little disturbance. Aerial predators such as gulls (*Larus spp.*) and Northern Ravens (*Corvus corax*) did not frequent this colony while the observers were present. The technique worked best while small to half-grown chicks were in the nest. Adults were less likely to tolerate the close presence of observers when there were only eggs, and larger chicks scramble from their nests if disturbed and may fall off the cliff.

The technique described here is a safe, low-disturbance method for capturing adult Great Cormorants at their nests. Behavioral differences may exist between cliff-nesting and ground-nesting birds regarding their tolerance to disturbance.

Acknowledgments

Research on both cormorant species was conducted during graduate studies by the author under the supervision of R.D. Morris, Brock University, St. Catharine's, Ontario. Field assistance during noosing operations was provided by Lawrence Gray and Ada Judson. Financial support was provided by the Prince Edward Island Fish and Wildlife Division, a National Research Council of Canada grant to R.D. Morris, and an Ontario Graduate Scholarship to the author. The manuscript was improved on the basis of comments by M.K. McNicholl, Daryl Guignon, and Ian MacQuarrie.

Literature Cited

- Cramp, S., and K.E.L. Simmons (eds.). 1977. The birds of the western Palearctic. Vol. 1. Oxford University Press, Oxford.
- Edgar, R.L. 1968. Catching colonial seabirds for banding. *Bird-Banding* 39:41-43.
- Godfrey, W.E. 1966. The birds of Canada. National Museum Canada Bull. 203.
- Hogan, G.G. 1979. Breeding parameters of Great Cormorants (*Phalacrocorax carbo carbo*) at mixed species colonies on Prince Edward Island, Canada. M.Sc. thesis. Brock University, St. Catharine's, Ontario.
- . 1983. P.E.I. cormorant census, 1983. Report to Fish and Wildlife Division, P.E.I. Department of Community and Cultural Affairs, Charlottetown.
- Lewis, H.F. 1929. The natural history of the Double-crested Cormorant [*Phalacrocorax auritus auritus* (Lesson)]. Ru-Mi-Lou Books, Ottawa.
- McNicholl, M.K. 1983. Use of noosing pole to capture Common Nighthawks. *NABB* 8:104-105.
- Tenaza, R.R. 1966. A method for trapping cormorants. *Bird-Banding* 37:207-208.

(Eastern)

