
January 1991

Field Observations

Field Observations Committee

Follow this and additional works at: <https://digitalcommons.usf.edu/ffn>

Recommended Citation

Field Observations Committee (1991) "Field Observations," *Florida Field Naturalist*. Vol. 19 : Iss. 1 , Article 8.

Available at: <https://digitalcommons.usf.edu/ffn/vol19/iss1/8>

This Note is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in Florida Field Naturalist by an authorized editor of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

Florida Field Naturalist 19(1): 25-30, 1991.

FIELD OBSERVATIONS

Prepared by the Field Observations Committee

LINDA COOPER, JIM COX (compiler, Florida Game and Fresh Water Fish Commission, 620 S. Meridian St., Tallahassee, FL 32399-1600), WAYNE HOFFMAN, PEGGY POWELL, and BILL PRANTY.

The sightings reported here are largely unsubstantiated field observations and have not been reviewed by the FOS Records Committee. As such, the observations should be considered tentative pending a more formal review. We encourage observers to submit appropriate records to the Records Committee *c/o* Jocie Baker (Secretary), 851 N. Surf Rd., #302, Hollywood, FL 33019.

Several conventions have been adopted to save space. A 3-character set of alpha-numeric initials is used to identify contributors within the accounts of individual species. A complete list of observers and corresponding initials is provided at the end. The list of observers is organized alpha-numerically by the 3-character initials. Names of observers for each report are drawn from a master list, and this procedure may result in occasional gaps in the alpha-numeric sequence (e.g., the listing of LA2 without a previous LA1).

Another convention adopted is the use of common names exclusively. Persons interested in scientific names should consult AOU (1983. Check-list of the North American Birds, 6th ed. Washington, D.C.: Am. Ornithol. Union) and revisions as published in *The Auk*. Other uncommon abbreviations used occasionally are: BBS, breeding bird survey; CBC, Christmas Bird Count; m. obs., many observers; NP, national park; NWR, national wildlife refuge; SP, state park; and SRA, state recreation area. Unless necessary, the counties of named locations are omitted.

The Field Observations Committee would like to thank everyone who contributed to this column. Please bring any unreported sightings to our attention by writing to Jim Cox, compiler.

Summer Report: June-August 1990

As has been the case with previous reports for 1990, drought conditions continue to influence distributions and abundances of birds around the state. Extra-range sightings of Snail Kites, certain wading birds, and waterfowl were reported from several northern areas of the state. One of the more noteworthy of these was a Roseate Spoonbill in Columbia Co. There were also high counts of Snail Kites in Polk and Osceola Cos., while no nesting was reported for many areas in the Everglades.

Southerly breeding attempts were noted for several songbirds. Indigo Buntings and Blue Grosbeaks were noted in new areas in central and southern Florida, and there were southerly breeding attempts reported for Hooded Warbler, Eastern Wood-Pee-wee, Belted Kingfisher, Mississippi Kite, and Field Sparrow. Several large concentrations of terns and skimmers were noted at new areas where breeding has not been reported. These areas should be followed closely in future years to monitor the potential development of colonies. A survey of coastal areas from Homosassa (Citrus Co.) to Steinhatchee (Taylor Co.) found no nesting colonies of terns or skimmers along this large coastal stretch.

Species Accounts

COMMON LOON: 1 in winter plumage at Alligator Point (Franklin Co.) on 23 Jun (DM1), somewhat late.

- NORTHERN GANNET:** immature at St. George Island (Franklin Co.) on 14 Jun (DM1), fairly late for area.
- BROWN PELICAN:** 550(!) on Apalachicola River spoil banks (Franklin Co.) on 3 Jul (DM1), large non-breeding aggregation for area.
- WHITE PELICAN:** scattered reports throughout period: high count of 86 at phosphate mines (Polk Co.) on 2 Jun (PF1); 43 at mines as late as 14 Jul (PF1, DF1); 48 at Merritt Island NWR on 29 Jun (DC3); 10 at St. Marks NWR on 19 Jul (CG2); 200 ± at Hickory Mound Impoundment (Taylor Co.) on 26 Jun (JC1), but only 60 ± on 26 Aug (JC3, KN1).
- MAGNIFICENT FRIGATEBIRD:** 200+ at Boca Grande Pass (Lee Co.) on 7 Jun (CF1, DF1), likely associated with new breeding colony off Ft. Myers; 3 reports along Big Bend coast with as many as 3 seen on 6 and 10 Jul (CG2, MB1, DM2); 50+ off Key West on 6 Aug (JO1).
- MASKED BOOBY:** 1 seen 25 miles off Port Canaveral on 10 Jun (DC4, m. obs.), rarely seen.
- LEAST BITTERN:** 1 at Wakulla Springs SP from early May through 27 Jul (DB3) was new for park.
- REDDISH EGRET:** 2 regular through Jun and Jul (DM1) at Lanark Reef (Franklin Co.); 1-2 immatures intermittently at St. Marks NWR from 14 Jun to end of period (CG2, RW2); 6 (2 adults, 4 immatures) at Hickory Mound Impoundment (Taylor Co.) on Aug 26 (JC3, KN1) and present through early Sep (JC1). Breeding attempt seems imminent along Big Bend coast.
- GLOSSY IBIS:** 423 (!) successful nests at Merritt Island NWR (DC3).
- GREAT BLUE HERON, White Morph:** 1 on coast of northern St. Johns Co. on 18 Aug (CH1); 1 at Alligator Point (Franklin Co.) for most of the report period (GS1).
- BLACK-CROWNED NIGHT-HERON:** increased numbers reported at St. Marks NWR (RW2); 12 adults, 2 sub-adults at Fiddlers Point (Wakulla Co.) on 23 Jun (DM1) suggest continued breeding nearby; 13 successful nests at Merritt Island NWR (DC3).
- ROSEATE SPOONBILL:** 1 at Alligator Lake (Columbia Co.) on 24 Jul represents a very rare interior sighting for north Florida (PS3, KG1); ca. 13 at phosphate mines (Polk Co., m. obs.) and 2 north of Interstate 4 (Polk Co.) throughout period (MR1), highest numbers ever for Polk Co.; 60 at Merritt Island NWR on 11 Aug (PF1, DF1) with 6 successful nests (DC3); immature at Carrabelle Beach (Franklin Co.) on 3 Jun (DM1), immature at St. Marks NWR throughout summer (m. obs.), and immature at Hickory Mounds Impoundment (Taylor Co.) on 4 Aug (DS1) were rare northern Gulf coast sightings.
- WOOD STORK:** high count of 60 ± at St. Marks NWR on 15 Jun (CG2); 4 seen in northern Okaloosa Co. on 3 Sep (SD1), rare in northwestern panhandle.
- BLACK-BELLIED WHISTLING-DUCK:** females with broods at phosphate mines in Hardee Co. on 31 Aug (LM2, TP1), first confirmed nesting in area; adults reported in reclaimed wetlands in Polk Co. all summer (SP1).
- FULVOUS WHISTLING-DUCK:** 10 at St. Marks NWR until late Jun with 3 remaining through end of period (CG1, DS1); 12 in Pensacola on 9 Jun (AF1); 38 at Crew's Lake (Pasco Co.) in early Jun (BP1, PY1, DR2).
- BLUE-WINGED TEAL:** female with 3 ducklings at Zellwood (Orange Co.) on 18 Aug (PF1, m.obs.), irregular nester; 2-3 from 2 Jun to 14 Jul at phosphate mines (Polk Co.), unusual in summer (PF1, DF1).
- NORTHERN SHOVELER:** 1 at phosphate mines (Polk Co.) on 2 and 9 Jun (PF1), late to leave.
- RING-NECKED DUCK:** pair at Crew's Lake (Pasco Co.) until early Jun when male left; female remained until mid Jul (BP1, DR2). Late group of 4 at phosphate mines (Polk Co.) on 2 Jun (PF1).
- RED-BREASTED MERGANSER:** DF1 and CF1 report 1 at Ft. Myers Beach on 8 Jun; 2 at Ft. DeSoto Park (Pinellas Co.) on 11 Jun; 2 at St. Marks NWR on 13 Jun; 1 at Merritt Island NWR on 24 Jun; and 1 in lower Keys on 26 Jun.

- RUDDY DUCK:** 5 at phosphate mines (Polk Co.) on 2 Jun (PF1), unusual in summer though a rare nester.
- OSPREY:** fledgling production at St. Marks NWR was up to 0.98 young per nest versus an alarmingly low 0.11 last year (JR4); 55 flying (mostly adults) at Lake Disston (Volusia Co.) on 17 Jun (HS1) may indicate a breeding aggregation in area.
- AMERICAN SWALLOW-TAILED KITE:** KM1 reports moderately successful reproduction this year. Fledgling rates of 16 monitored nests were ca. 50% as compared with 39% in 1988 and 81% in 1989. Few nests were found south of Lake Okeechobee, perhaps a result of drought conditions. The Corkscrew Swamp (Lee Co.) roost peaked at 317; the Okeechobee roost (Glades Co.) peaked around 25 Jul with 600-1000 (!). Four over south Jacksonville on 6 Jul (JC2) were uncommon for the area.
- BLACK-SHOULDERED KITE:** 2 nesting at Broward/Palm Beach Co. line on 9 Jul (DF1, CF1); 1 fledged 4 young (fide JB1); another nest fledged 3 young west of U.S. 27 near Sawgrass Fish Camp (JB1).
- SNAIL KITE:** sightings shifted north as a result of drought: Big Lake Tohopekaglia (Osceola Co.) had 87 birds, East Lake Tohopekaglia had 10 birds, and Lake Kissimmee (Polk Co.) had 72 birds. Most successful nesting limited to Lake Kissimmee where 58 nests were found (JR3); 2 nests on former phosphate lands near Lehigh Acres (Lee Co.) fledged young; nesting attempts along the lower St. Johns River failed. No nesting in Water Conservation Areas and fewer nest starts than usual at Lake Okeechobee (JR3). Single bird on Conservation Area 2A on 1 Aug (JB1) was only record for Everglades. Several sightings from other central Florida lakes.
- MISSISSIPPI KITE:** first confirmed nesting for Ocala NF and Marion Co. on 11 Jun (JS4, m. obs.).
- COOPER'S HAWK:** several reports for this secretive breeding species from around the state (m. obs.) lead to speculation that its numbers are increasing.
- SHORT-TAILED HAWK:** 3 sightings from Dixie Co.: 1 in Jena WMA on 6 Jun (BM2); 1 in California Swamp on 10 Jun (JC3, KN1); another at Jena WMA on 6 Jul (JR5, SR1).
- WILD TURKEY:** a female/immature tried "city life" for a while along heavily trafficked road near Bay Point (Bay Co.). The bird was first seen on 13 Jul and stayed in a woodlot across from a golf course for about 5 days (TM1, m. obs.).
- BLACK RAIL:** 2 called along Palm Beach/Broward Co. line on 30 Aug (JB1).
- SANDHILL CRANE:** 4 in extreme western Broward Co. on 27 Jun (JB1) may represent breeding attempts in new area.
- PIPING PLOVER:** 1 at St. George Island on 17 Jul (DM1) was early.
- SEMPALMATED PLOVER:** high count of 18 at Alligator Point (Franklin Co.) on 6 Jul (DM1); 12 at phosphate mines (Polk Co., PF1) on 9 Jun.
- BLACK-BELLIED PLOVER:** 1 in basic plumage in central Broward Co. on 11 Jun (JB1), probably late to leave; 1 in alternate plumage in Franklin Co. on 10 Jul (HS1, DM1), probably a returning migrant.
- LESSER GOLDEN PLOVER:** 1 in alternate plumage south of South Bay (Palm Beach Co.) on 25 Aug (DF1, CF1), infrequent migrant.
- BLACK-NECKED STILT:** 6 at Crew's Lake (Pasco Co.) on 6 Jun, disappeared the next day (PY1, DR2).
- AMERICAN AVOCET:** 2 in alternate plumage at phosphate mines (Polk Co.) on 2 Jun (PF1) seemed late; 3 in Homeland (Polk Co.) on 30 Aug (BC2, LC2) were early.
- WILLET:** fairly large aggregation of 185 at Lanark Reef (Franklin Co.) on 10 Jul (HS1, DM1).
- UPLAND SANDPIPER:** 11 south of South Bay (Palm Beach Co.) on 25 Aug (DF1, CF1), good number.
- LONG-BILLED CURLEW:** 1 on Shell Island (Bay Co.) on 26 Aug (JO2, LO1).
- MARbled GODWIT:** high count of 105 at Lanark Reef (Franklin Co.) on 10 Jul (HS1, DM1) with at least 30 present through summer.

- RED KNOT: 80-130 at Alligator Point throughout Jun-Jul (DM1), large numbers for summer.
- WHITE-RUMPED SANDPIPER: high count of 12 (!) at phosphate mines (Polk Co.) on 2 Jun (PF1); 1 at Hickory Mound Impoundment (Taylor Co.) on 10 Jun (DM1).
- PECTORAL SANDPIPER: 35 at sewage treatment ponds south of Tallahassee on 14 Jul (DM1, RW1), good concentration.
- DUNLIN: 2 at Lanark Reef (Franklin Co.) on 10 Jul (HS1, DM1), 1 in alternate plumage.
- STILT SANDPIPER: leucistic bird at Zellwood farms (Orange Co.) from 18-25 Aug (MB2, m. obs.); was it seen anywhere else during migration?
- SHORT-BILLED DOWITCHER: 150 ± at Lanark Reef (Franklin Co.) on 1 and 16 Jul (DM1), high summer counts for area.
- LONG-BILLED DOWITCHER: 1 in alternate plumage at St. Marks NWR on 14 Jul (NW1), heard calling in flight.
- AMERICAN WOODCOCK: several late summer reports around Tallahassee: 1 in southwest portion of town on 29 Jun (DB3); 3 at a small city park on 25 Aug (JC3).
- COMMON SNIBE: 1 at River Ranch (Polk Co.) on 30 Aug was very early (BC2).
- LAUGHING GULL: 4000+ adults and juveniles roosted at Lanark Reef on 17 Jul (DM1, RW1), site of only nesting colony in Franklin Co. The colony has grown to ca. 400 nests and appears to be so large as to inhibit nesting by Least Terns and Black Skimmers. Some of the gulls likely moved from St. Marks NWR as reduced numbers were reported there (CG2). In addition, 1 on nest and 1 feeding a downy juvenile at Wards Bank (Duval Co.) on 2 Jul (LB1), rare nesting report for area.
- GULL SPECIES: a confusing gull, which may be a Western, Lesser Black-backed, or hybrid gull, appeared (again) along the Pensacola waterfront on 30 Aug; known locally as "old one foot," the gull is estimated to be 16 years old (RD1).
- TERNS: aerial surveys along the Gulf coast from Homosassa (Citrus Co.) to Steinhatchee (Taylor Co.) found no colonies of terns or skimmers (JR3).
- GULL-BILLED TERN: 4 nests and 5 downy young at Hillsborough Co. phosphate mine (PF1, BC2) are first inland nesting records for region; 2 at St. Marks NWR on 11 Jul (HH1) were uncommon.
- SANDWICH TERN: 45 at Apalachicola Bay spoil bank (Franklin Co.) on 3 Jul and 60 at Lanark Reef (Franklin Co.) on 17 Jul (DM1), large non-breeding aggregations.
- FORSTER'S TERN: 500 roosted at Lanark Reef (Franklin Co.) on 17 Jul (DM1), high number.
- COMMON TERN: 1 at St. Marks NWR on 12 Jul (NW1), first fall report.
- LEAST TERN: North Florida: colony along St. George Island causeway (Franklin Co.) has grown from 150 to about 700+ pairs. Colony benefitted from habitat improvement, site posting, and educational brochures. Least tern colony at St. Marks NWR grew to 13 pairs and fledged 15 chicks (JR4) from specially constructed platforms. At least 6 small colonies produced young in Duval and St. Johns Counties, the largest being Wards Bank (Duval Co.) and Matanzas National Monument (St. Johns Co.) with 40 ± active nests each (LB1, PP1). South Florida: colony of 300+ nested (DS3) on a building in Lake Wales; 646 (!) at phosphate mines (Polk Co.) on 14 Jul (PF1, DF1) for record high count, but nesting in area limited to a small colony; 2 small colonies found on Merritt Island NWR on 20 Jun (DS2); 13 nesting colonies found in Broward Co. with 9 on rooftops and 4 at construction sites (JB1).
- BRIDLED TERN: 1 seen 25 miles off Port Canaveral (DC4, m. obs.) on 10 Jun, rarely reported.
- BLACK TERN: high count of 1,650 (!) at phosphate mines (Polk Co.) on 26 Aug (PF1); 175 roosting on Lanark Reef (Franklin Co.) on 1 Jul (DM1).
- BLACK SKIMMER: 820 reported at phosphate mines (Polk Co.) on 26 Aug (PF1), second highest interior count; 204 pairs nested on St. George Island causeway (see Least Tern).

- WHITE-WINGED DOVE: 1 seen along Broward/Collier Co. line (JB1) on 29 Jun, new for area.
- EURASIAN COLLARED-DOVE: careful observation of 1 at St. Marks NWR on 10 Jun (DJ1, SJ3), few reports for area.
- PSITTACINES: 2 *Amazona* spp. in Pensacola on 3 Aug were subsequently seen on several occasions (AF1); they were joined by 2 *Aratinga* spp. on 17 Aug.
- CHUCK-WILL'S-WIDOW: first migrant at Key West reported on 8 Aug (JO1).
- BELTED KINGFISHER: 1 at Lake Alfred (Polk Co.) on 14 Jun (PT2) was unusual, even though 1 nesting record for area; 1 at Merritt Island NWR on 6 Jul was rare (KB1, MH3); 1 in western Broward Co. on 25 Jul (JB1) and 1 at Key West on 30 Jul (JO1) were first "fall" reports for south Florida.
- RED-COCKADED WOODPECKER: a random sample of 50 of 186 colonies classified as active in 1989 by the U.S. Forest Service was re-sampled by FJ1, RW1, and others. They found 13 inactive, 14 with single birds, and 23 with breeding pairs/clans. The U.S. Forest Service re-visited areas that had not been studied since 1981 and found 8 active colonies not included in their 1989 inventory of active colonies (R. Costa and co-workers). The area studied by both groups contains about 30% of the active colonies in the Apalachicola National Forest.
- WOODPECKER: while conducting a BBS in Levy Co., JC3 had 6 species of woodpeckers calling at 1 3-minute census station (sans red-cockaded)!
- EASTERN KINGBIRD: an interesting recent phenomenon is nesting on metal power poles and transmission substations on former phosphate mines in Polk Co. (PF1, LC2); large flock of 500+ near Avon Park Bombing Range (Polk Co.) on 28 Aug (DF1, CF1).
- GRAY KINGBIRD: 13 pairs on Alligator Point (Franklin Co.) on 16-17 Jul (DM1), possibly more common in area than suspected.
- EASTERN WOOD-PEWEE: pair near Ocala on 8 Jun (JS4, WB1), nesting south of approximately Gainesville is rare.
- EASTERN PHOEBE: a late Jul nest at bridge near Laurel Hill (Okaloosa Co.) where first state record found (DW1).
- PURPLE MARTIN: roost of 1500-1800 at St. Marks NWR on 28 Jul (CG2); roosts of a few score began collecting around Key West ca. 30 Jul and increased to 1000+ by 29 Aug.
- BARN SWALLOW: first nest recorded in Highlands Co. on 6 Jun (FL1, MS2); 2 adults in Okeechobee Co. on 6 Jun (GM2); 6 juveniles, 2 adults at Key West on 3 Jun (JO1).
- BLUE-GRAY GNATCATCHER: 1st nest at Archbold Biological Station on 5 Jun (RM1).
- EASTERN BLUEBIRD: young hatched from a late nest near High Springs (Alachua Co.) on 8 Aug and fed through end of Aug (MA1).
- AMERICAN ROBIN: 1 sang from late Apr through 9 Jul in Gainesville (GK1); 1 at Mexico Beach (Bay Co.) on 9 Jun (m. obs.) was late.
- BLUE-WINGED WARBLER: 1 near Wakulla River (Wakulla Co.) on 24 Aug was early (JC1).
- GOLDEN-WINGED WARBLER: 1 near Wakulla River (Wakulla Co.) on 24 Aug also somewhat early (JC1).
- YELLOW WARBLER: 1 at St. Marks NWR on 12 Jul (NW1) and 1 in western Broward Co. on 28 Jul (JB1) were earliest reports.
- CHESTNUT-SIDED WARBLER: 1 in rural Alachua Co. on 21 Aug (JH2) and 1 at Saddle Creek Park (Polk Co.) on 26 Aug (PT2) were early.
- CAPE MAY WARBLER: 1 in western Broward Co. on 28 Jul was very early (JB1).
- PRAIRIE WARBLER: early stages of commercial pine plantations may be helping this species extend its range; recorded in new areas of northern Wakulla Co. (HS1) and northwestern Taylor Co. (JC3) at several times in early Jun.
- PALM WARBLER: early migrant at Myakka SP on 24 Aug (BC2, LC2).
- PROTHONOTARY WARBLER: early migrant near Venus (Highlands Co.) on 12 Jul (DS4); 11 at Saddle Creek Park (Polk Co.) on 21 Jul (PF1).

- SWAINSON'S WARBLER: 1 at San Felasco Hammock State Preserve (Alachua Co.) on 31 Aug (RR1); rarely seen here.
- KENTUCKY WARBLER: adult with juvenile in southern Suwannee Co. on 27 Jul (BP1) was far east of published breeding range.
- HOODED WARBLER: first confirmed breeding for Levy Co. on Lower Suwannee River NWR on 4 Jul (m. obs.).
- YELLOW-BREASTED CHAT: adults with fledglings ca. 10 miles south of Chiefland (Levy Co.) on 4 Jul (DH2, DF3), first confirmed nesting for Co.
- ROSE-BREASTED GROSBEAK: 1 at Saddle Creek Park (Polk Co.) on 25 Aug (PF1) was early.
- BLUE GROSBEAK: several southerly breeding season records: 5 singing males in Highlands and Okeechobee Cos. (BK1, MK1, GM2, DS4) from Jun-Jul; 8 males and 1 female in Brevard and Volusia Cos. on 13, 15 Jun (DS2); 5 birds at Zellwood (Orange Co.) on 4 Aug (DF1, CF1, PF1); 2 pairs in northeastern Pinellas Co. on 16 Jun (LH1).
- PAINTED BUNTING: 1 singing male 5 miles southwest of Ocala on 28 Jun (JS4); 9 singing males at Merritt Island NWR on 13 and 15 Jun (DS2); 2 males at Panama City on 30 Jun, 1 begging for food (RI1, AI1, m. obs.); male at St. Marks NWR on 2 Jul (BJ1), an early migrant or pioneer.
- INDIGO BUNTING: late migrants or possible breeders include: 1 singing male at old channel of Kissimmee River (Highlands Co.) on 6 Jun (FL1, GM2, MS2); 7 singing males at Merritt Island NWR on 12, 15 Jun (DS2); 6 at Zellwood (Orange Co.) on 4 Aug (PF1, DF1, CF1).
- FIELD SPARROW: singing male near Trenton on 8 and 20 Jun (Gilchrist Co.) was somewhat south (DM3, BM2, RC2).
- BOBOLINK: very late migrant at Merritt Island NWR on 15 Jun (DS2).
- NORTHERN ORIOLE: 1 at Key West on 9 Aug (JO1) was first of the season.
- SHINY COWBIRD: male at Eastpoint (Franklin Co.) on 23 Jun (DM1) and on 10 Jul (DM1, HS1); 10-12 regular now west of Apalachicola (RW1, HS1, DM1); common throughout period at Key West (JO1); pair at feeder (JB1) in Cooper City (Broward Co.).
- BROWN-HEADED COWBIRD: female at Key West on 20 Aug (JO1) was first of fall season.
- HOUSE FINCH: fledged young in Pensacola on 15 Jun (AF1); fledglings found at several locations around Tallahassee (m. obs.).

Observer abbreviations: Adams, Mary (MA1), Bryan, Dana (DB3), Baker, Jocie (JB1), Bennet, Ken (KB1), Bremer, Linda (LB1), Brothers, Michael (MB1), Biggs, Wes (WB1), Cooper, Buck (BC2), Cooley, Dwight (DC3), Click, Dan (DC4), Cavanagh, Jim (JC1), Cocks, Julie (JC2), Cox, James (JC3), Cooper, Linda (LC2), Duncan, Robert A. (RD1), Duncan, Scot (SD1), Forster, Ann (AF1), Ford, Clarice (CF1), Ford, Don (DF1), Fagan, Dorothy (DF3), Fellers, Paul J. (PF1), Geanangel, Chuck (CG1), Gidden, C.S. (CG2), Greenberg, Katie (KG1), Hough, C. Royce (CH1), Hill, Hugh (HH1), Hintermister, John (JH2), Hopkins, Larry (LH1), Harrel, Mary (MH3), Ingram, Ann (AI1), Ingram, Richard (RI1), Jordan, Blannie (BJ1), Jue, Dean (DJ1), James, Fran (FJ1), Jue, Sally (SJ3), Kittredge, Bruce (BK1), Kiltie, Gracie (GK1), Kittredge, Marion (MK1), Lohrer, Fred (FL1), Millsap, Brian (BM2) McNair, Doug (DM1), Mehlman, David (DM2), Miller, Dorothy (DM3), Myers, Ken (KM1), Menart, Tony (TM1), NeSmith, Katy (KN1), Ondrejko, Joe (JO1), Oswald, Joe (JO2), Oswald, Lois (LO1), Pranty, Bill (BP1), Powell, Peggy (PP1), Robinson, Don (DR2), Rodgers, Jim (JR3), Reinman, Joe (JR4), Rogers, John (JR5), Rowan, Rex (RR1), Rogers, Steve (SR1), Sandee, Daan (DS1), Simpson, David (DS2), Seamon, Greg (GS1), Stevenson, Henry (HS1), Sharpe, John (JS4), Stapleton, Martin (MS2), Southall, Pete (PS3), Timmer, Pete (PT2), Young, Paul (PY1), Ware, Don (DW1), Wamer, Noel (NW1), West, Rick (RW1), Will, Robin (RW2).