
January 1992

Loggerhead Shrikes Eat Crayfish

Reuven Yosef

Follow this and additional works at: <https://digitalcommons.usf.edu/ffn>

Recommended Citation

Yosef, Reuven (1992) "Loggerhead Shrikes Eat Crayfish," *Florida Field Naturalist*. Vol. 20 : Iss. 3 , Article 5.
Available at: <https://digitalcommons.usf.edu/ffn/vol20/iss3/5>

This Note is brought to you for free and open access by the Searchable Ornithological Research Archive at Digital Commons @ University of South Florida. It has been accepted for inclusion in Florida Field Naturalist by an authorized editor of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

resented by populations with low chances of continued existence, then additional oak scrub needs to be protected despite the current representation of oak scrub on conservation lands. On the other hand, these analyses do help to show the poor representation of upland cover types on conservation lands in comparison to wetland cover types. Increased attention needs to be given to many upland cover types.

LITERATURE CITED

- DAVIS, J. H. 1967. General map of natural vegetation of Florida. University of Florida, Gainesville.
- HARTMAN, B. J. 1978. Description of major terrestrial and wetland habitats of Florida, pages xvi-xix in *Rare and endangered biota of Florida*. Volume 2, Birds (H. W. Kale II, ed.). University Presses of Florida, Gainesville.
- KAUTZ, R. 1984. Criteria for evaluating impacts of development on wildlife habitats. Proc. Annu. Conf. Southeast Assoc. Fish and Wildl. Agenc. 38:121-136.
- KAUTZ, R., G. MAULDIN, AND T. GILBERT. 1991. Mapping Florida wildlife habitat using Landsat Thematic Mapper Imagery. Final Report. Florida Game and Fresh Water Fish Commission, Tallahassee.
- MULLER, J., D. HARDIN, D. JACKSON, S. GATEWOOD, AND N. CAIRE. 1989. Summary report on the vascular plants, animals, and plant communities endemic to Florida. Florida Game and Fresh Water Fish Commission, Nongame Wildlife Program Technical Report 7, Tallahassee.
- MYERS, R., AND J. EWELL (eds.). 1990. *Ecosystems of Florida*. University of Central Florida Press, Orlando.
- TYDAC. 1988. SPANS Spatial Analysis System. Version 4.30. TYDAC Technologies, Inc., Ottawa.

Fla. Field Nat. 20(3):75-76, 1992

LOGGERHEAD SHRIKES EAT CRAYFISH

REUVEN YOSEF

*Dept. of Zoology, Ohio State University
1735 Neil Avenue, Columbus, OH 43210, and
Archbold Biological Station
P.O. Box 2057, Lake Placid, Florida 33852*

Loggerhead Shrikes (*Lanius ludovicianus*) are opportunistic feeders that feed largely on invertebrates (Beal and McAtee 1912, Howell 1932, Craig 1978, Scott and Morrison 1990), although they may consume mammals when insects are scarce (Judd 1898, Kridelbaugh 1982). Graber et al. (1973) showed that shrikes adjust their diet according to prey availability, and even feed on road-kills (Robertson 1930).

John Condit, of the Ohio State University's Museum of Zoology, initially discovered that loggerhead shrikes caught and impaled crayfish (*Procambrus alleni*; Hobbs and Hobbs 1991) at the MacArthur Agro-ecology Center of the Archbold Biological Station, in February 1991. This occurred on the territory of a shrike that had a shallow canal (~ 1 m deep) flowing through its boundaries. It is unusual for shrikes to prey on crustaceans, and to date reports have implied that only isopods have been taken as prey (Scott and Morrison 1990).

In September and October 1991, the author found crayfish remains at the feeding and impaling sites of seven shrikes, all of which had shallow canals in their territories. Twenty three crayfish were found impaled, 18 of which were still alive and moved their legs and claws in an attempt to free themselves. All were impaled on barbed wire, ventral side down, immediately behind the legs at the juncture of the cephalothorax and the abdomen.

I saw shrikes eating crayfish on eight occasions. The shrikes removed the claws and walking feet of the crayfish, prior to further handling and ingestion. They reimpaled the crayfish and pulled at the abdomen so that all visceral organs were pulled out, and then the crayfish was consumed. The whole process usually took about 15 min (\bar{x} = 14.73, SD = 3.5, N = 8). Shrikes consumed the crayfish within three hours from when impaled. On three occasions Boat-tailed Grackles (*Quiscalus major*) stole the impaled crayfish.

Shrikes processed crayfish only at certain impaling sites along the fence-line. Five times I moved crayfish away from where they had been impaled, and the shrikes returned them to their original location. Shrikes impaled all crayfish within a 15 cm radius of a fence post, possibly because impaling them farther away from the fencepost requires the shrike to divide its attention between maintaining its balance and dismemberment of the prey.

I thank Mark Deyrup for help in identification of the crayfish. K. L. Bildstein and M. L. Morrison provided comments on an earlier draft of the manuscript, which improved it considerably. This is contribution No. 6 of the MacArthur Agro-ecology Center.

LITERATURE CITED

- BEAL, F. E. L., AND W. L. MCATEE. 1912. Food of some well-known birds of forest, farm, and garden. U.S. Dept. Agric. Farmer's Bull. 506:1-35.
- CRAIG, R. B. 1978. An analysis of the predatory behavior of the Loggerhead Shrike. Auk 95:221-234.
- GRABER, R. R., J. W. GRABER, AND E. L. KIRK. 1973. Illinois birds: Laniidae. III. Nat. Hist. Surv. Biol. Note 83.
- HOBBS, H. H., AND H. H. HOBBS III. 1991. An illustrated key to the crayfishes of Florida. Fla. Sci. 54:13-24.
- HOWELL, A. H. 1932. Florida bird life. Coward-McCann, Inc., New York.
- JUDD, S. D. 1898. The food of shrikes. U.S. Dept. Agric. Div. Biol. Surv. 9:15-26.
- KRIDELBAUGH, A. L. 1982. An ecological study of Loggerhead Shrikes in central Missouri. M.S. thesis, Univ. Missouri, Columbia.
- ROBERTSON, J. M. 1930. Roads and birds. Condor 32:142-146.
- SCOTT, T. A., AND M. L. MORRISON. 1990. Natural history and management of the San Clemente Loggerhead Shrike. Proc. of the Western Found. of Vert. Zool. Vol. 4:23-57.