

2018

Genome Sequence of PM2-Like Phage Cr39582, Induced from a Pseudoalteromonas sp. Isolated from the Gut of Ciona robusta

Brittany Leigh
University of South Florida

Mya Breitbart
University of South Florida, mya@usf.edu

Hanna Oksanen
University of Helsinki

Dennis Bamford
University of Helsinki

Larry Dishaw
University of South Florida

Follow this and additional works at: https://scholarcommons.usf.edu/msc_facpub


 Part of the [Life Sciences Commons](#)

Scholar Commons Citation

Leigh, Brittany; Breitbart, Mya; Oksanen, Hanna; Bamford, Dennis; and Dishaw, Larry, "Genome Sequence of PM2-Like Phage Cr39582, Induced from a Pseudoalteromonas sp. Isolated from the Gut of Ciona robusta" (2018). *Marine Science Faculty Publications*. 673.

https://scholarcommons.usf.edu/msc_facpub/673

This Article is brought to you for free and open access by the College of Marine Science at Scholar Commons. It has been accepted for inclusion in Marine Science Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact scholarcommons@usf.edu.



Genome Sequence of PM2-Like Phage Cr39582, Induced from a *Pseudoalteromonas* sp. Isolated from the Gut of *Ciona robusta*

Brittany A. Leigh,^{a,b*} Mya Breitbart,^a Hanna M. Oksanen,^c Dennis H. Bamford,^c Larry J. Dishaw^b

^aCollege of Marine Science, University of South Florida, St. Petersburg, Florida, USA

^bDepartment of Pediatrics, Children's Research Institute, University of South Florida, St. Petersburg, Florida, USA

^cMolecular and Integrative Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

ABSTRACT Phage Cr39582 was induced by mitomycin C from *Pseudoalteromonas* sp. strain Cr6751, isolated from a marine invertebrate gut. *Pseudoalteromonas* phage Cr39582 has 85% pairwise nucleotide identity with phage PM2 but lacks sequence homology in the spike protein. This report supports previous bioinformatic identification of corticoviral sequences within aquatic bacterial genomes.

Pseudoalteromonas virus PM2 is currently the only virus species of the *Corticoviridae* family (1). The classification is based on the presence of an internal membrane layer and a supercoiled circular double-stranded DNA (dsDNA) genome of approximately 10 kb (2, 3). PM2 is a virulent marine phage discovered off the coast of Chile in the 1960s (2, 4). Although sequences related to phage PM2 are widespread within the genomes of aquatic proteobacteria (5), it is unknown if these prophages can be experimentally induced (5). To date, seven genomes of virulent phages infecting *Pseudoalteromonas* strains and 31 prophages have been reported (6). Here, we describe the genome sequence of *Pseudoalteromonas* phage Cr39582, a prophage of *Pseudoalteromonas* sp. strain Cr6751 that contains a circular dsDNA genome of 10,584 bp with 20 predicted open reading frames (ORFs). Phage Cr39582 is only the second described PM2-like phage and a potential member of the family *Corticoviridae*; thus, its genome sequence increases our understanding of nontailed dsDNA viruses that may represent an abundant and ecologically important fraction of marine viral communities (7, 8).

Pseudoalteromonas sp. Cr6751 was isolated from the gut of the marine invertebrate *Ciona robusta*, as described previously (9). Bacterial identity was determined based on full-length 16S rRNA gene Sanger sequencing. The presence of active prophages was determined based on exposure of the bacterial culture to 0.5 $\mu\text{g ml}^{-1}$ mitomycin C (10). The culture supernatant was passed through a 0.22- μm filter, and the filtrate was stained with SYBR Gold to enumerate viral-like particles (VLPs) (11). The VLPs were purified by cesium chloride density gradient ultracentrifugation (1.23 g ml^{-1}) and imaged by transmission electron microscopy (TEM), revealing a nontailed viral particle (~45-nm capsid diameter) resembling phage PM2. DNA was extracted from the purified filtrate using the Qiagen MinElute virus spin kit and sequenced on the Illumina MiSeq platform (2 \times 250 bp). The resulting 71,931 reads were trimmed with Trimmomatic version 0.35.0 and assembled with SPAdes version 3.6.0 according to the iVirus pipeline (12). The assembled genome had 3,038-fold coverage and a G+C content of 48.2%.

ORFs were called using GLIMMER3 (13) and annotated using a BLASTx search against the GenBank nonredundant database. The genome of phage Cr39582 is syntenous with

Received 26 March 2018 Accepted 13 April 2018 Published 24 May 2018

Citation Leigh BA, Breitbart M, Oksanen HM, Bamford DH, Dishaw LJ. 2018. Genome sequence of PM2-like phage Cr39582, induced from a *Pseudoalteromonas* sp. isolated from the gut of *Ciona robusta*. *Genome Announc* 6:e00368-18. <https://doi.org/10.1128/genomeA.00368-18>.

Copyright © 2018 Leigh et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Brittany A. Leigh, brittany.a.leigh@vanderbilt.edu, or Larry J. Dishaw, ldishaw@health.usf.edu.

* Present address: Brittany A. Leigh, Department of Biological Sciences, Vanderbilt University, Nashville, Tennessee, USA.

phage PM2, demonstrating sequence homology throughout the genome with the exception of the ORF spanning positions 7982 to 9430, which is most similar to hypothetical *Pseudoalteromonas* sp. proteins. This region corresponds to the PM2 gene encoding the spike protein, which recognizes a host receptor that results in fusion with the bacterial membrane, ultimately allowing genome entry (14, 15). Phage Cr39582 was not able to form plaques on phage PM2's bacterial host, *Pseudoalteromonas espejiana* BAL-31, likely due to the dissimilarity between the two spike proteins. Full-genome alignment of phage Cr39582 to PM2 revealed ~85% overall identity. Pairwise nucleotide identities between Cr39582 and PM2 in the 5' and 3' regions flanking the spike protein gene were ~91% and ~96%, respectively.

Accession number(s). The phage Cr39582 genome sequence and the 16S rRNA gene sequence of the host *Pseudoalteromonas* sp. Cr6751 were deposited in GenBank under accession numbers [MG966533](#) and [MH014962](#), respectively.

ACKNOWLEDGMENTS

We thank Anthony Greco for imaging assistance on the TEM and Lauren McDaniel for her expertise and supply of the *P. espejiana* BAL-31 strain.

These studies were supported by a grant from the National Science Foundation (IOS-1456301) to L.J.D. and M.B. and by a National Science Foundation Graduate Research Fellowship (award number 1144244) to B.A.L. We acknowledge the use of the Instruct-HiLIFE Biocomplex Unit (University of Helsinki and Instruct-FI) and the Academy of Finland's support (grant 1306833) for the unit.

REFERENCES

- Oksanen HM; ICTV Report Consortium. 2017. ICTV virus taxonomy profile: *Corticoviridae*. *J Gen Virol* 98:888–889. <https://doi.org/10.1099/jgv.0.000795>.
- Männistö RH, Kivelä HM, Paulin L, Bamford DH, Bamford JK. 1999. The complete genome sequence of PM2, the first lipid-containing bacterial virus to be isolated. *Virology* 262:355–363. <https://doi.org/10.1006/viro.1999.9837>.
- Kivela HM, Kalkkinen N, Bamford DH. 2002. Bacteriophage PM2 has a protein capsid surrounding a spherical proteinaceous lipid core. *J Virol* 76:8169–8178. <https://doi.org/10.1128/JVI.76.16.8169-8178.2002>.
- Espejo RT, Canelo ES. 1968. Properties of bacteriophage PM2: a lipid-containing bacterial virus. *Virology* 34:738–747.
- Krupovic M, Bamford DH. 2007. Putative prophages related to lytic tailless marine dsDNA phage PM2 are widespread in the genomes of aquatic bacteria. *BMC Genomics* 8:236. <https://doi.org/10.1186/1471-2164-8-236>.
- Duhaime MB, Solonenko N, Roux S, Verberkmoes NC, Wichels A, Sullivan MB. 2017. Comparative omics and trait analyses of marine *Pseudoalteromonas* phages advance the phage OTU concept. *Front Microbiol* 8:1241. <https://doi.org/10.3389/fmicb.2017.01241>.
- Brum JR, Schenck RO, Sullivan MB. 2013. Global morphological analysis of marine viruses shows minimal regional variation and dominance of non-tailed viruses. *ISME J* 7:1738–1751. <https://doi.org/10.1038/ismej.2013.67>.
- Kauffman KM, Hussain FA, Yang J, Arevalo P, Brown JM, Chang WK, VanInsberghe D, Elsherbini J, Sharma RS, Cutler MB, Kelly L, Polz MF. 2018. A major lineage of non-tailed dsDNA viruses as unrecognized killers of marine bacteria. *Nature* 554:118–122. <https://doi.org/10.1038/nature25474>.
- Leigh B, Karrer C, Cannon JP, Breitbart M, Dishaw LJ. 2017. Isolation and characterization of a *Shewanella* phage-host system from the gut of the tunicate, *Ciona intestinalis*. *Viruses* 9:60. <https://doi.org/10.3390/v9030060>.
- Williamson SJ, Houchin LA, McDaniel L, Paul JH. 2002. Seasonal variation in lysogeny as depicted by prophage induction in Tampa Bay, Florida. *Appl Environ Microbiol* 68:4307–4314. <https://doi.org/10.1128/AEM.68.9.4307-4314.2002>.
- Patel A, Noble RT, Steele JA, Schwalbach MS, Hewson I, Fuhrman JA. 2007. Virus and prokaryote enumeration from planktonic aquatic environments by epifluorescence microscopy with SYBR Green I. *Nat Protoc* 2:269–276. <https://doi.org/10.1038/nprot.2007.6>.
- Bolduc B, Youens-Clark K, Roux S, Hurwitz BL, Sullivan MB. 2017. iVirus: facilitating new insights in viral ecology with software and community data sets imbedded in a cyberinfrastructure. *ISME J* 11:7–14. <https://doi.org/10.1038/ismej.2016.89>.
- Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* 27:4636–4641.
- Abrescia NGA, Grimes JM, Kivelä HM, Assenberg R, Sutton GC, Butcher SJ, Bamford JKH, Bamford DH, Stuart DI. 2008. Insights into virus evolution and membrane biogenesis from the structure of the marine lipid-containing bacteriophage PM2. *Mol Cell* 31:749–761. <https://doi.org/10.1016/j.molcel.2008.06.026>.
- Kivelä HM, Daugelavicius R, Hankkio RH, Bamford JKH, Bamford DH. 2004. Penetration of membrane-containing double-stranded-DNA bacteriophage PM2 into *Pseudoalteromonas* hosts. *J Bacteriol* 186:5342–5354. <https://doi.org/10.1128/JB.186.16.5342-5354.2004>.