
USF Patents

12-6-2016

Compositions including tricitiribine and methods of use thereof

Jin Q. Cheng

Said M. Sebt

Follow this and additional works at: https://digitalcommons.usf.edu/usf_patents

Recommended Citation

Cheng, Jin Q. and Sebt, Said M., "Compositions including tricitiribine and methods of use thereof" (2016).
USF Patents. 907.

https://digitalcommons.usf.edu/usf_patents/907

This Patent is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Patents by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.



US009511084B2

(12) **United States Patent**
Cheng et al.

(10) **Patent No.:** **US 9,511,084 B2**
(45) **Date of Patent:** ***Dec. 6, 2016**

(54) **COMPOSITIONS INCLUDING TRICIRIBINE
AND METHODS OF USE THEREOF**

16/2863 (2013.01); *A61K 2039/505* (2013.01);
C07K 2317/24 (2013.01); *C07K 2317/76*
(2013.01)

(71) Applicant: **University of South Florida**, Tampa,
FL (US)

(58) **Field of Classification Search**

None

See application file for complete search history.

(72) Inventors: **Jin Q. Cheng**, Tampa, FL (US); **Said
M. Sebti**, Tampa, FL (US)

(56) **References Cited**

(73) Assignee: **University of Florida**, Tampa, FL (US)

U.S. PATENT DOCUMENTS

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

4,123,524 A 10/1978 Townsend
5,681,970 A 10/1997 Didier et al.

(Continued)

This patent is subject to a terminal dis-
claimer.

FOREIGN PATENT DOCUMENTS

(21) Appl. No.: **14/514,311**

JP 2007-530705 11/2007
WO WO 2005/094322 10/2005

(Continued)

(22) Filed: **Oct. 14, 2014**

(65) **Prior Publication Data**

US 2015/0174149 A1 Jun. 25, 2015

OTHER PUBLICATIONS

Al-Janadi, Anas, et al.; "Activated Akt is Frequently Overexpressed
in Human Breast Cancer Cells and Its Blockade Can Inhibit
Growth," 2000 ASCO Annual Meeting, Proc Am Soc Clin Oncol,
19: 2000 (abstr 1879)

(Continued)

Related U.S. Application Data

(63) Continuation of application No. 13/463,576, filed on
May 3, 2012, now Pat. No. 8,901,086, which is a
continuation-in-part of application No. 12/992,556,
filed on May 18, 2011, now abandoned, which is a

(Continued)

Primary Examiner — Anna Pagonakis

(74) *Attorney, Agent, or Firm* — Seyfarth Shaw LLP

(51) **Int. Cl.**

A61K 31/7064 (2006.01)

A61K 31/282 (2006.01)

(Continued)



(57) **ABSTRACT**

This invention encompasses combination therapies includ-
ing TCN, TCN-P, TCN-PM and/or related compounds and
one or more additional anti-cancer agents, for example,
taxanes a molecule that modulates the HER2/neu (erbB2)
receptor, anthracyclin compounds, epidermal growth factor
receptor inhibitor compounds, one or more platinum com-
pounds and bortezomib and derivatives thereof and compo-
sitions with reduced toxicity for the treatment and preven-
tion of tumors, cancer, and other disorders associated with
abnormal cell proliferation.

(52) **U.S. Cl.**

CPC *A61K 31/7064* (2013.01); *A61K 31/282*
(2013.01); *A61K 31/337* (2013.01); *A61K*
31/4353 (2013.01); *A61K 31/7076* (2013.01);
A61K 33/24 (2013.01); *A61K 39/39558*
(2013.01); *A61K 45/06* (2013.01); *C07K*

15 Claims, 45 Drawing Sheets



Related U.S. Application Data

continuation-in-part of application No. 12/118,848, filed on May 12, 2008, now abandoned, which is a continuation-in-part of application No. 12/118,861, filed on May 12, 2008, now abandoned, which is a continuation-in-part of application No. 12/118,868, filed on May 12, 2008, now abandoned, which is a continuation-in-part of application No. 12/118,870, filed on May 12, 2008, now abandoned, which is a continuation-in-part of application No. 12/118,834, filed on May 12, 2008, now abandoned, which is a continuation-in-part of application No. 12/118,828, filed on May 12, 2008, now abandoned, which is a continuation-in-part of application No. 11/096,082, filed on Mar. 29, 2005, now Pat. No. 8,435,959.

(60) Provisional application No. 60/557,599, filed on Mar. 29, 2004.

(51) Int. Cl.

A61K 31/337 (2006.01)
A61K 31/4353 (2006.01)
A61K 31/7076 (2006.01)
A61K 33/24 (2006.01)
A61K 39/395 (2006.01)
A61K 45/06 (2006.01)
C07K 16/28 (2006.01)
A61K 39/00 (2006.01)

(56) References Cited**U.S. PATENT DOCUMENTS**

6,306,874 B1 * 10/2001 Fraley A61K 31/495
 514/300
 8,435,959 B2 * 5/2013 Cheng C07H 17/02
 514/23
 8,623,834 B2 * 1/2014 Cheng A61K 31/7064
 514/23
 8,901,086 B2 * 12/2014 Cheng A61K 31/7064
 514/23
 9,115,162 B2 * 8/2015 Cheng C07H 17/02
 2003/0096823 A1 * 5/2003 Asp A61K 39/39558
 514/252.11
 2015/0320784 A1 * 11/2015 Cheng C07H 17/02
 514/52

FOREIGN PATENT DOCUMENTS

WO WO 2008/060581 5/2008
 WO WO 2008/070100 6/2008
 WO WO 2008/094321 8/2008

OTHER PUBLICATIONS

Barnett, Stanley F., et al.; "The Akt/PKB Family of Protein Kinases: A Review of Small Molecule Inhibitors and Progress Towards Target Validation", Current Topics in Medicinal Chemistry, vol. 5, pp. 109-125, 2005.

Cheng, et al.; (Oncogene (1997), 14, 2793-2801).

Cheng, J.Q., et al.; "The Akt/PKB Pathway: Molecular Target for Cancer Drug Discovery," Oncogene 20051114 GB, vol. 24(50), pp. 7482-7492, 2005.

Dou, et al.; (IDrugs (2002), 5(8), 828-834) (Abstract).

Feun, et al.; (Cancer Research (1984), 44, 3608-3612).

Feun, Lynn G., et al.; "A Phase II Trial of Tricyclic Nucleoside Phosphate in Patients with Advanced Squamous Cell Carcinoma of the Cervix: A Gynecologic Oncology Group Study", American Journal of Clinical Oncology (Cancer Clinical Trials), vol. 16(6), pp. 506-508, 1993.

Goldenberg; (Clinical Therapeutics (1999), 21(2), 309-319.

Gura; (Science (1997), 278(4534), 1041-1042).

Herceptin (Trastuzumab) Prescribing Information, Jan. 2008.

Hoffman, et al.; (Cancer Chemother Pharmacol (1996), 37, 254-258).

Jetzt, et al.; (Cancer Research (2003), 63, 6697-6706).

Lan, Keng-Hsueh, et al.; "Mechanisms of Trastuzumab Resistance and Their Clinical Implications," Annals of the New York Academy of Sciences, vol. 1059, pp. 70-75, 2005.

Lu, Chien-Hsing, et al.; "Preclinical Testing of Clinically Applicable Strategies for Overcoming Trastuzumab Resistance Caused by PTEN Deficiency," Clinical Cancer Research, The American Association for Cancer Research, vol. 13(9), pp. 5583-5588, 2007.

Mittelman, et al.; (Cancer Treatment Reports (1983), 67(2), 159-162).

O'Connell, Michael J., et al.; "Phase II Clinical Trial of Tricyclic Nucleoside Phosphate for Advanced Colorectal Cancer 1,2", Cancer Treatment Reports, vol. 71(3), pp. 333-334, 1987.

Park, S.-S. et al., Oncology Report, 2007, vol. 18, pp. 139-143.

Schilcher, et al.; (Journal of Chromatography (1985), 337(1), 56-62) (Abstract).

Schilcher, et al.; (Cancer Research (1986), 46, 3147-3151).

Shedden, et al.; (Pharmaceutical Research (2003), 20, 843-847).

Sridhar, et al.; (The Lancet Oncology (2003), 4, 397-406).

Sun, et al.; (American Journal of Pathology (2001), 159(2), 431-437).

Yu, Dihua, et al.; "Strategies for Overcoming Trastuzumab Resistance Caused by PTEN Deficiency," Proceedings of the American Association for Cancer Research Annual Meeting, vol. 49, p. 159, 2008.

Waud, et al.; (Cancer Chemotherapy and Pharmacology (1991), 27, 456-463).

Yang, Lin, et al.; "Akt/Protein Kinase B Signaling Inhibitor-2, a Selective Small Molecule inhibitor of Akt Signaling with Antitumor Activity in Cancer Cells Overexpressing Akt", Cancer Research, vol. 64, pp. 4394-4399, 2004.

Office Action issued in co-pending Japanese Patent Application No. 2011-509454 mailed Feb. 12, 2013.

Extended European Search Report issued by the European Patent Office of Mar. 5, 2013 in European patent application No. 12197499.2.

Office Action issued in co-pending Japanese Patent Application No. 2011-509454 mailed Jan. 21, 2014.

Office Action in related co-pending U.S. Appl. No. 12/118,828 mailed Mar. 17, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,834 mailed Mar. 1, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,870 mailed Mar. 4, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,861 mailed Mar. 4, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,868 mailed Mar. 4, 2011.

Office Action in related co-pending U.S. Appl. No. 11/096,082 mailed Nov. 10, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,828 mailed Nov. 23, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,834 mailed Nov. 23, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,861 mailed Nov. 23, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,868 mailed Nov. 23, 2011.

Office Action in related co-pending U.S. Appl. No. 12/118,870 mailed Nov. 23, 2011.

Office Action in related co-pending U.S. Appl. No. 11/090,082 mailed Jun. 29, 2012.

Office Action in related co-pending U.S. Appl. No. 12/992,556 mailed Aug. 10, 2012.

Notice of Allowance in related co-pending U.S. Appl. No. 11/096,082 mailed Jan. 9, 2013.

Non-Final Office Action in related co-pending U.S. Appl. No. 13/453,807 mailed Feb. 26, 2013.


(56)

References Cited**OTHER PUBLICATIONS**


Non-Final Office Action in related co-pending U.S. Appl. No. 13/452,478 mailed Mar. 25, 2013.
Non-Final Office Action in related co-pending U.S. Appl. No. 13/453,778 mailed Mar. 8, 2013.
Non-Final Office Action in related co-pending U.S. Appl. No. 13/453,789 mailed Mar. 14, 2013.
Non-Final Office Action in co-pending U.S. Appl. No. 13/453,834 mailed Aug. 5, 2013.
Notice of Allowance in co-pending U.S. Appl. No. 13/453,807 mailed Sep. 24, 2013.
Notice of Allowance in co-pending U.S. Appl. No. 13/453,778 mailed Sep. 25, 2013.
Non-Final Office Action in co-pending U.S. Appl. No. 12/992,556 mailed Sep. 26, 2013.
Notice of Allowance in co-pending U.S. Appl. No. 13/453,789 mailed Oct. 28, 2013.
Notice of Allowance in co-pending U.S. Appl. No. 13/452,478 mailed Nov. 7, 2013.
Notice of Allowance in co-pending U.S. Appl. No. 13/453,834 mailed Apr. 9, 2014.
Non-Final Office Action issued in U.S. Appl. No. 13/463,576, now U.S. Pat. No. 8,901,086, mailed Apr. 25, 2013.
Final Office Action issued in U.S. Appl. No. 13/463,576, now U.S. Pat. No. 8,901,086, mailed Apr. 7, 2014.
Notice of Allowance issued in U.S. Appl. No. 13/463,576, now U.S. Pat. No. 8,901,086, mailed Jul. 11, 2014.
Notice of Allowance in U.S. Appl. No. 14/321,572 on Jul. 17, 2015.

Notice of Allowance in U.S. Appl. No. 14/092,323 on Sep. 28, 2015.
Schilcher, Rudolf B., et al.: "Reversed-Phase High-Performance Liquid Chromatographic Determination of Tricyclic Nucleotide and Tricyclic Nucleotide 5'- Phosphate in Biological Specimens," *Journal of Chromatography*, vol. 337, pp. 55-62, 1985.
Waud, William R., et al.: "Antitumor Drug Cross-Resistance in vivo in a Cisplatin-Resistant Murine P388 Leukemia," *Cancer Chemotherapy Pharmacology*, vol. 27, pp. 456-563, 1991.
Non-Final Office Action in U.S. Appl. No. 14/092,335 on Feb. 2, 2015.
Non-Final Office Action in U.S. Appl. No. 14/321,572 on Feb. 13, 2015.
Notice of Allowance in U.S. Appl. No. 14/091,073 on Apr. 13, 2015.
Notice Allowance in U.S. Appl. No. 13/864,953 on Apr. 17, 2015.
Non-Final Office Action in U.S. Appl. No. 14/789,347 on Feb. 25, 2016.
Non-Final Office Action in U.S. Appl. No. 14/800,286 on Mar. 9, 2016.
Non-Final Office Action in U.S. Appl. No. 14/832,417 on Mar. 1, 2016.
Non-Final Office Action in U.S. Appl. No. 14/881,777 on Feb. 25, 2016.
Non-Final Office Action in U.S. Appl. No. 14/886,882 on Feb. 26, 2016.
Non-Final Office Action in U.S. Appl. No. 14/998,352 on Apr. 22, 2016.
Decision of Refusal issued in Japanese Patent Application No. 2014-264781 on Apr. 26, 2016.


* cited by examiner



১
 ২
 ৩
 ৪



ॐ
ॐ
ॐ



ம
த
த
த

உ
த
உ




FIG. 2B





FIG. 2C






FIG. 2D




FIG. 2E




FIG. 2F




FIG. 3A




FIG. 3B-1




FIG. 3B-4




FIG. 3B-2




FIG. 3B-5




FIG. 3B-3




FIG. 3B-6




FIG. 3B-7




FIG. 3B-9




FIG. 3B-8




FIG. 3B-10




FIG. 3B-11




FIG. 3B-12




FIG. 3C-1




FIG. 3C-2




FIG. 3C-3




FIG. 3C-4




FIG. 3C-6




FIG. 3C-5




FIG. 3C-7




FIG. 4A




FIG. 4C




FIG. 4E




FIG. 4D




FIG. 4B-1




FIG. 4B-2




FIG. 4B-3




FIG. 4B-4




FIG. 4B-5

PIP3:	-	+	+	+	+
PDK1:	-	+	+	+	+
Akt:	+	+	+	+	+
API-2:	-	-	-	25	10 (μ M)



FIG. 5

atgaacgaogtagccattgtgaaggagggctgggtgcacaaacgaggggaatatattaa
1 -----+-----+-----+-----+-----+-----+-----+ 60
tacttgctgcctcggtasacacttccctcccgacggacgtgtttgctccctttatataattt

a M N D V A I V K E S W L H K R G E Y F K -

acctggcgggccacgctacttccctccctcaagaacgatggcaacctttattggctacaaaggaa
61 -----+-----+-----+-----+-----+-----+ 120
tggacgcgcgggtgcgatgaaggaggagttctttgctacogtggaaatatccgatgttccct

a T W R P R Y F L L K N D G T F I G Y K E -

cgccctcaggatgttggatcagcgagagtgccccactcaacaacttctcagtgggcacaatgc
121 -----+-----+-----+-----+-----+-----+ 180
ggcggagtcctacacctagtcgctctcaggggtgagttgttgaagagtcaccgtgttacg

a R F Q D V D Q R E S P L N N F S V A Q C -

PatI
|
cagctgatgaagacagagcgggccaaggcccaacacacctttatcctccgctgcctgcagtg
181 -----+-----+-----+-----+-----+-----+ 240
gtcgactacttctgtctcgcgggtccgggttgttggaatatagtagggacggacgctcacc

a Q L M K T E R P R P N T F I I R C L Q W -

accgcagtcattgagcgccaccttcctatgtagaacgcctgaggagcgggaagaatggggcc
241 -----+-----+-----+-----+-----+-----+ 300
tgytgcagtaactcgcggtggaagggtacatctttgaggactcctcgcccttcttaaccgg

a T T V I E R T F H V E T P E E R E E W A -

accgccattcagactgttggcggatggactcaagaggcaggaagaagagacgatggacttc
301 -----+-----+-----+-----+-----+-----+ 360
tggcggtaagtctgacaccggctacctgagttctccgtccttcttctctgctacctgaag

a T A I Q T V A D G L K R Q E E E T M D F -

cgatcaggctcaccacagtgacaactcaggggtgaagagatggaggtgtccctggccaaag
361 -----+-----+-----+-----+-----+-----+ 420
gctagtcggagtggggtcactgttgagtccccgacttctctacctccacagggacgggttc

a R S G S P S D N S G A E E M E V S L A K -

FIG. 6A

```
cccaagcacccgtgtgaccatgaacgagtttgagtacctgaaactactgggcaagggcacc
421 -----+-----+-----+-----+-----+-----+-----+ 480
gggttcgtggcacactgggtacttgcacaactcatggactttgatgacccgttcccggtg

a   P K H R V T M N E F E Y L K L L G K G T -

tttgggaagtgattctgtgtaaaagagaaggccacaggccgctactatgccatgaagatc
481 -----+-----+-----+-----+-----+-----+-----+ 540
aaaccctttcactaagaccattttctcttccgggtgtccggcgatgatacggtaacttctag

a   F G K V I L V K E K A T G R Y Y A M K I -

ctcaagaaggagggtcatcgctcgccaaggatgaggttgccacacgcttactgaaaacogt
541 -----+-----+-----+-----+-----+-----+-----+ 600
gagttcttccctcagtagcagcggttctactcacaacgggtgtgcgaatgacttttggca

a   L K K E V I V A K D E V A H T L T E N R -

      PetI
      |
gtcctgcagaaotctagggcatcccttcccttaogggccctcaagtactcattccagaccac
601 -----+-----+-----+-----+-----+-----+-----+ 660
caggacgttcttgagatccgttaggggaagggaatgccggggagtccatgagtaagggtctgggtg

a   V L Q N S R H P F L T A L K Y S F Q T H -

                                     XhoI
                                     |
gaaccgcctctgctttgtcctggagtatgccaaacgggggagagctcttcttccacctgtct
661 -----+-----+-----+-----+-----+-----+-----+ 720
ctggcgggagacgaacagtagacctacatcaggttgcccccgctcgagagagaagggtggacaga

a   D R L C F V M E Y A N G G E L F F H L S -

cgagagcgcggtgttctccgaggacccggcccgcttctatggtgaggagattgtgtctgcc
721 -----+-----+-----+-----+-----+-----+-----+ 780
gctctcgcgccacaagagggtctcctggccccgggagagataaccacgcctctaacacagacgg

a   R E R V F S E D R A R F Y G A E I V S A -

ctggactacttgcaactccgagaagaacgtggtgtaccgggacctgaagctagagaaacctc
781 -----+-----+-----+-----+-----+-----+-----+ 840
gacctgatgaacgtgagggctctcttcttgcaccacatggccctggacttcgatctcttggag

a   L D Y L H S E K N V V Y R D L K L E N L -
```

FIG. 6B

atgctggacaaggacgggcacatcaagataacggacttcgggctgtgcaggaggggatac
841 -----+-----+-----+-----+-----+-----+-----+ 900
taagacctgttcctgcccgtgtagtcttatattgcctgaagcccgacacgttcctcccttag
a M L D K D G H I K I T D F G L C K E G I ~
aaggatgggtgcactatgaagacattctgcyygaacgcgggagtacctggcccttgaggtg
901 -----+-----+-----+-----+-----+-----+ 960
ttcctaccacgggtgatacttctgttaagacgccttgccggcctcatggacccggggactccac
a K D G A T M K T F C G T P E Y L A P E V ~
ctggaagacaaacgactaaggccgtgcagtggaactgggtggggcctgggctgtgtatgtat
961 -----+-----+-----+-----+-----+-----+ 1020
gaccttctgttgctgatgcgggcacgtcacctgacccaccccccgcacccgcacagtacta
a L E D N D Y G R A V D W W G L G V V M Y ~
gagatgatgtgtggccgcctgccttctacaaacaggacccagagaagctgttcgagctg
1021 -----+-----+-----+-----+-----+-----+ 1080
ctctactacacacccggcggaaggagatgttggtcctgtgtctcttcgacaagctcgac
a E M M C G R L F F Y N Q D H E K L F E L ~
atcctcatggaggagatccgcttcccgcgacacactcggccctgaggccaaagtccctgctc
1081 -----+-----+-----+-----+-----+-----+ 1140
taggagtaacctcctctaggcgaaagggcgctgtgtagccgggactcgggttcagggaacag
a I L M E E I R F P R T L G P R A K S L L ~
tcgggctgtctcaagaaggacccctacacagaggctcgggtggggcctctgaggatgccaaag
1141 -----+-----+-----+-----+-----+-----+ 1200
aggcccgacagagtctctcctgggatgtgtctccgagccaccccccagagactcctacggttc
a S G L L K K D F T Q R L G G G S E D A K ~
gagatcatgcagcacccggttctttgccaaacatcgtgtggcaggatgtgtatgagaagag
1201 -----+-----+-----+-----+-----+-----+ 1260
ctctagtacgtcgtggccaagaaacgggttgttagcacacccgtcctacacatactctctctc
a E I M Q H R F F A N I V W Q D V Y E K K ~

FIG. 6C

```
ctgagcccaacctttcaagcccccagggtcacctctgagactgacaccagggtatttcgatgag
1261 -----+-----+-----+-----+-----+-----+ 1320
gactcgggttggaagttcggggtccagtgaggagactctgactgtggtccataaagctactc

a    L S P P F K P Q V T S E T D T R Y F D E -

gagttcacagctcagatgatcaccatcacgcgcctgatcaagatgacagcatggagtgt
1321 -----+-----+-----+-----+-----+-----+ 1380
ctcaagtgctcagtcctactagtggtagtgccggcggaotagttctactgtcgtacctcaca

a    E F T Q Q M I T I T P P D Q D D S M E C -

gtggacagtgagcgggagggccgcacttcccccagttctcctactgacccagtgggccagcc
1381 -----+-----+-----+-----+-----+-----+ 1440
cacctgtcactcgcctccggcgtgaaggggggtcaagaggatgagtcgggtcacccgtgtcgg

a    V D S E R R P H F F Q F S Y S A S G T A -

tga
1441 --- 1443
act

a    * -
```

FIG. 6D

ATGAATGAGGTGTCTGTCTCATCAAGAAGGCTGGCTCCACAAGCGTGGTGAATACATCAAG
328 ----- 387
TACTTACTCCACAGACAGTATGTTCTTCGACCGAGGTGTTCCGACCACCTTATGTAGTTC

a M N E V S V I K E G W L H K R G E Y I K -

ACCTGGAGGCCACGGTACTTCCTGCTGAAGAGCGACGGCTCCTTCATTGGGTACAAGGAG
388 ----- 447
TGGACCTCCGGTGGCATGAAGGACGACTTCTCGCTGCCGAGGAAGTAACCCATGTTCCCTC

a T W R P R Y F L L K S D G S F I G Y K E -

AGGCCCGAGGCCCTGATCAGACTCTACCCCCCTTAAACAACCTTCTCCGTAGCAGATGC
448 ----- 507
TCCGGGCTCCGGGCACTAGTCTEAGATGGGGGAATTGTTGAAGAGGCATCGTCTTACG

a R P E A P D Q T L P P L N N F S V A E C -

PstI
|
CAGCTGATGAAGACCGAGAGGCCCGGACCCACACCTTTGTCTATACGCTGGCTGCAGTGG
508 ----- 567
GTGCACTACTTCTGGCTCTCCGGCGCTGGGTTGTGGAACAGTATGCGACGGACGTCACC

a Q L M K T E R P R P N T F V I R C L Q W -

ACCACAGTCATCGAGAGGACCTTCCACGTGGATTCTCCAGACGAGAGGGAGCAGTGGATG
568 ----- 627
TGGTGTCACTAGCTCTCTCGAAGGTGCACCTAAGAGGTCTGCTCTCCCTCCTCACCTAC

a T T V I E R T F H V D S P D E R E E W M -

ApaI
|
CGGGCCATCCAGATGGTCCGCAACAGCCTCAAGCAGCGGGGCCCCAGGCGAGGACCCCATG
628 ----- 687
GCCCCGTAGGTCTACACGCGTTGTGCGAGTTCGTCGCCCCGGGTCCGCTCCTGGGGTAC

a R A I Q M V A N S L K Q R A P G E D F M -

GACTACAAGTGTGGCTCCCCAGTGACTCCTCCACGACTGAGGAGATGGAAGTGGCGGTC
688 ----- 747
CTGATGTTACACCGAGGGGCTCACTGAGGAGGTGCTGACTCCTCTACCTTCACCGCCAG

a D Y K C G S P S D S S T T E E M E V A V -

FIG. 7A




FIG. 7B

CTCATGCTGGACAAAGATGGCCACATCAAGATCACTGACTTTGGCCCTCTSCAAAGAGGGG
1168 ---+-----+-----+-----+-----+-----+----- 1227
GASTACGACCTGTTTCTACCGGTGTAGTTCCTAGTGACTGAAACCGGAGACCTTCTCCCG

a L M L D K D G H I K I T D F G L C K E G ~

ATCAGTGACGGGGCCACCATGAAAACCTTCTGTGGGACCCCGAGTACCTGGCGCCTGAG
1228 ---+-----+-----+-----+-----+-----+----- 1287
TAGTCACTGCCCCGGTGGTACTTTTGAAGACACCCTGGGGCCCTCATGGACCGCGGACTC

a I S D G A T M K T F C G T P E Y L A P E ~

GTGCTGGAGGACAATGACTATGSCCGGGCCGTGGACTGGTGGGGGCTGGGTGTGTGTCATG
1288 ---+-----+-----+-----+-----+-----+----- 1347
CACGACCTCCTGTACTGATACCGCCCGGCACCTGACCACCCCGACCCACACCACTAC

a V L E D N D Y G R A V D W W G L G V V M ~

TACGAGATGATGTGCGGCCCTGCCCCCTTCTACAACCGAGACCACGAGCGCCTCTTCGAG
1348 ---+-----+-----+-----+-----+-----+----- 1407
ATGCTCTACTACACGCCCGCGGACCGGAGATGTTGGTCTGCTGCTCGCGAGAGCTC

a Y E M M C G R L P F Y N Q D H E R L F E ~

SacI
|
CTCATCCTCATGGAAGAGATCCGCTTCCCGCCACGCTCAGCCCCGAGGCCAAGTCCCTG
1408 ---+-----+-----+-----+-----+-----+----- 1467
GAGTAGGAGTACCTTCTTAGGCGAAGGGCGCGTGCAGTCCGGGCTCCGGTTCAGGGAC

a L I L M E E I R F P R T L S F E A K S L ~

ApaI
|
CTTGCTGGCTGCTTAAGAAGGACCCCAAGCAGAGGCTTGGTGGGGGGCCAGCGATGCC
1468 ---+-----+-----+-----+-----+-----+----- 1527
GAACGACCCGACGAATTCCTTGGGGTTCGTCTCCGAACCAACCCCGGGTCCGTACGG

a L A G L L K K D P K Q R L G G G P S D A ~

AAGGAGGTCATGGAGCACAGGTTCTTCCTCAGCATCAACTGGCAGGACGTGGTCCAGAAG
1528 ---+-----+-----+-----+-----+-----+----- 1587
TTCCTCCAGTACCTCGTGTCCAAGAAGGAGTCTAGTTGACCGTCTGCACCAAGGCTCTC

a K E V M E H R F F L S I N W Q D V V Q K ~

FIG. 7C

SallI
|

AAGCTCCTGCCACCCTTCAAACCTCAGGTCACGTCGAGGTCGACACAAGGTACTTCGAT
1588 ---+-----+-----+-----+-----+-----+----- 1647
TTCGAGGACGGTGGGAAGTTTGGAGTCCAGTSCAGGCTCCAGCTGTGTTCCATGAAGCTA

a K L L P P F K P Q V T S E V D T R Y F D ~

GATGAATTTACCGCCCAAGTCCATCACAATCACACCCCTGACCGCTATGACAGCCTGGGC
1648 ---+-----+-----+-----+-----+-----+----- 1707
CTACTTAAATGGCGGCTCAGGTAGTGTGTTAGTGTGGGGACTGGCGATACTGTTCGGACCCG

a D E F T A Q S I T I T P P D R Y D S L G ~

TTACTGGAGCTGGACCAGCGGACCCACTTCCCCAGTTCTCTCTACTCGGCCAGCATCCGC
1708 ---+-----+-----+-----+-----+-----+----- 1767
AATGACCTCGACCTGGCTCGCCTGGGTGAAGCGGGTCAAGAGGATGAGCCGGTCTAGGCC

a L L E L D Q R T H F P Q F S Y S A S I R ~

GAGTGAGCAGTCTGCCACGCAGAGGACGACCGCTCGTGTCCATCACCCTGGGTGGTTT
1768 ---+-----+-----+-----+-----+-----+----- 1827
CTCACTCGTCAGACGGGTGCGTCTCCTGCGTGGAGCGACGGTAGTGGCGACCCACCAA

a E * A V C P R R G R T L A A I T A G W F ~

TTTACCCCTGCCCG
1828 ---+-----+----- 1842
AAATGGGGACGGGCC

a F T P A R ~

FIG. 7D

[illegible]

FIG. 8A

atgcctctacaacccatcatataaagaagacaaatgaatgattttgactatttgaaactac
421 -----+-----+-----+-----+-----+-----+ 480
taaggagatggttggttagtattttctttctgttacttactaaaaactgataaaactttgatg
a A S T T H H K R * T M N D F D Y L K L L -
taggtaaaaggcaacttttgggaagttatatttgggttgcagagaaggcaagtggaaaaact
481 -----+-----+-----+-----+-----+-----+ 540
atccattttcogtgaasaccctttcaataasaccaagctctcttccgttcaccttttatga
a G K G T F G K V I L V R E K A S G K Y Y -
atgcctatgaagattctgaagaagaagtcatttattgcagaaggatgaagtggcacacactc
541 -----+-----+-----+-----+-----+-----+ 600
taagataacttctaagscttctttcttcagtaataacgttttctacttccacgtgtgtgag
a A M K I L K K E V I I A K D E V A H T L -
taactgaagacagagtatttaagaacactagacatccctttttaacatccttgaatat
601 -----+-----+-----+-----+-----+-----+ 660
attgactttcgtctcataatttctgtgatctgttagggaaaaattgttaggaactttataa
a T E S R V L K N T R H P F L T S L K Y S -
ccttcagacaaaaagaccgttttgtgttttgtgatggaaatatgttaatgggggagagctgt
661 -----+-----+-----+-----+-----+-----+ 720
ggaaggtctgttttctggycaaacacaaacactaccttatacaattacccccgctcgaca
a F Q T K D R L C F V M E Y V N G G E L F -
ttttccatttgtgcagagagcgggtgttctctgaggaccgcacacgtttctatgtgtcag
721 -----+-----+-----+-----+-----+-----+ 780
aaaaggtaaaacagctctctctgcgcccaagagactcctggcgtgtgcasaagataccacgtc
a F H L S R E R V F S E D R T R F Y G A E -
aaattgtctctgccttggactatctacattccggaaagattgtgtacogtgcattcgaagt
781 -----+-----+-----+-----+-----+-----+ 840
tttaacagagacggaaacctgatagatgtaaggcctttctaacacatggcaactagagttca
a I V S A L D Y L H S G K I V Y R D L K L -

FIG. 8B

```

tggagaatctaatgctggcacaagatggtggccacataaaaaattacagatttttggacttttgca
841 -----+-----+-----+-----+-----+-----+-----+-----+ 900
acctotttagattacagacctgtttctacoggygtatttttaagtgtctaaaaacctgaaacct

a      E N L M L D K D G H I K I T D F G L C K -

aagaaggggatcacagatgcagcccccacatgaagacattctgtggcactccagaatatctgg
901 -----+-----+-----+-----+-----+-----+-----+ 960
ttcttccctsgtgtctacgtcggtggtactctctgtaagacacctgtgaggtctttatagacc

a      E G I T D A A T M K T F C G T F E Y L A -

caccagaggtgttagaagataatgactatggcccgagcagtagactggtggggccctagggg
961 -----+-----+-----+-----+-----+-----+-----+ 1020
gtgggtctccacaatctttctattactgtataccggctogtctcttgaccaccccggtacccc

a      P E V L E D N D Y G R A V D W W G L G V -

ttgtcatgtatgaaatgatgtgtgtgggaggttacctttctacaaccaggaccatgagaac
1021 -----+-----+-----+-----+-----+-----+-----+ 1080
aacagttacatactttactacacacccctcccaatggaagatgttgggtcctggtactctttg

a      V M Y E M M C G R L P F Y N Q D H E K L -

tttttgaattaatattaatggaagacattaaatttctctgaacactctctcttcagatgcaa
1081 -----+-----+-----+-----+-----+-----+-----+ 1140
aaaaacttaattataattacctttctgttaatttaaaggagcttgtgagagaagttctacgtt

a      F E L I L M E D I K F P R T L S S D A K -

                                     BamHI
                                     |
aatcattgcttttcagggctctttgataaaggatccaaataaaacgccttgggtggaggaccag
1141 -----+-----+-----+-----+-----+-----+-----+ 1200
ttagtaacgaaagtcoccgagaactatttctctaggtttattttgoggaaaccacctcctggtc

a      S L L S G L L I K D P N K R L G G G F D -

atgatgcasaagaattatgagacacagtttctctctctggagtaaaactggcagatgtat
1201 -----+-----+-----+-----+-----+-----+-----+ 1260
tactacgtttttctttaatactctgtgtcaagaagagacotcatttgaaccttctacata

a      D A K E I M R H S F P S G V N W Q D V Y -

```

FIG. 8C

HindIII
|
atgataaaaagcttgtagcctccttttaaacctcaagtaacatctgagacagatactagat
1261 -----+-----+-----+-----+-----+-----+ 1320
tactattttttcgaacatggagggaatttggagttcattgtagactctgtctatgatcta
a D K K L V P P F K P Q V T S E T D T R Y -
atcttgatgaagaatttacagctcagactattacaataacaccacctgaaaaatatgatg
1321 -----+-----+-----+-----+-----+-----+ 1380
taaaactactctttaaatgtcgagctctgataaatgttattgtggaggactttttatactao
a F D E E F T A Q T I T I T P P E K Y D E -
NotI
|
aggatgggtatggactgcctggacaatgagagggcgccgcatttcacctcaattttcctact
1381 -----+-----+-----+-----+-----+-----+ 1440
tcctaccatacctgaagtaacctgttaactctccgcggcgtaaaagggagttaaaaggatga
a D G M D C M D N E R R P H F P Q F S Y S -
ctgcaagtggacgagaaataagttctctttcattctgtacttcactgtcactcttcattta
1441 -----+-----+-----+-----+-----+-----+ 1500
gacgttcacctgtctcttattccagagaaagtaagacgatgaagtgcagtagaagttaat
a A S G R E + V S F I L L L H C H L Q F I -
ttactgaaa
1501 -----+ 1510
aatgactttt
a T E -

FIG. 8D

Synergistic effect of TCN and Taxol on apoptosis in
A2780S and OVCAR3 cells






FIG. 9

Drug	target	% growth inhibition in PTEN AS BT474.m1 cells				Ttzm + drug significantly better than either alone?		
		Ttzm only	drug only	Ttzm + drug	Ttzm + drug			
Triciribine	Akt	9.5	+/-4.2	19.2	+/-9.2	36.6	+/-3.3	yes (P<0.001)
KP 372-1	Akt	0	+/-4.1	36	+/-2.1	38.6	+/-3.2	no
4ADPIB	Akt	0.9	+/-9.7	23.7	+/-2.4	32.2	+/-3.4	no
Edelfosine	Akt	2.2	+/-9.6	41.8	+/-1.2	51.9	+/-2.3	no
RAD001	mTOR	10.9	+/-4.9	19.3	+/-2.5	33.7	+/-1.4	yes (P<0.05)
QLT 0267	ILK	20	+/-6.8	30.3	+/-8.4	44.8	+/-6.9	yes (P<0.05)

FIG. 10A






FIG. 11




FIG. 12A




FIG. 12B




FIG. 13A




FIG. 13B




FIG. 14A




FIG. 14B




FIG. 15




FIG. 16




FIG. 17A




FIG. 17B




FIG. 18A




FIG. 18B




FIG. 19A




FIG. 19B




FIG. 20B





FIG. 20C




FIG. 21A




FIG. 21B




FIG. 22A




FIG. 22B





FIG. 23C

HA-Aurora-A




FIG. 24A




FIG. 24B




FIG. 24C




FIG. 24D




FIG. 25




FIG. 26A




FIG. 26B







FIG. 26C



COMPOSITIONS INCLUDING TRICIRIBINE AND METHODS OF USE THEREOF

The present application is a continuation of U.S. patent application Ser. No. 13/463,576, filed May 3, 2012, which issued as U.S. Pat. No. 8,901,086, which is a continuation of U.S. patent application Ser. No. 12/992,556, filed May 18, 2011, which is abandoned, which is a continuation of U.S. patent application Ser. No. 12/118,848, filed May 12, 2008, which is abandoned, and is a continuation-in-part of U.S. patent application Ser. No. 11/096,082, filed Mar. 29, 2005, which issued as U.S. Pat. No. 8,435,959, which claims the benefit of U.S. provisional patent application No. 60/557,599, filed Mar. 29, 2004, each of which is incorporated herein by reference.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

This invention was made with government support CA077935 and CA089242 awarded by the National Institutes of Health. The government has certain rights to the invention.

FIELD OF THE INVENTION

The invention encompasses combination therapies including TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents, for example, one or more taxanes; a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof; anthracyclin compounds; epidermal growth factor receptor inhibitor compounds, particularly erlotinib-like compounds; one or more platinum compounds; and bortezomib or salts thereof. The invention also encompasses compositions with reduced toxicity for the treatment and prevention of tumors, cancer, and other disorders associated with abnormal cell proliferation.

The instant application contains a Sequence Listing which has been submitted in ASCII format, created Nov. 3, 2005, is named 065031-5001-US Sequence Listing.txt, and is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

Cancer is an abnormal growth of cells. Cancer cells rapidly reproduce despite restriction of space, nutrients shared by other cells, or signals sent from the body to stop reproduction. Cancer cells are often shaped differently from healthy cells, do not function properly, and can spread into many areas of the body. Abnormal growths of tissue, called tumors, are clusters of cells that are capable of growing and dividing uncontrollably. Tumors can be benign (noncancerous) or malignant (cancerous). Benign tumors tend to grow slowly and do not spread. Malignant tumors can grow rapidly, invade and destroy nearby normal tissues, and spread throughout the body.

Cancers are classified according to the kind of fluid or tissue from which they originate, or according to the location in the body where they first developed. In addition, some cancers are of mixed types. Cancers can be grouped into five broad categories, carcinomas, sarcomas, lymphomas, leukemias, and myelomas, which indicate the tissue and blood classifications of the cancer. Carcinomas are cancers found in body tissue known as epithelial tissue that covers or lines surfaces of organs, glands, or body structures. For example, a cancer of the lining of the stomach is called a carcinoma.

Many carcinomas affect organs or glands that are involved with secretion, such as breasts that produce milk. Carcinomas account for approximately eighty to ninety percent of all cancer cases. Sarcomas are malignant tumors growing from connective tissues, such as cartilage, fat, muscle, tendons, and bones. The most common sarcoma, a tumor on the bone, usually occurs in young adults. Examples of sarcoma include osteosarcoma (bone) and chondrosarcoma (cartilage). Lymphoma refers to a cancer that originates in the nodes or glands of the lymphatic system, whose job it is to produce white blood cells and clean body fluids, or in organs such as the brain and breast. Lymphomas are classified into two categories: Hodgkin's lymphoma and non-Hodgkin's lymphoma. Leukemia, also known as blood cancer, is a cancer of the bone marrow that keeps the marrow from producing normal red and white blood cells and platelets. White blood cells are needed to resist infection. Red blood cells are needed to prevent anemia. Platelets keep the body from easily bruising and bleeding. Examples of leukemia include acute myelogenous leukemia, chronic myelogenous leukemia, acute lymphocytic leukemia, and chronic lymphocytic leukemia. The terms myelogenous and lymphocytic indicate the type of cells that are involved. Finally, myelomas grow in the plasma cells of bone marrow. In some cases, the myeloma cells collect in one bone and form a single tumor, called a plasmacytoma. However, in other cases, the myeloma cells collect in many bones, forming many bone tumors. This is called multiple myeloma.

Thus, a combination of a TCN, TCN-P, TCN-PM and/or related compounds or a derivative thereof and an additional anti-cancer agent holds promise as a potential combination therapy for treating tumors, cancer, and abnormal cell proliferation.

3. SUMMARY OF THE INVENTION

The invention encompasses novel therapeutic regimens of triciribine ("TCN"), triciribine phosphate ("TCN-P"), triciribine phosphate monohydrate ("TCN-PM") and related compounds in combination with one or more additional anti-cancer agents, for example, one or more taxanes, a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, anthracyclin compounds, epidermal growth factor receptor inhibitor compounds, particularly erlotinib-like compounds, one or more platinum compounds and bortezomib or salts thereof.

The invention is based on the discovery that tumors or cancers, which overexpress Akt kinase are sensitive to the cytotoxic effects of TCN, TCN-P, TCN-PM and related compounds and a synergistic affect arises when administering a combination of these compounds and certain additional anti-cancer agents. The inventors have determined, contrary to the prior knowledge and experience, how to successfully use a combination of TCN, TCN-P, TCN-PM and related compounds and at least one additional anti-cancer agent to treat tumors and cancer by one or a combination of (i) administering TCN, TCN-P, TCN-PM and related compounds and at least one additional anti-cancer agent to patients who exhibit enhanced sensitivity to the triciribine; (ii) use of a described dosage level that minimizes the toxicity of the drugs but yet still exhibits efficacy; or (iii) use of a described dosage regimen that minimizes the toxicity of the drugs.

The inventions also encompasses methods useful to treat tumors and cancers that are particularly susceptible to the toxic effects of TCN, TCN-P, TCN-PM and/or related com-

pounds. In another embodiment, methods are provided for treating a tumor in a mammal, particularly a human that includes (i) obtaining a biological sample from the tumor; (ii) determining whether the tumor overexpresses an Akt kinase, and (iii) treating the tumor that overexpresses Akt kinase with TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein.

In another embodiment, the level of Akt kinase expression can be determined by assaying the tumor or cancer for the presence of a phosphorylated Akt kinase, for example, by using an antibody that can detect the phosphorylated form. In another embodiment, the level of Akt expression can be determined by assaying a tumor or cancer cells obtained from a subject and comparing the levels to a control tissue. In certain embodiments, the Akt can be overexpressed at least 2, 2.5, 3 or 5 fold in the cancer sample compared to the control. In certain embodiments, the overexpressed Akt kinase can be a hyperactivated and phosphorylated Akt kinase.

In another aspect of the invention, dosing regimens are provided that limit the toxic side effects of TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein. In another embodiment, such dosing regimens minimize or eliminate toxic side effects, including, but not limited to, hepatotoxicity, thrombocytopenia, hyperglycemia, vomiting, hypocalcemia, anemia, hypoalbuminemia, myelosuppression, hypertriglyceridemia, hyperamylasemia, diarrhea, stomatitis and/or fever.

In another embodiment, the administration of TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein, provides at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15, 20, or 25% of the subjects.

In another embodiment, a method is provided to treat a subject which has been diagnosed with a tumor by administering to the subject an effective amount of TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein according to a dosing schedule that includes administering the drug approximately one time per week for approximately three weeks followed by a one week period wherein the drug is not administered.

In another embodiment, methods are provided to treat tumor or cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein each one time per week. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be administered as a single bolus dose over a short period of time, for example, about 5, 10 or 15 minutes.

In further embodiments, dosing schedules are provided in which the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein are administered via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the continuous administration can be repeated at least once a week, once every two weeks and/or once a month. In other embodiments, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be administered at least once every three weeks. In further

embodiments, the compounds can be administered at least once a day for at least 2, 3, 4 or 5 days.

In further embodiments, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents as described herein can be administered to patients in an amount that is effective in causing tumor regression. The administration the TCN, TCN-P, TCN-PM and/or related compounds and at least one additional anti-cancer agent can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least about 2, 5, 10, 15, 20, 30 or 50 mg/m² of the TCN, TCN-P, TCN-PM and/or related compounds and at least about 0.1, 1, 2, 5, 10, 15, 20, 30 or 50 mg of one or more additional anti-cancer agents, as described herein can be administered to a subject. The administration of the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen can include administering less than 20 mg/m² of the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein. In one embodiment, less than 10 mg/m² of the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be administered once a week. In further embodiments, dosages of or less than 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of the TCN, TCN-P, TCN-PM and/or related compounds and at least about 0.1, 1, 2, 5, 10, 15, 20, 30 or 50 mg of one or more additional anti-cancer agents, as described herein can be administered to a subject. In another embodiment, less than 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be administered to a subject via continuous infusion for at least five days. In particular embodiments, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein as disclosed herein can be used for the treatment of pancreatic, prostate, colorectal and/or ovarian cancer.

In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be used to prevent and/or treat a carcinoma, sarcoma, lymphoma, leukemia, and/or myeloma. In other embodiments of the invention, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be used to treat solid tumors. In still further embodiments, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein and compositions disclosed herein can be used for the treatment of a tumor or cancer, such as, but not limited to cancer of the following organs or tissues: breast, prostate, bone, lung, colon, including, but not limited to colorectal, urinary, bladder, non-Hodgkin lymphoma, melanoma, kidney, renal, pancreas, pharynx, thyroid, stomach, brain, and/or ovaries.

In a particular embodiment, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be used for the treatment of pancreatic, breast, colorectal and/or ovarian cancer. In further embodiments of the invention, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as

described herein disclosed herein can be used in the treatment of angiogenesis-related diseases. In certain embodiments, methods are provided to treat leukemia via continuous infusion of, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes via continuous infusion for at least 24, 48, 72 or 96 hours. In other embodiments, the continuous infusion can be repeated, for example, at least once every two, three or four weeks.

In another embodiment, there is provided a method for the treatment of tumors, cancer, and others disorders associated with an abnormal cell proliferation in a host, the method including administering to the host an effective amount of the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein optionally in combination with a pharmaceutically acceptable carrier.

In one aspect, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein and compositions can be administered in combination and can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In other embodiments, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein can be used to treat tumors or cancers resistant to one or more drugs, including the embodiments of tumors or cancers and drugs disclosed herein. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein as disclosed herein is administered in an effective amount for the treatment of a patient with a drug resistant tumor or cancer, for example, multidrug resistant tumors or cancer including, but not limited to, those resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In certain embodiments, a method is provided including administering to a host in need thereof an effective amount of TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein, or pharmaceutical composition including a TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes, in an effective amount for the treatment of tumors, cancer, and others disorders associated with an abnormal cell proliferation in a host.

In another embodiment, a method for the treatment of a tumor or cancer is provided including an effective amount of a TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents disclosed herein, or a salt, isomer, prodrug or ester thereof, to an individual in need thereof, wherein the cancer is for example, carcinoma, sarcoma, lymphoma, leukemia, or myeloma. The compound, or salt, isomer, prodrug or ester thereof, is optionally provided in a pharmaceutically acceptable composition including the appropriate carriers, such as water, which is formulated for the desired route of administration to an individual in need thereof. Optionally the compound is administered in combination or alternation with at least one additional therapeutic agent for the treatment of tumors or cancer.

Also within the scope of the invention is the use of a TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more additional anti-cancer agents, as described herein or a salt, prodrug or ester thereof in the treatment of a tumor or cancer, optionally in a pharmaceutically acceptable carrier; and the use of a TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more addi-

tional anti-cancer agents, as described herein or a salt, prodrug or ester thereof in the manufacture of a medicament for the treatment of cancer or tumor, optionally in a pharmaceutically acceptable carrier.

4. BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A-FIG. 1D demonstrates the identification of API-2 (tricitriline) as a candidate of Akt inhibitor from the NCI Diversity Set. FIG. 1A illustrates the chemical structure of API-2 (tricitriline). FIG. 1B demonstrates that API-2 inhibits phosphorylation levels of AKT2 in AKT2-transformed NIH3T3 cells. Wild type AKT2-transformed NIH3T3 cells were treated with API-2 (1 μ M) for indicated times and subjected to immunoblotting analysis with anti-phospho-Akt-T308 and -S473 antibodies (top and middle panels). The bottom panel shows expression of total AKT2. In FIG. 1C, it is shown that API-2 inhibits three isoforms of Akt. HEK293 cells were transfected with HA-Akt1, -AKT2 and -AKT3 and treated with API-2 (1 μ M) or wortmannin (15 μ M) prior to EGF stimulation, the cells were lysed and immunoprecipitated with anti-HA antibody. The immunoprecipitates were subjected to in vitro kinase assay (top) and immunoblotting analysis with anti-phospho-Akt-T308 (bottom) antibody. Middle panel shows expression of transfected Akt1, AKT2 and AKT3. FIG. 1D illustrates that API-2 did not inhibit Akt in vitro. In vitro kinase assay of constitutively active AKT2 recombinant protein in a kinase buffer containing 1 μ M API-2 (lane 3).

FIG. 2A-FIG. 2F demonstrates that API-2 does not inhibit PI3K, PDK1 and the closely related members of AGC kinase family. FIG. 2A demonstrates an in vitro PI3K kinase assay. HEK293 cells were serum-starved and treated with API-2 (1 μ M) or Wortmannin (15 μ M) for 30 minutes prior to EGF stimulation. Cells were lysed and immunoprecipitated with anti-p110 α antibody. The immunoprecipitates were subjected to in vitro kinase assay using PI-4-P as substrate. FIG. 2B illustrates the effect of API-2 on in vitro PDK1 activation (top panel), closed circles show inhibition by API-2. Open circles show inhibition by the positive control staurosporine, which is a potent PDK1 inhibitor (IC₅₀=5 nM). Bottom panels are immunoblotting analysis of HEK293 cells that were transfected with Myc-PDK1 and treated with wortmannin or API-2 prior to EGF stimulation. The immunoblots were detected with indicated antibodies. FIG. 2C illustrates an immunoblot analysis of phosphorylation levels of PKC α with anti-phospho-PKC α T638 (top) and total PKC α (bottom) antibodies following treatment with API-2 or a non-selective PKC inhibitor Ro31-8220. FIG. 2D shows an in vitro SGK kinase assay. HEK293 cells were transfected with HA-SGK and treated with API-2 or wortmannin prior to EGF stimulation. In vitro kinase was performed with HA-SGK immunoprecipitates using MBP as substrate (top). Bottom panel shows the expression of transfected HA-SGK. FIG. 2E illustrates the results of a PKA kinase assay. Immuno-purified PKA was incubated in ADB buffer (Upstate Biotechnology Inc) containing indicated inhibitors (API-2 or PKAI) and substrate Kemptide. The kinase activity was quantified. In FIG. 2F, a western blot is shown. OVCAR3 cells were treated with API-2 for indicated times. Cell lysates were immunoblotted with indicated anti-phospho-antibodies (panels 1-4) and anti-actin antibody (bottom).

FIG. 3A-FIG. 3C demonstrates that API-2 inhibits Akt activity and cell growth and induces apoptosis in human cancer cells with elevated Akt. FIG. 3A is a western blot, following treatment with API-2, phosphorylation levels of

Akt were detected with anti-phospho-Akt-T308 antibody in indicated human cancer cell lines. The blots were reprobed with anti-total Akt antibody (bottom panels). In FIG. 3B, a cell proliferation assay is shown. Cell lines as indicated in the figure were treated with different doses of API-2 for 24 h and 48 h and then analyzed with CellTiter 96 Cell Proliferation Assay kit (Promega). FIG. 3C provides an apoptosis analysis. Cells were treated with API-2 and stained with annexin V and PI and analyzed by FACScan.

FIG. 4A-FIG. 4D shows that API-2 inhibits downstream targets of Akt and exhibits anti-tumor activity in cancer cell lines with elevated Akt in mouse xenograft. In FIG. 4A, it is demonstrated that API-2 inhibits Akt phosphorylation of tuberlin, Bad, AFX and GSK-3 β . Following treatment with API-2, OVAR3 cells were lysed and immunoblotted with indicated antibodies. FIG. 4B shows that API-2 inhibits tumor growth. Tumor cells were subcutaneously injected into nude mice with low level of Akt cells on left side and elevated level of Akt cells on right side. When the tumors reached an average size of about 100-150 mm³, animals were treated with either vehicle or 1 mg/kg/day API-2. Each measurement represents an average of 10 tumors. FIG. 4C illustrates a representation of the mice with OVCAR3 (right) and OVCAR5 (left) xenograft treated with API-2 or vehicle (control). FIG. 4D shows examples of tumor size (bottom) and weight (top) at the end of experiment. In FIG. 4E, immunoblot analysis of tumor lysates was performed with anti-phospho-Akt-S473 (top) and anti-AKT2 (bottom) antibodies in OVCAR-3-derived tumors that were treated (T3 and T4) and untreated (T1 and T2) with API-2.

FIG. 5 shows that API-2 (tricitiribine) inhibits Akt kinase activity in vitro. In vitro kinase assay was performed with recombinant of PDK1 and Akt in a kinase buffer containing phosphatidylinositol-3,4,5-P3 (PIP3), API-2 and histone H2B as substrate. After incubation of 30 min, the reactions were separated by SDS-PAGE and exposed in a film.

FIG. 6A-FIG. 6D provides the mRNA and amino acid sequence of human Akt1, restriction enzyme sites are also noted.

FIG. 7A-FIG. 7D provides the mRNA and amino acid sequence of human Akt2 restriction enzyme sites are also noted.

FIG. 8A-FIG. 8D provides the mRNA and amino acid sequence of human Akt3 restriction enzyme sites are also noted.

FIG. 9 shows the synergistic effect of a combination of tricitiribine and Taxol on different cancer cell lines.

FIG. 10A-FIG. 10C shows growth inhibition by the combination of trastuzumab and Akt/mTOR pathway inhibitors. PTEN anti-sense or non-specific oligonucleotide transfected BT474.m1 cells were treated with inhibitors of the Akt/mTOR pathway alone or in combination with trastuzumab and the relative cell growth was assessed. FIG. 10A shows a panel of Akt/mTOR inhibitors. Growth inhibition was assessed in PTEN AS transfected BT474.m1 cells. The doses shown are: Tricitiribine (TCN) 1 μ M; RAD001 0.2 nM; QLT0267 10 μ M; KP 372-1 0.05 μ M; 4ADPIB 5 μ M; Edelfosine 7.5 μ M; and trastuzumab (Ttzm) 2 μ g/ml. The standard deviation (SD) in the percent growth inhibition is indicated. Results shown are the combined data from 2-3 experiments with triplicates of each treatment within each experiment. FIG. 10B shows TCN inhibits cell growth in combination with trastuzumab. BT474.m1 cells were transfected with PTEN AS oligonucleotide or non-specific (NS) oligonucleotide, treated with trastuzumab and TCN, alone and in combination, at multiple doses of TCN and assayed for growth inhibition. Trastuzumab was administered at a

single concentration. FIG. 10C shows RAD001 inhibits cell growth in combination with trastuzumab. BT474.m1 cells were transfected with PTEN AS oligonucleotide or non-specific (NS) oligonucleotide, treated with trastuzumab and RAD001, alone and in combination, at multiple doses of RAD001 and assayed for growth inhibition. For FIGS. 10B and 10C: * indicates a significant difference in growth inhibition following combination treatment as compared to either trastuzumab or TCN/TAD001 alone. $P < 0.05$ was considered significant. Error bars depict the SEM.

FIG. 11 shows the synergistic effects on apoptosis. Twenty-four hours after plating, PTEN AS and NS transfected BT474.m1 cells were treated as indicated with trastuzumab (Ttzm), TCN and/or RAD001 at the following concentrations: trastuzumab 2 μ g/ml; tricitiribine 2.5 μ M; RAD001 0.4 nM. Apo-BrdU TUNEL assays were performed to assess apoptosis. The experiment was performed 3 times and the data shown is the mean apoptosis. Error bars depict the standard deviation. Trastuzumab+tricitiribine treatment significantly induced apoptosis ($p < 0.01$) as compared with all other treatments.

FIG. 12A-FIG. 12B shows inhibition of Akt and p70S6K activity. To evaluate the effects of these drugs on the Akt/mTOR pathway, PTEN AS and NS oligonucleotides were transfected into BT474.m1 cells. Two days later, the cells were treated for 2 hours with trastuzumab and tricitiribine (TCN) (FIG. 12A) or trastuzumab and RAD001 (FIG. 12B). Total cell lysates were collected, separated by SDS-PAGE and immunoblotted as indicated. The concentration of trastuzumab was 2 μ g/ml, tricitiribine was 2.5 μ M and RAD001 was 0.4 nM. The experiments were repeated at least twice to insure that results were reproducible.

FIG. 13A-FIG. 13B shows combination treatments inhibited tumor growth in a SCID mice xenograft model. SCID mice received BT474.m1 breast cancer cell xenografts in mammary fat pad. The xenografts grew for 3 weeks to generate tumors with an average size of 100-150 mm³. PTEN antisense oligonucleotides, trastuzumab, tricitiribine (FIG. 13A) and RAD001 (FIG. 13B) were administered. The tumors were measured twice weekly with calipers and tumor size was averaged for each treatment group. Error bars denote the standard error of the mean. * indicates a significant difference in growth inhibition following combination treatment as compared to either trastuzumab (Ttzm), TCN or DMSO alone. $P < 0.05$ was considered significant.

FIG. 14A-14B shows that cisplatin (CDDP) resistant cell lines endogenously express hyperactive Akt. FIG. 14A shows a graph of cell survival for two ovarian cancer cell lines (A2780S and OV2008) and their cisplatin resistant counterparts (A2780CP and C13, respectively) treated with increasing amounts of cisplatin. The cisplatin resistant variants, A2780CP and C13, demonstrated increased survival in response to cisplatin. FIG. 14B shows that a comparison of cisplatin sensitive ovarian cancer cells, A2780S and OV2008, to their cisplatin resistant counterparts. The upper panels illustrate hyperactive Akt in the cisplatin resistant cell lines C13 and A2780S. The lower panels illustrate comparable total amounts of Akt are present in each cell type.

FIG. 15 illustrates that tricitiribine (TCN) (API-2) overcomes cisplatin resistance. Cisplatin resistant cell lines, A2780S and C13, were treated with 0, 5, 10, and 20 μ M cisplatin. The white and striped bars represent treatment with cisplatin only. The black and grey bars represent treatment with cisplatin and 10 μ M TCN. The co-treatment with TCN decreased cell survival, indicating TCN overcomes cisplatin resistance.

FIG. 16 illustrates the synergistic effect seen with TCN and cisplatin on the cell survival of C13 ovarian cancer cells. C13 cells were treated with 0, 1, 5, 10 and 20 μ M TCN, with and without 10 μ M cisplatin. The combination of cisplatin and TCN synergistically reduced cell survival.

FIG. 17A-FIG. 17B illustrates that TCN enhances the ability of cisplatin to inhibit the growth of C13 ovarian cancer cells in nude mice xenografts. Nude mice received C13 xenografts, followed by treatment with vehicle (DMSO), cisplatin alone, TCN alone, or cisplatin and TCN, followed by analysis of tumor progression at 7, 14, 21, or 28 days post-inoculation. The left panel FIG. 17A shows a graph illustrating that the combination of cisplatin and TCN significantly reduces tumor growth. The right panel FIG. 17B depicts representative tumor mass in inoculated mice receiving the represented treatments.

FIG. 18A-FIG. 18B shows the rapid activation of Akt pathway by mTOR inhibitors in human ovarian cancer cells. The upper panels FIG. 18A show western blots of OV3 cell lysates following the treatment with 1 nM mTOR inhibitor rapamycin for the time indicated. Membranes were probed with antibodies specific to phosphorylated isoforms and to the total protein for comparison of Akt, p70S6K and FKHL1. The lower panels FIG. 18B illustrate western blots of MCF7 cell lysates following treatment with the mTOR inhibitor RAD001 for the time indicated. Membranes were probed with antibodies specific to phosphorylated isoforms and to the total protein for comparison of Akt and p70S6K.

FIG. 19A-FIG. 19B shows that activation of AKT pathway via mTOR inhibitors is attenuated by TCN. The left panel FIG. 19A illustrates the effect of TCN on Akt phosphorylation with rapamycin treatment in OV3 cells. The right panel FIG. 19B illustrates the same effect of TCN on Akt phosphorylation with RAD001 treatment in MCF7 cells.

FIG. 20A-FIG. 20C illustrates the enhanced effect on cell growth through the combination of TCN with RAD001 or rapamycin. The top left panel FIG. 20A is a graph illustrating cell number counts of DU-145 cells over six days treated with control, rapamycin alone, TCN alone, or TCN and rapamycin. The top right panel FIG. 20B is a graph illustrating cell number counts of MCF7 cells over six days treated with control, RAD001 alone, TCN alone, or TCN and RAD001. The bottom panel FIG. 20C is a graph illustrating cell number counts of OV3 cells over six days treated with control, rapamycin alone, TCN alone, or TCN and rapamycin.

FIG. 21A-FIG. 21B illustrates the induction of cell survival and chemoresistance through Aurora-A activation of Akt. The top panels in FIG. 21A show successfully transfection and expression of Aurora-A in A2780S, A2780CP and OV2008 ovarian cancer cell lines. The second panel in FIG. 21A is a graph illustrating that expression of Aurora-A in A2780S cells increases cell survival and resistance to paclitaxel. FIG. 21B is a graph illustrating that expression of Aurora-A in A2780S cells increases cell survival and resistance to cisplatin. The bottom panel (D) is a graph illustrating that expression of Aurora-A in OV2008 cells increases cell viability and resistance to cisplatin.

FIG. 22A-FIG. 22B shows the effect of TCN (API-2) on Aurora-A induced cisplatin resistance. FIG. 22A shows A2780S cells transfected with empty vector or Aurora-A. FIG. 22B illustrate OV2008 cells transfected with empty vector or Aurora-A, followed by treatment with cisplatin alone, API-2 alone or API-2 and cisplatin.

FIG. 23A-FIG. 23C illustrates the effect of Aurora-A on cytochrome c release and Akt activation. FIG. 23A shows

immunofluorescence of cytochrome c in A2780S cells transfected with empty vector or Aurora-A treated with or without cisplatin. Cisplatin induced release of cytochrome c is inhibited by expression of Aurora-A. FIG. 23B is a western blot showing expression of Aurora-A increases phosphorylation of Akt in cisplatin sensitive cells. FIG. 23C shows that increased concentrations of Aurora-A increase the phosphorylation and activation of Akt.

FIG. 24A-FIG. 24D shows a schematic cartoon of the roles API-2, Aurora-A, p53 and Akt play in cell survival.

FIG. 25 shows the effect of the Src family tyrosine kinase inhibitor dasatinib (Sprycel®) on Src phosphorylation and Akt phosphorylation in A375 cells. Inhibition of Src tyrosine kinases increases phosphorylation of Akt.

FIG. 26A-FIG. 26C shows bortezomib synergistically enhances the effect of API-2 (TCN) on the survival myeloma cells. H929 cells (multiple myeloma derived) were treated with 0, 2.5, 5, 10, 20, or 40 nM bortezomib alone or in the presence of 10 μ M API-2, or with 0, 2.5, 5, 10, 20, or 40 μ M API-2 alone. Cell survival rates were calibrated and graphed (FIG. 26B). U266 cells (multiple myeloma derived) were treated with 0, 2.5, 5, 10, 20, or 40 μ M API-2 alone or in the presence of 10 nM API-2, or with 0, 2.5, 5, 10, 20, or 40 nM bortezomib alone. Cell survival rates were calibrated and graphed (FIG. 26C).

FIG. 27A-FIG. 27C shows the synergistic effect of bortezomib and API-2 (TCN) on mantle cell lymphoma cells. Jeko-1 cells (mantle cell lymphoma derived) were treated with 0, 2.5, 5, 10, 20, 40, or 80 nM bortezomib, alone or in the presence of 5 μ M API-2, or with 0, 1.25, 2.5, 5, 10, 20, or 40 μ M API-2. Cell survival rates were calibrated and graphed.

5. DETAILED DESCRIPTION OF THE INVENTION

The inventors have determined, contrary to the prior art and experience, how to successfully use TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more taxanes to treat tumors and cancer by one or a combination of (i) administering TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane; (ii) using a described dosage level that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane but yet still exhibits efficacy; or (iii) using a described dosage regimen that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane.

The inventors have determined, contrary to the prior art and experience, how to successfully use TCN, TCN-P, TCN-PM and/or related compounds in combination with a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, to treat tumors and cancer by one or a combination of (i) administering TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, to patients, who according to a diagnostic test described below, exhibit enhanced sensitivity to the TCN, TCN-P, TCN-PM and/or related compounds and/or the trastuzumab or a salt thereof; (ii) using a described dosage level that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the trastuzumab or a salt thereof but yet still exhibits efficacy; or (iii) using a described dosage regimen that

11

minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the trastuzumab or a salt thereof.

The inventors have determined, contrary to the prior art and experience, how to successfully use TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more anthracycline analogs to treat tumors and cancer by one or a combination of (i) administering TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the TCN, TCN-P, TCN-PM and/or related compounds and/or the anthracycline analogs; (ii) using a described dosage level that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the anthracycline analogs but yet still exhibits efficacy; or (iii) using a described dosage regimen that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the anthracycline analogs.

The inventors have determined, contrary to the prior art and experience, how to successfully use TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more erlotinib-like compounds to treat tumors and cancer by one or a combination of (i) TCN, TCN-P, TCN-PM and/or related compounds and one or more erlotinib-like compounds to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the TCN, TCN-P, TCN-PM and/or related compounds and/or the erlotinib-like compounds; (ii) using a described dosage level that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the erlotinib-like compounds but yet still exhibits efficacy; or (iii) using a described dosage regimen that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the erlotinib-like compounds.

The inventors have determined, contrary to the prior art and experience, how to successfully use TCN, TCN-P, TCN-PM and/or related compounds in combination with one or more platinum compounds to treat tumors and cancer by one or a combination of (i) administering TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds only to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the TCN, TCN-P, TCN-PM and/or related compounds and/or the one or more platinum compounds; (ii) using a described dosage level that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the one or more platinum compounds but yet still exhibits efficacy; or (iii) using a described dosage regimen that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the one or more platinum compounds.

The inventors have determined, contrary to the prior art and experience, how to successfully use TCN, TCN-P, TCN-PM and/or related compounds in combination with bortezomib and derivatives thereof analogs to treat tumors and cancer by one or a combination of (i) administering TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs only to patients which according to a diagnostic test described below, exhibit enhanced sensitivity to the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs; (ii) using a described dosage level that minimizes the toxicity of the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs but yet still exhibits efficacy; or (iii) using a described dosage regimen that minimizes the toxicity of the

12

TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs.

5.1. DEFINITIONS

As used herein, the term “compounds of the invention” refers to compounds of formula I-XXIII, and combinations thereof.

As used herein, the terms “cancer” and “cancerous” refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth, i.e., proliferative disorders. Examples of such proliferative disorders include cancers such as carcinoma, lymphoma, blastoma, sarcoma, and leukemia, as well as other cancers disclosed herein. More particular examples of such cancers include breast cancer, prostate cancer, colon cancer, squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, pancreatic cancer, cervical cancer, ovarian cancer, liver cancer, e.g., hepatic carcinoma, bladder cancer, colorectal cancer, endometrial carcinoma, kidney cancer, and thyroid cancer.

Other non-limiting examples of cancers are basal cell carcinoma, biliary tract cancer; bone cancer; brain and CNS cancer; choriocarcinoma; connective tissue cancer; esophageal cancer; eye cancer; cancer of the head and neck; gastric cancer; intra-epithelial neoplasm; larynx cancer; lymphoma including Hodgkin's and Non-Hodgkin's lymphoma; melanoma; myeloma; neuroblastoma; oral cavity cancer (e.g., lip, tongue, mouth, and pharynx); pancreatic cancer; retinoblastoma; rhabdomyosarcoma; rectal cancer; cancer of the respiratory system; sarcoma; skin cancer; stomach cancer; testicular cancer; uterine cancer; cancer of the urinary system, as well as other carcinomas and sarcomas.

As used herein, the term “tumor” refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues. For example, a particular cancer may be characterized by a solid mass tumor. The solid tumor mass, if present, may be a primary tumor mass. A primary tumor mass refers to a growth of cancer cells in a tissue resulting from the transformation of a normal cell of that tissue. In most cases, the primary tumor mass is identified by the presence of a cyst, which can be found through visual or palpation methods, or by irregularity in shape, texture or weight of the tissue. However, some primary tumors are not palpable and can be detected only through medical imaging techniques such as X-rays (e.g., mammography), or by needle aspirations. The use of these latter techniques is more common in early detection. Molecular and phenotypic analysis of cancer cells within a tissue will usually confirm if the cancer is endogenous to the tissue or if the lesion is due to metastasis from another site.

The term alkyl, as used herein, unless otherwise specified, includes a saturated straight, branched, or cyclic, primary, secondary, or tertiary hydrocarbon of for example C₁ to C₂₄, and specifically includes methyl, trifluoromethyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, t-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, cyclohexylmethyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. The alkyl is optionally substituted, e.g., with one or more substituents such as halo (F, Cl, Br or I), (e.g., CF₃, 2-Br-ethyl, CH₂F, CH₂Cl, CH₂CF₃ or CF₂CF₃), hydroxyl (e.g. CH₂OH), amino (e.g., CH₂NH₂, CH₃NHCH₃ or CH₂N(CH₃)₂), alkylamino, arylamino, alkoxy, aryloxy, nitro, azido (e.g. CH₂N₃), cyano (e.g. CH₂CN), sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as neces-

13

sary, as known to those skilled in the art, for example, as taught in Greene, et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, Second Edition, 1991, hereby incorporated by reference.

The term lower alkyl, as used herein, and unless otherwise specified, refers to a C₁ to C₄ saturated straight, branched, or if appropriate, a cyclic (for example, cyclopropyl) alkyl group, including both substituted and unsubstituted forms.

The term alkylamino or arylamino includes an amino group that has one or two alkyl or aryl substituents, respectively.

The term amino acid includes naturally occurring and synthetic α , β , γ or δ amino acids, and includes but is not limited to, amino acids found in proteins, i.e. glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine. In a preferred embodiment, the amino acid is in the L-configuration. Alternatively, the amino acid can be a derivative of alanyl, valinyl, leucinyl, isoleuccinyl, prolinyl, phenylalaninyl, tryptophanyl, methioninyl, glycyl, serinyl, threoninyl, cysteinyl, tyrosinyl, asparaginyl, glutaminyl, aspartoyl, glutaroyl, lysinyl, argininyl, histidinyl, β -alanyl, β -valinyl, β -leucinyl, β -isoleuccinyl, β -prolinyl, β -phenylalaninyl, β -tryptophanyl, β -methioninyl, β -glycinyl, β -serinyl, β -threoninyl, β -cysteinyl, β -tyrosinyl, β -asparaginyl, β -glutaminyl, β -aspartoyl, β -glutaroyl, β -lysyl, β -argininyl or β -histidinyl. When the term amino acid is used, it is considered to be a specific and independent disclosure of each of the esters of a natural or synthetic amino acid, including but not limited to α , β , γ or δ glycine, alanine, valine, leucine, isoleucine, methionine, phenylalanine, tryptophan, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartate, glutamate, lysine, arginine and histidine in the D and L-configurations.

The term "protected" as used herein and unless otherwise defined includes a group that is added to an oxygen, nitrogen, sulfur or phosphorus atom to prevent its further reaction or for other purposes. A wide variety of oxygen and nitrogen protecting groups are known to those skilled in the art of organic synthesis (see Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, Inc., New York, N.Y., 1999).

The term aryl, as used herein, and unless otherwise specified, includes phenyl, biphenyl, or naphthyl, and preferably phenyl. The aryl group is optionally substituted with one or more moieties such as halo, hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate, either unprotected, or protected as necessary, as known to those skilled in the art, for example, as taught in Greene, et al., *Protective Groups in Organic Synthesis*, John Wiley and Sons, 3rd Ed., 1999.

The term alkaryl or alkylaryl includes an alkyl group with an aryl substituent. The term aralkyl or arylalkyl includes an aryl group with an alkyl substituent.

The term halo, as used herein, includes chloro, bromo, iodo, and fluoro.

The term acyl includes a carboxylic acid ester in which the non-carbonyl moiety of the ester group is selected from straight, branched, or cyclic alkyl or lower alkyl, alkoxyalkyl including methoxymethyl, aralkyl including benzyl, aryloxyalkyl such as phenoxymethyl, aryl including phenyl optionally substituted with halogen, C₁ to C₄ alkyl or C₁ to C₄ alkoxy, sulfonate esters such as alkyl or aralkyl sulfonyl including methanesulfonyl, the mono, di or triphosphate ester, trityl or monomethoxytrityl, substituted benzyl, tri-

14

alkylsilyl (e.g. dimethyl-t-butylsilyl) or diphenylmethylsilyl. Aryl groups in the esters optimally include a phenyl group. The term "lower acyl" refers to an acyl group in which the non-carbonyl moiety is lower alkyl.

As used herein, the term "substantially free of" or "substantially in the absence of" with respect to enantiomeric purity, refers to a composition that includes at least 85% or 90% by weight, preferably 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the designated enantiomer. In a preferred embodiment, in the methods and compounds of this invention, the compounds are substantially free of other enantiomers.

Similarly, the term "isolated" refers to a compound composition that includes at least 85% or 90% by weight, preferably 95% to 98% by weight, and even more preferably 99% to 100% by weight, of the compound, the remainder including other chemical species or enantiomers.

The term "independently" is used herein to indicate that the variable, which is independently applied, varies independently from application to application. Thus, in a compound such as RⁿXYR^m, wherein Rⁿ is "independently carbon or nitrogen," both Rⁿ can be carbon, both R^m can be nitrogen, or one Rⁿ can be carbon and the other R^m nitrogen.

The term "pharmaceutically acceptable salt or prodrug" is used throughout the specification to describe any pharmaceutically acceptable form (such as an ester, phosphate ester, salt of an ester or a related group) of a compound, which, upon administration to a patient, provides the compound. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. Pharmaceutically acceptable prodrugs refer to a compound that is metabolized, for example hydrolyzed or oxidized, in the host to form the compound of the present invention. Typical examples of prodrugs include compounds that have biologically labile protecting groups on a functional moiety of the active compound. Prodrugs include compounds that can be oxidized, reduced, aminated, deaminated, hydroxylated, dehydroxylated, hydrolyzed, dehydrolyzed, alkylated, dealkylated, acylated, deacylated, phosphorylated, dephosphorylated to produce the active compound.

The term "pharmaceutically acceptable esters" as used herein, unless otherwise specified, includes those esters of one or more compounds, which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of hosts without undue toxicity, irritation, allergic response and the like, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.

The term "subject" as used herein refers to an animal, preferably a mammal, most preferably a human. Mammals can include non-human mammals, including, but not limited to, pigs, sheep, goats, cows (bovine), deer, mules, horses, monkeys and other non-human primates, dogs, cats, rats, mice, rabbits or any other known or disclosed herein.


The term "TCN, TCN-P, TCN-PM and/or related compounds" includes compounds encompassed by the compounds of Formulas I to IV.

5.2. COMPOUNDS OF THE INVENTION


As used herein and unless otherwise indicated, the term "TCN, TCN-P, TCN-PM and/or related compounds," "tricitiribine," "tricitiribine phosphate," and "tricitiribine phosphate

15


monohydrate" and "tricitriline and related compounds" refers to compounds encompassed by the following structures:




Formula I



Formula II



Formula III



Formula IV

wherein each R_2' , R_3' and R_5' are independently hydrogen, optionally substituted phosphate or phosphonate (including mono-, di-, or triphosphate or a stabilized phosphate prod-
rug); acyl (including lower acyl); alkyl (including lower alkyl); amide, sulfonate ester including alkyl or arylalkyl; sulfonyl, including methanesulfonyl and benzyl, wherein the phenyl group is optionally substituted with one or more
substituents as for example as described in the definition of an aryl given herein; optionally substituted arylsulfonyl; a

16


lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound wherein R_2' , R_3' or R_5' is independently H or mono-, di- or tri-phosphate;

wherein R^x and R^y are independently hydrogen, optionally substituted phosphate; acyl (including lower acyl); amide, alkyl (including lower alkyl); aromatic, polyoxyalkylene such as polyethyleneglycol, optionally substituted arylsulfonyl; a lipid, including a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group. In one embodiment, the compound is administered as a 5'-phosphoether lipid or a 5'-ether lipid.

R_1 and R_2 each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl.

In one embodiment, R_2' and R_3' are hydrogen. In another embodiment, R_2' and R_5' are hydrogen. In yet another embodiment, R_2' , R_3' and R_5' are hydrogen. In yet another embodiment, R_2' , R_3' , R_5' , R_1 and R_2 are hydrogen.

In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds has the following structure:




Formula V

wherein R_3 is H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, NH_2 , NHR^4 , $N(R^4)_2$, aryl, alkoxyalkyl, aryloxyalkyl, or substituted aryl; and


each R^4 independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl, or aryl. In a subembodiment, R_3 is a straight chained C1-11 alkyl, iso-propyl, t-butyl, or phenyl.

In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds provided herein have the following structure:

17




Formula VI




Formula VII

In another embodiment, the tricyclic compounds provided herein have the following structure:



Formula VIII

In another embodiment, the tricyclic compounds provided herein have the following structure:



Formula IX

wherein R_6 is H, alkyl, (including lower alkyl) alkenyl, alkynyl, alkoxyalkyl, hydroxyalkyl, arylalkyl, cycloalkyl, NH_2 , NHR^4 , NR^4R^4 , CF_3 , CH_2OH , CH_2F , CH_2Cl , CH_2CF_3 , $C(Y^3)_3$, $C(Y^3)_2C(Y^3)$, $C(=O)OH$, $C(=O)OR^4$, $C(=O)-$


18

alkyl, $C(=O)-aryl$, $C(=O)-alkoxyalkyl$, $C(=O)NH_2$, $C(=O)NHR^4$, $C(=O)N(R^4)_2$, where each Y^3 is independently H or halo; and

each R^4 independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl, or aryl.

In a subembodiment, R_6 is ethyl, CH_2CH_2OH , or CH_2 -phenyl.

In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds provided herein have the following structure:




Formula X

wherein R_7 is H, halo, alkyl (including lower alkyl), alkenyl, alkynyl, alkoxy, alkoxyalkyl, hydroxyalkyl, cycloalkyl, nitro, cyano, OH , OR^4 , NH_2 , NHR^4 , NR^4R^4 , SH , SR^4 , CF_3 , CH_2OH , CH_2F , CH_2Cl , CH_2CF_3 , $C(Y^3)_3$, $C(Y^3)_2C(Y^3)$, $C(=O)OH$, $C(=O)OR^4$, $C(=O)-alkyl$, $C(=O)-aryl$, $C(=O)-alkoxyalkyl$, $C(=O)NH_2$, $C(=O)NHR^4$, $C(=O)N(R^4)_2$, or N_3 , where each Y^3 is independently H or halo; and

each R^4 independently is H, acyl including lower acyl, alkyl including lower alkyl such as but not limited to methyl, ethyl, propyl and cyclopropyl, alkenyl, alkynyl, cycloalkyl, alkoxy, alkoxyalkyl, hydroxyalkyl.

In a subembodiment, R_7 is methyl, ethyl, phenyl, chloro or NH_2 .


In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds provided herein have the following structures:




Formula XI

19

-continued



In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds provided herein have the following structure:



Formula XII

Formula XIII

20

In one embodiment, the taxane compound has the following structure:

5

Formula XV

10

15

20

25

30

35

40

wherein R^{11} each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, aryl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl.

In another embodiment, the taxane compound has the following structure:


30

Formula XVI

35

40

As used herein and unless otherwise indicated, the term “taxanes” refers to the following formula:



Formula XIV

50


55

wherein

R^{10} and R^{11} each are independently H, optionally substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkenyl, or alkynyl, aryl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkylsulfonyl.


60

Formula XVII



21


In another embodiment, the taxane compound has the following structure:



As used herein and unless otherwise indicated, the term “a molecule that modulates the HER2/neu (erbB2) receptor,” refers to compounds which are recombinant DNA-derived humanized monoclonal antibodies that selectively binds with high affinity in a cell-based assay to the extra-cellular domain of the human epidermal growth factor receptor 2 protein, HER2. In certain embodiments, the antibody is an IgG₁ kappa that contains human framework regions with the complementary-determining regions of a murine antibody (4D5) that binds to HER2.

As used herein and unless otherwise indicated, the term “trastuzumab” refers to trastuzumab or a salt thereof. In certain embodiment, the term trastuzumab refers to the compound with the DrugBank accession number BTID00098 and the trade name Herceptin, which is a recombinant humanized monoclonal antibody directed against the human epidermal growth factor receptor 2 (HER2). HER2 is over-expressed by many adenocarcinomas, particularly breast adenocarcinomas. Trastuzumab binds to HER2 on the cell surface of a tumor cell, thereby inducing an antibody-dependent cell-mediated cytotoxicity against tumor cells that overexpress HER2.

As used herein and unless otherwise indicated, the term “anthracycline analogs” refers to a compound of Formula XVIII or Formula XIX, wherein the compound of Formula XVIII has the following structure:



Formula XVIII

R₁, R₄, R₆, R₇ are each independently hydrogen, hydroxy, alkoxy;

R₂ and R₃ are each independently hydrogen, alkyl, alkoxy, hydroxy, halogen;


R₅ is hydrogen, an optionally substituted alkyl chained, branched or cyclic alkyl, hydroxy, alkoxy;

R₈ is CO-alkyl, CO-halogen substituted alkyl, CO-aryl or heteroaryl.

In other embodiments, the term “anthracycline compounds” refers to compound encompassed by Formula XIX having the following structure:

22


Formula XIX




R₁ and R₂ are each independently hydrogen, alkyl, alkoxy, hydroxy, halogen;

R₃ and R₄ are each independently hydrogen, optionally substituted alkyl chained, branched or cyclic alkyl, optionally substituted alkyl chained, branched or cyclic alcohol, optionally substituted alkyl chained, branched or cyclic amine.

In an illustrative embodiment of species encompassed by the term “anthracycline compounds of Formula XVIII or Formula XIX,” includes, but are limited to, a compound with any structure A-G:







Idarubicin




Daunorubicin

23

-continued


**24**

-continued



As used herein and unless otherwise indicated, the term “epidermal growth factor receptor inhibitor” refers to compounds that target the epidermal growth factor receptor (EGFR) tyrosine kinase, which is highly expressed and occasionally mutated in various forms of cancer. In certain embodiments, the compounds bind in a reversible fashion to the adenosine triphosphate (ATP) binding site of the receptor. In certain embodiments, two members of the EGFR family come together to form a homodimer. These then use the molecule of ATP to autophosphorylate each other, which causes a conformational change in their intracellular structure, exposing a further binding site for binding proteins that cause a signal cascade to the nucleus. By inhibiting the ATP, autophosphorylation is not possible and the signal is stopped. Illustrative examples of epidermal growth factor receptor inhibitor compounds include erlotinib-like compounds such as for example, gefitinib and erlotinib.

As used herein and unless otherwise indicated, the term “erlotinib-like compounds” refers to a compound of Formula XX:





Formula XX

wherein


each R_1 and R_2 is independently hydrogen, independently optionally substituted alkoxy, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl; and

each R_3 and R_4 is independently hydrogen, independently optionally substituted aromatic amine, heteroaromatic amine, or cyclic amine. In an illustrative embodiment, the erlotinib-like compounds include, but are limited to, the following structures:

25






As used herein and unless otherwise indicated, the term “platinum compounds” refers to a compound of formula XXI.



wherein:







R₁, R₂, R₃ and R₄ are each independently hydrogen, optionally substituted amine, aromatic amine, heteroaromatic amine, optionally substituted straight chained, branched or cyclic alkyl, aromatic, heteroaromatic or cyclic ring formed among R₁, R₂, R₃ and R₄.

In an illustrative embodiment, the platinum compounds include, but are limited to, platinum compounds of structures A-I:



26


-continued



As used herein and unless otherwise indicated, the term “bortezomib and derivatives thereof” refers to a compound of formula XXII:

27

Formula XXII




wherein

R_1 , R_2 , R_3 , R_4 , and R_5 each are independently H, optionally halogenated, substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkoxy, alkenyl, or alkynyl, aryl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkyl-sulfonyl.

In one embodiment, the bortezomib and derivatives thereof have the following structure:


Formula XXIII



wherein

R_1 , R_2 , R_3 , R_4 , and R_5 each are independently H, optionally halogenated, substituted straight chained, branched or cyclic alkyl (including lower alkyl), alkoxy, alkenyl, or alkynyl, aryl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, aralkyl-sulfonyl.

In another embodiment, the bortezomib and derivatives thereof analogs have the following structure:



28

Each of the compounds included herein is commercially available or can be made by syntheses known to those of ordinary skill in the art.

It is to be understood that the compounds disclosed herein may contain chiral centers. Such chiral centers may be of either the (R) or (S) configuration, or may be a mixture thereof. Thus, the compounds provided herein may be enantiomerically pure, or be stereoisomeric or diastereomeric mixtures. It is understood that the disclosure of a compound herein encompasses any racemic, optically active, polymorphic, or stereoisomeric form, or mixtures thereof, which preferably possesses the useful properties described herein, it being well known in the art how to prepare optically active forms and how to determine activity using the standard tests described herein, or using other similar tests which are well known in the art.

Examples of methods that can be used to obtain optical isomers of the compounds include the following:

(i) physical separation of crystals—a technique whereby macroscopic crystals of the individual enantiomers are manually separated. This technique can be used if crystals of the separate enantiomers exist, i.e., the material is a conglomerate, and the crystals are visually distinct;

(ii) simultaneous crystallization—a technique whereby the individual enantiomers are separately crystallized from a solution of the racemate, possible only if the latter is a conglomerate in the solid state;

(iii) enzymatic resolutions—a technique whereby partial or complete separation of a racemate by virtue of differing rates of reaction for the enantiomers with an enzyme

(iv) enzymatic asymmetric synthesis—a synthetic technique whereby at least one step of the synthesis uses an enzymatic reaction to obtain an enantiomerically pure or enriched synthetic precursor of the desired enantiomer;

(v) chemical asymmetric synthesis—a synthetic technique whereby the desired enantiomer is synthesized from an achiral precursor under conditions that produce asymmetry (i.e., chirality) in the product, which may be achieved using chiral catalysts or chiral auxiliaries;

(vi) diastereomer separations—a technique whereby a racemic compound is reacted with an enantiomerically pure reagent (the chiral auxiliary) that converts the individual enantiomers to diastereomers. The resulting diastereomers are then separated by chromatography or crystallization by virtue of their now more distinct structural differences and the chiral auxiliary later removed to obtain the desired enantiomer;

(vii) first- and second-order asymmetric transformations—a technique whereby diastereomers from the racemate equilibrate to yield a preponderance in solution of the diastereomer from the desired enantiomer or where preferential crystallization of the diastereomer from the desired enantiomer perturbs the equilibrium such that eventually in principle all the material is converted to the crystalline diastereomer from the desired enantiomer. The desired enantiomer is then released from the diastereomer;

(viii) kinetic resolutions—this technique refers to the achievement of partial or complete resolution of a racemate (or of a further resolution of a partially resolved compound) by virtue of unequal reaction rates of the enantiomers with a chiral, non-racemic reagent or catalyst under kinetic conditions;

(ix) enantiospecific synthesis from non-racemic precursors—a synthetic technique whereby the desired enantiomer is obtained from non-chiral starting materials and where the stereochemical integrity is not or is only minimally compromised over the course of the synthesis;

(x) chiral liquid chromatography—a technique whereby the enantiomers of a racemate are separated in a liquid mobile phase by virtue of their differing interactions with a stationary phase. The stationary phase can be made of chiral material or the mobile phase can contain an additional chiral material to provoke the differing interactions;

(xi) chiral gas chromatography—a technique whereby the racemate is volatilized and enantiomers are separated by virtue of their differing interactions in the gaseous mobile phase with a column containing a fixed non-racemic chiral adsorbent phase;

(xii) extraction with chiral solvents—a technique whereby the enantiomers are separated by virtue of preferential dissolution of one enantiomer into a particular chiral solvent;

(xiii) transport across chiral membranes—a technique whereby a racemate is placed in contact with a thin membrane barrier. The barrier typically separates two miscible fluids, one containing the racemate, and a driving force such as concentration or pressure differential causes preferential transport across the membrane barrier. Separation occurs as a result of the non-racemic chiral nature of the membrane which allows only one enantiomer of the racemate to pass through.

In some embodiments, tricitribine, tricitribine phosphate (TCN-P), tricitribine 5'-phosphate (TCN-P), or the DMF adduct of tricitribine (TCN-DMF) are provided. TCN can be synthesized by any technique known to one skilled in the art, for example, as described in Tetrahedron Letters, 1971. 49: p. 4757-4760. TCN-P can be prepared by any technique known to one skilled in the art, for example, as described in U.S. Pat. No. 4,123,524. The synthesis of TCN-DMF is described, for example, in INSERM, 1978. 81: p. 37-82. Other compounds related to TCN as described herein can be synthesized, for example, according to the methods disclosed in Gudmundsson K. S. et al., Nucleosides Nucleotides Nucleic Acids, 2001. 20(10-11): p. 1823-1830; Porcari A. R. et al., J Med Chem, 2000. 43(12): p. 2457-2463; Porcari A. R. et al., Nucleosides Nucleotides, 1999. 18(11-12): p. 2475-2497; Porcari A. R. et al., J Med Chem, 2000. 43(12): p. 2438-2448; Porcari A. R. et al., Nucleosides Nucleotides Nucleic Acids, 2003. 22(12): p. 2171-2193; Porcari A. R. et al., Nucleosides Nucleotides Nucleic Acids, 2004. 23(1-2): p. 31-39; Schweinsberg P. D. et al., Biochem Pharmacol, 1981. 30(18): p. 2521-2526; Smith K. L. et al., Bioorg Med Chem Lett, 2004. 14(13): p. 3517-3520; Townsend L. B. et al., Nucleic Acids Symp Ser, 1986. 1986(17): p. 41-44; and/or Wotring L. L. et al., Cancer Treat Rep, 1986. 70(4): p. 491-7, each of which is incorporated herein by reference.

5.3. PHARMACEUTICALLY ACCEPTABLE SALTS, HYDRATES AND PRODRUGS

In cases where compounds are sufficiently basic or acidic to form stable nontoxic acid or base salts, administration of the compound as a pharmaceutically acceptable salt may be appropriate. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium and magnesium, among numerous other acids well known in the pharmaceutical art. In particular, examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids, which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate,

benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Suitable inorganic salts may also be formed, including, sulfate, nitrate, bicarbonate, and carbonate salts.

In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds is a solvate, for example, tricitribine phosphate monohydrate (TCN-PM).

Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

Any of the nucleotides described herein can be administered as a nucleotide prodrug to increase the activity, bioavailability, stability or otherwise alter the properties of the nucleoside. A number of nucleotide prodrug ligands are known. In general, alkylation, acylation or other lipophilic modification of the mono, di or triphosphate of the nucleoside will increase the stability of the nucleotide. Examples of substituent groups that can replace one or more hydrogens on the phosphate moiety are alkyl, aryl, steroids, carbohydrates, including sugars, 1,2-diacylglycerol and alcohols. Many are described in R. Jones and N. Bischofberger, *Antiviral Research*, 27 (1995) 1-17. Any of these can be used in combination with the disclosed nucleosides to achieve a desired effect.

In one embodiment, the tricitribine or a related compound is provided as 5'-hydroxyl lipophilic prodrug. Nonlimiting examples of U.S. patents that disclose suitable lipophilic substituents that can be covalently incorporated into the nucleoside, preferably at the 5'-OH position of the nucleoside or lipophilic preparations, include U.S. Pat. Nos. 5,149,794 5,194,654 5,223,263 5,256,641 5,411,947 5,463,092 5,543,389 5,543,390 5,543,391 and 5,554,728, all of which are incorporated herein by reference.

Foreign patent applications that disclose lipophilic substituents that can be attached to the tricitribine or a related compound of the present invention, or lipophilic preparations, include WO 89/02733, WO 90/00555, WO 91/16920, WO 91/18914, WO 93/00910, WO 94/26273, WO/15132, EP 0 350 287, EP 93917054.4, and WO 91/19721, each of which is incorporated herein by reference.

Additional nonlimiting examples of derivatives of tricitribine or related compounds are those that contain substituents as described in the following publications. These derivatized tricitribine or related compounds can be used for the indications described in the text or otherwise as antiviral agents, including as anti-HIV or anti-HBV agents. Ho D. H. W., *Cancer Res.*, 1973 33: p. 2816-2820; Holy A. in *Advances in Antiviral Drug Design*, Vol. I, De Clercq (ed.), JAI Press, pp. 179-231; Hong C. I. et al. *Biochem Biophys Res Commun*, 1979. 88: p. 1223-1229; Hong C. I. et al., *J Med Chem*, 1980. 28: p. 171-177; Hostetler K. Y. et al., *J Biol Chem*, 1990. 266: p. 11714-11717; Hostetler K. Y. et al., *Antiviral Res*, 1994. 24: p. 59-67; Hostetler K. Y. et al., *Antimicrobial Agents Chemother*, 1994. 38: p. 2792-2797; Hunston R. N. et al., *J Med Chem*, 1984. 27: p. 440-444; Ji Y. H. et al., *J Med Chem*, 1990. 33: p. 2264-2270; Jones A. S. et al., *J Chem Soc Perkin Trans*, 1984. I: p. 1471-1474; Juodka B. A. and Smart J., *Coll Czech Chem Comm*, 1974. 39: p. 363-968; Kataoka S. et al., *Nucleic Acids Res Symp Ser*, 1989. 21: p. 1-2; Kataoka S. et al., *Heterocycles*, 1991. 32: p. 1351-1356; Kinchington D. et al., *Antiviral Chem Chemother*, 1992. 3: p. 107-112; Kodama K. et al., *Jpn J Cancer Res*, 1989. 80: p. 679-685; Korty M. and Engels J., *Naunyn-Schmiedeberg's Arch Pharmacol*, 1979. 10: p. 103-

111; Kumar A. et al., *J Med Chem*, 1990. 33: p. 2368-2375; LeBec C. and Huynh-dinh T., *Tetrahedron Lett*, 1991. 32: p. 6553-6556; Lichtenstein J. et al., *J Biol Chem*, 1960. 235: p. 457-465; Luethy J. et al., *Mitt Geg Lebensmittelunters Hyg*, 1981. 72: p. 131-133 (Chem. Abstr. 95, 127093); McGuigan C. et al., *Nucleic Acids Res*, 1989. 17: p. 6065-6075; McGuigan C. et al., *Antiviral Chem Chemother*, 1990. 1: p. 107-113; McGuigan C. et al., *Antiviral Chem Chemother*, 1990. 1: p. 355-360; McGuigan C. et al., *Antiviral Chem Chemother*, 1990. 1: p. 25-33; McGuigan C. et al., *Antiviral Res*, 1991. 15: p. 255-263; McGuigan C. et al., *Antiviral Res*, 1992. 17: p. 311-321; McGuigan C. et al., *Antiviral Chem Chemother*, 1993. 4: p. 97-101; McGuigan C. et al., *J Med Chem*, 1993. 36: p. 1048-1052, each of which is incorporated herein by reference.

Alkyl hydrogen phosphonate derivatives of the anti-HIV agent AZT may be less toxic than the parent nucleoside analogue. *Antiviral Chem. Chemother*. 5: 271-277; Meyer R. B. et al., *Tetrahedron Lett*, 1973. 269-272; Nagyvary J. et al., *Biochem Biophys Res Commun*, 1973. 55: p. 1072-1077; Namane A. et al., *J Med Chem*, 1992. 35: p. 3939-3044; Nargeot J. et al., *Natl. Acad. Sci. U.S.A.*, 1983. 80: p. 2395-2399; Nelson K. A. et al., *J Am Chem Soc*, 1987. 109: p. 4058-4064; Nerbonne J. M. et al., *Nature*, 1984. 301: p. 74-76; Neumann J. M. et al., *J Am Chem Soc*, 1989. 111: p. 4270-4277; Ohno R. et al., *Oncology*, 1991. 48: p. 451-455; Palomino E. et al., *J Med Chem*, 1989. 32: p. 622-625; Perkins R. M. et al., *Antiviral Res*, 1993. 20(Suppl. I): p. 84; Piantadosi C. et al., *J Med Chem*, 1991. 34: 1408-1414; Pompon A. et al., *Antiviral Chem Chemother*, 1994. 5: p. 91-98; Postemark T., *Annu Rev Pharmacol*, 1974. 14: p. 23-33; Prisbe E. J. et al., *J Med Chem*, 1986. 29: p. 671-675; Pucchi F. et al., *Antiviral Res*, 1993. 22: p. 155-174; Pugaeva V. P. et al., *Gig Trf Prof Zabol*, 1969. 13: p. 47-48 (Chem. Abstr. 72, 212); Robins R. K., *Pharm Res*, 1984. 11-18; Rosowsky A. et al., *J Med Chem*, 1982. 25: p. 171-178; Ross W., *Biochem Pharm*, 1961. 8: p. 235-240; Ryu E. K. et al., *J Med Chem*, 1982. 25: p. 1322-1329; Saffhill R. and Hume W. J., *Chem Biol Interact*, 1986. 57: p. 347-355; Saneyoshi M. et al., *Chem Pharm Bull*, 1980. 28: p. 2915-2923; Sastry J. K. et al., *Mol Pharmacol*, 1992. 41: p. 441-445; Shaw J. P. et al., 9th Annual AAPS Meeting, 1994. San Diego, Calif. (Abstract). Shuto S. et al., *Tetrahedron Lett*, 1987. 28: p. 199-202; Shuto S. et al., *Chem Pharm Bull*, 1988. 36: p. 209-217. One preferred phosphate prodrug group is the S-acyl-2-thioethyl group, also referred to as "SATE," each of which is incorporated herein by reference.

Additional examples of prodrugs that can be used are those described in the following patents and patent applications: U.S. Pat. Nos. 5,614,548, 5,512,671, 5,770,584, 5,962,437, 5,223,263, 5,817,638, 6,252,060, 6,448,392, 5,411,947, 5,744,592, 5,484,809, 5,827,831, 5,696,277, 6,022,029, 5,780,617, 5,194,654, 5,463,092, 5,744,461, 4,444,766, 4,562,179, 4,599,205, 4,493,832, 4,221,732, 5,116,992, 6,429,227, 5,149,794, 5,703,063, 5,888,990, 4,810,697, 5,512,671, 6,030,960, 2004/0259845, 6,670,341, 2004/0161398, 2002/082242, 5,512,671, 2002/0082242, and or PCT Publication Nos WO 90/11079, WO 96/39197, and/or WO 93/08807, each of which is incorporated herein by reference.

5.4. IN VIVO EFFICACY/DOSING REGIMENS

In another aspect of the invention, dosing regimens are provided that limit the toxic side effects of TCN, TCN-P, TCN-PM and/or related compounds. In one embodiment, such dosing regimens minimize the following toxic side

effects, including, but not limited to, hepatotoxicity, thrombocytopenia, hyperglycemia, vomiting, hypocalcemia, anemia, hypoalbuminemia, myelosuppression, hypertriglyceridemia, hyperamylasemia, diarrhea, stomatitis and/or fever.

In another embodiment, the administration TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes according to a dosing schedule that includes administering the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane is not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In another particular embodiment, about 1 to about 50 mg of one or more taxanes is administered. In a particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of one or more taxanes can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of one or more taxanes one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds as disclosed herein can be administered one time per week. In another particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of one or more taxanes can be administered one time per week.

In embodiments of the invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every

twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane can be administered according to any combination disclosed herein, for example, the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane are not administered.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of one or more taxanes can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of one or more taxanes either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds or related compounds can be administered once a week concurrently with less than about 30 mg of one or more taxanes. In another embodiment, less than 20 mg/m² of TCN or related compounds can be administered once a week and less than about 30 mg of one or more taxanes can be administered the following week.

In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30, 25, 20, 15, or mg of one or more taxanes can be administered to a subject. In another embodiment, less than 10 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of one or more taxanes can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

In another embodiment, the administration of TCN, TCN-P, TCN-PM and/or related compounds and a molecule that

modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, according to a dosing schedule that includes administering the TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, is not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 15 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In another particular embodiment, about 1 to about 50 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, is administered. In a particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds as disclosed herein can be administered one time per week. In another particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered one time per week.

In embodiments of the invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the

administration of the TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered according to any combination disclosed herein, for example, the TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the TCN, TCN-P, TCN-PM and/or related compounds and/or a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, are not administered.

The TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P, TCN-PM and/or related compounds and trastuzumab or a salt thereof can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered once a week concurrently with less than about 30 mg

of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof. In another embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered once a week and less than about 30 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered the following week.

In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 300, 250, 200, 150, or 100 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered to a subject. In another embodiment, less than 10 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds and less than about 300 mg of a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

In another embodiment, the administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs according to a dosing schedule that includes administering the TCN, TCN-P, TCN-PM and/or related compounds and/or the anthracycline analogs one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs are not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In another particular embodiment, about 1 to about 50 mg of an anthracycline derivative is administered. In a particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, or 100 mg of an anthracycline derivative can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of one or more anthracycline analogs one time per week. In par-

ticular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds as disclosed herein can be administered one time per week. In another particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, or 100 mg of an anthracycline derivative can be administered one time per week.

In embodiments of the present invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs can be administered according to any combination disclosed herein, for example, the TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs are not administered.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, or 100 mg of an anthracycline derivative can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than

about 30 mg of an anthracycline derivative either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered once a week concurrently with less than about 30 mg of an anthracycline derivative. In another embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered once a week and less than about 30 mg of an anthracycline derivative can be administered the following week.

In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30, 25, 20, 15, or 10 mg of an anthracycline derivative can be administered to a subject. In another embodiment, less than 10 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of an anthracycline derivative can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

In another embodiment, the administration of TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, according to a dosing schedule that includes administering the TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, is not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 15 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered according to a 42 day cycle. In another particular embodiment, about 1 to about 50 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof is administered. In a particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds as disclosed herein can be administered one time per week. In another particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, can be administered one time per week.

In embodiments of the present invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, can be administered according to any combination disclosed herein, for example, the TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the TCN, TCN-P, TCN-PM and/or related compounds and/or an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, are not administered.

The TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5,

4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered once a week concurrently with less than about 30 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof. In another embodiment, less than 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds can be administered once a week and less than about 30 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be administered the following week.

In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN or a related compound and less than about 300, 250, 200, 150, or 100 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be administered to a subject. In another embodiment, less than 10 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds and less than about 300 mg of an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

In another embodiment, the administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds provides at least a partial or complete response in vivo in at least 15-20% of the subjects. In particular embodiments, a partial response can be at least 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80 or 85% regression of the tumor. In other embodiments, this response can be evident in at least 15, 15, 20, 25, 30, 35, 40, 50, 55, 60, 65, 70, 75, 80, 85 or 90% of the subjects treated with the therapy. In further embodiments, such response rates can be obtained by any therapeutic regimen disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM or a related compound and one or more platinum compounds according to a dosing schedule that includes administering the TCN, TCN-P, TCN-PM and/or related compounds and/or the one or more platinum compounds one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds are not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of

the tumor is evident. In a particular embodiment, less than 12, less than 11 or less than 10 mg/m² of TCN, TCN-P, TCN-PM or a related compound can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 mg/m² of TCN, TCN-P, TCN-PM or a related compound can be administered according to a 42 day cycle. In another particular embodiment, about 1 to about 50 mg of a platinum compound is administered. In a particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90 or 100 mg/kg of a platinum compound can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of one or more platinum compounds one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds as disclosed herein can be administered one time per week. In another particular embodiment, 1, 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, or 100 mg/kg of a platinum compound can be administered one time per week.

In embodiments of the present invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds can be administered according to any combination disclosed herein, for example, the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds are not administered.

The TCN, TCN-P, TCM-PM and related compounds and one or more platinum compounds as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at

least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P, TCN-PM or a related compound disclosed herein can be administered to a subject. In certain embodiments, 10, 20, 50, 100, 150, 200, 250, 300, 350, or 400 mg of a platinum compound can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 100 mg of a platinum compound either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m² of TCN or related compounds can be administered once a week concurrently with less than about 100 mg of a platinum compound. In another embodiment, less than 20 mg/m² of TCN or related compounds can be administered once a week and less than about 100 mg of a platinum compound can be administered the following week.

In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN or a related compound and less than about 30, 25, 20, 50, or 100 mg of a platinum compound can be administered to a subject. In another embodiment, less than 10 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds and less than about 100 mg of a platinum compound can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

In other embodiments, methods are provided to treat a subject that has been diagnosed with cancer by administering to the subject an effective amount of TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs according to a dosing schedule that includes administering the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib or salt or derivatives thereof one time per week for three weeks followed by a one week period wherein the drug is not administered (i.e., via a 28 day cycle). In other embodiments, such 28 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident.

In further embodiments, a 42 day cycle is provided in which the compounds disclosed herein can be administered once a week for four weeks followed by a two week period in which the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof is not administered. In other embodiments, such 42 day cycles can be repeated at least 2, 3, 4, or 5 times or until regression of the tumor is evident. In a particular embodiment, less than 50, less than 25 or less than 10 mg/m² of TCN, TCN-P, TCN-PM or a related compound can be administered according to a 42 day cycle. In other particular embodiments, 2, 3, 4, 5, 6, 7, 8, 9, 10 or 11 mg/m² of TCN, TCN-P, TCN-PM or a related compound can be administered according to a 42 day cycle. In another particular embodiment, about 0.1 mg/m² to about 50 mg/m² of bortezomib or a derivative thereof is administered. In a particular embodiment, 0.1, 0.5, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg/m² of bortezomib or a salt thereof can be administered according to a 42 day cycle.

In another embodiment, methods are provided to treat cancer in a subject by administering to the subject a dosing regimen of 10 mg/m² or less of TCN, TCN-P, TCN-PM or a related compound and less than about 30 mg of bortezomib and derivatives thereof analogs one time per week. In particular embodiments, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 mg/m² of TCN, TCN-P, TCN-PM or a related compound as disclosed herein can be administered one time per week. In another particular embodiment, 0.1, 0.5, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg/m² of bortezomib or a derivative thereof can be administered one time per week.

In embodiments of the present invention, the compounds disclosed herein can be administered simultaneously as a single bolus dose over a short period of time, for example, about 5, 10, 15, 20, 30 or 60 minutes. In further embodiments, dosing schedules are provided in which the compounds are administered simultaneously via continuous infusion for at least 24, 48, 72, 96, or 120 hours. In certain embodiments, the administration of the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs via continuous or bolus injections can be repeated at a certain frequency at least: once a week, once every two weeks, once every three weeks, once a month, once every five weeks, once every six weeks, once every eight weeks, once every ten weeks and/or once every twelve weeks. The type and frequency of administrations can be combined in any manner disclosed herein to create a dosing cycle. The TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs can be repeatedly administered via a certain dosing cycles, for example as a bolus injection once every two weeks for three months. The dosing cycles can be administered for at least: one, two three, four five, six, seven, eight, nine, ten, eleven, twelve, eighteen or twenty four months. Alternatively, at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15 or 20 dosing cycles can be administered to a patient. The TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs can be administered according to any combination disclosed herein, for example, the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs can be administered once a week every three weeks for 3 cycles.

In further embodiments, the compounds can be administered separately at least once a day for at least 2, 3, 4, 5, 6, 7, 8, 9 or 10 days. Such administration can be followed by corresponding periods in which the TCN, TCN-P, TCN-PM and/or related compounds and/or the bortezomib and derivatives thereof analogs are not administered.

The TCN, TCN-P, TCN-PM and related compounds and bortezomib and derivatives thereof analogs as disclosed herein can be administered to patients in an amount that is effective in causing tumor regression. The administration of TCN, TCN-P, TCN-PM or related compounds and bortezomib and derivatives thereof analogs can provide at least a partial, such as at least 15, 20 or 30%, or complete response in vivo in at least 15-20% of the subjects. In certain embodiments, at least 2, 5, 10, 15, 20, 30 or 50 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein can be administered to a subject. In certain embodiments, at least about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 12, 15, 17, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 150, 165, 175, 200, 250, 300, or 350 mg/m² of TCN, TCN-P, TCN-PM or a related compound disclosed herein can be administered to

a subject. In certain embodiments, 1, 5, 10, 15, 20, 25, 30, 35, or 40 mg/m² of bortezomib can be administered to a subject.

The administration of the compound can be conducted according to any of the therapeutic regimens disclosed herein. In particular embodiments, the dosing regimen includes administering less than about 20 mg/m² of TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg of bortezomib either concurrently, sequentially, or conducted over a period of time. In one embodiment, less than 20 mg/m² of TCN or related compounds can be administered once a week concurrently with less than about 10 mg/m² of bortezomib. In another embodiment, less than 20 mg/m² of TCN or related compounds can be administered once a week and less than about 30 mg of bortezomib can be administered the following week.

In further embodiments, 2 mg/m², 5 mg/m², 10 mg/m², and/or 15 mg/m² of TCN or a related compound and less than about 30, 25, 20, 15, 10, 5, 1, 0.5, or 0.1 mg/m² of bortezomib or a salt or a derivative thereof can be administered to a subject. In another embodiment, less than 10 mg/m² of a TCN, TCN-P, TCN-PM and/or related compounds and less than about 30 mg/m² of bortezomib can be administered to a subject via continuous infusion for at least five days. The present invention provides for any combination of dosing type, frequency, number of cycles and dosage amount disclosed herein.

5.5. SCREENING OF PATIENT POPULATIONS

In another embodiment of the invention, methods are provided to identify cancers or tumors that are susceptible to the toxic effects of tricitriline (TCN) and related compounds. In one embodiment, methods are provided to treat a cancer or tumor in a mammal by (i) obtaining a biological sample from the tumor; (ii) determining whether the cancer or tumor overexpresses Akt kinase or hyperactivated and phosphorylated Akt kinase, and (iii) treating the cancer or tumor with tricitriline or a related compound as described herein. In one embodiment, the biological sample can be a biopsy. In other embodiments, the biological sample can be fluid, cells and/or aspirates obtained from the tumor or cancer.

The biological sample can be obtained according to any technique known to one skilled in the art. In one embodiment, a biopsy can be conducted to obtain the biological sample. A biopsy is a procedure performed to remove tissue or cells from the body for examination. Some biopsies can be performed in a physician's office, while others need to be done in a hospital setting. In addition, some biopsies require use of an anesthetic to numb the area, while others do not require any sedation. In certain embodiments, an endoscopic biopsy can be performed. This type of biopsy is performed through a fiberoptic endoscope (a long, thin tube that has a close-focusing telescope on the end for viewing) through a natural body orifice (i.e., rectum) or a small incision (i.e., arthroscopy). The endoscope is used to view the organ in question for abnormal or suspicious areas, in order to obtain a small amount of tissue for study. Endoscopic procedures are named for the organ or body area to be visualized and/or treated. The physician can insert the endoscope into the gastrointestinal tract (alimentary tract endoscopy), bladder (cystoscopy), abdominal cavity (laparoscopy), joint cavity (arthroscopy), mid-portion of the chest (mediastinoscopy), or trachea and bronchial system (laryngoscopy and bronchoscopy).

In another embodiment, a bone marrow biopsy can be performed. This type of biopsy can be performed either from

the sternum (breastbone) or the iliac crest hipbone (the bone area on either side of the pelvis on the lower back area). The skin is cleansed and a local anesthetic is given to numb the area. A long, rigid needle is inserted into the marrow, and cells are aspirated for study; this step is occasionally uncomfortable. A core biopsy (removing a small bone 'chip' from the marrow) may follow the aspiration.

In a further embodiment, an excisional or incisional biopsy can be performed on the mammal. This type of biopsy is often used when a wider or deeper portion of the skin is needed. Using a scalpel (surgical knife), a full thickness of skin is removed for further examination, and the wound is sutured (sewed shut with surgical thread). When the entire tumor is removed, it is referred to as an excisional biopsy technique. If only a portion of the tumor is removed, it is referred to as an incisional biopsy technique. Excisional biopsy is often the method usually preferred, for example, when melanoma (a type of skin cancer) is suspected.

In still further embodiments, a fine needle aspiration (FNA) biopsy can be used. This type of biopsy involves using a thin needle to remove very small pieces from a tumor. Local anesthetic is sometimes used to numb the area, but the test rarely causes much discomfort and leaves no scar. FNA is not, for example, used for diagnosis of a suspicious mole, but may be used, for example, to biopsy large lymph nodes near a melanoma to see if the melanoma has metastasized (spread). A computed tomography scan (CT or CAT scan) can be used to guide a needle into a tumor in an internal organ such as the lung or liver.

In other embodiments, punch shave and/or skin biopsies can be conducted. Punch biopsies involve taking a deeper sample of skin with a biopsy instrument that removes a short cylinder, or "apple core," of tissue. After a local anesthetic is administered, the instrument is rotated on the surface of the skin until it cuts through all the layers, including the dermis, epidermis, and the most superficial parts of the subcutis (fat). A shave biopsy involves removing the top layers of skin by shaving it off. Shave biopsies are also performed with a local anesthetic. Skin biopsies involve removing a sample of skin for examination under the microscope to determine if, for example, melanoma is present. The biopsy is performed under local anesthesia.

In particular embodiment, methods are provided to determine whether the tumor overexpresses an Akt kinase. Akt kinase overexpression can refer to the phosphorylation state of the kinase. Hyperphosphorylation of Akt can be detected according to the methods described herein. In one embodiment, a tumor biopsy can be compared to a control tissue. The control tissue can be a normal tissue from the mammal in which the biopsy was obtained or a normal tissue from a healthy mammal Akt kinase overexpression or hyperphosphorylation can be determined if the tumor biopsy contains greater amounts of Akt kinase and/or Akt kinase phosphorylation than the control tissue, such as, for example, at least approximately 1.5, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 5.5, 6, 7, 8, 9, or 10-fold greater amounts of Akt kinase than contained in the control tissue.

In one embodiment, the present invention provides a method to detect aberrant Akt kinase expression in a subject or in a biological sample from the subject by contacting cells, cell extracts, serum or other sample from the subjects or said biological sample with an immunointeractive molecule specific for an Akt kinase or antigenic portion thereof and screening for the level of immunointeractive molecule-Akt kinase complex formation, wherein an elevated presence of the complex relative to a normal cell is indicative of an aberrant cell that expresses or overexpresses Akt. In one

example, cells or cell extracts can be screened immunologically for the presence of elevated levels of Akt kinase.

In an alternative embodiment, the aberrant expression of Akt in a cell is detected at the genetic level by screening for the level of expression of a gene encoding an Akt kinase wherein an elevated level of a transcriptional expression product (i.e., mRNA) compared to a normal cell is indicative of an aberrant cell. In certain embodiments, real-time PCR as well as other PCR procedures can be used to determine transcriptional activity. In one embodiment, mRNA can be obtained from cells of a subject or from a biological sample from a subject and cDNA optionally generated. The mRNA or cDNA can then be contacted with a genetic probe capable of hybridizing to and/or amplifying all or part of a nucleotide sequence encoding Akt kinase or its complementary nucleotide sequence and then the level of the mRNA or cDNA can be detected wherein the presence of elevated levels of the mRNA or cDNA compared to normal controls can be assessed.

Yet another embodiment of the present invention contemplates the use of an antibody, monoclonal or polyclonal, to Akt kinase in a quantitative or semi-quantitative diagnostic kit to determine relative levels of Akt kinase in suspected cancer cells from a patient, which can include all the reagents necessary to perform the assay. In one embodiment, a kit utilizing reagents and materials necessary to perform an ELISA assay is provided. Reagents can include, for example, washing buffer, antibody dilution buffer, blocking buffer, cell staining solution, developing solution, stop solution, anti-phospho-protein specific antibodies, anti-Pan protein specific antibodies, secondary antibodies, and distilled water. The kit can also include instructions for use and can optionally be automated or semi-automated or in a form which is compatible with automated machine or software. In one embodiment, a phosphor-ser-473 Akt antibody that detects the activated form of AKT (Akt phosphorylated at serine 474) can be utilized as the antibody in a diagnostic kit. See, for example, Yuan et al. (2000) "Frequent Activation of AKT2 and induction of apoptosis by inhibition of phosphoinositide-3-OH kinase/Akt pathway in human ovarian cancer," *Oncogene* 19:2324-2330.

5.6. Akt KINASES

Akt, also named PKB3, represents a subfamily of the serine/threonine kinase. Three members, AKT1, AKT2, and AKT3, have been identified in this subfamily. Akt is activated by extracellular stimuli in a PI3K-dependent manner (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). Full activation of Akt requires phosphorylation of Thr308 in the activation loop and Ser473 in the C-terminal activation domain. Akt is negatively regulated by PTEN tumor suppressor. Mutations in PTEN have been identified in various tumors, which lead to activation of Akt pathway (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). In addition, amplification, overexpression and/or activation of Akt have been detected in a number of human malignancies (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999, Cheng, J. Q., and Nicosia, S. V. AKT signal transduction pathway in oncogenesis. In Schwab D, editor. *Encyclopedic Reference of Cancer*. Berlin Heidelberg and New York: Springer; 2001. pp 35-7). Ectopic expression of Akt, especially constitutively active Akt, induces cell survival and malignant transformation whereas inhibition of Akt activity stimulates apoptosis in a range of mammalian cells (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999, Cheng, J. Q., and Nicosia, S. V. AKT signal transduction pathway in oncogenesis. In

Schwab D, editor. *Encyclopedic Reference of Cancer*. Berlin Heidelberg and New York: Springer; 2001. pp 35-7, Sun, M., et al. *Am. J. Pathol.*, 159: 431-437, 2001, Cheng, J. Q., et al. *Oncogene*, 14: 2793-2801, 1997). Further, activation of Akt has been shown to associate with tumor invasiveness and chemoresistance (West, K. A., et al. *Drug Resist. Updat.*, 5: 234-248, 2002).

Activation of the Akt pathway plays a pivotal role in malignant transformation and chemoresistance by inducing cell survival, growth, migration, and angiogenesis. The present invention provides methods to determine levels of Akt kinase overexpression and/or hyperactivated and phosphorylated Akt kinase.

The Akt kinase can be any known Akt family kinase, or kinase related thereto, including, but not limited to Akt 1, Akt 2, Akt 3. The mRNA and amino acid sequences of human Akt1, Akt2, and Akt 3 are illustrated in FIGS. 6a-c, 7a-d, and 8a-c, respectively.

In another embodiment, the compositions of the invention including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes kills cancer or tumor cells which express Akt.

5.7. DIAGNOSTIC ASSAYS

Immunological Assays

In one embodiment, a method is provided for detecting the aberrant expression of an Akt kinase in a cell in a mammal or in a biological sample from the mammal, by contacting cells, cell extracts or serum or other sample from the mammal or biological sample with an immunointeractive molecule specific for an Akt kinase or antigenic portion thereof and screening for the level of immunointeractive molecule-Akt kinase complex formations and determining whether an elevated presence of the complex relative to a normal cell is present.

The immunointeractive molecule can be a molecule having specificity and binding affinity for an Akt kinase or its antigenic parts or its homologs or derivatives thereof. In one embodiment, the immunointeractive molecule can be an immunoglobulin molecule. In other embodiments, the immunointeractive molecules can be an antibody fragments, single chain antibodies, and/or deimmunized molecules including humanized antibodies and T-cell associated antigen-binding molecules (TABMs). In one particular embodiment, the antibody can be a monoclonal antibody. In another particular embodiment, the antibody can be a polyclonal antibody. The immunointeractive molecule can exhibit specificity for an Akt kinase or more particularly an antigenic determinant or epitope on an Akt kinase. An antigenic determinant or epitope on an Akt kinase includes that part of the molecule to which an immune response is directed. The antigenic determinant or epitope can be a B-cell epitope or where appropriate a T-cell epitope. In one embodiment, the antibody is a phosphor-ser 473 Akt antibody.

One embodiment of the present invention provides a method for diagnosing the presence of cancer or cancer-like growth in a mammal, in which aberrant Akt activity is present, by contacting cells or cell extracts from the mammal or a biological sample from the subject with an Akt kinase-binding effective amount of an antibody having specificity for the Akt kinase or an antigenic determinant or epitope thereon and then quantitatively or qualitatively determining the level of an Akt kinase-antibody complex wherein the presence of elevated levels of said complex compared to a normal cell is determined.

Antibodies can be prepared by any of a number of means known to one skilled in the art. For example, for the detection of human Akt kinase, antibodies can be generally but not necessarily derived from non-human animals such as primates, livestock animals (e.g. sheep, cows, pigs, goats, horses), laboratory test animals (e.g. mice, rats, guinea pigs, rabbits) and/or companion animals (e.g. dogs, cats). Antibodies may also be recombinantly produced in prokaryotic or eukaryotic host cells. Generally, antibody based assays can be conducted in vitro on cell or tissue biopsies. However, if an antibody is suitably deimmunized or, in the case of human use, humanized, then the antibody can be labeled with, for example, a nuclear tag, administered to a patient and the site of nuclear label accumulation determined by radiological techniques. The Akt kinase antibody can be a cancer targeting agent. Accordingly, another embodiment of the present invention provides deimmunized forms of the antibodies for use in cancer imaging in human and non-human patients.

In general, for the generation of antibodies to an Akt kinase, the enzyme is required to be extracted from a biological sample whether this be from animal including human tissue or from cell culture if produced by recombinant means. The Akt kinase can be separated from the biological sample by any suitable means. For example, the separation may take advantage of any one or more of the Akt kinase's surface charge properties, size, density, biological activity and its affinity for another entity (e.g. another protein or chemical compound to which it binds or otherwise associates). Thus, for example, separation of the Akt kinase from the biological fluid can be achieved by any one or more of ultra-centrifugation, ion-exchange chromatography (e.g. anion exchange chromatography, cation exchange chromatography), electrophoresis (e.g. polyacrylamide gel electrophoresis, isoelectric focussing), size separation (e.g., gel filtration, ultra-filtration) and affinity-mediated separation (e.g. immunoaffinity separation including, but not limited to, magnetic bead separation such as Dynabead (trademark) separation, immunochromatography, immuno-precipitation). The separation of Akt kinase from the biological fluid can preserve conformational epitopes present on the kinase and, thus, suitably avoids techniques that cause denaturation of the enzyme. In a further embodiment, the kinase can be separated from the biological fluid using any one or more of affinity separation, gel filtration and/or ultra-filtration.

Immunization and subsequent production of monoclonal antibodies can be carried out using standard protocols known in the art, such as, for example, described by Kohler and Milstein (Kohler and Milstein, *Nature* 256: 495-499, 1975; Kohler and Milstein, *Eur. J. Immunol.* 6(7): 511-519, 1976), Coligan et al. ("*Current Protocols in Immunology*, John Wiley & Sons, Inc., 1991-1997) or Toyama et al. ("*Monoclonal Antibody, Experiment Manual*", published by Kodansha Scientific, 1987). Essentially, an animal is immunized with an Akt kinase-containing biological fluid or fraction thereof or a recombinant form of Akt kinase by standard methods to produce antibody-producing cells, particularly antibody-producing somatic cells (e.g. B lymphocytes). These cells can then be removed from the immunized animal for immortalization. In certain embodiment, a fragment of an Akt kinase can be used to generate antibodies. The fragment can be associated with a carrier. The carrier can be any substance of typically high molecular weight to which a non- or poorly immunogenic substance (e.g. a hapten) is naturally or artificially linked to enhance its immunogenicity.

Immortalization of antibody-producing cells can be carried out using methods which are well-known in the art. For example, the immortalization may be achieved by the transformation method using Epstein-Barr virus (EBV) (Kozbor et al., *Methods in Enzymology* 121: 140, 1986). In another embodiment, antibody-producing cells are immortalized using the cell fusion method (described in Coligan et al., 1991-1997, *supra*), which is widely employed for the production of monoclonal antibodies. In this method, somatic antibody-producing cells with the potential to produce antibodies, particularly B cells, are fused with a myeloma cell line. These somatic cells may be derived from the lymph nodes, spleens and peripheral blood of primed animals, preferably rodent animals such as mice and rats. In a particular embodiment, mice spleen cells can be used. In other embodiments, rat, rabbit, sheep or goat cells can also be used. Specialized myeloma cell lines have been developed from lymphocytic tumours for use in hybridoma-producing fusion procedures (Kohler and Milstein, 1976, *supra*; Shulman et al., *Nature* 276: 269-270, 1978; Volk et al., *J. Virol.* 42(1): 220-227, 1982). Many myeloma cell lines can also be used for the production of fused cell hybrids, including, e.g. P3.times.63-Ag8, P3.times.63-AG8.653, P3/NS1-Ag-4-1 (NS-1), Sp2/0-Ag14 and S194/5.XXO.Bu.1. The P3.times.63-Ag8 and NS-1 cell lines have been described by Kohler and Milstein (1976, *supra*). Shulman et al. (1978, *supra*) developed the Sp2/0-Ag14 myeloma line. The S194/5.XXO.Bu.1 line was reported by Trowbridge (*J. Exp. Med.* 148(1): 3 β -323, 1978). Methods for generating hybrids of antibody-producing spleen or lymph node cells and myeloma cells usually involve mixing somatic cells with myeloma cells in a 10:1 proportion (although the proportion may vary from about 20:1 to about 1:1), respectively, in the presence of an agent or agents (chemical, viral or electrical) that promotes the fusion of cell membranes. Fusion methods have been described (Kohler and Milstein, 1975, *supra*; Kohler and Milstein, 1976, *supra*; Gefter et al., *Somatic Cell Genet.* 3: 231-236, 1977; Volk et al., 1982, *supra*). The fusion-promoting agents used by those investigators were Sendai virus and polyethylene glycol (PEG). In certain embodiments, means to select the fused cell hybrids from the remaining unfused cells, particularly the unfused myeloma cells, are provided. Generally, the selection of fused cell hybrids can be accomplished by culturing the cells in media that support the growth of hybridomas but prevent the growth of the unfused myeloma cells, which normally would go on dividing indefinitely. The somatic cells used in the fusion do not maintain long-term viability in *in vitro* culture and hence do not pose a problem. Several weeks are required to selectively culture the fused cell hybrids. Early in this time period, it is necessary to identify those hybrids which produce the desired antibody, so that they may subsequently be cloned and propagated. Generally, around 10% of the hybrids obtained produce the desired antibody, although a range of from about 1 to about 30% is not uncommon. The detection of antibody-producing hybrids can be achieved by any one of several standard assay methods, including enzyme-linked immunoassay and radioimmunoassay techniques as, for example, described in Kennet et al. (*Monoclonal Antibodies and Hybridomas: A New Dimension in Biological Analyses*, pp 376-384, Plenum Press, New York, 1980) and by FACS analysis (O'Reilly et al., *Biotechniques* 25: 824-830, 1998).

Once the desired fused cell hybrids have been selected and cloned into individual antibody-producing cell lines, each cell line may be propagated in either of two standard ways. A suspension of the hybridoma cells can be injected

into a histocompatible animal. The injected animal will then develop tumours that secrete the specific monoclonal antibody produced by the fused cell hybrid. The body fluids of the animal, such as serum or ascites fluid, can be tapped to provide monoclonal antibodies in high concentration. Alternatively, the individual cell lines may be propagated *in vitro* in laboratory culture vessels. The culture medium containing high concentrations of a single specific monoclonal antibody can be harvested by decantation, filtration or centrifugation, and subsequently purified.

The cell lines can then be tested for their specificity to detect the Akt kinase of interest by any suitable immuno-detection means. For example, cell lines can be aliquoted into a number of wells and incubated and the supernatant from each well is analyzed by enzyme-linked immunosorbent assay (ELISA), indirect fluorescent antibody technique, or the like. The cell line(s) producing a monoclonal antibody capable of recognizing the target LIM kinase but which does not recognize non-target epitopes are identified and then directly cultured *in vitro* or injected into a histocompatible animal to form tumours and to produce, collect and purify the required antibodies.

The invention encompasses, therefore, a method of detecting in a sample an Akt kinase or fragment, variant or derivative thereof including contacting the sample with an antibody or fragment or derivative thereof and detecting the level of a complex containing the antibody and Akt kinase or fragment, variant or derivative thereof compared to normal controls wherein elevated levels of Akt kinase is determined. Any suitable technique for determining formation of the complex may be used. For example, an antibody according to the invention, having a reporter molecule associated therewith, may be utilized in immunoassays. Such immunoassays include but are not limited to radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs) immunochromatographic techniques (ICTs), and Western blotting which are well known to those of skill in the art. Immunoassays can also include competitive assays. The present invention encompasses qualitative and quantitative immunoassays.

Suitable immunoassay techniques are described, for example, in U.S. Pat. Nos. 4,016,043; 4,424,279; and 4,018, 653. These include both single-site and two-site assays of the non-competitive types, as well as the traditional competitive binding assays. These assays also include direct binding of a labeled antigen-binding molecule to a target antigen.

The invention further provides methods for quantifying Akt protein expression and activation levels in cells or tissue samples obtained from an animal, such as a human cancer patient or an individual suspected of having cancer. In one embodiment, the invention provides methods for quantifying Akt protein expression or activation levels using an imaging system quantitatively. The imaging system can be used to receive, enhance, and process images of cells or tissue samples, that have been stained with AKT protein-specific stains, in order to determine the amount or activation level of AKT protein expressed in the cells or tissue samples from such an animal. In embodiments of the methods of the invention, a calibration curve of AKT1 and AKT2 protein expression can be generated for at least two cell lines expressing differing amounts of AKT protein. The calibration curve can then used to quantitatively determine the amount of AKT protein that is expressed in a cell or tissue sample. Analogous calibration curves can be made for activated AKT proteins using reagents specific for the acti-

vation features. It can also be used to determine changes in amounts and activation state of AKT before and after clinical cancer treatment.

In one particular embodiment of the methods of the invention, AKT protein expression in a cell or tissue sample can be quantified using an enzyme-linked immunoabsorbent assay (ELISA) to determine the amount of AKT protein in a sample. Such methods are described, for example, in U.S. Patent Publication No. 2002/0015974.

In other embodiments enzyme immunoassays can be used to detect the Akt kinase. In such assays, an enzyme is conjugated to the second antibody, generally by means of glutaraldehyde or periodate. The substrates to be used with the specific enzymes are generally chosen for the production of, upon hydrolysis by the corresponding enzyme, a detectable colour change. It is also possible to employ fluorogenic substrates, which yield a fluorescent product rather than the chromogenic substrates. The enzyme-labeled antibody can be added to the first antibody-antigen complex, allowed to bind, and then the excess reagent washed away. A solution containing the appropriate substrate can then be added to the complex of antibody-antigen-antibody. The substrate can react with the enzyme linked to the second antibody, giving a qualitative visual signal, which may be further quantitated, usually spectrophotometrically, to give an indication of the amount of antigen which was present in the sample. Alternatively, fluorescent compounds, such as fluorescein, rhodamine and the lanthanide, europium (EU), can be chemically coupled to antibodies without altering their binding capacity. When activated by illumination with light of a particular wavelength, the fluorochrome-labeled antibody adsorbs the light energy, inducing a state of excitability in the molecule, followed by emission of the light at a characteristic colour visually detectable with a light microscope. The fluorescent-labeled antibody is allowed to bind to the first antibody-antigen complex. After washing off the unbound reagent, the remaining tertiary complex is then exposed to light of an appropriate wavelength. The fluorescence observed indicates the presence of the antigen of interest. Immunofluorometric assays (IFMA) are well established in the art and are particularly useful for the present method. However, other reporter molecules, such as radioisotope, chemiluminescent or bioluminescent molecules can also be employed.

In a particular embodiment, antibodies to Akt kinase can also be used in ELISA-mediated detection of Akt kinase especially in serum or other circulatory fluid. This can be accomplished by immobilizing anti-Akt kinase antibodies to a solid support and contacting these with a biological extract such as serum, blood, lymph or other bodily fluid, cell extract or cell biopsy. Labeled anti-Akt kinase antibodies can then be used to detect immobilized Akt kinase. This assay can be varied in any number of ways and all variations are encompassed by the present invention and known to one skilled in the art. This approach can enable rapid detection and quantitation of Akt kinase levels using, for example, a serum-based assay.

In one embodiment, an Akt Elisa assay kit may be used in the present invention. For example, a Cellular Activation of Signaling ELISA kit for Akt S473 from SuperArray Bioscience can be utilized in the present invention. In one embodiment, the antibody can be an anti-pan antibody that recognizes Akt S473. Elisa assay kit containing an anti-Akt antibody and additional reagents, including, but not limited to, washing buffer, antibody dilution buffer, blocking buffer, cell staining solution, developing solution, stop solution, secondary antibodies, and distilled water.

Nucleotide Detection

In another embodiment, a method to detect Akt kinases is provided by detecting the level of expression in a cell of a polynucleotide encoding an Akt kinase. Expression of the polynucleotide can be determined using any suitable technique known to one skilled in the art. In one embodiment, a labeled polynucleotide encoding an Akt kinase can be utilized as a probe in a Northern blot of an RNA extract obtained from the cell. In other embodiments, a nucleic acid extract from an animal can be utilized in concert with oligonucleotide primers corresponding to sense and antisense sequences of a polynucleotide encoding the kinase, or flanking sequences thereof, in a nucleic acid amplification reaction such as RT PCR. A variety of automated solid-phase detection techniques are also available to one skilled in the art, for example, as described by Fodor et al. (*Science* 251: 767-777, 1991) and Kazal et al. (*Nature Medicine* 2: 753-759, 1996).

In other embodiments, methods are provided to detect akt kinase encoding RNA transcripts. The RNA can be isolated from a cellular sample suspected of containing Akt kinase RNA, e.g. total RNA isolated from human cancer tissue. RNA can be isolated by methods known in the art, e.g. using TRIZOL reagent (GIBCO-BRL/Life Technologies, Gaithersburg, Md.). Oligo-dT, or random-sequence oligonucleotides, as well as sequence-specific oligonucleotides can be employed as a primer in a reverse transcriptase reaction to prepare first-strand cDNAs from the isolated RNA. Resultant first-strand cDNAs can then amplified with sequence-specific oligonucleotides in PCR reactions to yield an amplified product.

Polymerase chain reaction or "PCR" refers to a procedure or technique in which amounts of a preselected fragment of nucleic acid, RNA and/or DNA, are amplified as described, for example, in U.S. Pat. No. 4,683,195. Generally, sequence information from the ends of the region of interest or beyond is employed to design oligonucleotide primers. These primers will be identical or similar in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific RNA sequences and cDNA transcribed from total cellular RNA. See generally Mullis et al. (*Quant. Biol.* 51: 263, 1987; Erlich, eds., *PCR Technology*, Stockton Press, NY, 1989). Thus, amplification of specific nucleic acid sequences by PCR relies upon oligonucleotides or "primers" having conserved nucleotide sequences wherein the conserved sequences are deduced from alignments of related gene or protein sequences, e.g. a sequence comparison of mammalian Akt kinase genes. For example, one primer is prepared which is predicted to anneal to the antisense strand and another primer prepared which is predicted to anneal to the sense strand of a cDNA molecule which encodes a Akt kinase. To detect the amplified product, the reaction mixture is typically subjected to agarose gel electrophoresis or other convenient separation technique and the relative presence of the Akt kinase specific amplified DNA detected. For example, Akt kinase amplified DNA may be detected using Southern hybridization with a specific oligonucleotide probe or comparing its electrophoretic mobility with DNA standards of known molecular weight. Isolation, purification and characterization of the amplified Akt kinase DNA can be accomplished by excising or eluting the fragment from the gel (for example, see references Lawn et al., *Nucleic Acids Res.* 2: 6103, 1981; Goeddel et al., *Nucleic Acids Res.* 8: 4057-1980), cloning the amplified product into a cloning site of a suitable vector, such as the pCRII vector (Invitrogen), sequencing the cloned insert and comparing the DNA sequence to the known sequence of

LIM kinase. The relative amounts of LIM kinase mRNA and cDNA can then be determined.

In one embodiment, real-time PCR can be used to determine transcriptional levels of Akt nucleotides. Determination of transcriptional activity also includes a measure of potential translational activity based on available mRNA transcripts. Real-time PCR as well as other PCR procedures use a number of chemistries for detection of PCR product including the binding of DNA binding fluorophores, the 5' endonuclease, adjacent liner and hairpin oligoprobes and the self-fluorescing amplicons. These chemistries and real-time PCR in general are discussed, for example, in Mackay et al., *Nucleic Acids Res* 30(6): 1292-1305, 2002; Walker, J. *Biochem. Mol. Toxicology*, 15(3): 121-127, 2001; Lewis et al., *J. Pathol.* 195: 66-71, 2001.

In an alternate embodiment, the aberrant expression of Akt can be identified by contacting a nucleotide sequences isolated from a biological sample with an oligonucleotide probe having a sequence complementary to an Akt sequences selected from the nucleotide sequences of FIG. 6a-c, 7a-d, or 8a-c, or fragment thereof, and then detecting the sequence by hybridizing the probe to the sequence, and comparing the results to a normal sample. The hybridization of the probe to the biological sample can be detected by labeling the probe using any detectable agent. The probe can be labeled for example, with a radioisotope, or with biotin, fluorescent dye, electron-dense reagent, enzyme, hapten or protein for which antibodies are available. The detectable label can be assayed by any desired means, including spectroscopic, photochemical, biochemical, immunochemical, radioisotopic, or chemical means. The probe can also be detected using techniques such as an oligomer restriction technique, a dot blot assay, a reverse dot blot assay, a line probe assay, and a 5' nuclease assay. Alternatively, the probe can be detected using any of the generally applicable DNA array technologies, including macroarray, microarray and DNA microchip technologies. The oligonucleotide probe typically includes approximately at least 14, 15, 16, 18, 20, 25 or 28 nucleotides that hybridize to the nucleotides selected from FIGS. 6a-c, 7a-d, and 8a-c, or a fragment thereof. It is generally not preferred to use a probe that is greater than approximately 25 or 28 nucleotides in length. The oligonucleotide probe is designed to identify an Akt nucleotide sequence.

Kinase Assays

The activity of the Akt kinases can be measured using any suitable kinase assay known in the art. For example, and not by way of limitation, the methods described in Hogg et al (*Oncogene* 1994 9:98-96), Mills et al (*J. Biol. Chem.* 1992 267:16000-006) and Tomizawa et al 2001 (*FEBS Lett.* 2001 492: 221-7), Schmandt et al, (*J. Immunol.* 1994, 152:96-105) can be used. Further serine, threonine and tyrosine kinase assays are described in Ausubel et al. (*Short Protocols in Molecular Biology*, 1999, unit 17.6).

Akt kinase assays can generally use an Akt polypeptide, a labeled donor substrate, and a receptor substrate that is either specific or non-specific for Akt. In such assays Akt transfers a labeled moiety from the donor substrate to the receptor substrate, and kinase activity is measured by the amount of labeled moiety transferred from the donor substrate to the receptor substrate. Akt polypeptide can be produced using various expression systems, can be purified from cells, can be in the form of a cleaved or uncleaved recombinant fusion protein and/or can have non-Akt polypeptide sequences, for example a His tag or .beta.-galactosidase at its N- or C-terminus. Akt activity can be assayed in cancerous cells lines if the cancerous cell lines are used as

a source of the Akt to be assayed. Suitable donor substrates for Akt assays include any molecule that is susceptible to dephosphorylation by Akt., such as, for example include gamma.-labeled ATP and ATP analogs, wherein the label is ³³P, ³²P, ³⁵S or any other radioactive isotope or a suitable fluorescent marker. Suitable recipient substrates for Akt assays include any polypeptide or other molecule that is susceptible to phosphorylation by Akt. Recipient substrates can be derived from fragments of in vivo targets of Akt. Recipient substrates fragments can be 8 to 50 amino acids in length, usually 10 to 30 amino acids and particularly of about 10, 12, 15, 18, 20 and 25 amino acids in length. Further recipient substrates can be determined empirically using a set of different polypeptides or other molecules. Targets of Recipient substrates for TTK can be capable of being purified from other components of the reaction once the reaction has been performed. This purification is usually done through a molecular interaction, where the recipient substrates is biotinylated and purified through its interaction with streptavidin, or a specific antibody is available that can specifically recognize the recipient substrates. The reaction can be performed in a variety of conditions, such as on a solid support, in a gel, in solution or in living cells. The choice of detection methods depends on type of label used for the donor molecule and may include, for example, measurement of incorporated radiation or fluorescence by autoradiography, scintillation, scanning or fluorography.

6. METHODS OF TREATMENT

The compounds and pharmaceutical compositions provided herein can be used in the treatment of a condition including tumors, cancer, and other disorders associated with abnormal cell proliferation. In one embodiment, the compounds of the present invention can be used to treat a carcinoma, sarcoma, lymphoma, leukemia, and/or myeloma. In other embodiments of the present invention, the compounds disclosed herein can be used to treat solid tumors.

The compounds of the present invention can be used for the treatment of cancer, such as, but not limited to cancer of the following organs or tissues: breast, prostate, lung, bronchus, colon, urinary, bladder, non-Hodgkin lymphoma, melanoma, kidney, renal, pancreas, pharynx, thyroid, stomach, brain, multiple myeloma, esophagus, liver, intrahepatic bile duct, cervix, larynx, acute myeloid leukemia, chronic lymphatic leukemia, soft tissue, such as heart, Hodgkin lymphoma, testis, small intestine, chronic myeloid leukemia, acute lymphatic leukemia, anus, anal canal, anorectal, thyroid, vulva, gallbladder, pleura, eye, nose nasal cavity, middle ear, nasopharynx, ureter, peritoneum, omentum, mesentery, and gastrointestinal, high grade glioma, glioblastoma, colon, rectal, pancreatic, gastric cancers, hepatocellular carcinoma; head and neck cancers, carcinomas; renal cell carcinoma; adenocarcinoma; sarcomas; hemangioendothelioma; lymphomas; leukemias, mycosis fungoides. In additional embodiments, the compounds of the invention can be used to treat skin diseases including, but not limited to, the malignant diseases angiosarcoma, hemangioendothelioma, basal cell carcinoma, squamous cell carcinoma, malignant melanoma and Kaposi's sarcoma, and the non-malignant diseases or conditions such as psoriasis, lymphangiogenesis, hemangioma of childhood, Sturge-Weber syndrome, verruca vulgaris, neurofibromatosis, tuberous sclerosis, pyogenic granulomas, recessive dystrophic epidermolysis bullosa, venous ulcers, acne, rosacea, eczema, molluscum contagious, seborrheic keratosis, and actinic keratosis.

Compositions including the compounds of the invention can be used to treat these cancers and other cancers at any stage from the discovery of the cancer to advanced stages. In addition, compositions including compounds of the invention can be used in the treatment of the primary cancer and metastases thereof.

In other embodiments of the invention, the compounds described herein can be used for the treatment of cancer, including, but not limited to, the cancers listed in Table 1 below.

TABLE 1

Types of Cancer	
Acute Lymphoblastic Leukemia, Adult	Hairy Cell Leukemia
Acute Lymphoblastic Leukemia, Childhood	Head and Neck Cancer
Acute Myeloid Leukemia, Adult	Hepatocellular (Liver) Cancer, Adult (Primary)
Acute Myeloid Leukemia, Childhood	Hepatocellular (Liver) Cancer, Childhood (Primary)
Adrenocortical Carcinoma	Hodgkin's Lymphoma, Adult
Adrenocortical Carcinoma, Childhood	Hodgkin's Lymphoma, Childhood
AIDS-Related Cancers	Hodgkin's Lymphoma During Pregnancy
AIDS-Related Lymphoma	Hypopharyngeal Cancer
Anal Cancer	Hypothalamic and Visual Pathway Glioma, Childhood
Astrocytoma, Childhood Cerebellar	Intraocular Melanoma
Astrocytoma, Childhood Cerebral	Islet Cell Carcinoma (Endocrine Pancreas)
Basal Cell Carcinoma	Kaposi's Sarcoma
Bile Duct Cancer, Extrahepatic	Kidney (Renal Cell) Cancer
Bladder Cancer	Kidney Cancer, Childhood
Bladder Cancer, Childhood	Laryngeal Cancer
Bone Cancer, Osteosarcoma/Malignant	Laryngeal Cancer, Childhood
Fibrous Histiocytoma	Leukemia, Acute Lymphoblastic, Adult
Brain Stem Glioma, Childhood	Leukemia, Acute Lymphoblastic, Childhood
Brain Tumor, Adult	Leukemia, Acute Myeloid, Adult
Brain Tumor, Brain Stem Glioma, Childhood	Leukemia, Acute Myeloid, Childhood
Brain Tumor, Cerebellar	Leukemia, Chronic Lymphocytic
Astrocytoma, Childhood	Leukemia, Chronic Myelogenous
Brain Tumor, Cerebral	Leukemia, B Cell
Astrocytoma/Malignant Glioma, Childhood	Lip and Oral Cavity Cancer
Brain Tumor, Ependymoma, Childhood	Liver Cancer, Adult (Primary)
Brain Tumor, Medulloblastoma, Childhood	Liver Cancer, Childhood (Primary)
Brain Tumor, Supratentorial	Lung Cancer, Non-Small Cell
Primitive	Lung Cancer, Small Cell
Neuroectodermal Tumors, Childhood	Lymphoma, AIDS-Related
Brain Tumor, Visual Pathway and Hypothalamic Glioma, Childhood	Lymphoma, Burkitt's
Brain Tumor, Childhood	Lymphoma, Cutaneous T-Cell, see Mycosis
Breast Cancer	Fungoides and Sézary Syndrome
Breast Cancer, Childhood	Lymphoma, Hodgkin's, Adult
Breast Cancer, Male	Lymphoma, Hodgkin's, Childhood
Bronchial Adenomas/Carcinoids, Childhood	Lymphoma, Hodgkin's During Pregnancy
Burkitt's Lymphoma	Lymphoma, Non-Hodgkin's, Adult
Carcinoid Tumor, Childhood	Lymphoma, Non-Hodgkin's, Childhood
Carcinoid Tumor, Gastrointestinal	Lymphoma, Non-Hodgkin's During Pregnancy
Carcinoma of Unknown Primary	Lymphoma, Primary Central Nervous System
Central Nervous System	Macroglobulinemia, Waldenström's
Lymphoma, Primary	Malignant Fibrous Histiocytoma of Bone/Osteosarcoma
Cerebellar Astrocytoma, Childhood	Medulloblastoma, Childhood
Cerebral Astrocytoma/Malignant Glioma, Childhood	Melanoma
Cervical Cancer	Melanoma, Intraocular (Eye)
Childhood Cancers	Merkel Cell Carcinoma
Chronic Lymphocytic Leukemia	Mesothelioma, Adult Malignant
Chronic Myelogenous Leukemia	Mesothelioma, Childhood
Chronic Myeloproliferative Disorders	Metastatic Squamous Neck Cancer with Occult Primary
Colon Cancer	Multiple Endocrine Neoplasia Syndrome, Childhood
Colorectal Cancer, Childhood	

TABLE 1-continued

Types of Cancer	
5 Cutaneous T-Cell Lymphoma, see Mycosis Fungoides and Sézary Syndrome	Multiple Myeloma/Plasma Cell Neoplasm
Endometrial Cancer	Mycosis Fungoides
Ependymoma, Childhood	Myelodysplastic Syndromes
Esophageal Cancer	Myelodysplastic/Myeloproliferative Diseases
Esophageal Cancer, Childhood	Myelogenous Leukemia, Chronic
Ewing's Family of Tumors	Myeloid Leukemia, Adult Acute
10 Extracranial Germ Cell Tumor, Childhood	Myeloid Leukemia, Childhood Acute
Extragenital Germ Cell Tumor	Myeloma, Multiple
Extrahepatic Bile Duct Cancer	Myeloproliferative Disorders, Chronic
Eye Cancer, Intraocular Melanoma	Nasal Cavity and Paranasal Sinus Cancer
Eye Cancer, Retinoblastoma	Nasopharyngeal Cancer
15 Gallbladder Cancer	Nasopharyngeal Cancer, Childhood
Gastric (Stomach) Cancer	Neuroblastoma
Gastric (Stomach) Cancer, Childhood	Non-Hodgkin's Lymphoma, Adult
Gastrointestinal Carcinoid Tumor	Non-Hodgkin's Lymphoma, Childhood
20 Germ Cell Tumor, Extracranial, Childhood	Non-Hodgkin's Lymphoma During Pregnancy
Germ Cell Tumor, Extragenital	Non-Small Cell Lung Cancer
Germ Cell Tumor, Ovarian	Oral Cancer, Childhood
Gestational Trophoblastic Tumor	Oral Cavity Cancer, Lip and Oropharyngeal Cancer
Glioma, Adult	Osteosarcoma/Malignant Fibrous
Glioma, Childhood Brain Stem	Histiocytoma of Bone
25 Glioma, Childhood Cerebral	Ovarian Cancer, Childhood
Astrocytoma	Ovarian Epithelial Cancer
Glioma, Childhood Visual Pathway and Hypothalamic	Ovarian Germ Cell Tumor
Skin Cancer (Melanoma)	Ovarian Low Malignant Potential Tumor
30 Skin Carcinoma, Merkel Cell	Pancreatic Cancer
Small Cell Lung Cancer	Pancreatic Cancer, Childhood
Small Intestine Cancer	Pancreatic Cancer, Islet Cell
Soft Tissue Sarcoma, Adult	Paranasal Sinus and Nasal Cavity Cancer
Soft Tissue Sarcoma, Childhood	Parathyroid Cancer
Squamous Cell Carcinoma, see Skin	Penile Cancer
35 Cancer (non-Melanoma)	Pheochromocytoma
Squamous Neck Cancer with Occult	Pineoblastoma and Supratentorial Primitive
Primary, Metastatic	Neuroectodermal Tumors, Childhood
Stomach (Gastric) Cancer	Pituitary Tumor
Stomach (Gastric) Cancer, Childhood	Plasma Cell Neoplasm/Multiple Myeloma
40 Supratentorial Primitive	Pleuropulmonary Blastoma
Neuroectodermal Tumors, Childhood	Pregnancy and Breast Cancer
T-Cell Lymphoma, Cutaneous, see Mycosis Fungoides and Sézary Syndrome	Pregnancy and Hodgkin's Lymphoma
45 Testicular Cancer	Pregnancy and Non-Hodgkin's Lymphoma
Thymoma, Childhood	Primary Central Nervous System Lymphoma
Thymoma and Thymic Carcinoma	Prostate Cancer
Thyroid Cancer	Rectal Cancer
Thyroid Cancer, Childhood	Renal Cell (Kidney) Cancer
50 Transitional Cell Cancer of the Renal Pelvis and Ureter	Renal Cell (Kidney) Cancer, Childhood
Trophoblastic Tumor, Gestational	Renal Pelvis and Ureter, Transitional Cell Cancer
Unknown Primary Site, Carcinoma of, Adult	Retinoblastoma
Unknown Primary Site, Cancer of, Childhood	Rhabdomyosarcoma, Childhood
55 Unusual Cancers of Childhood	Salivary Gland Cancer
Ureter and Renal Pelvis, Transitional Cell Cancer	Salivary Gland Cancer, Childhood
Urethral Cancer	Sarcoma, Ewing's Family of Tumors
Uterine Cancer, Endometrial	Sarcoma, Kaposi's
60 Uterine Sarcoma	Sarcoma, Soft Tissue, Adult
Vaginal Cancer	Sarcoma, Soft Tissue, Childhood
Visual Pathway and Hypothalamic Glioma, Childhood	Sarcoma, Uterine
Vulvar Cancer	Sézary Syndrome
Waldenström's Macroglobulinemia	Skin Cancer (non-Melanoma)
65 Wilms' Tumor	Skin Cancer, Childhood

In further embodiments of the present invention, the compounds disclosed herein can be used in the treatment of angiogenesis-related diseases.

Antiangiogenic small molecules include thalidomide, which acts in part by inhibiting NFkB, 2-methoxyestradiol, which influences microtubule activation and hypoxia inducing factor (HIF1a) activation, cyclo-oxygenase 2 (COX2) inhibitors, and low doses of conventional chemotherapeutic agents, including cyclophosphamide, taxanes, anthracycline analogs, and vinca alkaloids (vincristine, vinblastine) (D'Amato, R. J. et al. (1994) *Proc. Natl. Acad. Sci. U.S.A* 91, 3964-3968, D'Amato, R. J. et al. (1994) *Proc. Natl. Acad. Sci. U.S.A* 91, 4082-4085). In addition, certain tyrosine kinase inhibitors indirectly decrease angiogenesis by decreasing production of VEGF and other proangiogenic factors by tumor and stromal cells. These drugs include Herceptin, imatinib (Glivec), and Iressa (Bergers, G. et al. (2003) *Journal of Clinical Investigation* 111, 1287-1295, Ciardiello, F. et al. (2001) *Clinical Cancer Research* 7, 1459-1465, Plum, S. M. et al. (2003) *Clinical Cancer Research* 9, 4619-4626).

Recently, angiogenesis inhibitors have moved from animal models to human patients. Angiogenesis inhibitors represent a promising treatment for a variety of cancers. Recently, Avastin a high affinity antibody against vascular endothelial growth factor (VEGF), has been shown to prolong life as a single agent in advanced renal cell carcinoma and prolong life in combination with chemotherapy in advanced colon cancer (Yang, J. C. et al. (2003) *New England Journal of Medicine* 349, 427-434, Kabbinar, F. et al. (2003) *Journal of Clinical Oncology* 21, 60-65).

Angiogenesis-related diseases include, but are not limited to, inflammatory, autoimmune, and infectious diseases; angiogenesis-dependent cancer, including, for example, solid tumors, blood born tumors such as leukemias, and tumor metastases; benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; rheumatoid arthritis; psoriasis; eczema; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis; Osler-Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; and wound granulation. In addition, compositions of this invention can be used to treat diseases such as, but not limited to, intestinal adhesions, atherosclerosis, scleroderma, warts, and hypertrophic scars (i.e., keloids). Compositions of this invention can also be used in the treatment of diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochele minalia quintosa*), ulcers (*Helobacter pylori*), tuberculosis, and leprosy.

6.1. Treatment of Drug Resistant Tumors or Cancers

The invention provides compounds that can be used to treat drug resistant cancer, including the embodiments of cancers and the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more additional anti-cancer agents disclosed herein.

Multidrug resistance (MDR) occurs in human cancers and can be a significant obstacle to the success of chemotherapy. Multidrug resistance is a phenomenon whereby tumor cells in vitro that have been exposed to one cytotoxic agent develop cross-resistance to a range of structurally and functionally unrelated compounds. In addition, MDR can occur

intrinsically in some cancers without previous exposure to chemotherapy agents. Thus, in one embodiment, the present invention provides methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer, by administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents as disclosed herein. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents as disclosed herein can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

The invention encompasses compounds that can be used to treat drug resistant cancer, including the embodiments of cancers disclosed herein by administering TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes.

In one embodiment, the invention encompasses methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer by administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

The invention encompasses compounds that can be used to treat drug resistant cancer, including the embodiments of cancers disclosed herein by administering TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof.

In one embodiment, the invention encompasses methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer by administration of TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

The invention encompasses compounds that can be used to treat drug resistant cancer, including the embodiments of cancers disclosed herein by administering TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof.

In one embodiment, the invention encompasses methods for the treatment of a patient with a drug resistant cancer, for

example, multidrug resistant cancer by administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

The invention encompasses compounds that can be used to treat drug resistant cancer, including the embodiments of cancers disclosed herein by administering TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof.

In one embodiment, the invention encompasses methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer by administration of TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

The invention encompasses compounds that can be used to treat drug resistant cancer, including the embodiments of cancers disclosed herein by administering TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds.

In one embodiment, the invention encompasses methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer by administration of TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

The invention encompasses compounds that can be used to treat drug resistant cancer, including the embodiments of cancers disclosed herein by administering TCN, TCN-P, TCN-PM and/or related compounds and bortezomib or a salt thereof.

In one embodiment, the invention encompasses methods for the treatment of a patient with a drug resistant cancer, for example, multidrug resistant cancer by administration of TCN, TCN-P, TCN-PM and/or related compounds and bortezomib or a salt thereof. In certain embodiments, TCN, TCN-P, TCN-PM and/or related compounds and bortezomib or a salt thereof can be used to treat cancers that are resistant to taxol alone, rapamycin, tamoxifen, cisplatin, and/or gefitinib (iressa).

In one embodiment, TCN, TCN-P, TCN-PM and/or related compounds and bortezomib or a salt thereof can be used for the treatment of drug resistant cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein.

6.2. Combination Therapy

In certain embodiments, the TCN, TCN-P, TCN-PM and/or related compounds and one or more first additional anti-cancer agents of the invention can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more first additional anti-cancer agents and compositions thereof, when used in the treatment of solid tumors, can be administered in conjunction with the use of radiation.

In another embodiment of the present invention, the TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents and compositions disclosed herein can be combined with at least one second additional chemotherapeutic agent. The second additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more additional anti-cancer agents disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and taxanes and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopril, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angioinhibins, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimitotic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharomuricin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyproterone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cisplatin, carbonplatin, oxaliplatin, paraplatin) or nitroso-

61

reas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed below in Table 2.

TABLE 2

Chemotherapeutic Agents	
13-cis-Retinoic Acid	Neosar
2-Amino-6-Mercaptopurine	Neulasta
2-CdA	Neumega
2-Chlorodeoxyadenosine	Neupogen
5-fluorouracil	Nilandron
5-FU	Nilutamide
6-TG	Nitrogen Mustard
6-Thioguanine	Novaldex
6-Mercaptopurine	Novantrone
6-MP	Octreotide
Accutane	Octreotide acetate
Actinomycin-D	Oncospar
Adriamycin	Oncovin
Adrucil	Ontak
Agrylin	Onxal
Ala-Cort	Oprevelkin
Aldesleukin	Orapred
Alemtuzumab	Orasone
Alitretinoin	Oxaliplatin
Alkaban-AQ	Paclitaxel
Alkeran	Pamidronate
All-transretinoic acid	Panretin
Alpha interferon	Paraplatin
Altretamine	Pediapred
Amethopterin	PEG Interferon
Amifostine	Pegaspargase
Aminoglutethimide	Pegfilgrastim
Anagrelide	PEG-INTRON
Anandron	PEG-L-asparaginase
Anastrozole	Phenylalanine Mustard
Arabinosylcytosine	Platinol
Ara-C	Platinol-AQ
Aranesp	Prednisolone
Aredia	Prednisone
Arimidex	Prelone
Aromasin	Procarbazine
Arsenic trioxide	PROCRIT
Asparaginase	Proleukin
ATRA	Prolifeprospan 20 with
Avastin	Carmustine implant
BCG	Purinethol
BCNU	Raloxifene
Bevacizumab	Rheumatrex
Bexarotene	Rituxan
Bicalutamide	Rituximab
BiCNU	Roveron-A (interferon alfa-2a)
Blenoxane	Rubex
Bleomycin	Rubidomycin hydrochloride
Bortezomib	Sandostatatin
Busulfan	Sandostatatin LAR
Busulfex	Sargramostim
C225	Solu-Cortef
Calcium Leucovorin	Solu-Medrol
Campath	STI-571
Camptosar	Streptozocin
Camptothecin-11	Tamoxifen
Capecitabine	Targretin
Carac	Taxol
Carboplatin	Taxotere

62

TABLE 2-continued

Chemotherapeutic Agents		
	Carmustine	Temodar
5	Carmustine wafer	Temozolomide
	Casodex	Teniposide
	CCNU	TESPA
	CDDP	Thalidomide
	CeeNU	Thalomid
	Cerubidine	TheraCys
10	cetuximab	Thioguanine
	Chlorambucil	Thioguanine Tabloid
	Cisplatin	Thiophosphamide
	Citrovorum Factor	Thioplex
	Cladribine	Thiotepa
	Cortisone	TICE
15	Cosmegen	Toposar
	CPT-11	Topotecan
	Cyclophosphamide	Toremifene
	Cytadren	Trastuzumab
	Cytarabine	Tretinoin
	Cytarabine liposomal	Trexall
20	Cytosar-U	Trisenox
	Cytosan	TSPA
	Dacarbazine	VCR
	Dactinomycin	Velban
	Darbepoetin alfa	Velcade
	Daunomycin	VePesid
25	Daunorubicin	Vesanoid
	Daunorubicin hydrochloride	Viadur
	Daunorubicin liposomal	Vinblastine
	DaunoXome	Vinblastine Sulfate
	Decadron	Vincasar Pfs
	Delta-Cortef	Vincristine
	Deltasone	Vinorelbine
30	Denileukin difitox	Vinorelbine tartrate
	DepoCyt	VLB
	Dexamethasone	VP-16
	Dexamethasone acetate	Vumon
	dexamethasone sodium	Xeloda
	phosphate	Zanosar
35	Dexasone	Zevalin
	Dextrazoxane	Zinecard
	DHAD	Zoladex
	DIC	Zoledronic acid
	Diodex	Zometa
	Docetaxel	Gliadel wafer
40	Doxil	Glivec
	Doxorubicin	GM-CSF
	Doxorubicin liposomal	Goserelin
	Droxia	granulocyte - colony stimulating factor
	DTIC	Granulocyte macrophage colony
	DTIC-Dome	stimulating factor
	Duralone	Halotestin
45	Efidex	Herceptin
	Eligard	Hexadrol
	Ellence	Hexalen
	Eloxatin	Hexamethylmelamine
	Elspar	HMM
	Emcyt	Hycamtin
50	Epirubicin	Hydrea
	Epoetin alfa	Hydrocort Acetate
	Erbitux	Hydrocortisone
	Erwinia L-asparaginase	Hydrocortisone sodium phosphate
	Estramustine	Hydrocortisone sodium succinate
	Ethylol	Hydrocortone phosphate
55	Etopophos	Hydroxyurea
	Etoposide	Ibritumomab
	Etoposide phosphate	Ibritumomab Tiuxetan
	Eulexin	Idamycin
	Evista	Idarubicin
	Exemestane	Ifex
60	Fareston	IFN-alpha
	Faslodex	Ifosfamide
	Femara	IL-2
	Filgrastim	IL-11
	Floxuridine	Imatinib mesylate
	Fludara	Imidazole Carboxamide
	Fludarabine	Interferon alfa
65	Fluoroplex	Interferon Alfa-2b (PEG conjugate)
	Fluorouracil	Interleukin-2

TABLE 2-continued

Chemotherapeutic Agents	
Fluorouracil (cream)	Interleukin-11
Fluoxymesterone	Intron A (interferon alfa-2b)
Flutamide	Leucovorin
Folinic Acid	Leukeran
FUDR	Leukine
Fulvestrant	Leuprolide
G-CSF	Leurocristine
Gefitinib	Leustatin
Gemcitabine	Liposomal Ara-C
Gemtuzumab ozogamicin	Liquid Pred
Gemzar	Lomustine
Gleevec	L-PAM
Lupron	L-Sarcosin
Lupron Depot	Meticorten
Matulane	Mitomycin
Maxidex	Mitomycin-C
Mechlorethamine	Mitoxantrone
Mechlorethamine Hydrochlorine	M-Prednisol
Medralone	MTC
Medrol	MTX
Megace	Mustargen
Megestrol	Mustine
Megestrol Acetate	Mutamycin
Melphalan	Myleran
Mercaptopurine	Iressa
Mesna	Irinotecan
Mesnex	Isotretinoin
Methotrexate	Kidrolase
Methotrexate Sodium	Lanacort
Methylprednisolone	L-asparaginase
Mylocel	LCR
Letrozole	

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and a one or more taxanes and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

In another embodiment of the present invention, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopyr, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101,

CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimetabolic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharomuricin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyprotrone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplalin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as gramisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed in Table 2.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

In another embodiment of the present invention, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopyr,

anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimitotic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharmorubicin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyprotrone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplatin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed in Table 2.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline compounds of the invention can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline compounds and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

In another embodiment of the present invention, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline compounds and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline compounds disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline compounds and compositions, when

used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopryl, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimitotic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharmorubicin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyprotrone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplatin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed in Table 2.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

In another embodiment of the invention, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for

example, gefitinib, erlotinib or a salt thereof disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopryl, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlomethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimitotic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharmorubicin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuprorelin, goserelin, cyproterone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplalin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed in Table 2.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds of the invention can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

In another embodiment of the present invention, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed

herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopryl, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate (w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlomethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimitotic effects, such as those which target cytoskeletal elements, including podophylotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharmorubicin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuprorelin, goserelin, cyproterone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplalin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed below in Table 2.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs of the invention can be administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs and compositions thereof, when used in the treatment of solid tumors, can be administered the use of radiation.

In another embodiment of the present invention, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib

and derivatives thereof analogs and compositions disclosed herein can be combined with at least one additional chemotherapeutic agent. The additional agents can be administered in combination or alternation with the compounds disclosed herein. The drugs can form part of the same composition, or be provided as a separate composition for administration at the same time or a different time.

In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs disclosed herein can be combined with antiangiogenic agents to enhance their effectiveness, or combined with other antiangiogenic agents and administered together with other cytotoxic agents. In another embodiment, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs and compositions, when used in the treatment of solid tumors, can be administered with the agents selected from, but not limited to IL-12, retinoids, interferons, angiostatin, endostatin, thalidomide, thrombospondin-1, thrombospondin-2, captopril, anti-neoplastic agents such as alpha interferon, COMP (cyclophosphamide, vincristine, methotrexate and prednisone), etoposide, mBACOD (methotrexate, bleomycin, doxorubicin, cyclophosphamide, vincristine and dexamethasone), PRO-MACE/MOPP (prednisone, methotrexate w/leucovorin rescue), doxorubicin, cyclophosphamide, etoposide/mechlorethamine, vincristine, prednisone and procarbazine), vincristine, vinblastine, angiostatin, TNP-470, pentosan polysulfate, platelet factor 4, angiostatin, LM-609, SU-101, CM-101, Techgalan, thalidomide, SP-PG and radiation. In further embodiments, the compounds and compositions disclosed herein can be administered in combination or alternation with, for example, drugs with antimetabolic effects, such as those which target cytoskeletal elements, including podophyllotoxins or vinca alkaloids (vincristine, vinblastine); antimetabolite drugs (such as 5-fluorouracil, cytarabine, gemcitabine, purine analogues such as pentostatin, methotrexate); alkylating agents or nitrogen mustards (such as nitrosoureas, cyclophosphamide or ifosfamide); drugs which target DNA such as the anthracycline drugs adriamycin, doxorubicin, pharomycin or epirubicin; drugs which target topoisomerases such as etoposide; hormones and hormone agonists or antagonists such as estrogens, antiestrogens (tamoxifen and related compounds) and androgens, flutamide, leuporelin, goserelin, cyproterone or octreotide; drugs which target signal transduction in tumour cells including antibody derivatives such as herceptin; alkylating drugs such as platinum drugs (cis-platin, carbonplatin, oxaliplatin, paraplatin) or nitrosoureas; drugs potentially affecting metastasis of tumours such as matrix metalloproteinase inhibitors; gene therapy and antisense agents; antibody therapeutics; other bioactive compounds of marine origin, notably the didemnins such as aplidine; steroid analogues, in particular dexamethasone; anti-inflammatory drugs, including nonsteroidal agents (such as acetaminophen or ibuprofen) or steroids and their derivatives in particular dexamethasone; anti-emetic drugs, including 5HT-3 inhibitors (such as granisetron or ondasetron), and steroids and their derivatives in particular dexamethasone. In still further embodiments, the compounds and compositions can be used in combination or alternation with the chemotherapeutic agents disclosed in Table 2.

In certain embodiments, interferons (IFNs) can be used in combinations with the compounds of the invention. Suitable interferons include: interferon alpha-2a, interferon alpha-2b, pegylated interferon alpha, including interferon alpha-2a and interferon alpha 2b, interferon beta, interferon gamma, interferon tau, interferon omega, INFERGEN (interferon

alphacon-1) by InterMune, OMNIFERON (natural interferon) by Viragen, ALBUFERON by Human Genome Sciences, REBIF (interferon beta-1a) by Ares-Serono, Omega Interferon by BioMedicine, Oral Interferon Alpha by Amarello Biosciences, and interferon gamma, interferon tau, and/or interferon gamma-1b by InterMune.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextniguldipine, valsopodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In another embodiment TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextniguldipine, valsopodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In another embodiment TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline compounds as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextniguldipine, valsopodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In another embodiment TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer

known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextiguldipine, valspodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In another embodiment TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextiguldipine, valspodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In another embodiment TCN, TCN-P, TCN-PM and/or related compounds and bortezomib or a salt or derivative thereof as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer. Drug resistant cancers can include cancers of the colon, bone, kidney, adrenal, pancreas, liver and/or any other cancer known in the art or described herein. In one embodiment, the additional chemotherapeutic agent can be a P-glycoprotein inhibitor. In certain non-limiting embodiments, the P-glycoprotein inhibitor can be selected from the following drugs: verapamil, cyclosporin (such as cyclosporin A), tamoxifen, calmodulin antagonists, dexverapamil, dextiguldipine, valspodar (PSC 833), biricodar (VX-710), tariquidar (XR9576), zosuquidar (LY335979), laniquidar (R101933), and/or ONT-093.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and anthracycline compounds as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table

2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer.

In one embodiment TCN, TCN-P, TCN-PM and/or related compounds and bortezomib or a salt or derivative thereof as disclosed herein can be used in combination or alternation with additional chemotherapeutic agents, such as those described herein or in Table 2, for the treatment of drug resistant cancer, for example multiple drug resistant cancer.

7. PHARMACEUTICAL COMPOSITIONS

The compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The TCN, TCN-P, TCN-PM and/or related compounds and in combination with one or more taxanes may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with one or more taxanes.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

The compositions including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The TCN, TCN-P, TCN-PM and/or related compounds and in combination with a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be formulated as the sole pharmaceutically active ingredients in the composition or may be combined.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

The compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The TCN, TCN-P, TCN-PM and/or related compounds and in combination with one or more anthracycline analogs may be formulated as the sole

pharmaceutically active ingredient in the composition or may be combined with one or more anthracycline analogs.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

The compositions including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The TCN, TCN-P, TCN-PM and/or related compounds and in combination with an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be formulated as the sole pharmaceutically active ingredients in the composition or may be combined.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

The compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The TCN, TCN-P, TCN-PM and/or related compounds and in combination with one or more platinum compounds may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with one or more platinum compounds.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous, intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

The compositions including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs can optionally be administered with a pharmaceutical carrier or excipient. Pharmaceutical carriers suitable for administration of the compounds provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration. The TCN, TCN-P, TCN-PM and/or related compounds and in combination with bortezomib and derivatives thereof analogs may be formulated as the sole pharmaceutically active ingredient in the composition or may be combined with bortezomib and derivatives thereof analogs.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal, or parenteral (including subcutaneous, intramuscular, subcutaneous,

intravenous, intradermal, intraocular, intratracheal, intracisternal, intraperitoneal, and epidural) administration. Preferably the compositions are administered intravenously.

The compositions may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into association one or more compositions of the present invention and one or more pharmaceutical carriers or excipients.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes and compositions thereof can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel Introduction to Pharmaceutical Dosage Forms, Fourth Edition 1985, 126).

The TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, and compositions thereof can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel, Introduction to Pharmaceutical Dosage Forms, Fourth Ed., 1985, p. 126).

The TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs and compositions thereof can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel Introduction to Pharmaceutical Dosage Forms, Fourth Edition 1985, 126).

The TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof and compositions thereof can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel, Introduction to Pharmaceutical Dosage Forms, Fourth Ed., 1985, p. 126).

The TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds and compositions thereof can be formulated into suitable pharmaceutical

preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel Introduction to Pharmaceutical Dosage Forms, Fourth Ed., 1985, p. 126).

The TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs and compositions thereof can be formulated into suitable pharmaceutical preparations such as solutions, suspensions, tablets, dispersible tablets, pills, capsules, powders, sustained release formulations or elixirs, for oral administration or in sterile solutions or suspensions for parenteral administration, as well as transdermal patch preparation and dry powder inhalers. In one embodiment, the TCN, TCN-P, TCN-PM and/or related compounds described above are formulated into pharmaceutical compositions using techniques and procedures well known in the art (see, e.g., Ansel Introduction to Pharmaceutical Dosage Forms, Fourth Edition 1985, 126).

In the compositions, effective concentrations of one or more compounds or pharmaceutically acceptable derivatives thereof may be mixed with one or more suitable pharmaceutical carriers. The compounds of the invention may be derivatized as the corresponding salts, esters, enol ethers or esters, acetals, ketals, orthoesters, hemiacetals, hemiketals, acids, bases, solvates, hydrates or prodrugs prior to formulation. The concentrations of the compounds in the compositions are effective for delivery of an amount, upon administration, that treats, prevents, or ameliorates one or more of the symptoms of the target disease or disorder. In one embodiment, the compositions are formulated for single dosage administration. To formulate a composition, the weight fraction of compound is dissolved, suspended, dispersed or otherwise mixed in a selected carrier at an effective concentration such that the treated condition is relieved, prevented, or one or more symptoms are ameliorated.

Compositions suitable for oral administration may be presented as discrete units such as, but not limited to, tablets, caplets, pills or dragees capsules, or cachets, each containing a predetermined amount of one or more of the compositions; as a powder or granules; as a solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil emulsion or as a bolus, etc.

Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, or otherwise mixing a TCN, TCN-P, TCN-PM and/or related compounds and optional pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, glycols, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting agents, emulsifying agents, solubilizing agents, pH buffering agents, preservatives, flavoring agents, and the like, for example, acetate, sodium citrate, cyclodextrine derivatives, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, and other such agents. Methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 15th Edition, 1975.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or more TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or more TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or more TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or more TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having one or more TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

Compositions of the present invention suitable for topical administration in the mouth include for example, lozenges, having the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; pastilles, having TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs of the present invention in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes, having one or more of the compositions of the present invention administered in a suitable liquid carrier.

The tablets, pills, capsules, troches and the like can contain one or more of the following ingredients, or compounds of a similar nature: a binder; a lubricant; a diluent; a glidant; a disintegrating agent; a coloring agent; a sweetening agent; a flavoring agent; a wetting agent; an emetic coating; and a film coating. Examples of binders include microcrystalline cellulose, gum tragacanth, glucose solution, acacia mucilage, gelatin solution, molasses, poly-

nylpyrrolidine, povidone, croscopovidones, sucrose and starch paste. Lubricants include talc, starch, magnesium or calcium stearate, *lycopodium* and stearic acid. Diluents include, for example, lactose, sucrose, starch, kaolin, salt, mannitol and dicalcium phosphate. Glidants include, but are not limited to, colloidal silicon dioxide. Disintegrating agents include croscarmellose sodium, sodium starch glycolate, alginic acid, corn starch, potato starch, bentonite, methylcellulose, agar and carboxymethylcellulose. Coloring agents include, for example, any of the approved certified water soluble FD and C dyes, mixtures thereof; and water insoluble FD and C dyes suspended on alumina hydrate. Sweetening agents include sucrose, lactose, mannitol and artificial sweetening agents such as saccharin, and any number of spray dried flavors. Flavoring agents include natural flavors extracted from plants such as fruits and synthetic blends of compounds which produce a pleasant sensation, such as, but not limited to peppermint and methyl salicylate. Wetting agents include propylene glycol monostearate, sorbitan monooleate, diethylene glycol monolaurate and polyoxyethylene laural ether. Emetic-coatings include fatty acids, fats, waxes, shellac, ammoniated shellac and cellulose acetate phthalates. Film coatings include hydroxyethylcellulose, sodium carboxymethylcellulose, polyethylene glycol 4000 and cellulose acetate phthalate.

Compositions suitable for topical administration to the skin may be presented as ointments, creams, gels, and pastes, having one or more of the compositions administered in a pharmaceutical acceptable carrier.

Compositions for rectal administration may be presented as a suppository with a suitable base including, for example, cocoa butter or a salicylate.

Compositions suitable for nasal administration, when the carrier is a solid, include a coarse powder having a particle size, for example, in the range of 20 to 500 microns which is administered in the manner in which snuff is taken, (i.e., by rapid inhalation through the nasal passage from a container of the powder held close up to the nose). When the carrier is a liquid (for example, a nasal spray or as nasal drops), one or more of the compositions can be admixed in an aqueous or oily solution, and inhaled or sprayed into the nasal passage.

Compositions suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing one or more of the compositions and appropriate carriers.

Compositions suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The compositions may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules, and tablets of the kind previously described above.

Pharmaceutical organic or inorganic solid or liquid carrier media suitable for enteral or parenteral administration can be used to fabricate the compositions. Gelatin, lactose, starch, magnesium stearate, talc, vegetable and animal fats and oils, gum, polyalkylene glycol, water, or other known carriers may all be suitable as carrier media.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be used in combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be used in combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be used in combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be used in

combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be used in combination with one or more pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be used in combination with pharmaceutically acceptable carrier mediums and/or excipients. As used herein, "pharmaceutically acceptable carrier medium" includes any and all carriers, solvents, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants, adjuvants, vehicles, delivery systems, disintegrants, absorbents, preservatives, surfactants, colorants, flavorants, or sweeteners and the like, as suited to the particular dosage form desired.

Additionally, the compositions including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be combined with pharmaceutically acceptable excipients, and, optionally, sustained-release matrices, such as biodegradable polymers, to form therapeutic compositions. A "pharmaceutically acceptable excipient" includes a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

It will be understood, however, that the total daily usage of the compositions will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular host will depend upon a variety of factors, including for example, the disorder being treated and the severity of the

disorder; activity of the specific composition employed; the specific composition employed, the age, body weight, general health, sex and diet of the patient; the time of administration; route of administration; rate of excretion of the specific compound employed; the duration of the treatment; the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane used in combination or coincidental with the specific composition employed; the TCN, TCN-P, TCN-PM and/or related compounds and one or more molecules that modulate the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof used in combination or coincidental with the specific composition employed; the TCN, TCN-P, TCN-PM and/or related compounds and/or the anthracycline analogs used in combination or coincidental with the specific composition employed; an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof used in combination or coincidental with the specific composition employed; the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds used in combination or coincidental with the specific composition employed, the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more bortezomib and derivatives thereof used in combination or coincidental with the specific composition employed, and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the composition at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce

the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein refers to a physically discrete unit of the composition appropriate for the host to be treated. Each dosage should contain the quantity of composition calculated to produce the desired therapeutic affect either as such, or in association with the selected pharmaceutical carrier medium.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub-dose, or an appropriate fraction thereof, of the administered ingredient. For example, approximately 1-50 mg per day of a compound disclosed herein can reduce the volume of a solid tumor in mice.

The dosage will depend on host factors such as weight, age, surface area, metabolism, tissue distribution, absorption rate and excretion rate. In one embodiment, approximately 0.5 to 7 grams per day of a TCN, TCN-P, TCN-PM and/or related compounds disclosed herein may be administered to humans. Optionally, approximately 1 to 4 grams per day of the compound can be administered to humans. In certain embodiments 0.001-5 mg/day is administered to a human. The therapeutically effective dose level will depend on many factors as noted above. In addition, it is well within the skill of the art to start doses of the composition at relatively low levels, and increase the dosage until the desired effect is achieved.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho) esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (copolymers of lactic acid and glycolic acid).

The TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be

used. The liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microtine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with one or more compositions of the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (copolymers of lactic acid and glycolic acid).

The TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be formulated as aerosols for application, such as by inhalation.

These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microtine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with one or more compositions of the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

The TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microtine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally

with surgery, radiation, or chemotherapy combined with one or more compositions of the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (copolymers of lactic acid and glycolic acid).

The TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microtine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with one or more compositions of the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be used with a sustained-release matrix, which can be

made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

The TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The liposome can contain, in addition to one or more compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microtine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with one or more compositions of the present invention and then one or more compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

Compositions including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be used with a sustained-release matrix, which can be made of materials, usually polymers, which are degradable by enzymatic or acid-based hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix for example is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid), polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyr-

rolidone and silicone. A preferred biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

The TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically-acceptable and metabolizable lipid capable of forming liposomes can be used. The liposome can contain, in addition to compositions of the present invention, stabilizers, preservatives, excipients, and the like. Examples of lipids are the phospholipids and the phosphatidyl cholines (lecithins), both natural and synthetic. Methods to form liposomes are known in the art.

The TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be formulated as aerosols for application, such as by inhalation. These formulations for administration to the respiratory tract can be in the form of an aerosol or solution for a nebulizer, or as a microtine powder for insufflation, alone or in combination with an inert carrier such as lactose. In such a case, the particles of the formulation will, in one embodiment, have diameters of less than 50 microns, in one embodiment less than 10 microns.

Compositions including the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be used in combination with other compositions and/or procedures for the treatment of the conditions described above. For example, a tumor may be treated conventionally with surgery, radiation, or chemotherapy combined with compositions of the present invention and then compositions of the present invention may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize, inhibit, or reduce the growth of any residual primary tumor.

7.1. ADDITIONAL EMBODIMENTS

The pharmaceutical compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, *Remington's Pharmaceutical Sciences* (Martin E W [1995] Easton Pa., Mack Publishing Company, 19th ed.) describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations

of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The pharmaceutical compositions including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa. describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The pharmaceutical compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, *Remington's Pharmaceutical Sciences* (Martin E W [1995] Easton Pa., Mack Publishing Company, 19th ed.) describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The pharmaceutical compositions including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa. describes formulations which can be used in connection with the subject invention. Formulations

suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The pharmaceutical compositions including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, Remington's Pharmaceutical Sciences (Mack Publishing Company, Easton, Pa.) describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The pharmaceutical compositions including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations are described in a number of sources which are well known and readily available to those skilled in the art. For example, *Remington's Pharmaceutical Sciences* (Martin E W [1995] Easton Pa., Mack Publishing Company, 19th ed.) describes formulations which can be used in connection with the subject invention. Formulations suitable for administration include, for example, aqueous sterile injection solutions, which may contain antioxidants, buffers, bacteriostats, and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and nonaqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the condition of the sterile liquid carrier, for example, water for injections, prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powder, granules, tablets, etc. It should be understood that in addition to the ingredients particularly mentioned above, the

formulations of the subject invention can include other agents conventional in the art having regard to the type of formulation in question.

The methods of the present invention, for example, for inhibiting the growth of a cancerous cell, can be advantageously combined with at least one additional therapeutic method, including but not limited to chemotherapy, radiation therapy, therapy that selectively inhibits Ras oncogenic signaling, or any other therapy known to those of skill in the art of the treatment and management of cancer, such as administration of an anti-cancer agent.

Administration of TCN, TCN-P, TCN-PM and/or related compounds as a salt may be carried out. Examples of pharmaceutically acceptable salts are organic acid addition salts formed with acids which form a physiological acceptable anion, for example, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, alpha-ketoglutarate, and alpha-glycerophosphate. Suitable inorganic salts may also be formed, including hydrochloride, sulfate, nitrate, bicarbonate, and carbonate salts.

Pharmaceutically acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitable acid affording a physiologically acceptable anion. Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of carboxylic acids can also be made.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an

assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The

amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs can be formulated as pharmaceutical compositions and administered to a subject, such as a human or veterinary patient, in a variety of forms adapted to the chosen route of administration, i.e., orally or parenterally, by intravenous, intramuscular, topical or subcutaneous routes.

Thus, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs of the present invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle (i.e., carrier) such as an inert diluent or an assimilable edible carrier. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the compounds may be combined with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active agent. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2 to about 60% of the weight of a given unit dosage form. The amount of the active compound in such therapeutically useful compositions is such that an effective dosage level will be obtained.

The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to

otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the compounds of the invention, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the compounds of the invention may be incorporated into sustained-release preparations and devices.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active agents or their salts can

be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders including the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form must be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium including, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating compounds of the invention in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

For topical administration, the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

For topical administration, the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

For topical administration, the TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

For topical administration, the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof may be applied in pure-form, i.e., when they are liquids. However,

it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

For topical administration, the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

For topical administration, the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs may be applied in pure-form, i.e., when they are liquids. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Useful liquid carriers include water, alcohols or glycols or water-alcohol/glycol blends, in which the compounds of the invention can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user. Examples of useful dermatological compositions which can be used to deliver the compounds of the invention to the skin are disclosed in Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat. No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Woltzman (U.S. Pat. No. 4,820,508).

Useful dosages of the pharmaceutical compositions of the present invention can be determined by comparing their *in vitro* activity, and *in vivo* activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949.

In one non-limiting embodiment, the concentration of the active agent in a liquid composition, such as a lotion, can be from about 0.1-25 wt.-%, or from about 0.5-10 wt.-%. In one embodiment, the concentration in a semi-solid or solid composition such as a gel or a powder can be about 0.1-5 wt.-%, preferably about 0.5-2.5 wt.-%. In one embodiment, single dosages for injection, infusion or ingestion will generally vary between 5-1500 mg, and may be administered, i.e., 1-3 times daily, to yield levels of about 0.1-50 mg/kg, for adults. A non-limiting dosage of the present invention can be between 7.5 to 45 mg per day, administered orally, with appropriate adjustment for the body weight of an individual.

Accordingly, the invention includes a pharmaceutical composition including TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes or a pharma-

aceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the TCN, TCN-P, TCN-PM and/or related compounds and one or more taxanes in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of TCN, TCN-P, TCN-PM and/or related compounds (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

The invention also includes a pharmaceutical composition including TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab or a salt thereof, or a pharmaceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the TCN, TCN-P, TCN-PM and/or related compounds and a molecule that modulates the HER2/neu (erbB2) receptor, for example, trastuzumab, in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

The invention also includes a pharmaceutical composition including TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of TCN, TCN-P, TCN-PM and/or related compounds and one or more anthracycline analogs or a pharmaceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the TCN, TCN-P, TCN-PM and/or related compounds and

one or more anthracycline analogs in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

The invention also includes a pharmaceutical composition including TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the TCN, TCN-P, TCN-PM and/or related compounds and an erlotinib-like compound, for example, gefitinib, erlotinib or a salt thereof in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

The invention also includes a pharmaceutical composition including TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds or a pharmaceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the TCN, TCN-P, TCN-PM and/or related compounds and one or more platinum compounds in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

The invention also includes a pharmaceutical composition including TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof or pharmaceutically acceptable salts thereof, in combination with a pharmaceutically acceptable carrier. Pharmaceutical compositions adapted for oral, topical or parenteral administration, including an amount of TCN, TCN-P, TCN-PM and/or

related compounds and bortezomib and derivatives thereof analogs or a pharmaceutically acceptable salt thereof, constitute a preferred embodiment of the invention. The dose administered to a subject, particularly a human, in the context of the present invention should be sufficient to affect a therapeutic response in the patient over a reasonable time frame. One skilled in the art will recognize that dosage will depend upon a variety of factors including the condition of the animal, the body weight of the animal, as well as the severity and stage of the cancer.

A suitable dose is that which will result in a concentration of the TCN, TCN-P, TCN-PM and/or related compounds and bortezomib and derivatives thereof analogs in tumor tissue which is known to affect the desired response. The preferred dosage is the amount which results in maximum inhibition of cancer cell growth, without unmanageable side effects. Administration of API-2 (or a pharmaceutically acceptable salt thereof) can be continuous or at distinct intervals, as can be determined by a person of ordinary skill in the art.

Mammalian species which benefit from the disclosed methods for the inhibition of cancer cell growth, include, but are not limited to, primates, such as apes, chimpanzees, orangutans, humans, monkeys; domesticated animals (e.g., pets) such as dogs, cats, guinea pigs, hamsters, Vietnamese pot-bellied pigs, rabbits, and ferrets; domesticated farm animals such as cows, buffalo, bison, horses, donkey, swine, sheep, and goats; exotic animals typically found in zoos, such as bear, lions, tigers, panthers, elephants, hippopotamus, rhinoceros, giraffes, antelopes, sloth, gazelles, zebras, wildebeests, prairie dogs, koala bears, kangaroo, opossums, raccoons, pandas, hyena, seals, sea lions, elephant seals, otters, porpoises, dolphins, and whales. The terms "patient" and "subject" are used herein interchangeably and are intended to include such human and non-human mammalian species. Likewise, in vitro methods of the present invention can be earned out on cells of such mammalian species.

Patients in need of treatment using the methods of the present invention can be identified using standard techniques known to those in the medical profession.

The following examples are offered by way of illustration and not by way of limitation.

8. EXAMPLES

8.1. Example 1

In Vitro Screening

Cell Lines and NCI Diversity Set.

All cell lines can be purchased from ATCC or described previously (Cheng, J. Q., et al. *Oncogene*, 14: 2793-2801, 1997; West, K. A., et al. *Drug Resist. Updat.*, 5: 234-248, 2002; Satyamoorthy, K., et al. *Cancer Res.* 61: 7318-7324, 2001). The NCI Structural Diversity Set is a library of 1,992 compounds selected from the approximately 140,000-compound NCI drug depository. In-depth data on the selection, structures, and activities of these diversity set compounds can be found on the NCI Developmental Therapeutics Program web site.

Screening for Inhibition of Akt-Transformed Cell Growth.

AKT2 transformed NIH3T3 cells or LXSXN vector-transfected NIH3T3 control cells (Cheng, J. Q., et al. *Oncogene*, 14: 2793-2801, 1997) are plated into 96-well tissue culture plate. Following treatment with 5 μ M of NCI Diversity Set compound, cell growth can be detected with CellTier 96 One Solution Cell Proliferation kit (Promega). Compounds that

inhibit growth in AKT2-transformed but not LXSXN-transfected NIH3T3 cells are considered as candidates of Akt inhibitor and subjected to further analysis.

In Vitro Protein Kinase, Cell Survival and Apoptosis Assays.

In vitro kinase is performed as previously described (see, for example, Jiang, K., Coppola, et al. *Mol. Cell. Biol.*, 20:139-148, 2000). Cell survival is assayed with MTS (Promega). Apoptosis was detected with annexin V, which is performed as previously described (Jiang, K., Coppola, et al. *Mol. Cell. Biol.*, 20:139-148, 2000). Recombinant Akt and PDK1 are purchased from Upstate Biotechnology Inc.

Results

Identification of Small Molecule Inhibitor of Akt Signaling Pathway, API-2.

Frequent alterations of Akt has been detected in human cancer and disruption of Akt pathway induces apoptosis and inhibits tumor growth (Jetzt, A., et al. *Cancer Res.*, 63: 697-706, 2003). Thus, Akt is considered as an attractive molecular target for development of novel cancer therapeutics. To identify small molecule inhibitor(s) of Akt, a chemical library of 1,992-compounds from the NCI (the NCI Diversity Set) is evaluated for agents capable of inhibition of growth in AKT2-transformed but not empty vector LXSXN-transfected NIH3T3 cells. Repeated experiments showed that 32 compounds inhibited growth only in AKT2-transformed cells. The most potent of these compounds, API-2 (NCI identifier: NSC 154020), can suppress cell growth at a concentration of 50 nM. FIG. 1A shows the chemical structure of API-2, which is also known as triciribine (Schweinsberg, P. D., et al. *Biochem Pharmacol.*, 30: 2521-2526, 1981). The fact that API-2 inhibits selectively AKT-2 transformed cells over untransformed parental cells prompted us to determine whether API-2 is an inhibitor of AKT2 kinase.

To this end, AKT2 is immunoprecipitated with anti-AKT2 antibody from AKT-2 transformed NIH3T3 cells following treatment with API-2. AKT2 immunoprecipitates were immunoblotted with anti-phospho-Akt antibodies. As shown in FIG. 1B, API-2 significantly inhibited AKT2 phosphorylation at both threonine-309 and serine-474, which are required for full activation of AKT2 (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). As three isoforms of Akt share high homology and similar structure, the effect of API-2 on their kinase activities is evaluated. HEK293 cells are transfected with HA-Akt1, -AKT2 and -AKT3, serum-starved overnight and treated with API-2 for 60 min prior to EGF (50 ng/ml) stimulation. Triple experiments showed that API-2 suppressed EGF-induced kinase activity and phosphorylation of Akt1, AKT2 and AKT3 (FIG. 1C). However, kinase activity of recombinant constitutively active AKT2 (Myr-AKT2) is not inhibited by API-2 in an in vitro kinase reaction (FIG. 1D), suggesting that API-2 does not directly inhibit Akt in vitro and that API-2 neither functions as ATP competitor nor as the substrate competitor that binds to active site of Akt.

API-2 does not Inhibit Known Upstream Activators of Akt.

It has been well documented that Akt is activated by extracellular stimuli and intracellular signal molecules, such as active Ras and Src, through a PI3K-dependent manner. Therefore, API-2 inhibition of Akt could result from targeting upstream molecule(s) of Akt. As PI3K and PDK1 are direct upstream regulators of Akt (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999), whether API-2 inhibits PI3K and/or PDK1 is examined. HEK293 cells are serum-starved and then can be treated with API-2 or PI3K inhibitor, wortmannin, for 30 min prior to EGF stimulation. PI3K is

immunoprecipitated with anti-p110 α antibody. The immunoprecipitates are subjected to in vitro PI3K kinase assay using PI-4-P as a substrate. As shown in FIG. 2A, the EGF-induced PI3K activity is inhibited by wortmannin but not by API-2. To evaluate the effect of API-2 on PDK1, an assay in which recombinant PDK1 promotes the threonine-309 phosphorylation of AKT2 peptides is used in the presence of lipid vesicles containing phosphatidylinositol. As shown in FIG. 2B, the assay is potently inhibited by the control PDK1 inhibitor staurosporine (IC₅₀=5 nM). In contrast, API-2 displayed only 21% inhibition of the assay at the highest concentration tested (5.1 μ M). To further evaluate the effect of API-2 on PDK1 activation, the autophosphorylation level of PDK1 at serine-241, a residue that is phosphorylated by itself and is critical for its activity is examined (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999), following API-2 treatment of HEK293 cells. Triplicate experiments show that phosphorylation levels of PDK1 are not inhibited by API-2 (FIG. 2B). However, PI3K inhibitor wortmannin can inhibit EGF-stimulated PDK1 (FIG. 2B).

API-2 is Highly Selective for the Akt Over PKC, PKA, SGK, STAT, JNK, p38, and ERK Signaling Pathways.

Akt belongs to AGC (PKA/PKG/PKC) kinase family, which also include PKA, PKC, serum- and glucocorticoid-inducible kinase (SGK), p90 ribosomal S6 kinase, p70S6K, mitogen- and stress-activated protein kinase and PKC-related kinase. Among AGC kinase family, protein structures of PKA, PKC and SGK are more close to Akt kinase than other members. Therefore, next examined are the effects of API-2 on the enzymatic activities of these 3 kinases. HEK293 cells are transfected with HA-tagged PKA, PKC α or SGK. In vitro kinase assay and immunoblotting analysis show that the kinase activities of PKA and PKC α are inhibited by PKAI and Ro 31-8220, a PKC inhibitor, respectively, whereas API-2 exhibits no effect on their activities (FIGS. 2C and 2E). Further, serum-induced SGK kinase activity is attenuated by wortmannin but not by API-2 (FIG. 2D). In addition, it is determined whether API-2 has effect on other oncogenic survival pathways. Western blotting analyses with commercially available anti-phospho-antibodies reveals that phosphorylation levels of Stat3, JNK, p38 and Erk1/2 were not affected by API-2 treatment (FIG. 2F). These data indicate that API-2 specifically inhibits Akt signaling pathway.

API-2 Suppresses Cell Growth and Induces Apoptosis in Akt-Overexpressing/Activating Human Cancer Cell Lines.

The ability of API-2 to selectively inhibit the Akt pathway suggests that it should inhibit proliferation and/or induces apoptosis preferentially in those tumor cells with aberrant expression/activation of Akt. As activation of Akt in human malignancies commonly results from overexpression of Akt or PTEN mutations, API-2 is used to treat the cells that express constitutively active Akt, caused by overexpression of AKT2 (OVCAR3, OVCAR8, PANC1 and AKT2-transformed NIH3T3) or mutations of the PTEN gene (PC-3, LNCaP, MDA-MB-468), and cells that do not (OVCAR5, DU-145, T47D, COLO357 and LXSNIH3T3) as well as melanoma cells that are activated by IGF-1 to activate Akt or do not respond to growth stimulation by IGF-1 (Santamoorthy, K., et al. *Cancer Res.* 61: 7318-7324, 2001). Immunoblotting analysis showed that phosphorylation levels of Akt are inhibited by API-2 only in the cells expressing elevated Akt or responding to IGF-1 stimulation (FIG. 3A). Accordingly, API-2 inhibited cell growth to a much higher degree in Akt-overexpressing/activating cells as compared to those with low levels of Akt. As shown in FIG. 3B, API-2

treatment inhibited cell proliferation by approximate 50-60% in Akt-overexpressing/activating cell lines, LNCaP, PC-3, OVCAR3, OVCAR8, PANC1, MDA-MB-468, and WM35, whereas only by about 10-20% in DU145, OVCAR5, COLO357, T47D and WM852 cells, which exhibit low levels of Akt or do not respond to growth stimulation by IGF-1. Moreover, API-2 induces apoptosis by 8-fold (OVCAR3), 6-fold (OVCAR8), 6-fold (PANC1), and 3-fold (AKT2-NIH3T3). No significant difference of apoptosis is observed between API-2 and vehicle (DMSO) treatment in OVCAR5, COLO357 and LXSNIH3T3 cells. Thus, API-2 inhibits cell growth and induces apoptosis preferentially in cells that express aberrant Akt.

API-2 in Combination with Taxol Acts Synergistically to Suppress Cell Growth and Induce Apoptosis in Akt-Overexpressing/Activating Human Cancer Cell Lines.

The ability of API-2 and Taxol to selectively inhibit the Akt pathway suggests that it should inhibit proliferation and/or induces apoptosis preferentially in those tumor cells with aberrant expression/activation of Akt. As activation of Akt in human malignancies commonly results from overexpression of Akt or PTEN mutations, API-2 and Taxol in combination are used to treat the cells that express constitutively active Akt, caused by overexpression of AKT2 (OVCAR3, OVCAR8, PANC1 and AKT2-transformed NIH3T3) or mutations of the PTEN gene (PC-3, LNCaP, MDA-MB-468), and cells that do not (OVCAR5, DU-145, T47D, COLO357 and LXSNIH3T3) as well as melanoma cells that are activated by IGF-1 to activate Akt or do not respond to growth stimulation by IGF-1. (FIG. 3A). API-2 and Taxol inhibits cell growth to a much higher degree in Akt-overexpressing/activating cells as compared to those with low levels of Akt. As shown in FIG. 9, API-2/Taxol treatment inhibits cell proliferation by approximate 50-65% in Akt-overexpressing/activating cell lines, LNCaP, PC-3, OVCAR3, OVCAR8, PANC1, MDA-MB-468, and WM35, whereas only by about 10-20% in DU145, OVCAR5, COLO357, T47D and WM852 cells, which exhibit low levels of Akt or do not respond to growth stimulation by IGF-1. Thus, API-2 and Taxol acts synergistically to inhibit cell growth and induce apoptosis preferentially in cells that express aberrant Akt.

API-2 Inhibits Downstream Targets of Akt.

It has been shown that Akt exerts its cellular effects through phosphorylation of a number of proteins (Datta, S. R., et al. *Genes Dev.* 13: 2905-2927, 1999). More than 20 proteins have been identified as Akt substrates, including the members of Forkhead protein family (FKHR, AFX and FKHL1), tuberlin/TSC2, p70^{S6K}, GSK-3, p21^{WAF1/Cip1}, p27^{kip1}, MDM2, Bad, ASK1 and IKK α etc. It is next examined whether API-2 inhibits downstream targets of Akt. As anti-phospho-tuberlin, -Bad, -AFX, and -GSK-3 antibodies are commercially available, therefore, the effect of API-2 on their phosphorylation induced by Akt was determined. Following API-2 (1 μ M) treatment, OVCAR3 cells was lysed and immunoblotted with the individual anti-phospho-antibody. FIG. 4A shows that API-2 considerably inhibited the phosphorylation levels of tuberlin leading to stabilization and upregulation of tuberlin (Dan, H. C., et al. *J. Biol. Chem.*, 277: 35364-35370, 2002). The phosphorylation levels of Bad, GSK-3, and AFX are partially attenuated by API-2. These data suggest that API-2 induces cell death and cell growth arrest by inhibiting phosphorylation of its downstream targets. API-2 inhibition of Akt downstream targets at different degrees could be due to the fact that phosphorylation sites of these targets are also regulated by other kinase(s), for instance, Bad serine-136 is phosphory-

101

lated by PAK1 in addition to Akt (Schurmann, A., et al. Mol. Cell. Biol., 20: 453-461, 2000).

8.2 Example 2

Antitumor Activity in Nude Mouse Tumor Xenograft Model

Tumor cells can be harvested, suspended in PBS, and can be injected s.c. into the right and left flanks (2×10^6 cells/flank) of 8-week-old female nude mice as reported previously (Sun, J., Blaskovic, et al. Cancer Res., 59: 4919-4926, 1999). When tumors reach about 100-150 mm³, animals are randomized and dosed i.p. with 0.2 ml vehicle of the TCN, TCN-P, TCN-PM and/or related compounds and/or the taxane daily. Control animals receive DMSO (20%) vehicle, whereas treated animals can be injected with API-2 (1 mg/kg/day) in 20% DMSO.

API-2 Inhibits the Growth of Tumors in Nude Mice that Overexpress Akt. Frequent overexpression/activation and/or amplification of AKT1 and AKT2 in human ovarian and pancreatic cancer was shown (Cheng, J. Q., and Nicosia, S. V. AKT signal transduction pathway in oncogenesis. In Schwab D, Editor, Encyclopedic Reference of Cancer. Berlin Heidelberg and New York: Springer; 2001. pp 35-7) Inhibition of Akt pathway by inhibitors of PI3K, HSP70, Src and farnesyltransferase resulted in cell growth arrest and induction of apoptosis (Solit, D. B., et al. Cancer Res., 63: 2139-2144, 2003; Xu, W., et al. Cancer Res., 63: 7777-7784, 2003). A recent study showed that the tumor growth of xenografts with elevated Akt was also significantly inhibited by intratumoral injection of adenovirus of dominant negative Akt (Jetzt, A., et al. Cancer Res., 63: 697-706, 2003). Because API-2 inhibits Akt signaling and induces apoptosis and cell growth arrest only in cancer cells with elevated levels of Akt (FIG. 3), the growth of tumors with elevated levels of Akt should be more sensitive to API-2 than that of tumors with low levels of Akt in nude mice. To this end, s.c. Akt-overexpressing cells (OVCAR3, OVCAR8 and PANC-1) are s.c. implanted into the right flank, and those cell lines that express low levels of Akt (OVCAR5 and COLO357) into the left flank of mice. When the tumors reach an average size of about 100-150 mm³, the animals are randomized and treated i.p. with either vehicle or API-2 (1 mg/kg/day). As illustrated in FIG. 4B, OVCAR-5 and COLO357 tumors treated with vehicle grew to about 800-1,000 mm³ 49 days after tumor implantation. OVCAR3, OVCAR8 and PANC1 tumors treated with vehicle control grew to about 700-900 mm³ 49 days after tumor implantation. API-2 inhibited OVCAR3, OVCAR8 and PANC1 tumor growth by 90%, 88% and 80%, respectively. In contrast, API-2 has little effect on the growth of OVCAR5 and COLO357 cells in nude mice (FIGS. 4B-4D and data not shown). At dose 1 mg/kg/day, API-2 had no effects on blood glucose level, body weight, activity and food intake of mice. In treated tumor samples, Akt activity was inhibited by API-2 without change of total Akt content (FIG. 4E). Taken together, these results indicate that API-2 selectively inhibits the growth of tumors with elevated levels of Akt.

8.3 Example 3

TCN Directly Inhibits Wild Type Akt Kinase Activity

API-2 (TCN) can directly inhibit wild type Akt kinase activity induced by PDK1 in vitro (FIG. 1). This result

102

supports that API-2 is a direct Akt inhibitor and that the underlying mechanism may be API-2 binding to PH domain and/or threonine-308 of Akt. An in vitro kinase assay is performed with recombinant of PDK1 and Akt in a kinase buffer containing phosphatidylinositol-3,4,5-P3 (PIP3), API-2 and histone H2B as substrate. After incubation of 30 min, the reactions were separated by SDS-PAGE and exposed in a film.

8.4 Example 4

TCN is Effective in Cancer Resistant Cells

The effects of TCN (API-2) are tested in cisplatin, paclitaxel, and tamoxifen resistant A270CP, C-13, OVCAR433 and MCF7/TAM cells. API-2 overcame cisplatin, paclitaxel, and tamoxifen resistance in these cells.

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention, will be obvious to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of this invention.

8.5 Example 5

TCN Potentiates Growth Inhibition by Trastuzumab and Induces Apoptosis

Materials and Methods

Cell Lines and Cell Cultures. The tumorigenic BT474.m1 subline maintained in Dulbecco's modified Eagle's medium: Ham's F-12 medium (1:1) with 8-10% FBS.

Antibodies and Reagents. Trastuzumab was a gift from Genentech (San Francisco, Calif.). RAD001 (everolimus) was a gift from Novartis (East Hanover, N.J.). QLT0267 and KP 372-1 were gifts from QLT Inc. (Vancouver, BC). Triciribine (6-Amino-4-methyl-8-((3-D-ribofuranosyl)-4H, 8H-pyrrolo[4,3,2-de]pyrimido[4,5-c]pyridazine) was purchased from Berry & Associates, Inc. (Ann Arbor, Mich.). Edelfosine was purchased from Calbiochem (San Diego, Calif.). A selective Akt inhibitor, 4ADPIB (4-amino-2(3,4-dichloro-phenyl)-N-(1H-indazol-5-yl)-butyramide) (U.S. Pat. No. 6,919,340), was synthesized. PTEN antibodies were from Santa Cruz Biotechnology (Santa Cruz, Calif.). 3-actin antibodies were from Sigma (St. Louis, Mo.). All other antibodies were purchased from Cell Signaling Technology (Danvers, Mass.).

PTEN Antisense and Non-specific Oligonucleotide Transient Transfection. Antisense (AS) oligonucleotides specific for PTEN, control non-specific (NS) oligonucleotides and procedures for transfection were followed according to Nagata Y. et al., Cancer Cell, 2004. 6(2): p. 117-27.

Cell Proliferation Assay. PTEN AS/NS transfected BT474.m1 cells were plated 2500 cells/0.32 cm² well. Cells were treated with inhibitors of the Akt/mTOR pathway alone or in combination with trastuzumab as described for 5 days and viable cells were measured by MTS assay using the CellTiter 96 AQ nonradioactive cell proliferation assay kits according to the manufacturer's protocol (Promega, Madison, Wis.). Treated cells were compared to control DMSO treated BT474.m1 cells to calculate percentage of growth inhibition.

APO-BRDU TUNEL Assay. The PTEN AS and NS transfected BT474.m1 cells were plated in 6-well plates ($4-6 \times 10^5$ cells/well). Twenty-four hours after plating, the cells were treated as indicated for 72 hours with trastu-

zumab, TCN and/or RAD001. The floating and adherent cells were collected, labeled and stained using the APO-BRDU™ TUNEL assay kit (Phoenix Flow Systems, San Diego, Calif.) according to the manufacturer's protocol. Data was collected and analyzed using a FACScan flow cytometer and CellQuest Pro 4.02 software (Becton Dickinson, Franklin Lakes, N.J.). At least 10,000 events were examined.

SDS-PAGE and Immunoblot Analysis. Cells transfected with PTEN AS/NS oligonucleotides were treated as indicated. Immunoblotting was performed as described by Nagata Y. et al., *Cancer Cell*, 2004. 6(2): p. 117-27.

Results

Triciribine and RAD001 potentiate growth inhibition by trastuzumab in PTEN-deficient cells. To find a strategy to overcome trastuzumab resistance, particularly resistance caused by PTEN loss, 6 different small molecule inhibitors were tested, which directly or indirectly targeted the PI3K/Akt/mTOR signal transduction pathway, a major pathway activated by overexpression of ErbB2 and the loss of PTEN. The goal was to identify compounds that would exhibit synergistic effects with trastuzumab, preferably at a low dose of the compound in order to minimize toxicity. The drugs chosen targeted Akt, mTOR and integrin-linked kinase (ILK) (FIG. 10A). BT474.m1 cells are a tumorigenic subline of the BT474 breast cancer cell line and express high levels of ErbB2. When PTEN levels are decreased by transfection with PTEN antisense oligonucleotides (PTEN AS), BT474.m1 breast cancer cells become more resistant to the anti-proliferative effects of trastuzumab than cells with normal levels of PTEN and provide a good experimental model for breast cancers in which trastuzumab resistance is caused by PTEN loss (Nagata Y. et al., *Cancer Cell*, 2004. 6(2): p. 117-27 and FIGS. 9B and 9C). Nonspecific oligonucleotides (NS) were transfected as controls. Treatment with PTEN AS oligonucleotides effectively lowered PTEN levels (FIGS. 12A and 12B). Neither PTEN AS nor NS control oligonucleotides altered ErbB2 levels in the cells.

PTEN AS and NS-transfected BT474.m1 cells were treated with each of the 6 compounds or trastuzumab alone and in combination for 5 days and evaluated cell proliferation as compared to DMSO-treated control. Using growth inhibition as a biological endpoint, we compared the ability of each drug to exhibit cooperative effects with trastuzumab using doses of drug that resulted in ~20-40% growth inhibition when administered alone (FIG. 10A).

Almost all of the compounds displayed growth inhibitory effects, particularly at high concentrations and in cells with intact PTEN (FIG. 10A). However, two of the compounds, triciribine and RAD001, markedly enhanced growth inhibition in the PTEN AS cells when combined with trastuzumab as compared to trastuzumab or either compound alone (FIG. 10A). Triciribine (also called API-2), a compound that inhibits Akt activation, potentiated growth inhibition by trastuzumab over a 20-fold concentration range (FIG. 10B). The mTOR inhibitor RAD001 (everolimus) increased growth inhibition by trastuzumab when RAD001 was administered at low doses (<1 nM) (FIG. 10C). Strikingly, triciribine and RAD001 were able to cooperate with trastuzumab to inhibit cell growth at similar levels in the PTEN AS and NS cells (FIGS. 10B and 10C). In essence, triciribine and RAD001 were able to restore trastuzumab sensitivity to PTEN-deficient cells. Triciribine and RAD001 were also effective as single agents, both in PTEN AS and NS cells, at doses greater than 5 nM and 1.5 nM for triciribine and RAD001, respectively (FIGS. 10B and 10C).

A third compound, the ILK inhibitor QLT0267, potentiated growth inhibition by trastuzumab within a narrow dose range (~5-15 pM) (FIG. 10A). Because the dose range in which QLT0267 exhibited cooperative effects with trastuzumab was narrow, this compound further was not investigated further. At concentrations greater than 20 pM, QLT0267 had no cooperative effect with trastuzumab but significantly inhibited cell growth as a single agent.

Induction of Apoptosis Following Combination Treatment. To assess if growth inhibition was accompanied by apoptosis, we treated PTEN AS and NS transfected BT474.m1 cells with triciribine, RAD001 and trastuzumab, alone or combined, and quantified the levels of apoptosis (FIG. 10). RAD001 did not significantly induce apoptosis alone or in combination with trastuzumab. Although the number of TUNEL-positive, apoptotic cells increased slightly following treatment with trastuzumab or triciribine alone, this increase was not statistically significant. However, the combination of triciribine and trastuzumab, significantly induced apoptosis as compared with all other treatments in both PTEN AS and NS transfected cells (FIG. 11).

8.6 Example 6

TCN Inhibits Activation of Akt and mTOR Inhibition of Downstream Signaling Molecules

Immunoblot analysis verified that triciribine and RAD001 blocked activation of Akt and mTOR, important signaling molecules activated by ErbB2 and the targets of triciribine and RAD001 respectively. Phosphorylation of Akt on Thr308 and Ser473 was analyzed as an indicator of Akt activity and mTOR activity was assessed by the phosphorylation of p70S6K (70-kDa ribosomal protein S6 kinase), an mTOR target. After triciribine treatment, phosphorylation of Akt on both sites was substantially decreased (FIG. 12A). In PTEN-deficient cells, the levels of Akt phosphorylation following triciribine and trastuzumab combination treatment were similar to those seen in cells with intact PTEN (FIG. 12A, lanes 4 & 8). Thus, triciribine overcame the adverse effects of PTEN loss by effectively blocking Akt activation. RAD001 dramatically blocked phosphorylation of p70S6K (FIG. 12B). However, RAD001 combined with trastuzumab did not lower p70S6K phosphorylation beyond that seen with RAD001 alone (FIG. 12B). A feedback loop has been recently identified which results in Akt phosphorylation and activation following treatment with mTOR inhibitors, such as RAD001 (O'Reilly K. E. et al., *Cancer Res*, 2006. 66(3): p. 1500-8). It was also observed that feedback activation of Akt by RAD001 and combination therapy with trastuzumab and RAD001 eliminated Akt phosphorylation by this feedback loop (FIG. 12B, lanes 3 vs. 4), consistent with the notion that Akt activation following mTOR inhibition is dependent on upstream receptor tyrosine kinases (O'Reilly K. E. et al., *Cancer Res*, 2006. 66(3): p. 1500-8). In summary, both drugs inhibited their predicted target kinases and combination treatment had a larger inhibitory effect on the Akt/mTOR signaling pathway than any single agent, even in PTEN-deficient cells.

TCN and Trastuzumab Inhibit Tumor Growth in PTEN-Deficient Tumors

Materials and Methods

Xenograft Human Tumor Model in SCID Mice. Female, 6-week-old, severe combined immunodeficiency (SCID) mice were from Taconic Farms (Hudson, N.Y.). Tumor xenografts were performed as described in Nagata Y. et al., *Cancer Cell*, 2004, 6(2): p. 117-27. When the xenograft tumors reached the average size of 100-150 mm³, the mice were divided into 6 groups, each with 7 mice and an even distribution of tumor sizes, and treated as follows. PTEN antisense (30 pg) oligonucleotides were administered to each mouse weekly via intratumor injection. One week after PTEN AS oligonucleotide administration was initiated, drug treatment began. Trastuzumab was given at a dose of 0.5 mg/kg twice a week in 200 pL saline through intratumor injection at multiple sites. Triciribine was given at a dose of 0.5 mg/kg/day in 200 pL 20% DMSO saline solution through intraperitoneal (I.P.) injection. RAD001 was given via gavage at a dose of 1 pg/kg in 500 pL 5% glucose water twice a week. 20% DMSO saline solution (200 pL/day) was given through I.P. injection. The tumors were measured twice weekly with calipers and the volume of the tumors was calculated as: volume=length× width/2.

Statistical Analysis. One-way ANOVA was performed using GraphPad Prism 3.0 for Windows (Graph Pad Software, San Diego, Calif.).

Results

The earlier biological and molecular data were very promising, however, in vivo studies provide the most stringent test for therapeutic efficacy. Therefore, triciribine and RAD001 were tested in vivo. BT474.m1 cell xenografts were injected into the mammary fat pad of 6-week-old SCID mice. After tumors formed, the mice received PTEN AS weekly via intratumor injection. This protocol effectively models PTEN-deficient tumors in vivo (Nagata Y. et al., *Cancer Cell*, 2004, 6(2): p. 117-27). The mice were randomized into treatment groups receiving triciribine, RAD001, trastuzumab or DMSO alone or in combination. After treatment, the growth patterns of the tumors treated with DMSO, trastuzumab, RAD001, or triciribine alone were similar (FIGS. 13A and 13B). Growth of the tumors was not inhibited and the mice were euthanized after 3 weeks due to large tumor burdens. In contrast, combination treatment with triciribine and trastuzumab dramatically and significantly inhibited tumor growth (FIG. 13A). Many of the tumors actually decreased in size and four of 7 mice had no palpable tumors after 5 weeks of treatment. Following treatment with RAD001 and trastuzumab, tumor growth was relatively slower compared to RAD001 or trastuzumab alone (FIG. 13B). Thus combining trastuzumab with triciribine or RAD001 effectively inhibited ErbB2-overexpressing, PTEN-deficient human breast cancer xenografts in vivo.

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention will be obvious to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of this invention.

Immunoblot analysis verified that triciribine and RAD001 blocked activation of Akt and mTOR, important signaling molecules activated by ErbB2 and the targets of triciribine and RAD001 respectively. Phosphorylation of Akt on Thr308 and Ser473 was analyzed as an indicator of Akt

activity and mTOR activity was assessed by the phosphorylation of p70S6K (70-kDa ribosomal protein S6 kinase), an mTOR target. After triciribine treatment, phosphorylation of Akt on both sites was substantially decreased (FIG. 12A). In PTEN-deficient cells, the levels of Akt phosphorylation following triciribine and trastuzumab combination treatment were similar to those seen in cells with intact PTEN (FIG. 12A, lanes 4& 8). Thus, triciribine overcame the adverse effects of PTEN loss by effectively blocking Akt activation. RAD001 dramatically blocked phosphorylation of p70S6K (FIG. 12B). However, RAD001 combined with trastuzumab did not lower p70S6K phosphorylation beyond that seen with RAD001 alone (FIG. 12B). A feedback loop has been recently identified which results in Akt phosphorylation and activation following treatment with mTOR inhibitors, such as RAD001 (O'Reilly K. E. et al., *Cancer Res*, 2006, 66(3): p. 1500-8). It was also observed that feedback activation of Akt by RAD001 and combination therapy with trastuzumab and RAD001 eliminated Akt phosphorylation by this feedback loop (FIG. 12B, lanes 3 vs. 4), consistent with the notion that Akt activation following mTOR inhibition is dependent on upstream receptor tyrosine kinases (O'Reilly K. E. et al., *Cancer Res*, 2006, 66(3): p. 1500-8). In summary, both drugs inhibited their predicted target kinases and combination treatment had a larger inhibitory effect on the Akt/mTOR signaling pathway than any single agent, even in PTEN-deficient cells.

8.8 Example 8

Antitumor Activity in the Nude Mouse Tumor Xenograft Model

Tumor cells can be harvested, suspended in PBS, and can be injected s.c. into the right and left flanks (2×10⁶ cells/flank) of 8-week-old female nude mice as reported previously (Sun, J., Blaskovic, et al. *Cancer Res.*, 59: 4919-4926, 1999). When tumors reach about 100-150 mm³, animals are randomized and dosed i.p. with 0.2 ml vehicle of the TCN, TCN-P, TCN-PM and/or related compounds and/or anthracycline analogs daily. Control animals receive DMSO (20%) vehicle, whereas treated animals can be injected with API-2 (1 mg/kg/day) in 20% DMSO.

8.9 Example 9

Antitumor Activity in the Nude Mouse Tumor Xenograft Model

Tumor cells can be harvested, suspended in PBS, and can be injected s.c. into the right and left flanks (2×10⁶ cells/flank) of 8-week-old female nude mice as reported previously (Sun et al., 1999, *Cancer Res* 59: 4919-4926). When tumors reach about 100-150 mm³, animals are randomized and dosed i.p. with 0.2 ml vehicle of the TCN, TCN-P, TCN-PM and/or related compounds and/or one or more platinum compounds daily. Control animals receive DMSO (20%) vehicle, whereas treated animals can be injected with API-2 (1 mg/kg/day) in 20% DMSO.

API-2 Inhibits the Growth of Tumors in Nude Mice that Overexpress Akt. Frequent overexpression/activation and/or amplification of AKT1 and AKT2 in human ovarian and pancreatic cancer was shown (Cheng et al., AKT signal transduction pathway in oncogenesis, in Schwab D (ed.) *Encyclopedic Reference of Cancer*, Springer, pp 35-7) Inhibition of Akt pathway by inhibitors of PI3K, HSP70, Src and farnesyltransferase resulted in cell growth arrest and induc-

tion of apoptosis (Solit et al., 2003, *Cancer Res* 63: 2139-2144; Xu et al., 2003, *Cancer Res* 63: 7777-7784). A recent study showed that the tumor growth of xenografts with elevated Akt was also significantly inhibited by intratumoral injection of adenovirus of dominant negative Akt (Jetzt et al., 2003, *Cancer Res* 63: 697-706). Because API-2 inhibits Akt signaling and induces apoptosis and cell growth arrest only in cancer cells with elevated levels of Akt (FIG. 3), the growth of tumors with elevated levels of Akt should be more sensitive to API-2 than that of tumors with low levels of Akt in nude mice. To this end, s.c. Akt-overexpressing cells (OVCAR3, OVCAR8 and PANC-1) are s.c. implanted into the right flank, and those cell lines that express low levels of Akt (OVCAR5 and COLO357) into the left flank of mice. When the tumors reach an average size of about 100-150 mm³, the animals are randomized and treated i.p. with either vehicle or API-2 (1 mg/kg/day). As illustrated in FIG. 4B, OVCAR-5 and COLO357 tumors treated with vehicle grew to about 800-1,000 mm³ 49 days after tumor implantation. OVCAR3, OVCAR8 and PANC1 tumors treated with vehicle control grew to about 700-900 mm³ 49 days after tumor implantation. API-2 inhibited OVCAR3, OVCAR8 and PANC1 tumor growth by 90%, 88% and 80%, respectively. In contrast, API-2 has little effect on the growth of OVCAR5 and COLO357 cells in nude mice (FIGS. 4B-4D and data not shown). At dose 1 mg/kg/day, API-2 had no effects on blood glucose level, body weight, activity and food intake of mice. In treated tumor samples, Akt activity was inhibited by API-2 without change of total Akt content (FIG. 4E). Taken together, these results indicate that API-2 selectively inhibits the growth of tumors with elevated levels of Akt.

8.10 Example 10

TCN Overcomes Cisplatin Resistance in Cisplatin Resistant Ovarian Cancer Cells

Ovarian cancer cell lines sensitive to cisplatin, A2780S and OV2008, and their cisplatin resistant derivatives, A2780CP and C13, respectively, were treated with 0, 10, and 30 μ M cisplatin, followed by an assessment of the percentage of cells surviving the cisplatin. The resistance to cisplatin toxicity was confirmed in the A2780S and C13 cell lines (FIG. 14A). Lysates from all four cell types were resolved by SDS-PAGE and probed with antibodies to phosphorylated Akt and total Akt. The cisplatin resistant cell lines endogenously express significantly higher levels phosphorylated Akt (FIG. 14B).

It was next assessed what effects TCN would have on cisplatin resistance. A2780CP and C13 ovarian cancer cells were treated with 0, 5, 10 and 20 μ M, with or without 10 μ M TCN, followed by measurement of cell survival. The addition of TCN significantly lowered the resistance to cisplatin in both cell lines (FIG. 15). C13 cells were then treated with 0, 1, 5, 10 and 20 μ M TCN with or without 10 μ M cisplatin, followed by measurement of cell survival. The combination of TCN and cisplatin synergistically reduced cell survival (FIG. 16).

C13 cells were injected subcutaneously in nude mice, followed by treatment with vehicle (DMSO), cisplatin alone (2 mg/kg/day), TCN alone (2 mg/kg/day) or TCN and cisplatin. Tumor mass was assessed at weekly intervals for four weeks. The combination of cisplatin and TCN enhanced the ability to suppress the growth and progression of the tumor (FIG. 17).

8.11 Example 11

TCN Enhances the Effects of mTOR Inhibition

OV3 and MCF7 cells were treated with the mTOR inhibitors rapamycin and RAD001, respectively, over a 24 hour time course. Lysates were resolved by SDS-PAGE and transferred to a membrane for western blotting. Membranes were probed with antibodies to phospho-Akt, total Akt, phospho-p70S6K, and total p70S6K. Membranes with lysates derived from OV3 cells were additionally probed with antibodies to phospho-FKHRL 1 and total FKHRL 1. The use of mTOR inhibitors lead to rapid activation of Akt pathway in both cell lines (FIG. 18).

OV3 cells were then treated with either control, rapamycin, or rapamycin and TCN and MCF7 cells were treated with control, RAD001, or RAD001 and TCN. TCN markedly decreased the mTOR inhibitors' increase in Akt phosphorylation without a subsequent increase in p70S6K activity (FIG. 19). DU-145, MCF7 and OV3 cell growth rates were then assessed over 6 days in the presence of control, mTOR inhibitor, TCN, or TCN and mTOR inhibitor. The inhibition on cell growth by mTOR inhibition was enhanced by TCN (FIG. 20).

8.12 Example 12

TCN Overcomes Aurora-A Induced Cisplatin Resistance

Aurora-A is a serine/threonine kinase involved in progression of the cell cycle. The effect of Aurora-A on sensitivity to cisplatin was examined A2780S and A2780CP cells, as well as OV2008 cells, were transfected with empty vector or with a construct encoding an HA-tagged Aurora-A protein. Expression of Aurora-A was confirmed by western blotting lysates resolved by SDS-PAGE and transferred to a membrane (FIG. 21, top panels). The viability of cells was assessed in the presence of paclitaxel and cisplatin. A2780S and OV2008 cells transfected with Aurora-A showed improved viability over A2780S and OV2008 cells transfected with empty vector only (FIG. 21).

It was next assessed what effect Aurora-A exhibited on the induction of apoptosis induced by cisplatin. A2780S cells were transfected with empty vector or with a construct encoding the Aurora-A protein. Cells were then treated with cisplatin. Immunofluorescent staining (FIG. 22A) shows that expression of Aurora-A inhibits cytochrome c release. Western blot analysis revealed that introduction of Aurora-A increases phosphorylation of Akt, as well as Akt kinase activity (FIGS. 23B and C).

Next, A2780S and OV2008 cells, transfected with either empty vector or Aurora-A, were treated with either cisplatin, TCN, or cisplatin and TCN. Both A2780S and OV2008 cells expressing Aurora-A showed markedly decreased resistance to cisplatin in the presence of TCN (FIG. 23).

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention will be obvious to those skilled in the art from the foregoing detailed description of the invention. It is intended that all of these variations and modifications be included within the scope of this invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1

<211> LENGTH: 1443

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1) .. (1443)

<400> SEQUENCE: 1

```

atg aac gac gta gcc att gtg aag gag ggc tgg ctg cac aaa cga ggg      48
Met Asn Asp Val Ala Ile Val Lys Glu Gly Trp Leu His Lys Arg Gly
1          5          10          15

gaa tat att aaa acc tgg cgg cca cgc tac ttc ctc ctc aag aac gat      96
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Asn Asp
          20          25          30

ggc acc ttt att ggc tac aag gaa cgg cct cag gat gtg gat cag cga     144
Gly Thr Phe Ile Gly Tyr Lys Glu Arg Pro Gln Asp Val Asp Gln Arg
          35          40          45

gag tcc cca ctc aac aac ttc tca gtg gca caa tgc cag ctg atg aag     192
Glu Ser Pro Leu Asn Asn Phe Ser Val Ala Gln Cys Gln Leu Met Lys
          50          55          60

aca gag cgg cca agg ccc aac acc ttt atc atc cgc tgc ctg cag tgg     240
Thr Glu Arg Pro Arg Pro Asn Thr Phe Ile Ile Arg Cys Leu Gln Trp
          65          70          75          80

acc aca gtc att gag cgc acc ttc cat gta gaa acg cct gag gag cgg     288
Thr Thr Val Ile Glu Arg Thr Phe His Val Glu Thr Pro Glu Glu Arg
          85          90          95

gaa gaa tgg gcc acc gcc att cag act gtg gcc gat gga ctc aag agg     336
Glu Glu Trp Ala Thr Ala Ile Gln Thr Val Ala Asp Gly Leu Lys Arg
          100          105          110

cag gaa gaa gag acg atg gac ttc cga tca ggc tca ccc agt gac aac     384
Gln Glu Glu Glu Thr Met Asp Phe Arg Ser Gly Ser Pro Ser Asp Asn
          115          120          125

tca ggg gct gaa gag atg gag gtg tcc ctg gcc aag ccc aag cac cgt     432
Ser Gly Ala Glu Glu Met Glu Val Ser Leu Ala Lys Pro Lys His Arg
          130          135          140

gtg acc atg aac gag ttt gag tac ctg aaa cta ctg ggc aag ggc acc     480
Val Thr Met Asn Glu Phe Glu Tyr Leu Lys Leu Leu Gly Lys Gly Thr
          145          150          155          160

ttt ggg aaa gtg att ctg gta aaa gag aag gcc aca ggc cgc tac tat     528
Phe Gly Lys Val Ile Leu Val Lys Glu Lys Ala Thr Gly Arg Tyr Tyr
          165          170          175

gcc atg aag atc ctc aag aag gag gtc atc gtc gcc aag gat gag gtt     576
Ala Met Lys Ile Leu Lys Lys Glu Val Ile Val Ala Lys Asp Glu Val
          180          185          190

gcc cac acg ctt act gaa aac cgt gtc ctg cag aac tct agg cat ccc     624
Ala His Thr Leu Thr Glu Asn Arg Val Leu Gln Asn Ser Arg His Pro
          195          200          205

ttc ctt acg gcc ctc aag tac tca ttc cag acc cac gac cgc ctc tgc     672
Phe Leu Thr Ala Leu Lys Tyr Ser Phe Gln Thr His Asp Arg Leu Cys
          210          215          220

ttt gtc atg gag tat gcc aac ggg ggc gag ctc ttc ttc cac ctg tct     720
Phe Val Met Glu Tyr Ala Asn Gly Gly Glu Leu Phe Phe His Leu Ser
          225          230          235          240

cga gag cgc gtg ttc tcc gag gac cgg gcc cgc ttc tat ggt gcg gag     768
Arg Glu Arg Val Phe Ser Glu Asp Arg Ala Arg Phe Tyr Gly Ala Glu
          245          250          255

att gtg tct gcc ctg gac tac ttg cac tcg gag aag aac gtg gtg tac     816
Ile Val Ser Ala Leu Asp Tyr Leu His Ser Glu Lys Asn Val Val Tyr

```

-continued

260	265	270	
cgg gac ctg aag cta gag aac ctc atg ctg gac aag gac ggg cac atc			864
Arg Asp Leu Lys Leu Glu Asn Leu Met Leu Asp Lys Asp Gly His Ile			
275	280	285	
aag ata acg gac ttc ggg ctg tgc aag gag ggg atc aag gat ggt gcc			912
Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile Lys Asp Gly Ala			
290	295	300	
act atg aag aca ttc tgc gga acg ccg gag tac ctg gcc cct gag gtg			960
Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu Ala Pro Glu Val			
305	310	315	320
ctg gaa gac aac gac tac ggc cgt gca gtg gac tgg tgg ggg ctg ggc			1008
Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp Trp Gly Leu Gly			
325	330	335	
gtg gtc atg tat gag atg atg tgt ggc cgc ctg ccc ttc tac aac cag			1056
Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro Phe Tyr Asn Gln			
340	345	350	
gac cac gag aag ctg ttc gag ctg atc ctc atg gag gag atc cgc ttc			1104
Asp His Glu Lys Leu Phe Glu Leu Ile Leu Met Glu Glu Ile Arg Phe			
355	360	365	
ccg cgc aca ctc ggc cct gag gcc aag tcc ctg ctc tcc ggg ctg ctc			1152
Pro Arg Thr Leu Gly Pro Glu Ala Lys Ser Leu Leu Ser Gly Leu Leu			
370	375	380	
aag aag gac cct aca cag agg ctc ggt ggg ggc tct gag gat gcc aag			1200
Lys Lys Asp Pro Thr Gln Arg Leu Gly Gly Gly Ser Glu Asp Ala Lys			
385	390	395	400
gag atc atg cag cac cgg ttc ttt gcc aac atc gtg tgg cag gat gtg			1248
Glu Ile Met Gln His Arg Phe Phe Ala Asn Ile Val Trp Gln Asp Val			
405	410	415	
tat gag aag aag ctg agc cca cct ttc aag ccc cag gtc acc tct gag			1296
Tyr Glu Lys Lys Leu Ser Pro Pro Phe Lys Pro Gln Val Thr Ser Glu			
420	425	430	
act gac acc agg tat ttc gat gag gag ttc aca gct cag atg atc acc			1344
Thr Asp Thr Arg Tyr Phe Asp Glu Glu Phe Thr Ala Gln Met Ile Thr			
435	440	445	
atc acg ccg cct gat caa gat gac agc atg gag tgt gtg gac agt gag			1392
Ile Thr Pro Pro Asp Gln Asp Asp Ser Met Glu Cys Val Asp Ser Glu			
450	455	460	
cgg agg ccg cac ttc ccc cag ttc tcc tac tca gcc agt ggc aca gcc			1440
Arg Arg Pro His Phe Pro Gln Phe Ser Tyr Ser Ala Ser Gly Thr Ala			
465	470	475	480
tga			1443

<210> SEQ ID NO 2

<211> LENGTH: 480

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Asn Asp Val Ala Ile Val Lys Glu Gly Trp Leu His Lys Arg Gly
1 5 10 15

Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Asn Asp
20 25 30

Gly Thr Phe Ile Gly Tyr Lys Glu Arg Pro Gln Asp Val Asp Gln Arg
35 40 45

Glu Ser Pro Leu Asn Asn Phe Ser Val Ala Gln Cys Gln Leu Met Lys
50 55 60

Thr Glu Arg Pro Arg Pro Asn Thr Phe Ile Ile Arg Cys Leu Gln Trp
65 70 75 80

Thr	Thr	Val	Ile	Glu	Arg	Thr	Phe	His	Val	Glu	Thr	Pro	Glu	Glu	Arg
			85					90						95	
Glu	Glu	Trp	Ala	Thr	Ala	Ile	Gln	Thr	Val	Ala	Asp	Gly	Leu	Lys	Arg
			100					105					110		
Gln	Glu	Glu	Glu	Thr	Met	Asp	Phe	Arg	Ser	Gly	Ser	Pro	Ser	Asp	Asn
			115				120					125			
Ser	Gly	Ala	Glu	Glu	Met	Glu	Val	Ser	Leu	Ala	Lys	Pro	Lys	His	Arg
			130			135					140				
Val	Thr	Met	Asn	Glu	Phe	Glu	Tyr	Leu	Lys	Leu	Leu	Gly	Lys	Gly	Thr
					150					155					160
Phe	Gly	Lys	Val	Ile	Leu	Val	Lys	Glu	Lys	Ala	Thr	Gly	Arg	Tyr	Tyr
				165					170					175	
Ala	Met	Lys	Ile	Leu	Lys	Lys	Glu	Val	Ile	Val	Ala	Lys	Asp	Glu	Val
			180					185					190		
Ala	His	Thr	Leu	Thr	Glu	Asn	Arg	Val	Leu	Gln	Asn	Ser	Arg	His	Pro
							200					205			
Phe	Leu	Thr	Ala	Leu	Lys	Tyr	Ser	Phe	Gln	Thr	His	Asp	Arg	Leu	Cys
						215					220				
Phe	Val	Met	Glu	Tyr	Ala	Asn	Gly	Gly	Glu	Leu	Phe	Phe	His	Leu	Ser
					230					235					240
Arg	Glu	Arg	Val	Phe	Ser	Glu	Asp	Arg	Ala	Arg	Phe	Tyr	Gly	Ala	Glu
				245					250					255	
Ile	Val	Ser	Ala	Leu	Asp	Tyr	Leu	His	Ser	Glu	Lys	Asn	Val	Val	Tyr
				260				265					270		
Arg	Asp	Leu	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	Gly	His	Ile
							280					285			
Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Lys	Asp	Gly	Ala
						295					300				
Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	Pro	Glu	Val
					310					315					320
Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	Gly	Leu	Gly
				325					330					335	
Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	Tyr	Asn	Gln
				340				345					350		
Asp	His	Glu	Lys	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Glu	Ile	Arg	Phe
							360					365			
Pro	Arg	Thr	Leu	Gly	Pro	Glu	Ala	Lys	Ser	Leu	Leu	Ser	Gly	Leu	Leu
						375					380				
Lys	Lys	Asp	Pro	Thr	Gln	Arg	Leu	Gly	Gly	Gly	Ser	Glu	Asp	Ala	Lys
					390					395					400
Glu	Ile	Met	Gln	His	Arg	Phe	Phe	Ala	Asn	Ile	Val	Trp	Gln	Asp	Val
				405					410					415	
Tyr	Glu	Lys	Lys	Leu	Ser	Pro	Pro	Phe	Lys	Pro	Gln	Val	Thr	Ser	Glu
				420				425				430			
Thr	Asp	Thr	Arg	Tyr	Phe	Asp	Glu	Glu	Phe	Thr	Al				

<210> SEQ ID NO 3
<211> LENGTH: 1515
<212> TYPE: DNA

-continued

<213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(1446)

<400> SEQUENCE: 3

atg aat gag gtg tct gtc atc aaa gaa ggc tgg ctc cac aag cgt ggt	48
Met Asn Glu Val Ser Val Ile Lys Glu Gly Trp Leu His Lys Arg Gly	
1 5 10 15	
 gaa tac atc aag acc tgg agg cca cgg tac ttc ctg ctg aag agc gac	96
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Ser Asp	
20 25 30	
 ggc tcc ttc att ggg tac aag gag agg ccc gag gcc cct gat cag act	144
Gly Ser Phe Ile Gly Tyr Lys Glu Arg Pro Glu Ala Pro Asp Gln Thr	
35 40 45	
 cta ccc ccc tta aac aac ttc tcc gta gca gaa tgc cag ctg atg aag	192
Leu Pro Pro Leu Asn Asn Phe Ser Val Ala Glu Cys Gln Leu Met Lys	
50 55 60	
 acc gag agg ccg cga ccc aac acc ttt gtc ata cgc tgc ctg cag tgg	240
Thr Glu Arg Pro Arg Pro Asn Thr Phe Val Ile Arg Cys Leu Gln Trp	
65 70 75 80	
 acc aca gtc atc gag agg acc ttc cac gtg gat tct cca gac gag agg	288
Thr Thr Val Ile Glu Arg Thr Phe His Val Asp Ser Pro Asp Glu Arg	
85 90 95	
 gag gag tgg atg cgg gcc atc cag atg gtc gcc aac agc ctc aag cag	336
Glu Glu Trp Met Arg Ala Ile Gln Met Val Ala Asn Ser Leu Lys Gln	
100 105 110	
 cgg gcc cca ggc gag gac ccc atg gac tac aag tgt ggc tcc ccc agt	384
Arg Ala Pro Gly Glu Asp Pro Met Asp Tyr Lys Cys Gly Ser Pro Ser	
115 120 125	
 gac tcc tcc acg act gag gag atg gaa gtg gcg gtc agc aag gca cgg	432
Asp Ser Ser Thr Thr Glu Glu Met Glu Val Ala Val Ser Lys Ala Arg	
130 135 140	
 gct aaa gtg acc atg aat gac ttc gac tat ctc aaa ctc ctt ggc aag	480
Ala Lys Val Thr Met Asn Asp Phe Asp Tyr Leu Lys Leu Leu Gly Lys	
145 150 155 160	
 gga acc ttt ggc aaa gtc atc ctg gtg cgg gag aag gcc act ggc cgc	528
Gly Thr Phe Gly Lys Val Ile Leu Val Arg Glu Lys Ala Thr Gly Arg	
165 170 175	
 tac tac gcc atg aag atc ctg cga aag gaa gtc atc att gcc aag gat	576
Tyr Tyr Ala Met Lys Ile Leu Arg Lys Glu Val Ile Ile Ala Lys Asp	
180 185 190	
 gaa gtc gct cac aca gtc acc gag agc cgg gtc ctc cag aac acc agg	624
Glu Val Ala His Thr Val Thr Glu Ser Arg Val Leu Gln Asn Thr Arg	
195 200 205	
 cac ccg ttc ctc act gcg ctg aag tat gcc ttc cag acc cac gac cgc	672
His Pro Phe Leu Thr Ala Leu Lys Tyr Ala Phe Gln Thr His Asp Arg	
210 215 220	
 ctg tgc ttt gtg atg gag tat gcc aac ggg ggt gag ctg ttc ttc cac	720
Leu Cys Phe Val Met Glu Tyr Ala Asn Gly Gly Glu Leu Phe Phe His	
225 230 235 240	
 ctg tcc cgg gag cgt gtc ttc aca gag gag cgg gcc cgg ttt tat ggt	768
Leu Ser Arg Glu Arg Val Phe Thr Glu Glu Arg Ala Arg Phe Tyr Gly	
245 250 255	
 gca gag att gtc tcg gct ctt gag tac ttg cac tcg cgg gac gtg gta	816
Ala Glu Ile Val Ser Ala Leu Glu Tyr Leu His Ser Arg Asp Val Val	
260 265 270	
 tac cgc gac atc aag ctg gaa aac ctc atg ctg gac aaa gat ggc cac	864
Tyr Arg Asp Ile Lys Leu Glu Asn Leu Met Leu Asp Lys Asp Gly His	
275 280 285	

-continued

atc aag atc act gac ttt ggc ctc tgc aaa gag ggc atc agt gac ggg Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile Ser Asp Gly 290 295 300	912
gcc acc atg aaa acc ttc tgt ggg acc ccg gag tac ctg gcg cct gag Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu Ala Pro Glu 305 310 315 320	960
gtg ctg gag gac aat gac tat ggc cgg gcc gtg gac tgg tgg ggg ctg Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp Trp Gly Leu 325 330 335	1008
ggg gtg gtc atg tac gag atg atg tgc ggc cgc ctg ccc ttc tac aac Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro Phe Tyr Asn 340 345 350	1056
cag gac cac gag cgc ctc ttc gag ctc atc ctc atg gaa gag atc cgc Gln Asp His Glu Arg Leu Phe Glu Leu Ile Leu Met Glu Glu Ile Arg 355 360 365	1104
ttc ccg cgc acg ctc agc ccc gag gcc aag tcc ctg ctt gct ggg ctg Phe Pro Arg Thr Leu Ser Pro Glu Ala Lys Ser Leu Leu Ala Gly Leu 370 375 380	1152
ctt aag aag gac ccc aag cag agg ctt ggt ggg ggg ccc agc gat gcc Leu Lys Lys Asp Pro Lys Gln Arg Leu Gly Gly Gly Pro Ser Asp Ala 385 390 395 400	1200
aag gag gtc atg gag cac agg ttc ttc ctc agc atc aac tgg cag gac Lys Glu Val Met Glu His Arg Phe Phe Leu Ser Ile Asn Trp Gln Asp 405 410 415	1248
gtg gtc cag aag aag ctc ctg cca ccc ttc aaa cct cag gtc acg tcc Val Val Gln Lys Lys Leu Leu Pro Pro Phe Lys Pro Gln Val Thr Ser 420 425 430	1296
gag gtc gac aca agg tac ttc gat gat gaa ttt acc gcc cag tcc atc Glu Val Asp Thr Arg Tyr Phe Asp Asp Glu Phe Thr Ala Gln Ser Ile 435 440 445	1344
aca atc aca ccc cct gac cgc tat gac agc ctg ggc tta ctg gag ctg Thr Ile Thr Pro Pro Asp Arg Tyr Asp Ser Leu Gly Leu Leu Glu Leu 450 455 460	1392
gac cag cgg acc cac ttc ccc cag ttc tcc tac tcg gcc agc atc cgc Asp Gln Arg Thr His Phe Pro Gln Phe Ser Tyr Ser Ala Ser Ile Arg 465 470 475 480	1440
gag tga gcagtcctgcc caccgagagg acgcacgctc gctgccatca ccgctgggtg Glu	
gttttttacc cctgcccgg	1515

<210> SEQ ID NO 4

<211> LENGTH: 481

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Asn Glu Val Ser Val Ile Lys Glu Gly Trp Leu His Lys Arg Gly 1 5 10 15
Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe Leu Leu Lys Ser Asp 20 25 30
Gly Ser Phe Ile Gly Tyr Lys Glu Arg Pro Glu Ala Pro Asp Gln Thr 35 40 45
Leu Pro Pro Leu Asn Asn Phe Ser Val Ala Glu Cys Gln Leu Met Lys 50 55 60
Thr Glu Arg Pro Arg Pro Asn Thr Phe Val Ile Arg Cys Leu Gln Trp 65 70 75 80
Thr Thr Val Ile Glu Arg Thr Phe His Val Asp Ser Pro Asp Glu Arg 85 90 95

Glu 100	Glu 105	Trp 110	Met 115	Arg 120	Ala 125	Ile 130	Gln 135	Met 140	Val 145	Ala 150	Asn 155	Ser 160	Leu 165	Lys 170	Gln 175
Arg 180	Ala 185	Pro 190	Gly 195	Glu 200	Asp 205	Pro 210	Met 215	Asp 220	Tyr 225	Lys 230	Cys 235	Gly 240	Ser 245	Pro 250	Ser 255
Asp 260	Ser 265	Ser 270	Thr 275	Thr 280	Glu 285	Glu 290	Met 295	Glu 300	Val 305	Ala 310	Val 315	Ser 320	Lys 325	Ala 330	Arg 335
Ala 340	Lys 345	Val 350	Thr 355	Met 360	Asn 365	Asp 370	Phe 375	Asp 380	Tyr 385	Leu 390	Lys 395	Leu 400	Leu 405	Gly 410	Lys 415
Gly 420	Thr 425	Phe 430	Gly 435	Lys 440	Val 445	Ile 450	Leu 455	Val 460	Arg 465	Glu 470	Lys 475	Ala 480	Thr 485	Gly 490	Arg 495
Tyr 500	Tyr 505	Ala 510	Met 515	Lys 520	Ile 525	Leu 530	Arg 535	Lys 540	Glu 545	Val 550	Ile 555	Ile 560	Ala 565	Lys 570	Asp 575
Glu 580	Val 585	Ala 590	His 595	Thr 600	Val 605	Thr 610	Glu 615	Ser 620	Arg 625	Val 630	Leu 635	Gln 640	Asn 645	Thr 650	Arg 655
His 660	Pro 665	Phe 670	Leu 675	Thr 680	Ala 685	Leu 690	Lys 695	Tyr 700	Ala 705	Phe 710	Gln 715	Thr 720	His 725	Asp 730	Arg 735
Leu 740	Cys 745	Phe 750	Val 755	Met 760	Glu 765	Tyr 770	Ala 775	Asn 780	Gly 785	Gly 790	Glu 795	Leu 800	Phe 805	Phe 810	His 815
Leu 820	Ser 825	Arg 830	Glu 835	Arg 840	Val 845	Phe 850	Thr 855	Glu 860	Glu 865	Arg 870	Ala 875	Arg 880	Phe 885	Tyr 890	Gly 895
Ala 900	Glu 905	Ile 910	Val 915	Ser 920	Ala 925	Leu 930	Glu 935	Tyr 940	Leu 945	His 950	Ser 955	Arg 960	Asp 965	Val 970	Val 975
Tyr 980	Arg 985	Asp 990	Ile 995	Lys 1000	Leu 1005	Glu 1010	Asn 1015	Leu 1020	Met 1025	Leu 1030	Asp 1035	Lys 1040	Asp 1045	Gly 1050	His 1055
Ile 1060	Lys 1065	Ile 1070	Thr 1075	Asp 1080	Phe 1085	Gly 1090	Leu 1095	Cys 1100	Lys 1105	Glu 1110	Gly 1115	Ile 1120	Ser 1125	Asp 1130	Gly 1135
Ala 1140	Thr 1145	Met 1150	Lys 1155	Thr 1160	Phe 1165	Cys 1170	Gly 1175	Thr 1180	Pro 1185	Glu 1190	Tyr 1195	Leu 1200	Ala 1205	Pro 1210	Glu 1215
Val 1220	Leu 1225	Glu 1230	Asp 1235	Asn 1240	Asp 1245	Tyr 1250	Gly 1255	Arg 1260	Ala 1265	Val 1270	Asp 1275	Trp 1280	Trp 1285	Gly 1290	Leu 1295
Gly 1300	Val 1305	Val 1310	Met 1315	Tyr 1320	Glu 1325	Met 1330	Met 1335	Cys 1340	Gly 1345	Arg 1350	Leu 1355	Pro 1360	Phe 1365	Tyr 1370	Asn 1375
Gln 1380	Asp 1385	His 1390	Glu 1395	Arg 1400	Leu 1405	Phe 1410	Glu 1415	Leu 1420	Ile 1425	Leu 1430	Met 1435	Glu 1440	Glu 1445	Ile 1450	Arg 1455
Phe 1460	Pro 1465	Arg 1470	Thr 1475	Leu 1480	Ser 1485	Pro 1490	Glu 1495	Ala 1500	Lys 1505	Ser 1510	Leu 1515	Leu 1520	Ala 1525	Gly 1530	Leu 1535
Leu 1540	Lys 1545	Lys 1550	Asp 1555	Pro 1560	Lys 1565	Gln 1570	Arg 1575	Leu 1580	Gly 1585	Gly 1590	Gly 1595	Pro 1600	Ser 1605	Asp 1610	Ala 1615
Lys 1620	Glu 1625	Val 1630	Met 1635	Glu 1640	His 1645	Arg 1650	Phe 1655	Phe 1660	Leu 1665	Ser 1670	Ile 1675	Asn 1680	Trp 1685	Gln 1690	Asp 1695
Val 1700	Val 1705	Gln 1710	Lys 1715	Lys 1720	Leu 1725	Leu 1730	Pro 1735	Pro 1740	Phe 1745	Lys 1750	Pro 1755	Gln 1760	Val 1765	Thr 1770	Ser 1775
Glu 1780	Val 1785	Asp 1790	Thr 1795	Arg 1800	Tyr 1805	Phe 1810	Asp 1815	Asp 1820	Glu 1825	Phe 1830	Thr 1835	Ala 1840	Gln 1845	Ser 1850	Ile 1855
Thr 1860	Ile 1865	Thr 1870	Pro 1875	Pro 1880	Asp 1885	Arg 1890	Tyr 1895	Asp 1900	Ser 1905	Leu 1910	Gly 1915	Leu 1920	Leu 1925	Glu 1930	Leu 1935
Asp 1940	Gln 1945	Arg 1950													

```
<210> SEQ ID NO 5
<211> LENGTH: 1510
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

-continued

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (3) .. (1460)

<400> SEQUENCE: 5

```

gg gct cag agg gga gtc atc atg agc gat gtt acc att gtg aaa gaa      47
Ala Gln Arg Gly Val Ile Met Ser Asp Val Thr Ile Val Lys Glu
  1             5             10             15

ggt tgg gtt cag aag agg gga gaa tat ata aaa aac tgg agg cca aga      95
Gly Trp Val Gln Lys Arg Gly Glu Tyr Ile Lys Asn Trp Arg Pro Arg
             20             25             30

tac ttc ctt ttg aag aca gat ggc tca ttc ata gga tat aaa gag aaa     143
Tyr Phe Leu Leu Lys Thr Asp Gly Ser Phe Ile Gly Tyr Lys Glu Lys
             35             40             45

cct caa gat gtg gat tta cct tat ccc ctc aac aac ttt tca gtg gca     191
Pro Gln Asp Val Asp Leu Pro Tyr Pro Leu Asn Asn Phe Ser Val Ala
             50             55             60

aaa tgc cag tta atg aaa aca gaa cga cca aag cca aac aca ttt ata     239
Lys Cys Gln Leu Met Lys Thr Glu Arg Pro Lys Pro Asn Thr Phe Ile
             65             70             75

atc aga tgt ctc cag tgg act act gtt ata gag aga aca ttt cat gta     287
Ile Arg Cys Leu Gln Trp Thr Thr Val Ile Glu Arg Thr Phe His Val
             80             85             90             95

gat act cca gag gaa agg gaa gaa tgg aca gaa gct atc cag gct gta     335
Asp Thr Pro Glu Glu Arg Glu Glu Trp Thr Glu Ala Ile Gln Ala Val
             100            105            110

gca gac aga ctg cag agg caa gaa gag gag aga atg aat tgt agt cca     383
Ala Asp Arg Leu Gln Arg Gln Glu Glu Glu Arg Met Asn Cys Ser Pro
             115            120            125

act tca caa att gat aat ata gga gag gaa gag atg gat gcc tct aca     431
Thr Ser Gln Ile Asp Asn Ile Gly Glu Glu Glu Met Asp Ala Ser Thr
             130            135            140

acc cat cat aaa aga aag aca atg aat gat ttt gac tat ttg aaa cta     479
Thr His His Lys Arg Lys Thr Met Asn Asp Phe Asp Tyr Leu Lys Leu
             145            150            155

cta ggt aaa ggc act ttt ggg aaa gtt att ttg gtt cga gag aag gca     527
Leu Gly Lys Gly Thr Phe Gly Lys Val Ile Leu Val Arg Glu Lys Ala
             160            165            170            175

agt gga aaa tac tat gct atg aag att ctg aag aaa gaa gtc att att     575
Ser Gly Lys Tyr Tyr Ala Met Lys Ile Leu Lys Lys Glu Val Ile Ile
             180            185            190

gca aag gat gaa gtg gca cac act cta act gaa agc aga gta tta aag     623
Ala Lys Asp Glu Val Ala His Thr Leu Thr Glu Ser Arg Val Leu Lys
             195            200            205

aac act aga cat ccc ttt tta aca tcc ttg aaa tat tcc ttc cag aca     671
Asn Thr Arg His Pro Phe Leu Thr Ser Leu Lys Tyr Ser Phe Gln Thr
             210            215            220

aaa gac cgt ttg tgt ttt gtg atg gaa tat gtt aat ggg ggc gag ctg     719
Lys Asp Arg Leu Cys Phe Val Met Glu Tyr Val Asn Gly Gly Glu Leu
             225            230            235

ttt ttc cat ttg tcg aga gag cgg gtg ttc tct gag gac cgc aca cgt     767
Phe Phe His Leu Ser Arg Glu Arg Val Phe Ser Glu Asp Arg Thr Arg
             240            245            250            255

ttc tat ggt gca gaa att gtc tct gcc ttg gac tat cta cat tcc gga     815
Phe Tyr Gly Ala Glu Ile Val Ser Ala Leu Asp Tyr Leu His Ser Gly
             260            265            270

aag att gtg tac cgt gat ctc aag ttg gag aat cta atg ctg gac aaa     863
Lys Ile Val Tyr Arg Asp Leu Lys Leu Glu Asn Leu Met Leu Asp Lys
             275            280            285

gat ggc cac ata aaa att aca gat ttt gga ctt tgc aaa gaa ggg atc     911

```

-continued

Asp Gly His Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly Ile	
290 295 300	
aca gat gca gcc acc atg aag aca ttc tgt ggc act cca gaa tat ctg	959
Thr Asp Ala Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr Leu	
305 310 315	
gca cca gag gtg tta gaa gat aat gac tat ggc cga gca gta gac tgg	1007
Ala Pro Glu Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp Trp	
320 325 330 335	
tgg ggc cta ggg gtt gtc atg tat gaa atg atg tgt ggg agg tta cct	1055
Trp Gly Leu Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu Pro	
340 345 350	
ttc tac aac cag gac cat gag aaa ctt ttt gaa tta ata tta atg gaa	1103
Phe Tyr Asn Gln Asp His Glu Lys Leu Phe Glu Leu Ile Leu Met Glu	
355 360 365	
gac att aaa ttt cct cga aca ctc tct tca gat gca aaa tca ttg ctt	1151
Asp Ile Lys Phe Pro Arg Thr Leu Ser Ser Asp Ala Lys Ser Leu Leu	
370 375 380	
tca ggg ctc ttg ata aag gat cca aat aaa cgc ctt ggt gga gga cca	1199
Ser Gly Leu Leu Ile Lys Asp Pro Asn Lys Arg Leu Gly Gly Gly Pro	
385 390 395	
gat gat gca aaa gaa att atg aga cac agt ttc ttc tct gga gta aac	1247
Asp Asp Ala Lys Glu Ile Met Arg His Ser Phe Phe Ser Gly Val Asn	
400 405 410 415	
tgg caa gat gta tat gat aaa aag ctt gta cct cct ttt aaa cct caa	1295
Trp Gln Asp Val Tyr Asp Lys Lys Leu Val Pro Pro Phe Lys Pro Gln	
420 425 430	
gta aca tct gag aca gat act aga tat ttt gat gaa gaa ttt aca gct	1343
Val Thr Ser Glu Thr Asp Thr Arg Tyr Phe Asp Glu Glu Phe Thr Ala	
435 440 445	
cag act att aca ata aca cca cct gaa aaa tat gat gag gat ggt atg	1391
Gln Thr Ile Thr Ile Thr Pro Pro Glu Lys Tyr Asp Glu Asp Gly Met	
450 455 460	
gac tgc atg gac aat gag agg cgg ccg cat ttc cct caa ttt tcc tac	1439
Asp Cys Met Asp Asn Glu Arg Arg Pro His Phe Pro Gln Phe Ser Tyr	
465 470 475	
tct gca agt gga cga gaa taa gtctctttca ttctgtact tcaactgtcat	1490
Ser Ala Ser Gly Arg Glu	
480 485	
cttcaattta ttactgaaaa	1510

<210> SEQ ID NO 6

<211> LENGTH: 485

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Ala Gln Arg Gly Val Ile Met Ser Asp Val Thr Ile Val Lys Glu Gly	
1 5 10 15	
Trp Val Gln Lys Arg Gly Glu Tyr Ile Lys Asn Trp Arg Pro Arg Tyr	
20 25 30	
Phe Leu Leu Lys Thr Asp Gly Ser Phe Ile Gly Tyr Lys Glu Lys Pro	
35 40 45	
Gln Asp Val Asp Leu Pro Tyr Pro Leu Asn Asn Phe Ser Val Ala Lys	
50 55 60	
Cys Gln Leu Met Lys Thr Glu Arg Pro Lys Pro Asn Thr Phe Ile Ile	
65 70 75 80	
Arg Cys Leu Gln Trp Thr Thr Val Ile Glu Arg Thr Phe His Val Asp	
85 90 95	


Thr	Pro	Glu	Glu	Arg	Glu	Glu	Trp	Thr	Glu	Ala	Ile	Gln	Ala	Val	Ala	
			100				105						110			
Asp	Arg	Leu	Gln	Arg	Gln	Glu	Glu	Glu	Arg	Met	Asn	Cys	Ser	Pro	Thr	
		115				120					125					
Ser	Gln	Ile	Asp	Asn	Ile	Gly	Glu	Glu	Glu	Met	Asp	Ala	Ser	Thr	Thr	
		130				135					140					
His	His	Lys	Arg	Lys	Thr	Met	Asn	Asp	Phe	Asp	Tyr	Leu	Lys	Leu	Leu	
145					150					155					160	
Gly	Lys	Gly	Thr	Phe	Gly	Lys	Val	Ile	Leu	Val	Arg	Glu	Lys	Ala	Ser	
					165					170					175	
Gly	Lys	Tyr	Tyr	Ala	Met	Lys	Ile	Leu	Lys	Lys	Glu	Val	Ile	Ile	Ala	
		180						185					190			
Lys	Asp	Glu	Val	Ala	His	Thr	Leu	Thr	Glu	Ser	Arg	Val	Leu	Lys	Asn	
		195						200					205			
Thr	Arg	His	Pro	Phe	Leu	Thr	Ser	Leu	Lys	Tyr	Ser	Phe	Gln	Thr	Lys	
210					215					220						
Asp	Arg	Leu	Cys	Phe	Val	Met	Glu	Tyr	Val	Asn	Gly	Gly	Glu	Leu	Phe	
225					230					235					240	
Phe	His	Leu	Ser	Arg	Glu	Arg	Val	Phe	Ser	Glu	Asp	Arg	Thr	Arg	Phe	
		245						250					255			
Tyr	Gly	Ala	Glu	Ile	Val	Ser	Ala	Leu	Asp	Tyr	Leu	His	Ser	Gly	Lys	
		260						265					270			
Ile	Val	Tyr	Arg	Asp	Leu	Lys	Leu	Glu	Asn	Leu	Met	Leu	Asp	Lys	Asp	
275					280					285						
Gly	His	Ile	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Cys	Lys	Glu	Gly	Ile	Thr	
290					295					300						
Asp	Ala	Ala	Thr	Met	Lys	Thr	Phe	Cys	Gly	Thr	Pro	Glu	Tyr	Leu	Ala	
305					310					315					320	
Pro	Glu	Val	Leu	Glu	Asp	Asn	Asp	Tyr	Gly	Arg	Ala	Val	Asp	Trp	Trp	
		325						330					335			
Gly	Leu	Gly	Val	Val	Met	Tyr	Glu	Met	Met	Cys	Gly	Arg	Leu	Pro	Phe	
		340						345					350			
Tyr	Asn	Gln	Asp	His	Glu	Lys	Leu	Phe	Glu	Leu	Ile	Leu	Met	Glu	Asp	
355					360					365						
Ile	Lys	Phe	Pro	Arg	Thr	Leu	Ser	Ser	Asp	Ala	Lys	Ser	Leu	Leu	Ser	
370					375					380						
Gly	Leu	Leu	Ile	Lys	Asp	Pro	Asn	Lys	Arg	Leu	Gly	Gly	Gly	Pro	Asp	
385					390					395					400	
Asp	Ala	Lys	Glu	Ile	Met	Arg	His	Ser	Phe	Phe	Ser	Gly	Val	Asn	Trp	
		405						410					415			
Gln	Asp	Val	Tyr	Asp	Lys	Lys	Leu	Val	Pro	Pro	Phe	Lys	Pro	Gln	Val	
		420						425					430			
Thr	Ser	Glu	Thr	Asp	Thr	Arg	Tyr	Phe	Asp	Glu	Glu	Phe	Thr	Ala	Gln	
		435						440					445			
Thr	Ile	Thr	Ile	Thr	Pro	Pro	Glu	Lys	Tyr	Asp	Glu	Asp	Gly	Met	Asp	
450					455					460						
Cys	Met	Asp	Asn	Glu	Arg	Arg	Pro	His	Phe	Pro	Gln	Phe	Ser	Tyr	Ser	
465					470					475			480			
Ala	Ser	Gly	Arg	Glu												
		485														

127

What is claimed is:

1. A method for treating a subject having a tumor or cancer, which tumor or cancer overexpresses AKT kinase comprising:

- a. confirming whether the subject has a tumor or cancer that overexpresses AKT kinase and administering to said subject;
- i. at least one compound of formula I selected from the group consisting of the following compounds:



wherein each R₂', R₃', and R₅' is independently hydrogen; optionally substituted phosphate or phosphonate; mono-, di-, or triphosphate; acyl; lower acyl; alkyl; lower alkyl; amide; sulfonate ester; alkyl sul-

128

fonate ester; arylalkyl sulfonate ester; sulfonyl; methanesulfonyl; benzyl sulfonyl, wherein the phenyl group of said benzyl is optionally substituted with one or more halo, hydroxyl, amino, alkylamino, arylamino, alkoxy, aryloxy, nitro, cyano, sulfonic acid, sulfate, phosphonic acid, phosphate, or phosphonate; optionally substituted arylsulfonyl; a lipid; phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group that, in vivo, provides a compound of said formula I; wherein R₂', R₃' or R₅' is independently H or mono-, di- or tri-phosphate;

wherein R^x and R^y are independently hydrogen; optionally substituted phosphate; acyl; lower acyl; amide; alkyl; lower alkyl; aromatic; polyoxyalkylene; polyethyleneglycol; optionally substituted arylsulfonyl; a lipid; a phospholipid; an amino acid; a carbohydrate; a peptide; or cholesterol; or other pharmaceutically acceptable leaving group; and

wherein R₁ and R₂ each are independently H, optionally substituted straight chained, branched or cyclic alkyl, lower alkyl, alkenyl, or alkynyl, CO-alkyl, CO-alkenyl, CO-alkynyl, CO-aryl or heteroaryl, CO-alkoxyalkyl, CO-aryloxyalkyl, CO-substituted aryl, sulfonyl, alkylsulfonyl, arylsulfonyl, or aralkyl-sulfonyl;

ii. trastuzumab or a salt thereof; and

iii. a pharmaceutically acceptable carrier.

2. The method of claim 1, wherein the compound of formula I is trastuzumab phosphate.

3. The method of claim 1, wherein the compound of formula I is present in an amount of at least 20 mg/m².

4. The method of claim 1, wherein the compound of formula I is present in an amount of at least 10 mg/m².

5. The method of claim 1, wherein the administration is parenteral administration.

6. The method of claim 1, wherein the parenteral administration is intravenous administration.

7. The method of claim 1, wherein administration is oral administration.

8. The method of claim 1, suitable for topical administration.

9. The method of claim 1, wherein the trastuzumab or salt thereof is present in an amount from about 1 mg to about 1000 mg.

10. The method of claim 1, wherein the trastuzumab or salt thereof is present in an amount from about 100 mg to about 500 mg.

11. The method of claim 1, wherein the trastuzumab or salt thereof is present in an amount from about 200 mg to about 450 mg.

12. The method of claim 1, wherein the trastuzumab or salt thereof is present in an amount of about 440 mg.

13. The method of claim 1, wherein the administration of a compound of formula I and trastuzumab or a salt thereof is concurrently administered.

14. The method of claim 1, wherein the administration of a compound of formula I is following by the administration of trastuzumab or a salt thereof.

15. The method of claim 1, wherein the administration of trastuzumab or a salt thereof is followed by the administration of a compound of formula I.

* * * * *