

1993

## Wing-length Change in the First Postnuptial Molt of Gambel's White-crowned Sparrows

Carl D. Barrentine

Max W. Lincoln

Charles E. Corchran

Phillip M. Walters

Follow this and additional works at: <https://digitalcommons.usf.edu/nabb>

---

### Recommended Citation

Barrentine, Carl D.; Lincoln, Max W.; Corchran, Charles E.; and Walters, Phillip M. (1993) "Wing-length Change in the First Postnuptial Molt of Gambel's White-crowned Sparrows," *North American Bird Bander*. Vol. 18 : Iss. 4 , Article 3.

Available at: <https://digitalcommons.usf.edu/nabb/vol18/iss4/3>

This Contents is brought to you for free and open access by the Searchable Ornithological Research Archive at Digital Commons @ University of South Florida. It has been accepted for inclusion in North American Bird Bander by an authorized editor of Digital Commons @ University of South Florida. For more information, please contact [digitalcommons@usf.edu](mailto:digitalcommons@usf.edu).

# Wing-length Change in the First Postnuptial Molt of Gambel's White-crowned Sparrows

**Carl D. Barrentine**  
Integrated Studies/Biology  
Campus Box 7117  
University of North Dakota  
Grand Forks, ND 58202-7117

**Max W. Lincoln**  
Coyote Creek Riparian Station  
P.O. Box 1027  
Alviso, CA 95002

**Charles E. Corchran**  
7214 E. Camino Vecino  
Tucson, AZ 85715

**Phillip M. Walters**  
2001 W. Rudasill Road  
Tucson, AZ 85705

## INTRODUCTION

As an age group, adult Gambel's White-crowned Sparrows (*Zonotrichia leucophrys gambelii*) have longer wings than immatures (Fugle and Rothstein 1985, Mewaldt and King 1986). This wing-length difference is a consequence of the first postnuptial molt, which occurs on the breeding grounds and precedes the second fall migration (Chilgren 1978, Morton *et al.* 1969). Mewaldt (1973) described first postnuptial wing-length change in inclusive age samples of White-crowned Sparrows (*Z.l. pugetensis* and *Z.l. gambelii*). Changes in wing-length associated with this molt have been noted in other *Zonotrichia* (Piper and Wiley 1991), and are well-documented in other passerine species (Alatalo *et al.* 1984, Stewart 1963).

In this study, we describe first postnuptial wing-length change in samples of Gambel's White-crowned Sparrows from California and Arizona. We use measurements obtained from inclusive age samples of sparrows (*i.e.*, paired immature and adult wing-chord measurements from the *same* birds) to compare wing-length change at three widely-separated geographic locations.

## METHODS

We measured wing-length (unflattened chord, recorded to the nearest 1.0 mm using a steel rule with bend-of-wing stop at zero) in immature (HY, SY) and adult (AHY, ASY) White-crowned Sparrows. Sparrows were captured and banded at

three locations: Alviso, California (37° 26'N, 121° 56'W), Bakersfield, California (35° 21'N, 119° 09'W), and Tucson, Arizona (32° 10'N, 110° 40'W). Site descriptions and sampling methods are published elsewhere (Barrentine *et al.* 1990).

Measurements reported in this study were obtained from banded immature sparrows that were recaptured as returning adults in a subsequent winter season, 1985-90. That is, we reference paired immature and adult wing-length measurements from the *same* birds. All measurements were made at the time of first capture and first recapture for immatures and returning adults, respectively.

We compare mean wing-length change for samples at each location using two-tailed, paired-sample *t* tests (reject  $H_0: \mu_a - \mu_i = 0$  if  $|t| > t_{0.05(2),v}$ ). The Kruskal-Wallis test is used to test the  $H_0$ : wing-length change is the same for the three sample locations (reject  $H_0: \mu_1 = \mu_2 = \mu_3$  if  $|H_c| > \chi^2_{0.05,2}$ ). Statistical procedures used in this study are described by Zar (1984).

## RESULTS AND DISCUSSION

Paired immature and adult wing-length measurements were obtained from a total of 507 White-crowned Sparrows (Table 1). Mean wing-length increase in the first postnuptial molt was statistically significant ( $P < 0.001$ ) for samples at each location (Table 2), and this increase was similar for all three samples ( $H_c = 2.29$ ,  $df = 2$ ,  $P > 0.25$ ,

Table 1.	Wing-length frequency distributions for inclusive age samples (immature and adult) of White-crowned Sparrows at Alviso, CA; Bakersfield, CA; and Tucson, AZ.							
	LOCATION							
	Alviso, California		Bakersfield, Calif.		Tucson, Arizona		Combined Locs	
Wing length (mm)	Immat.	Adult	Immat.	Adult	Immat.	Adult	Immat.	Adult
	N	N	N	N	N	N	N	N
82		2				1		3
81		0				4		4
80	1	3		5	2	15	3	23
79	3	10	2	9	6	28	11	47
78	6	9	9	25	22	28	37	62
77	8	17	17	18	21	22	46	57
76	24	21	22	18	32	26	78	65
75	17	17	22	12	33	27	72	56
74	19	14	15	13	23	31	57	58
73	13	10	15	14	41	28	69	52
72	9	13	12	7	35	25	56	45
71	9	9	7	5	18	9	34	23
70	13	5	6	1	13	2	32	8
69	6	1	1	1	1	1	8	3
68	4	1					4	1
x (mm)	73.77	75.03	74.58	75.76	74.28	75.64	74.22	75.51
SD (mm)	2.79	2.80	2.28	2.48	2.45	2.76	2.52	2.72
N	132		128		247		507	

Kruskal-Wallis test, corrected for tied ranks). Overall, mean wing-length increased by 1.3 mm (or 1.7 %) in the first postnuptial molt.

Our observations of wing-length change compare favorably with the findings of three other studies. Mewaldt (1973) found mean wing-length increases of 1.1 to 1.5 mm (changes ranged from -2 mm to +5 mm) in inclusive age samples of *Z.l. gambelii* and *Z.l. pugetensis* from San Jose, California. Fugle and Rothstein (1985) found that the mean wing-length of adults was 1.4 mm (or 1.9 %) longer than immatures in mutually exclusive age samples of *Z.l. gambelii* from Santa Barbara, California. Finally, Mewaldt and King (1986) found that the mean wing-length of adult *Z.l. gambelii* was 1.2

and 1.4 mm (or 1.6 and 1.9%) longer than immatures in mutually exclusive age samples from Tucson, Arizona, and San Jose, California, respectively.

## SUMMARY

We document wing-length change in the first postnuptial molt of 507 Gambel's White-crowned Sparrows wintering at Alviso and Bakersfield, California, and Tucson, Arizona. We use paired immature and adult wing-length measurements from the *same* birds to show that (1) mean wing-length increases significantly (1.7%) in the first postnuptial molt, and that (2) this increase in mean wing-length is similar for birds sampled at the three widely-sepa-

rated geographic locations. This study corroborates and extends earlier observations of wing-length change for this subspecies.

## ACKNOWLEDGMENTS

We thank the 60+ volunteers at the Coyote Creek Riparian Station, Alviso, California. Shelby J. Barrentine assisted at the Environmental Studies Area, California State University, Bakersfield, California. Timothy Baird, Ross Chapin, Kristin Coleman, Robert Cote, Carol de Waard, Donald Lamm, Janice Luepke, and Jean Scheibe assisted at the Tanque Verde Ranch, Tucson, Arizona. We also thank John B. Dunning, Martin L. Morton, Kerry Paul Reese, Michael Rigney, and Robert C. Tweit for their helpful comments on earlier versions of this manuscript.

<b>Table 2.</b> Frequency distribution of wing-length change in first postnuptial molt for inclusive age samples of White-crowned Sparrows at Alviso, CA; Bakersfield, CA; and Tucson, AZ.				
	<b>LOCATION</b>			
<b>Wing-length change (mm)</b>	<b>Alviso, CA</b>	<b>Bakersfield, CA</b>	<b>Tucson, AZ</b>	<b>Comb. Loc.</b>
	<b>N</b>	<b>N</b>	<b>N</b>	<b>N</b>
+5	3		3	6
+4	11	3	12	26
+3	17	10	34	61
+2	22	33	71	126
+1	34	49	59	142
0	27	27	45	99
-1	14	6	17	37
-2	4		6	10
$\bar{x}$ (mm)	+1.26	+1.18	+1.36	+1.29
SD(mm)	1.63	1.08	1.42	1.40
N	132	128	247	507
t	8.87	12.41	15.08	20.81

(P) (P<0.001) (P<0.001) (P<0.001) (P<0.001)

## LITERATURE CITED

- Alatalo, R.V., L. Gustafsson and A. Lundberg. 1983. Why do passerine birds have shorter wings than older birds? *Ibis* 126:410-415.
- Barrentine, C.D., M.W. Lincoln, L.R. Mewaldt, C.E. Corchran and P.M. Walters. 1990. Comparative age and sex ratios in Gambel's White-crowned Sparrows in relation to year and latitude *N. Am. Bird Bander* 15:57-60.
- Chilgren, J.D. 1978. Effects of photoperiod and temperature on postnuptial molt in captive White-crowned Sparrows. *Condor* 80:222-229.
- Fugle, G.N. and S.I. Rothstein. 1985. Age- and sex-related variation in size and crown plumage brightness in wintering White-crowned Sparrows. *J. Field Ornithol.* 56:356-368.
- Mewaldt, L.R. 1973. Wing-length and age in White-crowned Sparrows. *W. Bird Bander* 48:54-56.
- \_\_\_\_\_ and J.R. King. 1986. Estimation of sex ratio from wing-length in birds when sexes differ in size but not coloration. *J. Field Ornithol.* 57:155-167.
- Morton, M.L., J.R. King, and D.S. Farner. 1969. Postnuptial and postjuvenile molt in White-crowned Sparrows in Alaska. *Condor* 71:376-385.
- Piper, W.H. and R.H. Wiley. 1991. Effects of laparotomies on wintering White-throated Sparrows and the usefulness of wing chord as a criterion for sexing. *J. Field Ornithol.* 62:40-45.
- Stewart, I.F. 1963. Variation of wing length with age. *Bird Study* 10:1-9.
- Zar, J.H. 1984. *Biostatistical analysis*, second edition. Prentice-Hall, Englewood Cliffs, N.J.