

1994

Diet-induced Color Variation in the White-throated Sparrow

Elizabeth W. Brooks

Follow this and additional works at: <https://digitalcommons.usf.edu/nabb>

Recommended Citation

Brooks, Elizabeth W. (1994) "Diet-induced Color Variation in the White-throated Sparrow," *North American Bird Bander*. Vol. 19 : Iss. 1 , Article 2.

Available at: <https://digitalcommons.usf.edu/nabb/vol19/iss1/2>

This Contents is brought to you for free and open access by the Searchable Ornithological Research Archive at Digital Commons @ University of South Florida. It has been accepted for inclusion in North American Bird Bander by an authorized editor of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

Diet-induced Color Variation in the White-throated Sparrow

Elizabeth W. Brooks
1435 Waterwells Road
Alfred Station, NY 14803

In recent years, many banders have observed Cedar Waxwings (*Bombycilla cedrorum*) with orange terminal tail bands. Incidence of this phenomenon was reported by Parkes and Wood (1988) and theories for a diet-based cause were proposed by Hudon and Brush (1989) and Brush (1990). Mulvihill et al. (1992), in an analysis of waxwings banded at Powdermill Nature Reserve since the early 1960's, provided evidence that consumption of the berries of Morrow's Honeysuckle (*Lonicera morrowii*) or Tartarian Honeysuckle (*L. tatarica*) was responsible for orange variant waxwing plumage. They also observed this phenomenon in two other berry-eating species. Six out of 12 adult (AHY) and three out of 16 hatching-year (HY) Yellow-breasted Chats (*Icteria virens*) banded during a five-year period exhibited bright orange incoming or freshly molted breast feathers that contrasted with any yellow unmolted breast feathers. In addition, two adult Kentucky Warblers (*Oporornis formosus*) encountered at Powdermill exhibited orange-yellow coloration on the superciliary, chin, throat and breast.

Since White-throated Sparrows (*Zonotrichia albicollis*) also consume berries in the fall (Bent 1968), including rhodoxanthin-containing honeysuckle fruit (pers. obs.), it should be expected that this species might exhibit the same chemically induced feather pigment aberrations as waxwings. Parkes (pers. comm.) has seen two specimens of xanthochroistic White-throated Sparrows in the collection at the Carnegie Museum of Natural History. CM 154167, a male netted at Elkins, WV, in October 1981, has its entire underparts washed with pale yellow which is intensified on the throat, and its superciliary line is a dirty yellow. A second specimen, CM 111070, collected at the mouth of Partridge Creek, James Bay, Ontario, as a HY female in October 1931, exhibits an odd orange-buff color on its throat, middle of the belly, and undertail coverts.

Seemingly more similar to the waxwing phenomenon were eight White-throated Sparrows showing orange lores. These birds, banded in western New York State, were among over 2700 White-throated Sparrows banded during fall at three locations between 1985-1993. Details of the orange-lore White-throated Sparrow bandings appear in Table 1.

Table 1. Banding data on eight White-throated Sparrows with orange lores.

Band Number	Age/Sex	Date Banded	Banding Location	Bander
1431-41013	HY-U	10-08-90	Alfred, NY	Brooks
1431-40930	AHY-U	10-19-90	Alfred, NY	Brooks
1421-87226	HY-U	10-20-90	Clarkson, NY	Symonds
1431-41964	U-U	10-09-91	Braddock Bay, NY	Brooks
1431-41994	U-U	10-10-91	Braddock Bay, NY	Brooks
1431-41719	HY-U	10-25-91	Braddock Bay, NY	Skelly
1481-09563	HY-U	09-09-93	Braddock Bay, NY	Brooks
1481-64315	AHY-M	10-08-93	Braddock Bay, NY	Brooks

This phenomenon was not observed in any of the 950 White-throated Sparrows banded during spring at the three western New York locations.

DISCUSSION

The eight examples detailed in Table 1 suggest that the same diet-induced feather pigment changes observed in Cedar Waxwing rectrices may occur in the yellow lores of White-throated Sparrows in fall. Banders should be alert for the possibility of this color variant in White-throated Sparrows and in other berry-consuming species they handle. It might be helpful, too, if banders attempt to describe color variations by using the Munsell Soil Color Charts (1990) or Smithe's color guide (1975, 1981).

ACKNOWLEDGMENTS

My appreciation to Sharon Skelly and William Symonds for sharing information on White-throated Sparrows they banded; to Charles R. Smith for useful suggestions on an earlier draft of this paper; and to Kenneth C. Parkes for his helpful response to my questions.

LITERATURE CITED

- Bent, A.C. 1968. Life histories of North American cardinals, grosbeaks, buntings, towhees, finches, sparrows, and allies. U.S. Natl. Mus. Bull. 237, Part 3. Smithsonian Institution Press, Washington, DC.
- Brush, A.H. 1990. A possible source for the rhodoxanthin in some Cedar Waxwing tails. *J. Field Ornithol.* 61:355.
- Hudon, J. and A.H. Brush. 1989. Probable dietary basis of a color variant of the Cedar Waxwing. *J. Field Ornithol.* 60:361-368.
- Mulvihill, R.S., K.C. Parkes, R.C. Leberman, and D.S. Wood. 1992. Evidence supporting a dietary basis for orange-tipped rectrices in the Cedar Waxwing. *J. Field Ornithol.* 63:212-216.
- Munsell Soil Color Charts. 1990. Macbeth Division of Kollmorgen Instruments Corp., Baltimore, MD.
- Parkes, K.C. and D.S. Wood. 1988. *Orange-tipped rectrices in the Cedar Waxwing: a preliminary report*. Abstracts of presented papers, 100th stated meeting of the Wilson Ornithological Society, No. 92.
- Smithe, F.B. 1975 and 1981. *Naturalist's color guide*. American Mus. of Natural History, NY.

