

---

January 2016

## A New Longevity Record for the Northern Caracara (*Caracara cheriway*) in Florida

Joan L. Morrison

Brian K. Schmidt

Steven M. McGehee

Follow this and additional works at: <https://digitalcommons.usf.edu/ffn>

---

### Recommended Citation

Morrison, Joan L.; Schmidt, Brian K.; and McGehee, Steven M. (2016) "A New Longevity Record for the Northern Caracara (*Caracara cheriway*) in Florida," *Florida Field Naturalist*: Vol. 44 : Iss. 1 , Article 5. Available at: <https://digitalcommons.usf.edu/ffn/vol44/iss1/5>

This Note is brought to you for free and open access by Digital Commons @ University of South Florida. It has been accepted for inclusion in Florida Field Naturalist by an authorized editor of Digital Commons @ University of South Florida. For more information, please contact [digitalcommons@usf.edu](mailto:digitalcommons@usf.edu).

**A NEW LONGEVITY RECORD FOR THE NORTHERN  
CARACARA (*Caracara cheriway*) IN FLORIDA**

JOAN L. MORRISON<sup>1</sup>, BRIAN K. SCHMIDT<sup>2</sup>, AND STEVEN M. MCGEHEE<sup>3</sup>

<sup>1</sup>*Department of Biology, Trinity College, 300 Summit St., Hartford, Connecticut 06106*

*E-mail: joan.morrison@trincoll.edu*

<sup>2</sup>*9016 S. Bay Drive, Haines City, Florida 33844*

<sup>3</sup>*2449 Pocahontas Place, St. Louis, Missouri 63144*

Raptors, at least those with larger body mass, are known to be relatively long-lived, but obtaining longevity records for wild birds is difficult unless individuals are marked or can be tracked for multiple years. Here we report a resighting that establishes a longevity record for the Northern Caracara (*Caracara cheriway*, hereafter caracara). This raptor occurs in Florida as an isolated and resident population in the south-central region (Morrison and Dwyer 2012). In this region, caracaras are most often found nesting on cattle ranches, which typically include large acreages of improved pastures, open grasslands that are intensely managed as forage for cattle (Morrison and Humphrey 2001). Breeding pairs of caracaras exhibit strong fidelity to a nest site and territory, even to a nest tree (Morrison 1999).

During the mid-1990s and again in the mid-2000s, this population was intensively studied, and over 500 caracaras, both adults and nestlings, were marked. Resighting and recovery information over the years has allowed estimation of annual survival for sub-adults (Dwyer et al. 2012) and for adults and juveniles (Morrison 2003). Adult annual survival is relatively high, 0.88 for males and 0.91 for females (Morrison 2003).

On 7 March 1994, JLM and SMM banded an adult breeding caracara at its nest (N 27° 28', W 85° 01') along NW 240th St., in northern Okeechobee County, Florida. This bird was sexed as male, using DNA obtained from a blood sample. JLM and SMM monitored nesting success in this territory, through 2000. During these 7 years, this male remained in this territory and, with its mate, successfully fledged a total of 11 young.

On 15 March 2015, BKS came upon a banded adult caracara feeding on road kill within 200 m of the site where the adult male caracara was banded in 1994. BKS observed the caracara for several minutes and watched it take food to a cabbage palm (*Sabal palmetto*), which presumably contained the caracara's nest. The caracara returned to the road kill within a few minutes, and at this time, BKS was able to obtain photographs of the caracara. On 16 March 2015, BKS submitted a banding report, along with several photographs showing the US Fish and Wildlife Service (USFWS) band, to the USGS Bird Banding Laboratory (BBL). Verification of this caracara's identity was made on 16 March 2015, when the BBL submitted the photographs to JLM, who confirmed the caracara's identity using numbers visible on the USFWS band. Since caracaras attain adult plumage at a minimum of 3 years of age (Nemeth and Morrison 2002), this male was at least 3 years old when it was banded; thus, when photographed in March 2015, it was at least 24 years old.

Previous longevity records for wild caracaras have been reported as 9 years (Klimkiewicz and Fitcher 1989) and 22 years (J. Layne, unpubl. data). No details are available for these records although they presumably are resightings of marked individuals. The lifespan of captive caracaras has been recorded as 15 years (Bent 1938),

21 years (J. Layne, unpubl. data), and 37 years (Newton 1979). For comparison, longevity records for other raptors resident in Florida include Snail Kite (*Rostrhamus sociabilis*), 14 years, and Red-shouldered Hawk (*Buteo lineatus*), 22 years (Lutmerding 2015).

Florida's population of the Northern Caracara is federally and state listed as Threatened. Perhaps as few as 500 nesting pairs may remain in the state, and the population is believed to be in decline because of ongoing loss of its favored pasture- and grassland habitats (Morrison and Humphrey 2001). All available habitat suitable for nesting appears to be occupied, and a large population of non-breeding floaters is continually searching for and evaluating prospective territories and breeding opportunities (Dwyer et al. 2013). The landscape in this male caracara's territory is typical of caracara nesting habitat and includes improved pasture, scattered cabbage palms and live oaks (*Quercus virginiana*), and seasonally flooded wetlands. This landscape has not changed since this caracara was banded in 1994, so it is likely that this individual has remained there and attempted nesting, annually, since it was banded.

Individual birds that live longest typically produce more offspring over their lifetimes, thereby have a greater probability of producing recruits and thus influencing long-term population persistence (Newton 1985, Espie and Oliphant 2000, Herényi et al. 2012). A management strategy focused on retaining suitable nesting habitat will likely benefit Florida's caracara population by enhancing survival and reproductive success of breeding individuals. However, the strong site fidelity and sedentary behavior typical of this species, combined with a long lifespan, suggest that breeding pairs may maintain residence on a territory even if habitat quality is declining, as long as they can forage successfully and attempt to breed there. Once breeding adults are gone from a territory, though, they might not be replaced if the altered landscape is no longer attractive to new potential breeders. In this case, a long lifespan may pose an additional conservation challenge for this species in that a decline in population size may not be noticed until it becomes drastic.

#### ACKNOWLEDGMENTS

We gratefully acknowledge support of personnel at the MacArthur Agro-Ecology Research Center and Archbold Biological Station, Lake Placid, Florida for their continued support of caracara research over the decades and the many field assistants who have helped with observations, nest searching, and tracking activities. Comments from T. Murphy improved the manuscript. This is Contribution No. 166 from the MacArthur Agro-Ecology Research Center of Archbold Biological Station.

#### LITERATURE CITED

- BENT, A. C. 1938. Life Histories of North American Birds of Prey. Part 2. U.S. National Museum Bulletin 170.
- DWYER, J. F., J. D. FRASER, AND J. L. MORRISON. 2012. Within-year survival of non-breeding Crested Caracaras. *Condor* 114:295–301.
- DWYER, J. F., J. D. FRASER, AND J. L. MORRISON. 2013. Range sizes and habitat use of non-breeding Crested Caracaras in Florida. *Journal of Field Ornithology* 84:223–233.
- ESPIE, R. H. M., AND L. W. OLIPHANT. 2000. Age-dependent breeding performance in merlins (*Falco columbarius*). *Ecology* 81:3404–3415.
- HERÉNYI, M., G. HEGYI, L. GARAMSZEGI, R. HARGITAI, G. MICHL, B. ROSIVALL, AND J. TÖRÖK. 2012. Lifetime offspring production in relation to breeding lifespan, attractiveness, and mating status in male collared flycatchers. *Oecologia* 170:935–942.
- KLIMKIEWICZ, M. K., AND A. G. FUTCHER. 1989. Longevity records of North American birds. Supplement 1. *Journal of Field Ornithology* 60:469–494.

- LUTMERDING, J. A. 2015. Longevity records of North American Birds. USGS Bird Banding Laboratory, Patuxent Wildlife Research Center. <[http://www.pwrc.usgs.gov/BBL/longevity/Longevity\\_main.cfm](http://www.pwrc.usgs.gov/BBL/longevity/Longevity_main.cfm)>. Accessed 26 December 2015.
- MORRISON, J. L. 1999. Breeding biology and productivity of Florida's Crested Caracaras. *Condor* 101:505–517.
- MORRISON, J. L. 2003. Age-specific survival of Florida's Crested Caracaras. *Journal of Field Ornithology* 74:321–330.
- MORRISON, J. L. AND S. R. HUMPHREY. 2001. Conservation value of private lands for Crested Caracaras in Florida. *Conservation Biology* 15:675–684.
- MORRISON, J. L., AND J. F. DWYER. 2012. Crested Caracara (*Caracara cheriway*) in The Birds of North America Online (A. Poole, Ed.). Cornell Lab of Ornithology, Ithaca, New York. <<http://bna.birds.cornell.edu/bna/species/249>>, doi:10.2173/bna.249>.
- NEMETH, N. M., AND J. L. MORRISON. 2002. Natal dispersal of the Crested Caracara in Florida. *Journal of Raptor Research* 36:203–206.
- NEWTON, I. 1979. *Population Ecology of Raptors*. Buteo Books, Vermillion, South Dakota.
- NEWTON, I. 1985. Lifetime reproductive output of female sparrowhawks. *Journal of Animal Ecology* 54:241–253.