

2020

Genome Sequence of *Vibrio natriegens* Phage vB_VnaS-AQKL99

Noah Yonas
University of South Florida

Paige Boleman
University of South Florida

Y Nguyen
University of South Florida

Makenzie Kerr
University of South Florida, burrowsm@usf.edu

Kema Malki
University of South Florida

See next page for additional authors

Follow this and additional works at: https://digitalcommons.usf.edu/msc_facpub

 Part of the [Life Sciences Commons](#)

Scholar Commons Citation

Yonas, Noah; Boleman, Paige; Nguyen, Y; Kerr, Makenzie; Malki, Kema; Greco, Anthony M.; and Breitbart, Mya, "Genome Sequence of *Vibrio natriegens* Phage vB_VnaS-AQKL99" (2020). *Marine Science Faculty Publications*. 1359.

https://digitalcommons.usf.edu/msc_facpub/1359

This Article is brought to you for free and open access by the College of Marine Science at Digital Commons @ University of South Florida. It has been accepted for inclusion in Marine Science Faculty Publications by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact scholarcommons@usf.edu.

Authors

Noah Yonas, Paige Boleman, Y Nguyen, Makenzie Kerr, Kema Malki, Anthony M. Greco, and Mya Breitbart



Genome Sequence of *Vibrio natriegens* Phage vB_VnaS-AQKL99

Noah Yonas,^a Paige Boleman,^a Y Nguyen,^a  Makenzie Kerr,^a  Kema Malki,^a Anthony M. Greco,^a  Mya Breitbart^a

^aCollege of Marine Science, University of South Florida, Saint Petersburg, Florida, USA

Noah Yonas and Paige Boleman contributed equally to this work. Author order was determined based on the amount of writing performed.

ABSTRACT *Vibrio natriegens* is a naturally occurring marine bacterium that is emerging as a microbiological model system. Here, we describe Aquatic Killer 99 (AQKL99), a novel phage that infects *Vibrio natriegens* 14048. The genome of the phage is 58,464 bp long, has a GC content of 45.9%, and contains 51 protein-coding genes.

Vibrio spp. are Gram-negative proteobacteria that inhabit saline and estuarine environments; however, several species are human pathogens (1). The characterization of phages infecting *Vibrio* spp. is of broad interest for the fields of ecology, evolution, human health, and biotechnology (2). *Vibrio natriegens* is emerging as a microbiological model system due to its rapid doubling time (<10 min) under standard laboratory conditions and the availability of tools for genetic manipulation (3, 4). Here, we describe the isolation and genome annotation of Aquatic Killer 99 (AQKL99), a novel phage that infects *Vibrio natriegens* 14048.

In June 2019, 3 liters of surface seawater from Bayboro Harbor (Tampa Bay, FL, USA) was prefiltered through a 0.8- μ m filter, and viruses were concentrated using iron chloride flocculation (5). A double-layer plaque assay on *Vibrio natriegens* 14048 on tryptic soy agar with 10 g per L sodium chloride at 25°C led to the isolation of a novel phage, which was subsequently plaque purified three times (6). Phage lysates were adsorbed to a copper grid, negatively stained with 2% (wt/vol) uranyl acetate (7), and viewed on a Hitachi 7100 transmission electron microscope equipped with a Gatan Orius high-resolution digital camera. The morphology of phage AQKL99 is consistent with that of members of the family *Siphoviridae*, with an average capsid diameter of 74 \pm 18 nm and a long, flexible tail averaging 131 \pm 38 nm, based on pixel measurements relative to the scale bar of 40 phage particles (Fig. 1). Phage AQKL99 was unable to produce plaques on 53 *Vibrio* colonies isolated from the same seawater sample on thiosulfate-citrate-bile salts-sucrose (TCBS) agar, suggesting a narrow host range. However, neither the identity nor the uniqueness of these *Vibrio* isolates was determined.

Phage DNA was isolated from a 0.2- μ m-filtered overnight lysate with the Qiagen MinElute virus spin kit and was sent to the Microbial Genome Sequencing (MiGS) Center for library preparation with the Illumina Nextera kit and sequencing on the NextSeq 550 platform. Sequences were processed using FastQC v0.11.5 (8) with default parameters, assembled using SPAdes v3.11.1 (9), and evaluated using QUAST v5.0.1 (10) to reveal a single contig with \sim 2,900 \times coverage. This contig, which was presumed to represent the complete linear genome of phage AQKL99, is 58,464 bp long with a GC content of 45.9%. As determined with Geneious Prime v2020.1.2 (11), phage AQKL99 shares 75% genome-wide pairwise identity with *Vibrio* phage VhaS-tm (GenBank accession number [KX198614](https://www.ncbi.nlm.nih.gov/nuccore/KX198614)), which was isolated from oyster tissue and can prevent *Vibrio harveyi* from causing disease in greenlip abalone (*Haliotis laevis*) (12).

Open reading frames (ORFs) were predicted through Geneious Prime v2020.1.2 using default parameters, which resulted in the identification and annotation of 51

Citation Yonas N, Boleman P, Nguyen Y, Kerr M, Malki K, Greco AM, Breitbart M. 2020. Genome sequence of *Vibrio natriegens* phage vB_VnaS-AQKL99. Microbiol Resour Announc 9:e00967-20. <https://doi.org/10.1128/MRA.00967-20>.

Editor Simon Roux, DOE Joint Genome Institute

Copyright © 2020 Yonas et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mya Breitbart, mya@usf.edu.

Received 17 August 2020

Accepted 9 September 2020

Published 1 October 2020




FIG 1 Transmission electron micrograph of *Vibrio* phage vB_VnaS-AQKL99.

protein-coding genes, mostly transcribed in the reverse direction (11). Amino acid sequence similarity searches were performed against the nonredundant database using BLASTp with default parameters and an E value cutoff value of <0.001 (13). Approximately 75% of the genes are of unknown function; however, the phage AQKL99 genome encodes several putative structural proteins (major capsid protein, minor head protein, portal protein, tail assembly protein, tail length tape measure protein, and large and small terminase subunits), DNA replication proteins (DNA helicase, DNA polymerase, DNA replicative clamp, and primase), and other notable proteins (autolysin *N*-acetylmuramoyl-L-alanine amidase, AAA family ATPase, and multimodular transpeptidase-transglycosylase). Interestingly, the phage AQKL99 genome encodes three proteins putatively involved in deazaguanine DNA modification (QueC, QueD, and QueD), which is a common nucleotide substitution used to protect phage DNA from host restriction enzymes (14).

Data availability. The genome sequence and associated metadata for *Vibrio* phage vB_VnaS-AQKL99 are available in GenBank under the accession number [MT795651.1](https://www.ncbi.nlm.nih.gov/nuclseq/MT795651.1). Raw reads are available in the SRA database under the accession number [SRX9058260](https://www.ncbi.nlm.nih.gov/sra/SRX9058260).

ACKNOWLEDGMENTS

This work was supported by funding to M.B. from the National Science Foundation (award OCE-1722761). N.Y. was part of the 2020 National Summer Undergraduate Research Project (NSURP). K.M. was supported by a National Science Foundation Graduate Research Fellowship (award 3900101301).

REFERENCES

1. Turner JW, Good B, Cole D, Lipp EK. 2009. Plankton composition and environmental factors contribute to *Vibrio* seasonality. *ISME J* 3:1082–1092. <https://doi.org/10.1038/ismej.2009.50>.
2. Comeau A, Chan A, Suttle C. 2006. Genetic richness of vibriophages isolated in a coastal environment. *Environ Microbiol* 8:1164–1176. <https://doi.org/10.1111/j.1462-2920.2006.01006.x>.
3. Weinstock MT, Heseck ED, Wilson CM, Gibson DG. 2016. *Vibrio natriegens* as a fast-growing host for molecular biology. *Nat Methods* 13:849–851. <https://doi.org/10.1038/nmeth.3970>.
4. Dalia TN, Hayes CA, Stolyar S, Marx CJ, McKinlay JB, Dalia AB. 2017. Multiplex genome editing by natural transformation (MuGENT) for synthetic biology in *Vibrio natriegens*. *ACS Synth Biol* 6:1650–1655. <https://doi.org/10.1021/acssynbio.7b00116>.
5. John S, Mendez C, Deng L, Poulos B, Kauffman A, Kern S, Brum J, Polz M, Boyle E, Sullivan M. 2011. A simple and efficient method for concentration of ocean viruses by chemical flocculation. *Environ Microbiol Rep* 3:195–202. <https://doi.org/10.1111/j.1758-2229.2010.00208.x>.
6. Kropinski AM, Mazzocco A, Waddell TE, Lingohr E, Johnson RP. 2009. Enumeration of bacteriophages by double agar overlay plaque assay. *Methods Mol Biol* 501:69–76. https://doi.org/10.1007/978-1-60327-164-6_7.
7. Ackermann H-W. 2009. Basic phage electron microscopy. *Methods Mol Biol* 501:113–126. https://doi.org/10.1007/978-1-60327-164-6_12.
8. Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. Babraham Institute, Cambridge, United Kingdom. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
9. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS,

- Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyskin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
10. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
11. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
12. Wang Y, Barton M, Elliott L, Li X, Abraham S, O'Dea M, Munro J. 2017. Bacteriophage therapy for the control of *Vibrio harveyi* in greenlip abalone (*Haliotis laevis*). *Aquaculture* 473:251–258. <https://doi.org/10.1016/j.aquaculture.2017.01.003>.
13. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
14. Hutinet G, Kot W, Cui L, Hillebrand R, Balamkundu S, Gnanakalai S, Neelakandan R, Carstens AB, Fa Lui C, Tremblay D, Jacobs-Sera D, Sassanfar M, Lee Y-J, Weigele P, Moineau S, Hatfull GF, Dedon PC, Hansen LH, de Crécy-Lagard V. 2019. 7-Deazaguanine modifications protect phage DNA from host restriction systems. *Nat Commun* 10:5442. <https://doi.org/10.1038/s41467-019-13384-y>.