



Available online at scholarcommons.usf.edu/ijis

International Journal of Speleology

Official Journal of Union Internationale de Spéléologie



Ecophysiological responses of two closely related epigean and hypogean *Niphargus* species to hypoxia and increased temperature: Do they differ?

Tatjana Simčič ^{1*} and Boris Sket ²

¹ Department of Organisms and Ecosystems Research, National Institute of Biology, Večna pot 111, SI-1000 Ljubljana, Slovenia

² Biotechnical Faculty, University of Ljubljana, Večna pot 111, SI-1000 Ljubljana, Slovenia

Abstract: Ecological performance of animals depends on physiological and biochemical processes that are adjusted to the environment. The responses to hypoxia or anoxia have been frequently studied in subterranean aquatic organisms in order to find potential adaptations to restrict oxygen conditions occurring in the underground habitats. However, some previous studies have compared phylogenetic distant epigean and hypogean species or the epigean and hypogean populations of the same species due to little chance to compare closely related epigean and hypogean species. Therefore, in this study, we compared the effects of exposure to hypoxia, followed by reoxygenation, and increased temperature on oxygen consumption, potential metabolic activity, and antioxidant activities in closely related epigean and hypogean species: *Niphargus zagrebensis* and *N. stygius*. Oxygen consumption of *N. stygius* increased similarly during post-hypoxic recovery at 10 and 20°C (approx. 5-times), while *N. zagrebensis* increased its oxygen consumption for 9.7 and 4.4-times at 10 and 20°C, respectively. We observed higher exploitation of metabolic potential for current oxygen consumption during reoxygenation in *N. zagrebensis* than *N. stygius*. Exposure to hypoxia and subsequent reoxygenation at 20°C increased catalase (CAT) activity in *N. stygius*, but not in *N. zagrebensis*. We observed increased glutathione reductase activity in both *Niphargus* species. We concluded that respiratory and antioxidant responses to severe hypoxia and increased temperature differed between closely related epigean and hypogean *Niphargus* species. Hypogean *Niphargus* species possess physiological and biochemical characteristics that are advantageous in temperature stable subterranean environments which support inhabiting of species that have low energetic demands, while epigean *Niphargus* species can successfully inhabit specific surface habitats.

Keywords: *Niphargus stygius*, *Niphargus zagrebensis*, hypoxia adaptation, temperature, metabolic activity
Received 19 November 2020; Revised 13 March 2021; Accepted 13 March 2021

Citation: Simčič, T., Sket, B., 2021. Ecophysiological responses of two closely related epigean and hypogean *Niphargus* species to hypoxia and increased temperature: Do they differ? International Journal of Speleology, 50(2), 111-120.
<https://doi.org/10.5038/1827-806X.50.2.2369>

INTRODUCTION

Survival, growth and reproduction of organisms depend on physiological and biochemical processes that are adjusted to the environment. Aquatic arthropods differ in tolerance to hypoxia or anoxia. Many species, including numerous crustaceans, exhibit oxygen debts that are repaid upon return to normoxia, but other invertebrates do not (Herreid, 1980; Ellington, 1983; Zou et al., 1996). Oxygen debt, i.e., the additional oxygen consumption by the animal in the recovery period from anaerobic stress, is generally interpreted as needed in order to meet the increased energy demands for disposal of

end products, including oxidation of anaerobic end products for energy, converting end products into storage products, such as glycogen, and regenerating the phosphagen and ATP stores depleted during severe hypoxia (Herreid, 1980).

Many aquatic subterranean organisms have to cope with periodic oxygen depletion in their habitats with sometimes rapid switches from normoxia to hypoxia or even anoxia. Numerous previous studies reported that hypogean species are better adapted to low oxygen content and are better equipped to remain aerobic under hypoxia than epigean ones (Hervant et al., 1995, 1996, 1997b, 1998). Nevertheless, Culver & Poulson (1971) pointed out

*Tatjana.Simcic@nib.si

that phylogenetic effects should be taken into account when comparing the biological responses of epigeal and hypogean species. Unfortunately, there is little opportunity to compare closely related epigeal and hypogean species. Therefore, an alternative approach comparing biological traits among epigeal and hypogean populations of the same species has been used (Biswas, 1991; see review of Malard & Hervant, 1999). For example, the locomotory and ventilatory activities, oxygen consumption, and the intermediary and energy metabolism modifications of a spring and a cave population of the aquatic amphipod crustacean *Gammarus minus* Say were investigated in normoxia, severe hypoxia ($P_{O_2} < 0.03$ kPa), and subsequent recovery to compare the reactions of both populations to these experimental conditions, and the degree of adaptation to hypoxia (Hervant et al., 1999a). Despite their different origins, both populations of *G. minus* presented identical responses in all experimental conditions. Thus, it was assumed that a high resistance to hypoxia is not general characteristic of hypogean organisms but is more related to oxygen availability of particular subterranean habitats (Hervant & Malard, 2019).

In Slovenia, genus *Niphargus* (Amphipoda, Niphargidae) is represented by closely related stygobiotic *Niphargus stygius* (Schoedte) and semisubterranean *Niphargus zagrebensis* S. Karaman that invaded into epigeal waters from the subterranean habitats (Karaman, 2019). Both species are morphologically similar since *N. stygius* is not strongly troglomorphic, while *N. zagrebensis* is considered as a member of an epigeal species group exhibiting the same degree of a fictitious troglomorphy (Sket, 2008). This distribution of species gives us a unique opportunity to compare the responses of both species to severe hypoxia and subsequent recovery in normoxia in order to test the previous findings obtained in distantly related species or the epigeal and hypogean populations of the same species.

Besides the immediate harmful effect of anaerobiosis, an increased production of reactive oxygen species (ROS) occurs during reoxygenation due to the recovery of activity of the mitochondrial respiratory chain (Hervant & Malard, 2019). Due to the damaging effects of ROS overproduction animals have developed both non-enzymatic and enzymatic antioxidant mechanisms. To prevent oxidative stress, animals developed anti-oxidative enzymes, including catalase (CAT) and glutathione reductase (GR) (Lushchak et al., 2001; Lushchak & Bagnyukova, 2006; Issartel et al., 2009; Lawniczak et al., 2013; Vranković et al. 2017). CAT is a very important enzyme in protecting the cell from oxidative damage by ROS, and it has one of the highest turnover numbers of all enzymes. It catalyzes the decomposition of hydrogen peroxide to water and oxygen (Zamocky et al., 2008). GR is the enzyme critical for the reduction of oxidized form of glutathione (GSSG) responsible for maintaining the supply of reduced glutathione (GSH) involved in neutralization of free radicals (Couto et al., 2016). Moreover, oxygen content has been reported as an environmental factor that correlated to CAT activity and GSH content in

zebra mussel (*Dreissena polymorpha* (Pallas)) from the two ecosystems (Wojtal-Frankiewicz et al., 2017). Previous study on the groundwater crustacean *Niphargus rhenorhodanensis* Schellenberg revealed an overactivation of superoxide dismutase (SOD) and glutathione peroxidase (GPx) activities after exposure to drastic variations in oxygen level compared to the control group (Lawniczak et al., 2013), but information about the antioxidant responses of the closely related epigeal *Niphargus* species is still lacking.

While the basis of hypoxia and temperature as interlinked stressors in aquatic organisms has been proven (Pörtner & Knust, 2007; Pörtner, 2010), increased temperature was also included in the present study. Because physiological and biochemical rates of ectothermic organisms are highly temperature-dependent (Hochachka & Somero, 2002), their behavioural and ecological performance, and even fitness, can be influenced by body temperature (Huey & Kingsolver, 1989). The increase in temperature stimulates all metabolic processes in accordance with known thermodynamic principles. For example, it enhances oxygen consumption and, therefore, may increase ROS production as side products of intensified metabolism, resulting in oxidative stress (Lushchak, 2011). Moreover, higher thermal sensitivity of hypogean *N. stygius* than epigeal *N. zagrebensis* that resulted in increased oxygen consumption as a consequence of stress response (Simčič & Sket, 2019) could enhance ROS production in subterranean species at increased temperatures.

An animal's energy demand is usually quantified by its metabolic rate. Measurement of the entire organism's oxygen consumption gives meaningful information on metabolic rates. For example, oxygen consumption has been frequently used in comparative studies of epigeal and hypogean organisms in relation to different environmental factors (e.g., Culver & Poulson, 1971; Hervant et al., 1995, 1996, 1997a,b, 1998, 1999a; Issartel et al., 2005; Simčič et al., 2005, 2010; Mezek et al., 2010), while measurement of respiratory electron transport system (ETS) activity has been used sporadically (Simčič et al., 2005, 2010; Mezek et al., 2010; Simčič & Sket, 2019). However, measurements of oxygen consumption can provide information about metabolic intensity under particular conditions, but the question remains as to what the measured energy demands mean, with regard to the whole metabolic capacity, for a single species. Therefore, ETS activity has been measured to estimate potential metabolic activity, i.e., the value of oxygen consumption that would occur if all enzymes functioned maximally (Muskó et al., 1995). Moreover, the ratio between oxygen consumption and potential metabolic activity (R/ETS ratio) has been frequently used as an estimator of the exploitation of metabolic potential for current metabolic activity in relation to various environmental conditions and reflects the plasticity and fitness of organisms (e.g., Muskó et al., 1995; Fanslow et al., 2001; Simčič et al., 2005; Lukančič et al., 2010; Simčič et al., 2015).

The goals of the present study were to: (i) investigate the effects of hypoxia exposure and increased

temperature on respiratory response during post-hypoxic recovery in closely related epigeal and hypogean species; (ii) estimate the exploitation of metabolic potential for current metabolic demands during reoxygenation; (iii) explore the activity of the antioxidant enzymes displayed by the epigeal and hypogean crustaceans during reoxygenation after exposure to severe hypoxia at different temperatures. To reach these goals, we measured oxygen consumption in normoxia and during normoxic recovery after exposure to severe hypoxia in closely related subterranean amphipod *N. stygius* and epigeal amphipod *N. zagrebensis* at 10 and 20°C. Moreover, ETS activity was determined to estimate the ratio between oxygen consumption and potential metabolic activity (R/ETS ratio). The activity of two antioxidant enzymes, i.e., catalase (CAT) and glutathione reductase (GR) was measured in both *Niphargus* species.

MATERIALS AND METHODS

Collection and maintenance of animals

Niphargus zagrebensis S. Karaman 1950 (syn. *N. valachicus* z., *N. elegans* z.) is eyeless and only feebly pigmented semisubterranean species that occurs in epigeal waters where it is mixed with strongly pigmented epigeal species (Karaman, 2019). Our sample was collected in stagnant or slowly flowing water in puddles and ditches with loamy bottoms and rich deposits of fallen leaves in a lowland oak forest Krakovski gozd at Kostanjevica (mean body mass \pm SE: 26.4 \pm 3.1 mg wet mass, n = 20). As access to caves or interstitial water is here mainly absent, *N. zagrebensis* is considered as an epigeal species in the present study.

Niphargus stygius (Schioedte 1847) is strongly stygobiotic species. The sample in this study was taken from the cave Unška koliševka at Planina (NE of Postojna, Slovenia) (30.1 \pm 2.6 mg, n = 20). The habitats were small puddles of percolated water in artificially reshaped karst caves (past military galleries). Specimens of both species used in the study were males of a similar body mass (t = 1.12, d.f. = 38, p > 0.05). The mean yearly temperature (i.e., also the permanent temperature of ground waters) in 2005-2014 was 10-12°C at Novo mesto (close to Krakovski gozd) and 9-11°C at Postojna. The mean monthly temperatures in January and July varied from -1.7 to 1.4°C and 20.5 to 22.8°C in Novo mesto, but from -2.3 to 4.7°C and 18.5 to 22.3°C in Postojna (Statistical office, 2018).

Specimens were collected using a hand net. They were stored in plastic bottles and transported to the laboratory, where they were kept in aquaria in constant darkness at 10°C (\pm 0.5°C) in a thermo-regulated chamber for three weeks. Aquaria were previously filled with chemically controlled (synthetic) and aerated water that was prepared by adding 2940 mg CaCl \times 2 H₂O, 1230 mg MgSO₄ \times 7 H₂O, 650 mg NaHCO₃ and 60 mg KCl into 10 L of bi-distilled water (ISO standard 6341, 1996). During this period of acclimation to laboratory conditions, animals were

fed *ad libitum* with a commercial food (Sera crabs natural™). Water in aquaria was changed twice a week.

After acclimation to laboratory conditions, individuals of each species were transferred into two aquaria containing synthetic water and food. One aquarium with experimental animals was kept at 10°C, while the water temperature in second one was gradually (i.e., 1°C/12 h) increased to 20°C. After 5 days of acclimation to the newly reached temperature, animals were separately transferred from aquaria into 100 mL beaker containing 50 mL of synthetic water of adequate temperature, i.e., 10 or 20°C. Animals were starved for 24 hours before oxygen consumption measurements were taken.

Oxygen consumption measurement in normoxia and during post-hypoxic recovery

Oxygen consumption was estimated by the closed bottle method (Lampert, 1984). First, ground-glass stoppered bottles were filled with synthetic and aerated water from the same, well-mixed, container to measure oxygen consumption in normoxia. The experimental bottles (n = 5) received animals, while two bottles served as controls. All bottles were stoppered and kept in the dark at 10 or 20°C. The concentration of dissolved oxygen in the experimental and control bottles was measured with a 4-Channel fiber optic oxygen meter (PreSens OXY-4, Germany) for the first time after 30 min, and a second time at the end of the incubation period (i.e., after 3 h). Oxygen consumption was expressed as mg O₂ per g of wet weight (mg O₂ g⁻¹WW h⁻¹). After normoxia measurements, the same animals were introduced into experimental bottles filled with deoxygenated water that was generated by bubbling pure nitrogen gas. Water replacement was carried out according to a procedure that ensured minimal stress for the animals. First, single animal was gently sucked into a wide glass dropper along with a minimal volume of water, from where it was slowly released into a bottle filled with deoxygenated water. All bottles were stoppered and kept in the dark at 10 or 20°C. The concentration of dissolved oxygen in bottles was below 0.3 mg L⁻¹ during hypoxia. The selected incubation time for the exposure to hypoxia was longer than that for normoxia measurement in order to perceive the consequences of hypoxia in *N. stygius*, but also it was short enough to ensure the survival of all specimens of *N. zagrebensis*. After 5 h of incubation, the deoxygenated water in the bottles was quickly replaced with air saturated water. Animals were transferred in a small volume of water using a wide glass dropper, as described above. Oxygen concentration was measured during post-hypoxic recovery after 15 min, 30 min, 1, 1.5, 2, 3, 4, 5, 6, 7, and 8 h at 10 and 20°C.

After oxygen consumption measurements, animals were placed between two sheets of filter paper and gently squeezed to remove the water from their surface. They were then placed on a pre-weighed piece of aluminium foil and weighed on an electrobalance (Sartorius BP 210 S) with 0.1 mg accuracy.

Procedure before measurements of ETS, CAT and GR activities

Animals ($n = 5$) were incubated in deoxygenated water (oxygen concentration was below 0.3 mg L^{-1}) for 5 h at 10 or 20°C . At the end of incubation in hypoxic conditions, the deoxygenated water was replaced with aerated water according to the procedure described above in "Oxygen consumption measurement in normoxia and during post-hypoxic recovery" subsection. After 1 h of post-hypoxic recovery, individuals were immediately frozen in liquid nitrogen and stored at -80°C , prior to the preparation of the homogenate for the measurements of ETS, CAT and GR activities and determination of protein concentration. For controls, the individuals of each species ($n = 5$) were kept in normoxic conditions at each experimental temperature and were frozen and stored as described above.

ETS activity

Respiratory electron transport system (ETS) activity was measured using the method originally proposed by Packard (1971) and improved by G.-Tóth (1999). The ETS activity of each animal was determined using the procedure described in Simčić & Sket (2019). The rate of tetrazolium dye reduction to formazan was converted to equivalent oxygen as described by Kenner & Ahmed (1975).

Catalase (CAT) activity

CAT activity was measured according to the method proposed by Aebi (1984). The procedure used in the present study was previously described in detail in Simčić et al. (2015). Enzyme activities were expressed as enzyme units (U), where one U was defined as the amount of CAT that degrades one μmol of hydrogen peroxide in 1 min. These results were divided by the total amount of protein to give the specific CAT activity per mg protein.

Glutathione reductase (GR) activity

GR activity was determined according to the method proposed by Carlberg & Mannervik (1985), using the procedure described in Simčić et al. (2015). One U was defined as the amount of GR that degrades 1 μmol of NADPH in 1 min. The specific GR activity per mg protein was calculated.

Protein concentration

Protein concentration was determined using a commercial Pierce™ BCA protein assay kit (Thermo Scientific, USA) according to the manufacturer's recommendations. Absorbance of the samples was measured using a Lambda UV/Vis spectrophotometer (PerkinElmer, USA).

Data analysis

Values are presented as means \pm SE. The data were first tested for normality of distributions (Shapiro-Wilk test) and homogeneity of variances (Levene's test), and log transformed when necessary. Body masses of the two species were compared by Student's t-test. The differences between values were investigated

using one-way ANOVAs. When significant differences in ANOVAs were found, the *post hoc* Tukey's test was performed to find out which means differ significantly. All statistical analyses were conducted with SPSS 20.0 (SPSS Inc. Chicago, Illinois, USA).

RESULTS

Oxygen consumption in normoxia and during post-hypoxic recovery

Oxygen consumption during reoxygenation varied significantly - from $1.16 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ to $0.12 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ in *N. zagrebensis* (ANOVA, $F_{10,44} = 8.61$; $p < 0.001$), and from $0.38 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ to $0.06 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ in *N. stygius* (ANOVA, $F_{10,44} = 5.46$; $p < 0.001$) at 10°C (Fig. 1a). Similarly, both species had significantly increased oxygen consumption after a half an hour in normoxic recovery period at 10°C when compared to oxygen consumption in normoxia (Control), but with different increases. Oxygen consumption was, compared to the control, nearly 5-times (4.9) higher in hypogean *N. stygius*, and almost 10-times (9.7) higher in epigeal *N. zagrebensis*. After reaching the maximal value, oxygen consumption of both species during normoxic recovery period was decreasing for several hours, until it was similar to the control value (after 2 h of normoxic recovery in *N. stygius* and 5 h in *N. zagrebensis*).

Significant variations in oxygen consumption during reoxygenation were also observed in *N. zagrebensis* (ANOVA, $F_{10,44} = 7.36$; $p < 0.001$) and *N. stygius* (ANOVA, $F_{10,44} = 15.70$; $p < 0.001$) at 20°C , where values ranged from $1.30 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ to $0.34 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ and from $0.76 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$ to $0.13 \text{ mg O}_2 \text{ g}^{-1} \text{ h}^{-1}$,




Fig. 1. Oxygen consumption in normoxia (Cont) and during subsequent normoxic recovery after severe hypoxia ($0.3 \text{ mg O}_2 \text{ L}^{-1}$; Hypo) measured in epigeal *Niphargus zagrebensis* and in hypogean *Niphargus stygius* in darkness at 10°C (a) and 20°C (b). Values are means \pm SE for $n = 5$. Full symbols differ significantly from control (Cont) — $p < 0.05$ (ANOVA, Tukey's HSD test).

respectively (Fig. 1b). Different patterns of subsequent normoxic recovery after severe hypoxia were observed at 20°C than at 10°C for *N. zagrebensis*. *N. stygius* had, similarly, the highest oxygen consumption at 0.5 h of recovery period, but *N. zagrebensis* reached the highest oxygen consumption after 1 h of recovery. Oxygen consumption rate increased for 4.6-times in hypogean *N. stygius* (0.5 h) and 4.4-times in epigeal *N. zagrebensis* (1 h). After reaching maximal value, oxygen consumption of *N. stygius* decreased gradually for several hours, but for *N. zagrebensis* a steep decrease in oxygen consumption was observed.

ETS activity

Significant differences in ETS activity between temperatures and oxygen conditions (i.e., in normoxia as control and 1 h after post-hypoxic recovery in normoxia) were observed for *N. zagrebensis* ($F_{3,16} = 23.83$, $p < 0.001$, Fig. 2a) and *N. stygius* ($F_{3,16} = 195.16$, $p < 0.001$, Fig. 2b). Tukey *post hoc* test showed that ETS activity of *N. zagrebensis* was similar in normoxia and during post-hypoxic recovery within each temperature, but it differed significantly between temperatures, where it was higher at 20°C. In *N. stygius*, significant differences were observed between temperatures and oxygen treatments. Higher ETS activities were obtained at higher temperature and during post-hypoxic recovery, respectively.




Fig. 2. Respiratory electron transport system (ETS) activity in normoxia (Control) and after 1 h of subsequent normoxic recovery from severe hypoxia ($< 0.3 \text{ mg O}_2 \text{ L}^{-1}$) determined in epigeal *Niphargus zagrebensis* (a) and hypogean *Niphargus stygius* (b) at 10°C and 20°C. Bars labelled with different letters differ significantly — $p < 0.05$ (ANOVA, Tukey's HSD test). Values are means \pm SE for $n = 5$.

R/ETS ratio

The R/ETS ratio was calculated to illustrate the exploitation of metabolic potential for actual metabolic activity. *N. stygius* and *N. zagrebensis* had similar R/ETS ratios at 10°C (12 and 13%, respectively) and at 20°C (8%) in normoxia (Control) before exposure to hypoxic stress (Fig. 3). During post-hypoxic recovery at 10°C, the R/ETS ratio of *N. zagrebensis* showed an extreme increase in oxygen consumption and exploitation of metabolic potential for current metabolic activity at 0.5 h of normoxic recovery (i.e., 115%), whereas that of *N. stygius* increased just up to 40%. At 20°C, maximal value of R/ETS ratio was 30% for *N. stygius* and 47% for *N. zagrebensis*.




Fig. 3. Estimation of the exploitation of electron transport system (ETS) activity (metabolic potential) for actual metabolic activity (R/ETS ratio, in percentages) in *Niphargus zagrebensis* and *N. stygius* at 10°C (a) and 20°C (b) in normoxia (Cont) and during normoxic recovery after severe hypoxia. Mean values of oxygen consumption rates and ETS activities were used for the calculation of ratios.

Response of antioxidant enzymes

Hypoxic stress, followed by 1 h of post-hypoxic recovery in normoxia and increased temperature, showed significant modification in CAT activity ($F_{3,16} = 40.84$, $p < 0.001$, Fig. 4) and GR activity ($F_{3,16} = 14.33$, $p < 0.001$, Fig. 5) for *N. stygius*, where higher activities of CAT and GR were measured in animals during post-hypoxic recovery in normoxia compared to control animals at 20°C. Similarly, *N. zagrebensis* showed significant variation in CAT activity ($F_{3,16} = 3.84$, $p < 0.05$) and GR activity ($F_{3,16} = 3.62$, $p < 0.05$). Tukey *post hoc* test showed that *N. zagrebensis* had significantly higher CAT activity during normoxic recovery after severe hypoxia at 20°C than at 10°C. Moreover, GR activity of *N. zagrebensis* was higher during subsequent post-hypoxic recovery compared to the control at 20°C.




Fig. 4. Catalase (CAT) activity measured in *Niphargus zagrebensis* (a) and *N. stygius* (b) at 10 and 20°C in normoxia (Control) and after 1 h of normoxic recovery from severe hypoxia (<0.3 mg O₂ L⁻¹). Bars labelled with different letters differ significantly — $p < 0.05$ (ANOVA, Tukey's HSD test). Values are means \pm SE for $n = 5$.




Fig. 5. Glutathion reductase (GR) activity measured in *Niphargus zagrebensis* (a) and *N. stygius* (b) at 10 and 20°C in normoxia (Control) and after 1 h of normoxic recovery from severe hypoxia (<0.3 mg O₂ L⁻¹). Bars labelled with different letters differ significantly — $p < 0.05$ (ANOVA, Tukey's HSD test). Values are means \pm SE for $n = 5$.

DISCUSSION

While oxygen level and temperature have been proven as interlinked stressors in aquatic organisms (Pörtner & Knust, 2007; Pörtner, 2010) and, moreover, they both have significant effects on oxygen consumption and ROS formation, the interaction of both factors expectedly resulted in different respiratory and antioxidant responses to hypoxia-reoxygenation and increased temperature in epigean and hypogean *Niphargus* species, namely *N. zagrebensis* and *N. stygius*. For both species, post-hypoxic recovery resulted in increased oxygen consumption. The higher increase of oxygen consumption in epigean than hypogean species is in accord with the findings of previous studies, where distantly related epigean and hypogean species were compared (Hervant et al., 1995, 1996, 1998). For example, Hervant et al. (1998) reported that the main explanation for the lower oxygen debt shown by hypogean organisms is the lower energetic expenditures noticed during hypoxia, partly due to a decrease in locomotory and ventilatory activities. Moreover, oxygen consumption of *N. stygius* increased similarly during recovery at 10 and 20°C, i.e., approximately for five-times of normoxic rates, but *N. zagrebensis* increased its oxygen consumption relatively less at 20°C than at 10°C. However, it is ecologically very important for organisms to recover quickly and completely from hypoxic or anoxic stress when oxygen is once more available (Hervant & Malard,

2019). Previous studies revealed faster replenishment of ATP levels in hypogean than in epigean species during post-hypoxic recovery (Hervant et al., 1995, 1996, 1997b).

Moreover, observations of the hypogean *Stenasellus virei* Dollfus, *Niphargus virei* Chevreux, *N. rhenorhodanensis* and the epigean *Asellus aquaticus* Linne and *Gammarus fossarum* Koch showed that part of the end products was excreted and part was metabolized (Hervant et al., 1995, 1996, 1997b, 1999b). Excretion is an important mechanism for the disposal of lactate during aerobic recovery in epigean species. This is a costly strategy because of a loss of energy-rich carbon chains. In contrast, hypogean crustaceans preferentially use glycconeogenesis to convert lactate into glycogen stores (Hervant & Malard, 2019). Thus, the gradual decrease of oxygen consumption in *N. stygius* after reaching its maximal value during normoxic recovery could indicate a subsequent turnover of the anaerobic end products. But in *N. zagrebensis*, the maximal oxygen consumption observed after 0.5 and 1 h of post-hypoxic recovery period at 10°C and 20°C, respectively, is followed by a rapid decrease at both temperatures. Therefore, it is assumed that the reason for the quick decrease in oxygen consumption and relatively low oxygen consumption in *N. zagrebensis* at higher temperatures is likely the excretion of anaerobic end products instead of being converted, as evidenced by epigean species in

previous studies (Hervant et al., 1995, 1996, 1997b, 1999b).

As expected, higher ETS activities were obtained at a higher temperature in both species, while differences between species were observed in response to hypoxic stress, where *N. stygius* had higher ETS activity after exposure to hypoxic stress compared to controls, but *N. zagrebensis* did not. Le Moullac et al. (2007) reported that 20 days of hypoxia significantly increased the activity of ETS in the oyster *Crassostrea gigas* (Thunberg). The authors assumed that the stimulation of the ETS activity, observed in molluscs, was linked to the up-regulation of some genes of the respiratory chain, as was previously observed in anoxia tolerant vertebrates (Krivoruchko & Storey, 2015). Many species have developed mechanisms that allow them to compensate for periods of low oxygen. These mechanisms improve oxygen delivery to tissues and/or to increase ATP production by oxygen-independent means (e.g., glycolysis) to compensate for the reduced ATP output by oxygen-dependent pathways in the mitochondria (Krivoruchko & Storey, 2015). Hervant et al. (1999b) reported that the metabolic response of *N. virei* during post-hypoxic recovery mainly exhibited the characteristics of vertebrates, where a strategy of lactate removal was quite different from that observed in epigeal crustaceans, favouring lactate supported gluco- and glycogenogenesis and rapid glycogen replenishment instead of rapid lactate removal via oxidative pathways. Thus, it seems that hypoxic tolerant *N. stygius* exhibited stimulation of ETS activity during hypoxic stress in order to provide sufficient ATP production, while *N. zagrebensis* was incapable of increasing its metabolic potential.

The ratio between oxygen consumption and ETS activity (R/ETS) is an important index of organisms' metabolism (Muskó et al., 1995; Fanslow et al., 2001; Simčič et al., 2005; Lukančič et al., 2010; Simčič et al., 2015; Simčič & Sket, 2019). The values of the present study, measured in normoxic conditions before exposure to severe hypoxia, are in agreement with those of Simčič et al. (2005) reported for hypogean *Niphargus* species, where exploitation of metabolic potential ranged from 12 to 20% in *N. krameri* and two populations of *N. stygius*. It seems that closely related epigeal and hypogean *Niphargus* species in normoxia possess a similar ratio, while a distantly related epigeal species, namely *Gammarus fossarum*, used a higher percentage (i.e., 40%) of metabolic potential for actual metabolic activity (Simčič et al., 2005). Lower exploitation of metabolic potential is an advantage in variable environments, where organisms could immediately use the existing enzyme machinery for increased metabolic activity during recovery under favourable conditions (Fanslow et al., 2001; Simčič et al., 2005; Žagar et al., 2015). Moreover, the relatively low ratio (i.e., 9%) of *Nereis virens* was explained by its low activity and intermittent exposure to anoxic conditions (Cammen et al., 1990). Stimulated ETS activity under severe hypoxia in *N. stygius*, observed in the present study, probably additionally supported quick and efficient post-hypoxic recovery. Thus, the exploitation of metabolic potential in *N. stygius*, which

increased up to 40% at 10°C, was a metabolically more favourable condition than the high R/ETS ratio in *N. zagrebensis* at 10°C which indicated exploitation of the whole metabolic potential. A similar pattern with higher exploitation of metabolic potential in *N. zagrebensis* than *N. stygius* was also observed at 20°C. Lower exploitation of metabolic potential at 20°C than at 10°C is in accord with the results of the previous studies, where R/ETS ratio decreased with increasing temperature due to different response of oxygen consumption and ETS activity to temperature change (Bamstedt, 1980; Muskó et al., 1995; Simčič & Brancelj, 1997; Simčič, 2005; Simčič & Sket, 2019). Furthermore, relatively lower R/ETS ratios in *N. stygius* exhibited at the end of the recovery period than they were displayed in normoxia before exposure to hypoxia (Control) indicated on lower energetic expenditures.

During hypoxia and subsequent reoxygenation increased ROS levels are expected to activate antioxidant defenses (Hermes-Lima et al., 2015). The results of the present study showed a higher activity of antioxidant enzyme CAT in *N. stygius* during post-hypoxic recovery than it was in control at 20°C. These results are in accord with the findings of the previous studies on crustaceans where increased CAT activities in hypoxia and reoxygenation were reported (Gorokhova et al., 2010, 2013; Trasviña-Arenas et al., 2013). Moreover, increased GR activity that catalyzes the reduction of GSSG to GSH was observed in both *Niphargus* species during post-hypoxic recovery at 20°C. Nevertheless, hypogean *N. stygius* had, under hypoxia and subsequent reoxygenation, increased activity of the primary antioxidant enzyme, i.e., CAT, while epigeal *N. zagrebensis* did not. Lawniczak et al. (2013) similarly reported the increased GPx activity in subterranean amphipod *N. rhenorhodanensis* after 24 h exposure to anoxia. If the antioxidant potential is high enough, ROS level returns to the initial level before oxidative stress could occur. However, when the efficiency of the antioxidant system does not counterbalance enhanced ROS production, the presence of excess ROS in cells causes oxidative damage of lipids, proteins, and nucleic acids (Lushchak et al., 2001). Thus, lower CAT activity in *N. zagrebensis* than *N. stygius* at 20°C indicates a lower response to hypoxia in former species. Consequently, it is assumed that a decline in ETS activity of *N. zagrebensis* at 20°C could be caused by a less efficient antioxidant defense that resulted in the oxidative damages of cellular structures. However, to test this assumption, some parameters of oxidative damage, e.g., lipid peroxidation or protein carbonylation, ought to be measured.

In this study, we observed higher activities of CAT and GR during post-hypoxic recovery in comparison to controls at 20°C, but not at 10°C. Bagnyukova et al. (2007) claimed that the activities of primary antioxidant enzymes – superoxide dismutase (SOD) and CAT – were unaffected in organs of goldfish *Carassius auratus*, whereas glutathione-dependent enzymes, such as also GR, were elevated after more prolonged exposure to high temperature. In the

present study, the exposure of organisms of both *Niphargus* species solely to higher temperatures did not increase CAT nor GR activities, whereas combined effects of high temperature and hypoxia with subsequent normoxic recovery resulted in increased CAT and GR activities of both species. As reported by Vinagre et al. (2012), an oxidative stress response is not directly correlated to temperature. It is lowest at the optimal temperature, and it increases outside this species' upper and lower optimum thermal limits. In addition, as 10°C is an average habitat temperature of both species and a permanent one for *N. stygius*, a higher temperature (i.e., 20°C) likely contributes to overall environmental stress. This explanation is in agreement with the reports of Gorokhova et al. (2013), who found that co-exposure of amphipod *Monoporeia affinis* (Lindstrom) to both fluctuating hypoxia and contaminants caused greater increase in antioxidants and lipid peroxidation, and slowed recovery from hypoxia as indicated by CAT and glutathione redox state (GSH/GSSG ratio). We, therefore, assumed that temperature-stable subterranean habitats represent an advantage for hypogean animals that have to face frequent exposure to variable oxygen concentrations.

In conclusion, closely related epigeal and hypogean *Niphargus* species have, in fact, similar oxygen consumption in normoxia, but they respond differently to exposure to environmental stressors, such as severe hypoxia and increased temperature. Thus, as reported previously by Hervant et al. (1998), the lower energetic expenditures noticed during hypoxia due to a decrease in locomotory and ventilatory activities contributed to lower oxygen debt in hypogean species. Moreover, the consistent respiratory response of *N. stygius* to hypoxia at 10 and 20°C, i.e., gradual decreasing of oxygen consumption during repayment of oxygen debt as a result of metabolization of anaerobic end products, and enhanced antioxidant enzyme activities after exposure to severe hypoxia and higher temperature, are probably related to some physiological and biochemical traits of stygobitic species that enable *N. stygius* to cope successfully with specific conditions in subterranean environments. However, a high resistance to hypoxia, followed by the activation of antioxidant defenses during reoxygenation, is not universally found in subterranean species (e.g., Issartel et al., 2009). Thus, additional studies on the mechanisms involved in hypoxia tolerance of different groundwater animals are needed to improve understanding of adaptation to restrict oxygen conditions.

ACKNOWLEDGEMENTS

The authors thank Jennifer C. Ellis for suggestions on an earlier draft of this manuscript and the English revision of the text and the reviewers for constructive comments. This study was financially supported by the Slovenian Research Agency (Research Program P1-0255).

Authorship statement: TS and BS designed and directed the study. BS collected and determined

the animals. TS performed the measurements and analyzed the data. TS wrote the paper with substantial input from BS.

REFERENCES

- Aebi, H., 1984. Catalase *in vitro*. Methods in Enzymology 105, 121–126.
[https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3)
- Bagnyukova, T.V., Lushchak, O.V., Storey, K.B., Lushchak, V.I., 2007. Oxidative stress and antioxidant defense responses by goldfish tissues to acute change of temperature from 3 to 23°C. Journal of Thermal Biology, 32, 227–234.
<https://doi.org/10.1016/j.jtherbio.2007.01.004>
- Bamstedt, U., 1980. ETS activity as an estimator of respiratory rate of zooplankton populations. The significance of variations in environmental factors. Journal of Experimental Marine Biology and Ecology, 42, 267–283.
[https://doi.org/10.1016/0022-0981\(80\)90181-1](https://doi.org/10.1016/0022-0981(80)90181-1)
- Biswas, J., 1991. Metabolic efficiency and regulation of body weight: a comparison between life in hypogean and epigeal ecosystems. International Journal of Speleology, 20, 15–22.
<https://doi.org/10.5038/1827-806X.20.1.2>
- Cammen, L.M., Corwin, S., Christensen, J.P., 1990. Electron transport system (ETS) activity as a measure of benthic macrofaunal metabolism. Marine Ecology Progress Series, 65, 171–182.
<https://doi.org/10.3354/meps065171>
- Carlberg, C., Mannervik, B., 1985. Glutathione reductase. In: Meister, A. (Ed.). Methods in Enzymology, vol 113. Academic Press, New York, p. 488–495.
- Couto N., Wood J., Barber, J., 2016. The role of glutathione reductase and related enzymes on cellular redox homeostasis network. Free Radical Biology and Medicine, 95, 27–42.
<https://doi.org/10.1016/j.freeradbiomed.2016.02.028>
- Culver, D.C., Poulson, T.L., 1971. Oxygen consumption and activity in closely related amphipod populations from cave and surface habitats. American Midland Naturalist, 85, 74–84. <https://doi.org/10.2307/2423913>
- Ellington, W.R., 1983. The recovery from anaerobic metabolism in invertebrates. Journal of Experimental Zoology, 228, 431–444.
<https://doi.org/10.1002/jez.1402280305>
- Fanslow, D.L., Nalepa, T.F., Johengen, T.H., 2001. Seasonal changes in the respiratory electron transport system (ETS) and respiration rate of the zebra mussel, *Dreissena polymorpha* in Saginaw Bay, Lake Huron. Hydrobiologia, 448, 61–70.
- Gorokhova E., Löf, M., Halldórsson, H.P., Tjärnlund, U., Lindström, M., Elfving, T., Sundelin, B., 2010. Single and combined effects of hypoxia and contaminated sediments on the amphipod *Monoporeia affinis* in laboratory toxicity bioassays based on multiple biomarkers. Aquatic Toxicology, 99, 263–274.
<https://doi.org/10.1016/j.aquatox.2010.05.005>
- Gorokhova, E., Löf, M., Reutgard, M., Lindström, M., Sundelin, B., 2013. Exposure to contaminants exacerbates oxidative stress in amphipod *Monoporeia affinis* subjected to fluctuating hypoxia. Aquatic Toxicology, 127, 46–53.
<https://doi.org/10.1016/j.aquatox.2012.01.022>
- G.-Tóth, L., 1999. Aktivität des Elektronentransportsystems. In: von Tümpling, W., Friedrich, G. (Eds.), Biologische Gewässeruntersuchung. Methoden der Biologischen Wasseruntersuchung 2, Gustav Fischer Verl., Jena, Stuttgart, Lübeck, Ulm, p. 465–473.

- Hermes-Lima, M., Moreira, D.C., Rivera-Ingraham, G.A., Giraud-Billoud, M., Genaro-Mattos, T.C., Campos, É.G., 2015. Preparation for oxidative stress under hypoxia and metabolic depression: Revisiting the proposal two decades later. *Free Radical Biology and Medicine*, 89, 1122–1143.
<https://doi.org/10.1016/j.freeradbiomed.2015.07.156>
- Herreid, C., 1980. Hypoxia in invertebrates. *Comparative Biochemistry & Physiology A*, 67, 311–320.
[https://doi.org/10.1016/S0300-9629\(80\)80002-8](https://doi.org/10.1016/S0300-9629(80)80002-8)
- Hervant, F., Mathieu, J., Garin, D., Freminet, A., 1995. Behavioral, ventilatory, and metabolic responses to severe hypoxia and subsequent recovery of the hypogean *Niphargus rhenorhodanensis* and the epigeal *Gammarus fossarum* (Crustacea: Amphipoda). *Physiological Zoology*, 68, 223–244.
<https://doi.org/10.1086/physzool.68.2.30166501>
- Hervant, F., Mathieu, J., Garin, D., Freminet, A., 1996. Behavioral, ventilatory, and metabolic responses of hypogean amphipod *Niphargus virei* and the epigeal isopod *Asellus aquaticus* to severe hypoxia and subsequent recovery. *Physiological Zoology*, 69, 1277–1300.
<https://doi.org/10.1086/physzool.69.6.30164261>
- Hervant, F., Mathieu, J., Barré, H., Simon, K., Pinon, C., 1997a. Comparative study on the behavioural, ventilatory, and respiratory responses of hypogean and epigeal crustaceans to long-term starvation and subsequent feeding. *Comparative Biochemistry & Physiology A*, 118, 1277–1283.
[https://doi.org/10.1016/S0300-9629\(97\)00047-9](https://doi.org/10.1016/S0300-9629(97)00047-9)
- Hervant, F., Mathieu, J., Messana, G., 1997b. Locomotory, ventilatory and metabolic responses of the subterranean *Stenasellus virei* (Crustacea, Isopoda) to severe hypoxia and subsequent recovery. *Comptes rendus de l'Académie des Sciences Paris, Sciences de la vie / Life Sciences*, 320, 139–148.
[https://doi.org/10.1016/S0764-4469\(97\)85005-6](https://doi.org/10.1016/S0764-4469(97)85005-6)
- Hervant, F., Mathieu, J., Messana, G., 1998. Oxygen consumption and ventilation in declining oxygen tension and post-hypoxic recovery in epigeal and hypogean crustaceans. *Journal of Crustacean Biology*, 18, 717–727.
<https://doi.org/10.1163/193724098X00593>
- Hervant, F., Mathieu, J., Culver, D.C., 1999a. Comparative responses to severe hypoxia and subsequent recovery in closely related amphipod populations (*Gammarus minus*) from cave and surface habitats. *Hydrobiologia*, 392, 197–204.
<https://doi.org/10.1023/A:1003511416509>
- Hervant, F., Garin D., Mathieu, J., Freminet, A., 1999b. Lactate metabolism and glucose turnover in the subterranean crustacean *Niphargus virei* during post-hypoxic recovery. *Journal of Experimental Biology*, 202, 579–592.
<https://jeb.biologists.org/content/202/5/579.article-info>
- Hervant, F., Malard, F., 2019. Adaptations: Low oxygen. In: White, W.B., Culver, D.C., Pipan, T. (Ed.), *Encyclopedia of caves* (3rd Ed.). Elsevier/Academic Press, London, p. 8–15.
<https://doi.org/10.1016/B978-0-12-814124-3.00002-9>
- Hochachka, P.W., Somero, G.N., 2002. *Biochemical adaptation: Mechanism and process in physiological evolution*. Oxford University Press, Oxford, 480 p.
- Huey, R.B., Kingsolver, J.G., 1989. Evolution of thermal sensitivity of ectotherms. *Trends in Ecology & Evolution*, 4, 131–135.
[https://doi.org/10.1016/0169-5347\(89\)90211-5](https://doi.org/10.1016/0169-5347(89)90211-5)
- ISO-standard 6341 (1996) Determination of the inhibition of the mobility of *Daphnia magna* Straus (Cladocera, Crustacea)-Acute toxicity test (3rd ed.). Geneva: International Organization for Standardization.
- Issartel, J., Hervant, F., Voituron, Y., Renault, D., Vernon, Ph., 2005. Behavioural, ventilatory and respiratory responses of epigeal and hypogean crustaceans to different temperatures. *Comparative Biochemistry & Physiology A*, 141, 1–7.
<https://doi.org/10.1016/j.cbpb.2005.02.013>
- Issartel, J., Hervant, F., de Fraipont, M., Clobert, J., Voituron, Y., 2009. High anoxia tolerance in the subterranean salamander *Proteus anguinus* without oxidative stress nor activation of antioxidant defenses during reoxygenation. *Journal of Comparative Physiology B*, 179, 543–551.
<https://doi.org/10.1007/s00360-008-0338-9>
- Karaman, G.S., 2019. The subterranean species *Niphargus zagrebensis* S. Kar. 1950 (Fam. Niphargidae) on Balkan (Contribution to the knowledge of the Amphipoda 312). *Agriculture & Forestry*, 65, 67–89.
<https://doi.org/10.17707/AgricultForest.65.4.07>
- Kenner, R.A., Ahmed, S.I., 1975. Measurements of electron transport activities in marine phytoplankton. *Marine Biology*, 33, 119–127.
<https://doi.org/10.1007/BF00390716>
- Krivoruchko, A., Storey, K.B., 2015. Turtle anoxia tolerance: Biochemistry and gene regulation-Review. *Biochimica et Biophysica Acta*, 1850, 1188–1196.
<https://doi.org/10.1016/j.bbagen.2015.02.001>
- Lampert, W., 1984. The measurement of respiration. In: Downing, J.A., Rigler, F.H. (Eds.) *A manual on methods for the assessment of secondary productivity in fresh water*. IPB Handbook 17, Blackwell Scientific Publications, Oxford, p. 413–468.
- Lawniczak, M., Romestaing, C., Roussel, D., Maazouzi, C., Renault, D., Hervant, F., 2013. Preventive antioxidant responses to extreme oxygen level fluctuation in a subterranean crustacean. *Comparative Biochemistry & Physiology A* 165, 299–303.
<https://doi.org/10.1016/j.cbpa.2013.03.028>
- Le Moullac, G., Quéau, I., Le Souchu, P., Pouvreau, S., Moal, J., Le Coz, J.R., Samain, J.F., 2007. Metabolic adjustments in the oyster *Crassostrea gigas* according to oxygen level and temperature. *Marine Biology Research*, 3, 357–366.
<https://doi.org/10.1080/17451000701635128>
- Lukančić, S., Žibrat, U., Mezek, T., Jerebic, A., Simčić, T., Brancelj, A., 2010. A new method for early assessment of effects of exposing two non-target crustacean species, *Asellus aquaticus* and *Gammarus fossarum*, to pesticides, a laboratory study. *Toxicology & Industrial Health*, 26, 217–228.
<https://doi.org/10.1177/0748233710362379>
- Lushchak, V.I., 2011. Environmentally induced oxidative stress in aquatic animals. *Aquatic Toxicology*, 101, 13–30.
<https://doi.org/10.1016/j.aquatox.2010.10.006>
- Lushchak, V.I., Lushchak, L.P., Mota, A.A., Hermes-Lima, M., 2001. Oxidative stress and antioxidant defenses in goldfish *Carassius auratus* during anoxia and reoxygenation. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 280, R100–R107.
<https://doi.org/10.1152/ajpregu.2001.280.1.R100>
- Lushchak, V.I., Bagnyukova, T.V., 2006. Effects of different environmental oxygen levels on free radical processes in fish. *Comparative Biochemistry & Physiology B*, 144, 283–289.
<https://doi.org/10.1016/j.cbpb.2006.02.014>

- Malard, F., Hervant, F., 1999. Oxygen supply and the adaptations of animals in groundwater. *Freshwater Biology*, 41, 1–30.
<https://doi.org/10.1046/j.1365-2427.1999.00379.x>
- Mezek, T., Simčič, T., Arts, M.T., Brancelj A., 2010. Effect of fasting on hypogean (*Niphargus stygius*) and epigeal (*Gammarus fossarum*) amphipods: a laboratory study. *Aquatic Ecology*, 44, 397–408.
<https://doi.org/10.1007/s10452-009-9299-7>
- Muskó, I.B., G.-Tóth, L, Szábo, E., 1995. Respiration and respiratory electron transport system (ETS) activity of two amphipods: *Corophium curvispinum* G. O. Sars and *Gammarus fossarum* Koch. *Polish Archives of Hydrobiology*, 42, 547–558.
- Packard, T.T., 1971. The measurement of respiratory electron transport activity in marine phytoplankton. *Journal of Marine Research*, 29, 235–244.
- Pörtner, H.-O., 2010. Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. *Journal of Experimental Biology*, 213, 881–893.
<https://doi.org/10.1242/jeb.037523>
- Pörtner, H.-O., Knust, R., 2007. Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science*, 315, 95–97.
<https://doi.org/10.1126/science.1135471>
- Simčič, T., 2005. Respiratory electron transport system (ETS) activity and respiration rate in cold-stenothermal and eurythermal chironomid larvae from high-mountain lakes. *Archiv für Hydrobiologie*, 162, 399–415.
<https://doi.org/10.1127/0003-9136/2005/0162-0399>
- Simčič, T., Brancelj, A., 1997. Electron transport system (ETS) activity and respiration rate in five *Daphnia* species at different temperatures. *Hydrobiologia*, 360, 117–125.
https://doi.org/10.1007/978-94-011-4964-8_13
- Simčič, T., Lukančič, S., Brancelj, A., 2005. Comparative study of electron transport system activity and oxygen consumption of amphipods from caves and surface habitats. *Freshwater Biology*, 50, 494–501.
<https://doi.org/10.1111/j.1365-2427.2005.01339.x>
- Simčič, T., Pajk, F., Brancelj, A., 2010. Electron transport system activity and oxygen consumption of two amphibious isopods, epigeal *Ligia italica* Fabricius and hypogean *Titanethes albus* (Koch), in air and water. *Marine and Freshwater Behaviour and Physiology*, 43, 149–156.
<https://doi.org/10.1080/10236244.2010.483052>
- Simčič, T., Jesenšek, D., Brancelj, A., 2015. Effects of increased temperature on metabolic activity and oxidative stress in the first life stages of marble trout (*Salmo marmoratus*). *Fish Physiology and Biochemistry*, 41, 1005–1014.
<https://doi.org/10.1007/s10695-015-0065-6>
- Simčič, T., Sket, B., 2019. Comparison of some epigeal and troglolobiotic animals regarding their metabolism intensity. Examination of a classical assertion. *International Journal of Speleology*, 48(2), 133–144.
<https://doi.org/10.5038/1827-806X.48.2.2251>
- Sket, B., 2008. Can we agree on an ecological classification of subterranean animals. *Journal of Natural History*, 42, 1549–1563.
<https://doi.org/10.1080/00222930801995762>
- Statistical office RS, 2018. Average annual and monthly air temperatures (°C) by meteorological stations, Slovenia, annual data until 2014 - in Slovene (<http://pxweb.stat.si/pxweb/Dialog/Saveshow.asp>) [accessed February 6, 2018]
- Trasviña-Arenas, C.H., Garcia-Triana, A., Peregrino-Uriarte, A.B., Yepiz-Plascencia, G., 2013. White shrimp *Litopenaeus vannamei* catalase: Gene structure, expression and activity under hypoxia and reoxygenation. *Comparative Biochemistry & Physiology B*, 164, 44–52.
<https://doi.org/10.1016/j.cbpb.2012.10.004>
- Vinagre, C., Madeira, D., Narciso, L., Cabral, H.N., Diniz, M., 2012. Effect of temperature on oxidative stress in fish: Lipid peroxidation and catalase activity in the muscle of juvenile seabass, *Dicentrarchus labrax*. *Ecological Indicators*, 23, 274–279.
<https://doi.org/10.1016/j.ecolind.2012.04.009>
- Vranković, J., Borković-Mitić, S., Ilić, B., Radulović, M., Milošević, S., Makarov, S., Mitić, B., 2017. Bioaccumulation of metallic trace elements and antioxidant enzyme activities in *Apfelbeckia insculpta* (L. Koch, 1867) (Diplopoda: Callipodida) from the cave Hadži-Prodanova Pećina (Serbia). *International Journal of Speleology*, 46(1), 99–108.
<https://doi.org/10.5038/1827-806X.46.1.1981>
- Wojtal-Frankiewicz, A., Bernasińska, J., Frankiewicz, P., Gwoździński, K., Jurczak, T., 2017. The role of environmental factors in the induction of oxidative stress in zebra mussel (*Dreissena polymorpha*). *Aquatic Ecology*, 51, 289–306.
<https://doi.org/10.1007/s10452-017-9617-4>
- Zamocky, M., Furtmüller, P.G., Obinger, C., 2008. Evolution of catalases from bacteria to humans. *Antioxidants & Redox Signaling*, 10, 1527–1548.
<https://doi.org/10.1089/ars.2008.2046>
- Zou, E., Du, N., Ali, W., 1996. The effects of severe hypoxia on lactate and glucose concentrations in the blood of the Chinese freshwater crab *Eriocheir sinensis* (Crustacea: Decapoda). *Comparative Biochemistry & Physiology A*, 114, 105–109.
[https://doi.org/10.1016/0300-9629\(95\)02101-9](https://doi.org/10.1016/0300-9629(95)02101-9)
- Žagar, A., Simčič, T., Carretero, M.A., Vrezec, A., 2015. The role of metabolism in understanding the altitudinal segregation pattern of two potentially interacting lizards. *Comparative Biochemistry & Physiology A*, 179, 1–6.
<https://doi.org/10.1016/j.cbpa.2014.08.018>