

2020

A trace bound for integer-diagonal positive semidefinite matrices

Lon Mitchell

University of South Florida St. Petersburg, lonmitchell@usf.edu

Follow this and additional works at: https://digitalcommons.usf.edu/fac_publications

 Part of the [Mathematics Commons](#)

Recommended Citation

Mitchell, L. (2020). A trace bound for integer-diagonal positive semidefinite matrices. *Special Matrices*, 8(1), 14–16. <https://doi.org/10.1515/spma-2020-0002>

This Article is brought to you for free and open access by the USF Faculty Publications at Digital Commons @ University of South Florida. It has been accepted for inclusion in USF St. Petersburg campus Faculty Publications by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact scholarcommons@usf.edu.

Note

Open Access

Lon Mitchell*

A trace bound for integer-diagonal positive semidefinite matrices

<https://doi.org/10.1515/spma-2020-0002>

Received September 27, 2019; accepted December 2, 2019

Abstract: We prove that an n -by- n complex positive semidefinite matrix of rank r whose graph is connected, whose diagonal entries are integers, and whose non-zero off-diagonal entries have modulus at least one, has trace at least $n + r - 1$.

Keywords: positive semidefinite matrices, integer-diagonal, trace

MSC: 15B48, 15A15, 05C50

1 Introduction

The graph of an n -by- n Hermitian matrix $M = (m_{ij})$ has vertex set $\{1, 2, \dots, n\}$ and edge set $\{ij \mid i < j, m_{ij} \neq 0\}$. As part of their work on the Schur-Siegel-Smyth problem for totally positive algebraic integers, James McKee and Pavlo Yatsyna [3] proved that an n -by- n positive definite matrix S whose entries are integers and whose graph is connected must have trace at least $2n - 1$. As a consequence, 2 is the smallest limit point of the absolute trace (which for an n -by- n matrix is the trace divided by n) of such matrices.

The integer entries are important to McKee and Yatsyna's proof: since S is positive definite, it can be factored as $S = B^T B$, and thus viewed as the Gram matrix of the columns x_1, x_2, \dots, x_n of B . In a minimal-trace connected counterexample, we can assume without loss of generality that x_1 is a unit vector. Then the Gram matrix of $x_1, x'_2, x_3, \dots, x_n$, where $x'_2 = x_2 - (x_1^T x_2)x_1$, still has integer entries and eventually provides a contradiction.

Are the integer entries necessary? In this note, we prove a generalization for complex positive semidefinite matrices and show that while the diagonal entries must be integers, the off-diagonal non-zero entries need only have modulus at least 1. A generalization of McKee and Yatsyna's absolute trace result follows as a corollary.

In addition to standard tools and definitions from matrix analysis [2] and graph theory [1], one fact we will use repeatedly is that, because the sum of a positive definite matrix and a positive semidefinite matrix is still positive definite, adding a positive number to a diagonal entry of a positive definite matrix results in another positive definite matrix. Also note that an empty graph on a single vertex is connected.

2 New Results

Lemma 1. An n -by- n complex positive definite matrix whose graph is a tree, whose diagonal entries are integers, and whose non-zero off-diagonal entries have modulus at least one, has trace at least $2n - 1$.

*Corresponding Author: Lon Mitchell: Lon Mitchell, University of South Florida St. Petersburg, Florida, USA, E-mail: lon-mitchell@usfsp.edu

Proof. Proceed by induction on n , noting the result is true for $n = 1$. Assume that the result is true for all k -by- k matrices where $1 \leq k < n$, and let $M = (m_{ij})$ be an n -by- n positive definite matrix whose graph is a tree with vertices labeled v_1, v_2, \dots, v_n corresponding to the rows of M . Assume for the sake of eventual contradiction that the trace of M , $\text{tr } M$, is less than $2n - 1$.

Since the graph G of M is a tree, it has a pendant vertex (a vertex of degree one). Without loss of generality, we can assume vertex v_1 has unique neighbor v_2 . If the diagonal element m_{11} of M is greater than 1, then applying the induction hypothesis to M_{11} , the matrix obtained from M by deleting the first row and column, yields $\text{tr } M \geq 2 + \text{tr } M_{11} \geq 2 + 2(n-1) - 1 = 2n - 1$, a contradiction. So we may assume $m_{11} = 1$. Since v_1 is pendant,

$$M = \begin{bmatrix} 1 & \bar{\alpha}e_1^* \\ \alpha e_1 & M_{11} \end{bmatrix},$$

where α is a complex number with $|\alpha| \geq 1$ and e_1 is the standard basis vector.

Consider next the Schur complement $M' = M_{11} - |\alpha|^2 e_1 e_1^T$, which is an $(n-1)$ -by- $(n-1)$ positive definite matrix. All off-diagonal elements of M' remain unchanged from the corresponding entries of M , so the graph of M' is a tree. All main-diagonal elements of M' also remain unchanged with the exception of $m'_{11} = m_{22} - |\alpha|^2 \leq m_{22} - 1$.

Since m'_{11} may not be an integer, let M'' be the matrix obtained from M' by replacing m'_{11} with $m''_{11} = m_{22} - 1$. Since $m''_{11} \geq m'_{11}$, M'' is also positive definite. Further, its graph is a tree, its diagonal entries are integers, and its non-zero off-diagonal entries have modulus at least one. Finally, $\text{tr } M'' = \text{tr } M - 2 < 2n - 3$, a contradiction of the induction hypothesis. Thus $\text{tr } M \geq 2n - 1$. \square

Theorem 1. An n -by- n complex positive definite matrix whose graph is connected, whose diagonal entries are integers, and whose non-zero off-diagonal entries have modulus at least one, has trace at least $2n - 1$.

Proof. Proceed by induction on n , noting the result is true for $n = 1$. Assume that the result is true for all k -by- k matrices where $1 \leq k \leq n - 1$, and let M be an n -by- n positive definite matrix whose diagonal entries are integers, whose graph is connected, and whose non-zero off-diagonal entries have modulus at least one.

Assume for the sake of eventual contradiction that $\text{tr } M < 2n - 1$. By adding to a diagonal entry if needed, we can assume that we have a matrix M with the above-mentioned properties and with $\text{tr } M = 2n - 2$.

Let G be the graph of M and let m_v be the diagonal entry of M corresponding to vertex v in G . For each vertex v of G , let $c(v)$ be the number of connected components of $G \setminus v$.

Suppose first that there is a vertex v of G such that $m_v > c(v)$. Consider $M(v)$, the matrix obtained from M by removing the row and column corresponding to v . Applying the induction hypothesis to the principal submatrices $M_1, M_2, \dots, M_{c(v)}$ of $M(v)$ corresponding to the connected components $C_1, C_2, \dots, C_{c(v)}$ of $G \setminus v$, we find that

$$\begin{aligned} \text{tr } M &= m_v + \text{tr } M(v) = m_v + \sum_{i=1}^{c(v)} \text{tr } M_i \\ &\geq m_v + \sum_{i=1}^{c(v)} (2|C_i| - 1) = m_v - c(v) + 2(n-1) \geq 2n - 1. \end{aligned}$$

Thus we must have that $m_v \leq c(v)$ for each vertex v .

Let T be a spanning tree of G . Since T is a tree on n vertices, it has $n - 1$ edges, and so

$$\sum_{v \in G} d_T(v) = 2(n-1)$$

where $d_T(v)$ is the degree of the vertex v in T . Since $d_T(v) \geq c(v) \geq m_v$ for each v but

$$\sum_{v \in G} d_T(v) = 2(n-1) = \text{tr } M = \sum_{v \in G} m_v,$$

we must have $d_T(v) = c(v) = m_v$ for each v .

For any vertex v of G , because $d_T(v) = c(v)$, there is a bijective correspondence between the neighbors of v in T and the connected components of $G \setminus v$. Thus, if vertices v_i and v_j are not adjacent in T , then they belong to different connected components of $G \setminus w$ for any vertex w on a path between them in T , and so are not adjacent in G either. So, in fact, $G = T$, and Lemma 1 requires $\text{tr } M \geq 2n - 1$, contradicting our earlier assumption. Thus $\text{tr } M \geq 2n - 1$. \square

Corollary 1. The smallest limit point of the set of absolute traces of matrices satisfying the conditions of Theorem 1 is 2.

Remark. The matrices $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\begin{bmatrix} 1.0 & 0.5 \\ 0.5 & 1.0 \end{bmatrix}$, and $\begin{bmatrix} 1.1 & 1.0 \\ 1.0 & 1.1 \end{bmatrix}$ show that none of the conditions of Theorem 1 can be removed.

Theorem 2. An n -by- n complex positive semidefinite matrix of rank r whose graph is connected, whose diagonal entries are integers, and whose non-zero off-diagonal entries have modulus at least one, has trace at least $n + r - 1$.

Proof. Proceed by induction on the nullity. The nullity zero case is Theorem 1. Assume the result is true for all nullities less than some $k > 0$. Let M be an n -by- n complex positive semidefinite matrix of nullity $k = n - r$ whose graph is connected, whose diagonal entries are integers, and whose non-zero off-diagonal entries have modulus at least one.

Consider M as the Gram matrix of linearly dependent vectors x_1, x_2, \dots, x_n in \mathbb{C}^n . Let l be such that x_l is in the span of the other vectors, and let y be a unit vector in \mathbb{C}^n orthogonal to each x_i . Then the Gram matrix M' of $x_1, x_2, \dots, x_{l-1}, x_l + y, x_{l+1}, \dots, x_n$ is equal to M except for an increase of 1 in the m_{ll} main-diagonal element, so its graph is connected, its diagonal entries are integers, and the non-zero off-diagonal entries have modulus at least one. The nullity of M' is $k - 1$, so by the induction hypothesis and by construction, $\text{tr } M = \text{tr } M' - 1 \geq (n + (r + 1) - 1) - 1 = n + r - 1$. \square

Corollary 2. An n -by- n complex positive semidefinite matrix of rank r whose graph has s connected components, whose diagonal entries are integers, and whose non-zero off-diagonal entries have modulus at least one, has trace at least $n + r - s$.

Acknowledgement: Publication of this article was funded by the University of South Florida St. Petersburg's Open Access Publication Fund.

References

- [1] Reinhard Diestel, *Graph theory*, 5th ed., Graduate Texts in Mathematics, 173, Springer, Berlin, 2018.
- [2] Roger A. Horn, Charles R. Johnson, *Matrix analysis*, 2nd ed., Cambridge University Press, Cambridge, 2013.
- [3] James McKee, Pavlo Yatsyna, *A trace bound for positive definite connected integer symmetric matrices*, Linear Algebra Appl., 444, 227–230, 2014.