November 2018

U–Pb-dated flowstones restrict South African early hominin record to dry climate phases

Robyn Pickering

Andy I.R. Herries

Jon D Woodhead

John C. Hellstrom

Helen E. Green

See next page for additional authors

Follow this and additional works at: https://digitalcommons.usf.edu/kip_articles

Recommended Citation

Pickering, Robyn; Herries, Andy I.R.; Woodhead, Jon D; Hellstrom, John C.; Green, Helen E.; Paul, Bence; Ritzman, Terrence; Strait, David S.; Schoville, Benjamin J.; and Hancox, Phillip J., "U–Pb-dated flowstones restrict South African early hominin record to dry climate phases" (2018). *KIP Articles*. 5511.
https://digitalcommons.usf.edu/kip_articles/5511

This Article is brought to you for free and open access by the KIP Research Publications at Digital Commons @ University of South Florida. It has been accepted for inclusion in KIP Articles by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

Creator

Robyn Pickering, Andy I.R. Herries, Jon D Woodhead, John C. Hellstrom, Helen E. Green, Bence Paul, Terrence Ritzman, David S. Strait, Benjamin J. Schoville, and Phillip J. Hancox

U–Pb–dated flowstones restrict South African early hominin record to dry climate phases

Robyn Pickering^{1,2*}, Andy I. R. Herries^{3,4}, Jon D. Woodhead⁵, John C. Hellstrom⁵, Helen E. Green⁵, Bence Paul⁵, Terrence Ritzman^{2,6,7}, David S. Strait⁷, Benjamin J. Schoville^{2,8} & Phillip J. Hancox⁹

The Cradle of Humankind (Cradle) in South Africa preserves a rich collection of fossil hominins representing *Australopithecus*, *Paranthropus* and *Homo*¹. The ages of these fossils are contentious^{2–4} and have compromised the degree to which the South African hominin record can be used to test hypotheses of human evolution. However, uranium–lead (U–Pb) analyses of horizontally bedded layers of calcium carbonate (flowstone) provide a potential opportunity to obtain a robust chronology⁵. Flowstones are ubiquitous cave features and provide a palaeoclimatic context, because they grow only during phases of increased effective precipitation^{6,7}, ideally in closed caves. Here we show that flowstones from eight Cradle caves date to six narrow time intervals between 3.2 and 1.3 million years ago. We use a kernel density estimate to combine 29 U–Pb ages into a single record of flowstone growth intervals. We interpret these as major wet phases, when an increased water supply, more extensive vegetation cover and at least partially closed caves allowed for undisturbed, semi-continuous growth

of the flowstones. The intervening times represent substantially drier phases, during which fossils of hominins and other fossils accumulated in open caves. Fossil preservation, restricted to drier intervals, thus biases the view of hominin evolutionary history and behaviour, and places the hominins in a community of comparatively dry-adapted fauna. Although the periods of cave closure leave temporal gaps in the South African fossil record, the flowstones themselves provide valuable insights into both local and pan-African climate variability.

The early hominin fossil record in South Africa is best represented by deposits preserved in a series of dolomite caves 40 km northwest of Johannesburg (Fig. 1). These sites, which are collectively known as the Cradle of Humankind World Heritage Site, have produced hominin fossils attributed to at least five taxa: *Australopithecus africanus*, *Australopithecus sediba*, *Paranthropus robustus*, *Homo naledi*¹ and a poorly understood collection of early Pleistocene fossils that we refer to as ‘early *Homo*’. Historically, the hominin fossil record in South Africa

Fig. 1 | Map of the localities and analyses of the flowstone ages versus geographic variables. a, Topographical map showing major hominin localities in East and South Africa. **b**, The Cradle shown in detail with locations of the cave sites (in bold with filled circles, U–Pb ages). **c**, Photograph of Bolt's Farm deposits shows stacking of fossil-bearing

sediments and thick flowstone layers. **d**, U–Pb flowstone ages plotted against site elevation, latitude and longitude reveal no simple relationship, suggesting these factors are not forcing the mode of cave deposition (sediment or flowstone). $n = 29$; diamonds, individual ages; whisker, 2σ errors.

¹Department of Geological Sciences, University of Cape Town, Cape Town, South Africa. ²Human Evolution Research Institute, University of Cape Town, Cape Town, South Africa. ³The Australian Archaeomagnetism Laboratory, Department of Archaeology and History, La Trobe University, Melbourne, Victoria, Australia. ⁴Centre for Anthropological Research, University of Johannesburg, Johannesburg, South Africa. ⁵School of Earth Sciences, The University of Melbourne, Melbourne, Victoria, Australia. ⁶Department of Neuroscience, Washington University School of Medicine, St. Louis, MO, USA. ⁷Department of Anthropology, Washington University in St. Louis, St. Louis, MO, USA. ⁸School of Social Science, University of Queensland, Brisbane, Queensland, Australia. ⁹Evolutionary Science Institute, University of the Witwatersrand, Johannesburg, South Africa. *e-mail: robyn.pickering@uct.ac.za

Table 1 | All U–Pb ages and site information for the Cradle caves

Site	Sample	U–Pb age		U–Pb age		%	Stratigraphy	
		²⁰⁴ Pb	2σ	²⁰⁶ Pb	2σ			
Bolt's Farm	WP160-6L	ND	ND	1.752	0.149	8.5	WP160	Top flowstone
	WP160.2	ND	ND	2.269	0.075	3.3	WP160	Basal flowstone
	BFMC-6	ND	ND	1.383	0.050	3.6	Pit 7	Top flowstone
	BFMC-1	ND	ND	1.776	0.073	4.1	Pit 7	Base of middle flowstone
	AV03	ND	ND	2.668	0.304	11.4	Pit 14	Top flowstone
Cooper's ¹⁵	CD-1	1.526	0.088	1.375	0.113	8.2	Unit 1	Basal flowstone
Drimolen	DN09	ND	ND	1.789	0.104	5.8	Main quarry	Top flowstone
	DN26	ND	ND	1.962	0.107	5.5	Main quarry	FS in middle of seds
	DN39A	ND	ND	2.673	0.103	3.9	Main quarry	Basal flowstone
	DMK5 ²⁷	ND	ND	2.664	0.392	14.7	Makondo	Basal flowstone
Haasgat	HG1	ND	ND	1.686	0.236	14.0		Middle flowstone
Hoogland	HL1	ND	ND	3.145	0.243	7.7		Basal flowstone
Malapa ^{15,20}	M-1	ND	ND	2.062	0.021	1.0	Pit 1	Basal flowstone
	M6	ND	ND	2.048	0.140	6.8	Pit 2	Top flowstone Pit 2
	M5	ND	ND	2.067	0.161	7.8	Pit 1	Top flowstone in Pit 1
Sterkfontein ^{5,14,15}	OE-13	1.810	0.060	1.784	0.090	5.0	Member 5	Base of MB5
	OE-14	2.014	0.055	2.030	0.061	3.0	Member 4	FS capping open air MB4
	BH4-9	2.650	0.300	2.645	0.183	6.9	Member 4	FS at base of MB4 in borehole 4
	BH1-8	2.830	0.344	2.800	0.140	5.0	Member 4	FS at base of MB4 in borehole 1
	BH1-15	2.800	0.280	2.747	0.172	6.3	Member 2	FS at base of borehole 1
	SB-1	2.347	0.101	2.275	0.176	7.7	Silberberg	FS below STW573
	STA09	2.170	0.065	ND	ND	ND	Silberberg	FS2C, associated with STW573
	STA12	2.110	0.060	ND	ND	ND	Silberberg	FS2C, associated with STW573
	SKA3	2.250	0.075	ND	ND	ND	Silberberg	FS2C, associated with STW573
	STA15	2.240	0.080	ND	ND	ND	Silberberg	FS2B, below STW537
Swartkrans ¹⁵	SWK-9	ND	ND	1.706	0.069	4.0	Member 1	FS capping Lower Bank
	SWK-5	ND	ND	1.800	0.005	0.3	Member 1	FS capping Hanging Remnant
	SWK-7	ND	ND	2.248	0.052	2.3	Member 1	FS base Hanging Remnant
	SWK-12	ND	ND	2.249	0.077	3.4	Member 1	FS base Lower Bank

²⁰⁴Pb and ²⁰⁶Pb, U–Pb ages determined using ²⁰⁴Pb and ²⁰⁶Pb, respectively. *n* = 29, errors on individual ages at 2σ; new ages in bold. Ages are given as Ma. FS, flowstone; seds, sediments. ND, not determined. Indicated data obtained from previous studies^{5,14,15,20,27}.

featured prominently in hypotheses about early hominin diversity, biogeography and evolutionary history, many of which posit a causal relationship between changing climatic conditions and human evolution^{8,9}. However, testing these hypotheses has been hampered by the imprecision of and disagreement about the ages of the deposits, such that more recent assessments of climate forcing in early human evolution have focused on the East African record^{10–13}. The paucity of absolute ages for the Cradle sites and lack of correlation between geological members in different caves has prevented the South African record from entering these discussions. Direct dating of the cave deposits^{2,5,14,15} is changing this picture.

Cave carbonates, or speleothems, are ubiquitous features at the Cradle caves and hold the key to understanding the ages of the fossils and the formational history of sites. Speleothems form from drip waters that percolate through the bedrock into caves, which if completely, or at least partially, closed to incoming sediment, can lead to the accumulation of horizontally bedded layers of flowstone that are up to several metres thick. The dominant control over speleothem formation is a positive water balance, driven by an increase in regional effective precipitation⁶. Therefore, the presence of speleothems, particularly in the form of massive flowstone layers, is indicative of increased cave drip water and more broadly reflects wetter conditions outside the cave. The onset, and termination, of a flowstone is therefore indicative of the crossing of a considerable threshold in the surface hydroclimate. Caves are well-known to be dynamic systems, often subject to multiple opening and closure events^{6,16}. In the Cradle context, increased effective

precipitation, coupled with a strong decrease in the flux of externally derived clastic sediment into the caves (that is, completely or partially closed caves) leads to flowstone formation⁷. A shift to a drier hydroclimate, with less vegetation, more open environments and increased surface erosion, favours the opening up of caves and deposition of sediment within them^{7,16}. The latter also explains the cessation of major flowstone formation during these sedimentation intervals. Given the apparent climatic forcing of these two modes of cave deposition, we argue that they are mutually exclusive, meaning that it is unlikely that sediment and flowstone formation occurs concomitantly. The Cradle is a small area (approximately 10 × 15 km²; Fig. 1), so shifts in local climate conditions should be simultaneous, with flowstone forming in different caves synchronously. Indeed, all Cradle sites preserve alternating stacks of flowstones and sediment (Extended Data Fig. 1), indicating that conditions conducive to the formation of both these deposits occurred repeatedly. Here we argue for a simple, binary, Cradle-wide control on these sedimentary facies, with caves being either open (accumulating sediments) or closed (growing flowstone), driven by changes in the hydroclimate.

The U–Pb method for dating speleothems¹⁷ has progressed in the last decade and now allows routine precise age determination of materials that are a few hundred thousand years to hundreds of millions of years old. Cradle flowstone U–Pb ages (Table 1) have uncertainties of at best around 1% (±20 thousand years on a date of 2.062 million years ago (Ma)), or at worst 15% (±390 thousand years on a date of 2.664 Ma), making them comparable to Ar–Ar ages used to date East African

Fig. 2 | U-Pb chronology of the Cradle. **a**, **b**, U-Pb ages plotted against time and by site (**a**), and summed together into a histogram and kernel density estimate (**b**). $n = 29$; diamonds, individual ages; whiskers, 2σ errors (**a**); open circles, individual ages (**b**). Peaks of the kernel density estimates represent FGIs (1–6, from oldest to youngest); troughs, phases of open caves with sediment- and fossil-accumulation (SED1–SED6). **c**, Cave

fills dating to each sediment- and fossil-accumulation interval are noted; proposed age ranges for the Cradle hominins are included. **d**, No simple relationship between the flowstone record and the stacked, global benthic $\delta^{18}\text{O}$ record²⁸. Numbers in brackets, number of flowstones dated at each cave site. MB, member. ? denotes age uncertainties.

hominin sites, which typically have uncertainties¹⁸ of 1–2%. Even with this precision, attributing individual flowstones to known records of climate variability is almost impossible, because the age uncertainties are greater than the period of climatic fluctuation. Therefore, we use a kernel density estimate¹⁹ to sum the Cradle-wide dataset of 29 U-Pb ages and their associated individual errors to produce a single, composite record of flowstone growth intervals (FGIs) (Fig. 2). Our record is one of the most complete chronologies for the Cradle produced by a single method and contextualizes the caves as localities within a single evolving landscape. We can now investigate the duration and frequency of the FGIs, constrain the intervening periods during which caves were open to receive sediments and bones, and correlate the fossil-bearing deposits from different caves (see Supplementary Video 1).

We acknowledge that not all of the Cradle caves were dated in this study; while material older than 3.2 Ma may exist, we have dated the basal flowstones at each site, suggesting that the major time of Cradle flowstone and sediment accumulation occurred between 3.2 and 1.3 Ma (Fig. 2). This range is not compatible with the cosmogenic nuclide burial date estimate of approximately 3.6 Ma for the hominin fossil STW573³, and we favour the reinterpretation of this burial date to around 2.8 Ma⁴ or possibly even more recent, given the reversed palaeomagnetic signal of the flowstone². There are younger U-Th-dated cave deposits in the Cradle, at Gladysvale⁷, Rising Star²⁰ and Plover's Lake²¹,

but these represent minor deposits and thin flowstones that are only a few centimetres thick, compared to the major flowstones and thick sedimentary layers of the older deposits (Extended Data Fig. 1). This probably reflects an increase in aridity, whereas the thick flowstones of the terminal Pliocene and early Pleistocene indicate an overall wetter climate. We see no simple spatial relationship between the ages of major flowstones and longitude, latitude or elevation (Fig. 1). This indicates that cave location is not the dominant factor in determining the age of the deposits; we argue instead that a changing hydroclimate (repeated wet-dry cycles) provides a better explanation.

We identify six FGIs, which were numbered 1–6 from the earliest (3.19–3.08 Ma) to the most recent (1.41–1.32 Ma) period (Fig. 2), that represent wetter periods and correspond to predominantly closed caves. The amplitude of the kernel density estimate is a function of the number of flowstones that formed during a FGI; the intervals during which the most flowstone was deposited, FGI3 (2.28–2.17 Ma) and FGI5 (1.82–1.63 Ma), consisted of 13 and 7 flowstones forming in five separate caves, respectively (Fig. 2). These extended periods of flowstone formation support our model, which predicts that all of the Cradle caves experienced the same external climatic conditions. Our model predicts that the periods between successive FGIs (3.08–2.83, 2.62–2.29, 2.17–2.12, 2.00–1.82, 1.63–1.41 and less than 1.32 Ma) were drier times during which the fossil-bearing clastic sedimentary

units (Fig. 2) accumulated in open caves. We argue that the entire early Cradle fossil record is restricted to these limited time intervals. Comparing these intervals to previously published ages for the fossils and their surrounding sediments can test this hypothesis; there is good corroboration overall^{2,4,22,23} (see Supplementary Information).

Our results have several implications for the interpretation of the South African hominin fossil record. First, the record is discontinuous, with substantial gaps represented by FG11–FG16. These discontinuities suggest that any anagenetic change within hominin lineages across sedimentary periods will appear punctuated. Similarly, gradual trends in faunal extinction or speciation will appear as sudden, correlated changes. This makes it impossible to falsify hypotheses of punctuated equilibrium²⁴ and turnover pulses⁹. Moreover, our ability to observe pivotal milestones that pertain to the origin of *Homo* and advances in tool technology are temporally restricted. Second, the record is biased towards representing drier-adapted plant and animal communities. Although palaeoenvironments may have shifted on average from more mesic to more arid over the time period during which the Cradle sites were accumulating sediments²⁵, the wettest periods are still missing as the caves were closed during speleothem formation. This bias is likely to be manifested in direct measures of hominin behaviour, such as dental microwear, phytoliths that are preserved in dental calculus and isotopes. Moreover, the inability to observe behaviours during wet periods constrains our ability to evaluate hypotheses of hominin adaptation using the Cradle record. Third, Plio-Pleistocene South Africa evidently experienced marked climatic cyclicity over timescales that cannot easily be explained by insolation due to Milankovitch cycles. These wet and dry periods do not obviously correspond to climate cycles in East Africa (Extended Data Fig. 2). Future assessments of hominin adaptations and palaeobiology need to account for this complexity. Moreover, climatic cyclicity has important implications for the biogeography of hominins and other mammals insofar as habitat theory²⁶ predicts that as vegetation zones shift, so will the faunal communities that access those habitats. Lastly, some hominin taxa (for example, *A. africanus*, *P. robustus* and early *Homo*) are found during dry periods that unequivocally straddle wet periods, indicating either that these species were ecological generalists or that they vacated the Cradle landscape during wet periods only to return at a later, drier time.

Although the closed caves produce unresolvable gaps in the hominin record, the compensation is that the flowstones themselves have enormous potential as archives of both local and pan-African palaeoclimate variation. The work presented here correlates sedimentary units across sites, establishes age ranges for hominins, facilitates comparisons with the East African record and provides a single, independent chronological framework upon which data can judiciously be leveraged to answer the questions that are central to the study of early human evolution.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, statements of data availability and associated accession codes are available at <https://doi.org/10.1038/s41586-018-0711-0>.

Received: 3 April 2018; Accepted: 12 September 2018;

Published online 21 November 2018.

1. Wood, B. & Boyle, E. K. Hominin taxic diversity: fact or fantasy? *Am. J. Phys. Anthropol.* **159**, 37–78 (2016).
2. Herries, A. I. R. & Shaw, J. Palaeomagnetic analysis of the Sterkfontein palaeocave deposits: implications for the age of the hominin fossils and stone tool industries. *J. Hum. Evol.* **60**, 523–539 (2011).
3. Granger, D. E. et al. New cosmogenic burial ages for Sterkfontein Member 2 *Australopithecus* and Member 5 Oldowan. *Nature* **522**, 85–88 (2015).
4. Kramers, J. D. & Dirks, P. H. G. M. The age of fossil StW573 ('Little Foot'): an alternative interpretation of ²⁶Al/¹⁰Be burial data. *S. Afr. J. Sci.* **113**, 2016-0085 (2017).
5. Pickering, R., Kramers, J. D., Partridge, T., Kodolanyi, J. & Pettke, T. U–Pb dating of calcite–aragonite layers in speleothems from hominin sites in South Africa by MC-ICP-MS. *Quat. Geochronol.* **5**, 544–558 (2010).
6. Ayliffe, L. K. et al. 500 ka precipitation record from southeastern Australia: evidence for interglacial relative aridity. *Geology* **26**, 147–150 (1998).

7. Pickering, R. et al. Stratigraphy, U–Th chronology, and palaeoenvironments at Gladysvale Cave: insights into the climatic control of South African hominin-bearing cave deposits. *J. Hum. Evol.* **53**, 602–619 (2007).
8. Dart, R. *Australopithecus africanus*: the man-ape of South Africa. *Nature* **115**, 195–199 (1925).
9. Vrba, E. S. in *Hominid Evolution. Past, Present and Future* (ed. Tobias, P. V.) 195–200 (Alan R. Liss, New York, 1985).
10. Potts, R. Variability selection in hominid evolution. *Evol. Anthropol.* **7**, 81–96 (1998).
11. Behrensmeyer, A. K., Todd, N. E., Potts, R. & McBrinn, G. E. Late Pliocene faunal turnover in the Turkana basin, Kenya and Ethiopia. *Science* **278**, 1589–1594 (1997).
12. Trauth, M. H., Larrasoana, J. C. & Mudelsee, M. Trends, rhythms and events in Plio-Pleistocene African climate. *Quat. Sci. Rev.* **28**, 399–411 (2009).
13. deMenocal, P. B. Climate and human evolution. *Science* **331**, 540–542 (2011).
14. Walker, J., Cliff, R. A. & Latham, A. G. U–Pb isotopic age of the StW 573 hominid from Sterkfontein, South Africa. *Science* **314**, 1592–1594 (2006).
15. Pickering, R., Kramers, J. D., Hancox, P. J., de Ruiter, D. J. & Woodhead, J. D. Contemporary flowstone development links early hominin bearing cave deposits in South Africa. *Earth Planet. Sci. Lett.* **306**, 23–32 (2011).
16. Brain, C. K. in *Paleoclimate and Evolution, with Emphasis on Human Origins* (eds Vrba, E. S. G. et al.) 451–458 (Yale Univ. Press, New Haven, 1995).
17. Woodhead, J. et al. U–Pb geochronology of speleothems by MC-ICPMS. *Quat. Geochronol.* **1**, 208–221 (2006).
18. McDougall, I. et al. New single crystal ⁴⁰Ar/³⁹Ar ages improve time scale for deposition of the Omo Group, Omo–Turkana Basin, East Africa. *J. Geol. Soc. Lond.* **169**, 213–226 (2012).
19. Vermeech, P. On the visualisation of detrital age distributions. *Chem. Geol.* **312–313**, 190–194 (2012).
20. Dirks, P. H. G. M. et al. The age of *Homo naledi* and associated sediments in the Rising Star Cave, South Africa. *eLife* **6**, e24231 (2017).
21. de Ruiter, D. J., Brophy, J. K., Lewis, P. J., Churchill, S. E. & Berger, L. R. Faunal assemblage composition and palaeoenvironment of Plovers Lake, a Middle Stone Age locality in Gauteng Province, South Africa. *J. Hum. Evol.* **55**, 1102–1117 (2008).
22. Gibbon, R. J. et al. Cosmogenic nuclide burial dating of hominin-bearing Pleistocene cave deposits at Swartkrans, South Africa. *Quat. Geochronol.* **24**, 10–15 (2014).
23. Herries, A. I. R. et al. in *Paleobiology of Australopithecus (Vertebrate Paleobiology and Paleoanthropology Series)* (eds Reed, K. E. et al.) 21–40 (Springer Netherlands, Dordrecht, 2013).
24. Eldredge, N. & Gould, S. J. in *Models in Paleobiology* (ed. Schopf, T. J. M.) 82–115 (Freeman, Cooper & Co, San Francisco, 1972).
25. Vrba, E. S. in *Ancestors: The Hard Evidence* (ed. Delson, E.) 63–71 (Alan R. Liss, New York, 1985).
26. Vrba, E. S. Mammals as a key to evolutionary theory. *J. Mamm.* **73**, 1–28 (1992).
27. Herries, A. I. R. et al. Geochronological and 3D visualisation approaches for contextualising in-situ fossil bearing palaeokarst in South Africa: a case study from the ~2.61 Ma Drimolen Makondo. *Quat. Int.* **483**, 90–110 (2018).
28. Lisiecki, L. E. & Raymo, M. E. A Pliocene–Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography* **20**, PA1003 (2005).

Acknowledgements We thank D. Braun, R. Potts, B. Wood and W. L. S. Jor for their insightful discussion. Site access granted by C. Steininger, R. Clarke, T. Pickering, C. Menter, S. Potze, J. Adams and L. Berger; permits provided by South African Heritage Resource Agency. This work was supported by Australian Research Council DECRA DE120102504 (to R.P.), University of Melbourne McKenzie Post-Doctoral Fellowship 0023249 (to R.P.), Australian Research Council Future Fellowship FT120100399 (to A.I.R.H.) and Discovery Project DP170100056 (to A.I.R.H. and D.S.S.) and National Science Foundation Grant BCS 0962564 (to A.I.R.H.).

Reviewer information Nature thanks C. Feibel, T. Rasbury and the other anonymous reviewer(s) for their contribution to the peer review of this work.

Author contributions R.P. devised the project, conducted the U–Pb dating, drafted and revised the manuscript; A.I.R.H. assisted in the project design, fieldwork and site access; J.D.W., J.C.H., H.E.G. and B.P. assisted with U–Pb and ²³⁴U/²³⁸U analysis, data processing and discussion; T.R. and D.S.S. contributed to the hominin discussion; B.J.S. assisted with data visualization, including Supplementary Video 1 and discussion; P.J.H. contributed to the conceptual design of the project and provided field support. All authors contributed equally to the paper.

Competing interests The authors declare no competing interests.

Additional information

Extended data is available for this paper at <https://doi.org/10.1038/s41586-018-0711-0>.

Supplementary information is available for this paper at <https://doi.org/10.1038/s41586-018-0711-0>.

Reprints and permissions information is available at <http://www.nature.com/reprints>.

Correspondence and requests for materials should be addressed to R.P.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

METHODS

Flowstone samples were carefully selected from each field site based on a visual assessment of their level of preservation. The association of flowstones with fossil-bearing sediments was carefully noted and ideally flowstones above and below fossil-rich layers were sampled to provide bracketing ages (Extended Data Fig. 1). The petrography of all major flowstone layers was evaluated using thin sections and standard transmitted light microscopy. All samples were pre-screened for uranium (and in some cases lead) concentrations and distributions using either passive radiation imaging with a FUJIFilm BAS-1800 beta-scanner⁵ or in situ laser ablation ICP-MS (inductively coupled plasma mass spectrometer). Layers with at least 100 ng g⁻¹ of U were selected and up to 15 small 0.5-cm³ blocks were manually cut out using a handheld dentist drill. These blocks were etched in weak HCl to remove surface contamination and all subsequent handling was performed in a Class 350 clean laboratory. Samples were spiked with a mixed ²³⁵U–²⁰⁵Pb tracer, and ion chromatography column chemistry was used to separate and concentrate U and Pb before measurement by MC-ICP-MS (multi-collector inductively coupled plasma mass spectrometer), following previously published protocols¹⁷. ²³⁴U/²³⁸U was similarly determined from separate sample dissolutions using established protocols²⁹.

Instrumental mass bias effects were monitored and corrected using NIST SRM 981 reference material in the case of Pb, and the sample's internal (¹³⁷.88) ²³⁸U/²³⁵U ratio for U. Instrument data files were processed initially using an in-house-designed importer, operating within the Iolite environment³⁰, which considers all data and reference material analyses obtained throughout a particular analytical session and permits a variety of corrections for instrumental mass bias and drift. The resulting data, now corrected for instrumental effects, were then blank-corrected and isotope-dilution calculations performed using previously described³¹ software.

Age plots were generated using Tera–Wasserberg constructs; the slope of this line and its intercept with an iteratively calculated disequilibrium concordia derived from measured ²³⁴U/²³⁸U values were used to calculate a final age¹⁷. All errors are quoted as $\pm 2\sigma$.

Our 25 U–Pb ages and four ages that have previously been published¹⁴ were summed together into a simple frequency histogram and a kernel density estimate curve, with a linear transformation, a bandwidth of 0.03 and 45 bins, using a previously published program¹⁹. There are five new U–Pb ages from Bolt's Farm, four from Drimolen²⁷ and one each from Haasgat, Hoogland and Malapa (Table 1 and Supplementary Table 1). Flowstone U–Pb ages from Sterkfontein, Swartkrans, Cooper's and Malapa have previously been published^{5,14,15,29,32,33}, but here we recalculate the ²³⁸U–²⁰⁴Pb ages for Sterkfontein and Cooper's using Tera–Wasserberg concordia plots to avoid using isotope ratios that include ²⁰⁴Pb and we therefore improve the precision by up to 50%.

We created a video (Supplementary Video 1) as a visualization of the age data on the landscape through time. We used ESRI ArcMap 10.4 (<http://desktop.arcgis.com/en/arcmap>) and Filmora v.8.6.1 (<https://filmora.wondershare.net/video-editor>) software. The underlying digital elevation model in the video was generated in ArcMap using the US Geological Survey 1-arcsec (30-m) Shuttle Radar Topography Mission dataset (<https://lta.cr.usgs.gov/SRTM1Arc>). The underlying

digital elevation model in Fig. 1 was generated in ArcMap using the global 30-arcsec digital elevation model (1 km, <https://lta.cr.usgs.gov/GTOPO30>). The site abbreviations are: BF, Bolt's Farm; CD, Cooper's Cave; DM, Drimolen; GND, Gondolin; GV, Gladysvale; HG, Haasgat; HL, Hoogland; KD, Kromdraai; MP, Malapa; RS, Rising Star; STK, Sterkfontein; SWK, Swartkrans. The animation proceeds in 1,000-year time intervals, starting at 3.2 Ma, using the Time Slider control feature in ArcGIS. When the displayed time interval contains the 2σ age of a sampled speleothem, a blue marker is displayed at that sample location. The size of the marker is relative to the magnitude of the error estimate on the sample, such that samples with smaller errors appear as larger circles than samples with larger errors. The colour of the marker is relative to the proximity of the displayed time slice to the mean age of the sample, such that the marker is displayed as the darkest shade of blue when the map is displaying a time slice that contains the mean age of the sample. The marker colour is progressively lighter towards the tail ends of the age distribution before and after the mean age (up to 2σ). The marker-shape file is displayed with 50% transparency to allow multiple samples from the same site to be visible. The time intervals containing wetter FGI1–6 and drier sedimentary units (SEDs) 1–6 are also displayed.

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this paper.

Data availability

The authors declare that all data supporting the findings of this study are available within the paper and the Supplementary Information (see Supplementary Information and Supplementary Table 1).

29. Pickering, R. et al. *Australopithecus sediba* at 1.977 Ma and implications for the origins of the genus *Homo*. *Science* **333**, 1421–1423 (2011).
30. Paton, C., Hellstrom, J., Paul, B., Woodhead, J. & Hergt, J. Iolite: freeware for the visualisation and processing of mass spectrometric data. *J. Anal. At. Spectrom.* **26**, 2508–2518 (2011).
31. Schmitz, M. D. & Schoene, B. Derivation of isotope ratios, errors, and error correlations for U–Pb geochronology using ²⁰⁵Pb–²³⁵U–(²³³U)-spiked isotope dilution thermal ionization mass spectrometric data. *Geochim. Geophys. Geosyst.* **8**, Q08006 (2007).
32. Dirks, P. et al. Geological setting and age of *Australopithecus sediba* from Southern Africa. *Science* **328**, 205–208 (2010).
33. de Ruiter, D. J. et al. New *Australopithecus robustus* fossils and associated U–Pb dates from Cooper's Cave (Gauteng, South Africa). *J. Hum. Evol.* **56**, 497–513 (2009).
34. Wynn, J. G. Influence of Plio-Pleistocene aridification on human evolution: evidence from paleosols of the Turkana Basin, Kenya. *Am. J. Phys. Anthropol.* **123**, 106–118 (2004).
35. Trauth, M. H., Maslin, M. A., Deino, A. & Strecker, M. R. Late Cenozoic moisture history of east Africa. *Science* **309**, 2051–2053 (2005).
36. Maslin, M. A. & Trauth, M. H. in *The First Humans: Origins of the Genus Homo* (eds Grine, F. E. et al.) 151–158 (Springer Netherlands, Dordrecht, 2009).
37. Potts, R. & Faith, J. T. Alternating high and low climate variability: the context of natural selection and speciation in Plio-Pleistocene hominin evolution. *J. Hum. Evol.* **87**, 5–20 (2015).

Extended Data Fig. 1 | Field photographs showing U–Pb-dated flowstone from the indicated sites, all of which record some variation of an alternating stack of flowstones and fossil-bearing sediments. a, The basal flowstone from Cooper's Cave. b, c, Flowstones at Haasgat are preserved in the now-deroofed section of the deposits (b) and inside the cave (c). d, The flowstone capping the MB1 Lower Bank at Swartkrans.

e, f, Flowstone from Bolt's Farm Pit 7 at the base (e) and top (f) of the sequence. g, Flowstone capping Member 4 at Sterkfontein. h, Flowstone underlying fossil bearing sediments at Malapa. i, Flowstone sandwiched between fossil-bearing sediments at Drimolen. j, Massive flowstone at the base of Member 4 at Sterkfontein is exposed in a borehole. k, Flowstone underlying fossil-bearing sediments at Hoogland.

Extended Data Fig. 2 | U-Pb ages plotted against time and by site, additional un-U-Pb dated Cradle sites and non-Cradle hominin cave sites included. a–e, A variation of Fig. 2. U-Pb ages of Cradle sites are shown, with Cradle sites not dated with U-Pb (Gondolin and Kromdraai) included, as well as the non-Cradle hominin cave sites (Makapansgat and Taung) shown for comparison. All U-Pb ages are plotted against time and by site, $n = 29$, diamonds represent individual ages and 2σ errors

are shown as whiskers. Also included here are four records of climate and variability derived from orbital parameters for East African sites (d), specifically arid phases from soil carbonates³⁴, periods of deep rift valley lakes³⁵, phases of extreme climate variability³⁶ and key phases of variability as described previously³⁷. Again, there is no clear relationship between these records and the new South African flowstone record. Indicated data were obtained from previous studies^{28,34–37}.

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see [Authors & Referees](#) and the [Editorial Policy Checklist](#).

Statistical parameters

When statistical analyses are reported, confirm that the following items are present in the relevant location (e.g. figure legend, table legend, main text, or Methods section).

n/a | Confirmed

- ☐ ☒ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☐ ☒ An indication of whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☒ ☐ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☒ ☐ A description of all covariates tested
- ☒ ☐ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☐ ☒ A full description of the statistics including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☒ ☐ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☒ ☐ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated
- ☐ ☒ Clearly defined error bars
State explicitly what error bars represent (e.g. SD, SE, CI)

Our web collection on [statistics for biologists](#) may be useful.

Software and code

Policy information about [availability of computer code](#)

Data collection

U and Pb isotope data were collected using protocols outlines in Woodhead et al (2006) and Pickering et al. (2010). In more detail, isotope ratios were collected using the machine software that comes standard with Nu Instruments MC-ICP-MS, Pb isotopes were corrected using Lolite (Paton et al., 2011), spike corrections done using Schmitz and Schoene (2007).

Paton, C., Hellstrom, J., Paul, B., Woodhead, J. & Hergt, J. Lolite: Freeware for the Visualisation and Processing of Mass Spectrometric Data. Vol. 26 (12) 2508-2518 (2011).

Schmitz, M. D. & Schoene, B. Derivation of isotope ratios, errors, and error correlations for U-Pb geochronology using 205Pb-235U-(233U)-spiked isotope dilution thermal ionization mass spectrometric data. *Geochemistry, Geophysics, Geosystems* 8, doi:Q08006 10.1029/2006GC001492. (2007).

Woodhead, J. et al. U-Pb geochronology of speleothems by MC-ICPMS. *Quaternary Geochronology* 1, 208-221, doi:10.1016/j.quageo.2006.08.002 (2006).

Pickering, R., Kramers, J. D., Partridge, T., Kodolanyi, J. & Pettke, T. U-Pb dating of calcite–aragonite layers in speleothems from hominin sites in South Africa by MC-ICP-MS. *Quaternary Geochronology* 5, 544-558,

Data analysis

The 29 U-Pb ages were summed together into a single record using a kernel density estimate - we used the free software by Vermeesch (2012).
 Vermeesch, P. On the visualisation of detrital age distributions. *Chemical Geology* 312–313, 190–194 (2012).
 Our data visualisation, presented as supplementary video 1, was created using ESRI ArcGIS 10.4 and Filmora v8.6.1

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors/reviewers upon request. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Research [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A list of figures that have associated raw data
- A description of any restrictions on data availability

All isotope concentration and ratio data needed to calculate U-Pb ages are provided in extended data Table 1.

There are no restrictions on data availability

Field-specific reporting

Please select the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☐ Behavioural & social sciences ☒ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/authors/policies/ReportingSummary-flat.pdf](https://www.nature.com/authors/policies/ReportingSummary-flat.pdf)

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description

Uranium-lead dating of cave carbonates from hominin fossil bearing caves in South Africa. Individual U-Pb ages (n=29) summed together with a kernel density estimate. One replicate was run, SB1 and STA15 were taken from the same flowstone.

Research sample

Based on my field experience, flowstone rock samples were selected and collected from 8 cave locations in the Cradle of Humankind, South Africa. The sites were chosen based on the importance of the associated fossil record and ease of access/permission to sample. We present 29 U-Pb ages. 13 of these ages are new, 16 previously published but recalculated here. Previous papers include:
 de Ruiter, D. J. et al. New Australopithecus robustus fossils and associated U-Pb dates from Cooper's Cave (Gauteng, South Africa). *Journal Of Human Evolution* 56, 497–513, doi:10.1016/j.jhevol.2009.01.009 (2009).
 Dirks, P. et al. Geological Setting and Age of Australopithecus sediba from Southern Africa. *Science* 328, 205–208, doi:10.1126/science.1184950 (2010).
 Pickering, R. et al. Australopithecus sediba at 1.977 Ma and Implications for the Origins of the Genus Homo. *Science* 333, 1421–1423, doi:10.1126/science.1203697 (2011).
 Pickering, R. & Kramers, J. D. A re-appraisal of the stratigraphy and new U-Pb dates at the Sterkfontein hominin site, South Africa. *Journal Of Human Evolution* 59, 70–86 (2010).
 Pickering, R., Kramers, J. D., Hancox, P. J., de Ruiter, D. J. & Woodhead, J. D. Contemporary flowstone development links early hominin bearing cave deposits in South Africa. *Earth and Planetary Science Letters* 306, 23–32 (2011).
 Pickering, R., Kramers, J. D., Partridge, T., Kodolanyi, J. & Pettke, T. U-Pb dating of calcite–aragonite layers in speleothems from hominin sites in South Africa by MC-ICP-MS. *Quaternary Geochronology* 5, 544–558, doi:10.1016/j.quageo (2010).
 Herries, A.I.R. et al. Geoarchaeological and 3D visualisation approaches for contextualising in-situ fossil bearing palaeokarst in South Africa: A case study from the ~2.61 Ma Drimolen Makondo. *Quaternary International*, doi.org/10.1016/j.quaint.2018.01.001 (in press)
 Walker, J., Cliff, R. A. & Latham, A. G. U-Pb isotopic age of the StW 573 hominid from Sterkfontein, South Africa. *Science* 314, 1592–1594, doi:10.1126/science.1132916 (2006).

Sampling strategy

Not every single cave in the Cradle is represented here, but all major hominin bearing deposits (Sterkfontein, Swartkrans, Drimolen, Coopers, Malapa) and a few non-hominin caves are (Haasgat, Hoogland, Bolt's Farm). This is the largest geochronological collection ever for these sites. This is a geological study, our sample sizes are small, around 2 samples per site is normal.

Data collection

Careful fieldwork included noting the relationship between the flowstones samples for dating and the fossil bearing sediments in a field note book and with photographs. RP did the bulk of the sample collection herself with a geological hammer, with field assistance from AIRH. All U-Pb dating was done at the University of Bern and the University of Melbourne by RP.

Timing and spatial scale

Sample collection in the Cradle began in 2005 as part of RP's PhD work, continued with yearly field campaigns until 2012. Break in fieldwork since 2012 is mainly due to RP having two babies (2013;2016) and associated time away from work. All U-Pb dating samples were run between 2005 and 2015.

Data exclusions

All collected U-Pb data are included in this paper, one sample taken from Haasgat could not provide a final age solution but is included anyway in the spirit of transparency.

Reproducibility	<p>We do not explore the reproducibility of the U-Pb dating in this contribution, but have at length done so in previous publications (see below). The same flowstone sample split between labs and dated by different people using slightly different protocols gave final ages in close agreement.</p> <p>Dirks, P. et al. Geological Setting and Age of Australopithecus sediba from Southern Africa. Science 328, 205-208, doi:10.1126/science.1184950 (2010)</p> <p>Woodhead, J. & Pickering, R. Beyond 500ka: Progress and prospects in the U-Pb chronology of speleothems, and their application to studies in palaeoclimate, human evolution, biodiversity and tectonics. Chemical Geology 322-323, 290-299 (2012))</p>
Randomization	We dated all the flowstones collected, there was no need nor scope for randomization
Blinding	We did not undertake any blinding, samples were collected and analyzed with full knowledge of where they were from etc. This is standard practice in this type of work.
Did the study involve field work?	☒ Yes ☐ No

Field work, collection and transport

Field conditions	Field work was done over a period of 10 years, the exact conditions of collection are not relevant to these geological samples
Location	Cradle of Humankind, South Africa, 25°50'S - 26°S, 27°70'E - 28°10'E, 1410-1550 mamsl
Access and import/export	The rock samples did not require South African Heritage Agency (SAHRA) permits as they are not fossils, artefacts or meteorites but all sampling was carried out under the umbrella of the SAHRA existing permits held for each site. The relevant permit holders are acknowledged in the paper. A full list of permit numbers and reports is available online from SAHRA.
Disturbance	The study did not cause any disturbance. All samples were collected with a view to minimizing the damage to the cave site.

Reporting for specific materials, systems and methods

Materials & experimental systems

n/a	Involved in the study
☒	☐ Unique biological materials
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology
☒	☐ Animals and other organisms
☒	☐ Human research participants

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging