

January 2000

New data concerning the age of Mesozoic limestone from Scărișoara (Bihor Mountains)

Bogdan P. Onac

Ioan I. Bucur

Follow this and additional works at: https://digitalcommons.usf.edu/kip_articles

Recommended Citation

Onac, Bogdan P. and Bucur, Ioan I., "New data concerning the age of Mesozoic limestone from Scărișoara (Bihor Mountains)" (2000). *KIP Articles*. 5791.
https://digitalcommons.usf.edu/kip_articles/5791

This Article is brought to you for free and open access by the KIP Research Publications at Digital Commons @ University of South Florida. It has been accepted for inclusion in KIP Articles by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

NEW DATA CONCERNING THE AGE OF MESOZOIC LIMESTONE FROM SCĂRIȘOARA (BIHOR MOUNTAINS)

IOAN I. BUCUR¹, BOGDAN P. ONAC²

ABSTRACT. Microfacies and micropalaeontologic studies on some limestone samples collected within the Scărișoara Glacier Cave are presented. The investigation pointed out that the age of the limestone around Scărișoara Cave is of Upper Jurassic (Kimmeridgian) age and not Triassic as it was considered in most of the studies.

KEYWORDS: Mesozoic, limestone, microfacies, Scărișoara Glacier Cave, Bihor Mts., Romania

Geographic and Geologic Setting

Scărișoara Glacier Cave is located in the central area of the Bihor Massif (Apuseni Mountains). The area is a karst plateau bounded on the east and west by Ordâncușa and Gârda Seacă valleys, respectively (Racoviță, 1927; Rusu et al., 1970) (Fig. 1).

Access to this region is provided by National Road 75 from both Oradea (125 km) and Cluj (146 km), connecting with a county road (18 km) at Gârda de Sus. This later road is not accessible during winter and early springtime.

A series of tourists reports and descriptions referring to the Scărișoara Glacier Cave (middle of the last century) were followed by two scientific papers. Both of them represented the results of an Austrian expedition in the Bihor Mountains financed by Archduke Albrecht (Schimdl, 1863). The first paper, published by geologist Peters in 1861, contains informations on the age of the limestone and its tectonic setting. Based on his investigations, Peters (1861) uphold a Jurassic-Neocomian age for the limestone around the Scărișoara Cave. The well-known geographer Adolf Schimdl wrote the second paper, which was printed in 1863. This book contains a geology review of the Bihor Mountains, and also a more detailed description of the cave.

After Racoviță's first monographic study on the Scărișoara Glacier Cave (1927), a number of articles were published by the researchers of the Speological Institute in Cluj as a result of an extremely detailed and systematic study on climatology, glaciology, and underground ecology. All their results, as well as some new investigations were compiled and presented in a recent monographic study (Racoviță and Onac, 2000).

¹ "Babeş-Bolyai" University, Department of Geology, 1 Kogălniceanu str., 3400 Cluj-Napoca, Romania.

² "Babeş-Bolyai" University, Department of Mineralogy 1 Kogălniceanu str. and Speological Institute "Emil Racoviță", Clinicilor 5, 3400 Cluj-Napoca, Romania.




Fig. 1. Location of the Scărișoara Glacier Cave

The geological setting of the Ocoale - Ghețar Plateau is rather simple. The entire plateau is developed on Mesozoic sedimentary rocks belonging to the Bihor Unit ("autochthonous"). To the west, the Bihor Unit is overthrust by the Permian detrital formation developed in a Verrucano-type facies (purplishred colored quartzite sandstone, conglomerates, and shales). This latter one is part of the Gârda Nappe (Balintoni, 1997).

Triassic vs Jurassic age of the limestone from Scărișoara Cave

In 1964, the Geological Survey of Romania has published the geological map 1:100000, Arieșeni sheet, on which according to Bleahu and Dimitrescu the limestone around the Scărișoara Cave is of Triassic age (Ladinian and Carnian). Bleahu & Dimitrescu (1964) presented these carbonate rocks as being massive reef limestone, locally dolomitized, and developed on Wetterstein facies.

After this map was released, all papers focussing on the geology of the Scărișoara Plateau considered the limestone around Scărișoara Cave to be of Triassic age.

The first ones to question the Triassic vs Jurassic age of these carbonates were Silvestru and Ghergari (1994) who sampled the limestones along the natural section provided by the cave. These samples were sent off to one of us (I.I. Bucur) to be

analyzed with respect to their microfossil content. This preliminary study undertaken by Bucur on ten thin sections advocate for an Upper Jurassic age of the limestone.

Recently, a new set of limestone samples was collected from various locations of the cave. The sampled sites are presented in Fig. 2.

Microfacies and microfossils of the limestone from Scărișoara Cave

The limestone samples we investigated show minor microfacies varieties. Most of them are relatively coarse biopelsparites (bioclastic-peloidal grainstones) (Pl. I, Fig. 2-5), composed of skeletal fragments: plates and spines of echinoderms, bivalves and gastropods, worm tubes, scleractinian corals, bryozoans, and agglutinate foraminifera.

NEW DATA CONCERNING THE AGE OF LIMESTONE FROM SCĂRIȘOARA




Fig. 2. Longitudinal cross-section through the Scărișoara Glacier Cave with the location of the limestone samples

Dasycladales, Bryopsidales and Rhodophyte algae are also present as well as "*Tubiphytes*" *morroneensis* CRESCENTI and large fragments of microbial micrites of trombolithic type. Micrite peloids have both elongated-oval and irregular contour. Intraclasts of pelmicrite sediment are rare. Sparry calcite is the normally pore-filling cement.

Another microfacies type is the coral boundstones (Pl. I, Fig. 1), made up of colonial scleractinian corals having their external surface covered by microbial crusts (including the "*Tubiphytes*" type) or by incrustated foraminifera. The voids between corals are filled with micrite sediment in which dasyclad algae are sometimes present.

The third type of facies is represented by well-packed bioclastic-peloidal grainstones (Pl. I, fig. 6) showing elongated, flattened and oriented peloids. These appearances are most probable due to tectonic strain.

The limestone samples we analyzed are typical for a shallow marine carbonate shelf. Most probably they were deposited in its external zone (shelf edge). The microfacies of the limestone samples from Scărișoara are very alike to some sequences within the Cornet limestone (Padurea Craiului Mountains) also of Upper Jurassic age.

Most of the species identified within the micropalaeontological assemblage are characteristic for the Tethysian Upper Jurassic. The following foraminifera were determined: *Andersenolina alpina* (LEUPOLD) (Pl. II, fig. 5), *Troglotella incrustans* WERNLI & FOOKES (Pl. II, fig. 4) and *Labyrinthina mirabilis* WEYNSCHENK (Pl. II, fig. 1-3). Calcareous algae are represented by: *Salpingoporella annulata* CAROZZI (Pl. II, fig. 6), *Salpingoporella pygmaea* (GUÉMBEL) (Pl. II, Fig. 7, 8; Pl. III, Fig. 1), *Linoporella capriotica* (OPPENHEIMER) (Pl. III, fig. 5), ?*Suppiluliumaella delphica* (CARRAS), *Nipponophycus ramosus* YABE & TOYAMA (Pl. III, fig. 3, 4), *Thaumatoporella parvovesiculifera* RAINERI and *Solenopora jurassica* NICHOLSON (Pl. III, fig. 2).

Generally speaking, the assemblage is characteristic for the Upper Jurassic shallow water deposits of the Tethys Basin (Bernier, 1984; Senowbari-Daryan et al., 1994). From a biostratigraphic point of view, the most important species is *Labyrinthina mirabilis*. This species was previously cited by Fourcade & Neumann (1965) from Kimmeridgian; Septfontaine (1981, 1988) from Oxfordian-Lower Tithonian;

Pelissier et al. (1984) from Upper Oxfordian-Lower Kimmeridgian; Heinz & Isenschmid (1988) from Upper Oxfordian-Kimmeridgian, Septfontaine et al. (1991) from Oxfordian-Kimmeridgian and by Luperto-Sinni & Masse (1994) from Kimmeridgian. Synthesizing all the knowledge over this foraminifer, Bassoullet (1997) consider that this species is characteristic for the Late Upper Oxfordian and Lower Tithonian (upper part, up to the base of the Middle Tithonian).

We can conclude that the age of the limestone around the Scărișoara Glacier Cave is definitely Upper Jurassic and not Triassic. Furthermore, the presence of the foraminifer *Labyrinthina mirabilis* allows us to precise the Kimmeridgian age of this limestone.

REFERENCES

1. Balintoni, I. (1997) – *Geotectonica regiunilor metamorfe din Romania*. Ed. Carpatice, 176 p., Cluj-Napoca.
2. Bassoullet J. P. (1997) – *Foraminifères. Les grands foraminifères*. In: Groupe français d'étude du Jurassique – Biostratigraphie du Jurassique ouest-européen et méditerranéen: zonations parallèles et distribution des invertébrés et microfossiles – Cariou, E. & Hantzpergue P. (coord.) – Bull. Centre. Rech. Elf. Explor. Prod., Mém. 17, p.293-304, 3 tabl., Pau.
3. Bernier P. (1984) – *Les formations carbonatées du Kimmeridgien et du Portlandien dans le Jura méridional. Stratigraphie, micropaléontologie, sédimentologie*. Docum. Lab. Géol. Lyon, 92 (1-2), p.1-803, 218 figs., 21 tabl., 36 pls., Lyon.
4. Bleahu, M. & Dimitrescu, R. (1964) – *Harta geologică a R.P.R., foaia Arieșeni* (1:100.000). Institutul Geologic, București.
5. Fourcade E. & Neumann M. (1965) – *A propos des genres Labyrinthina Weynschenk, 1951 et Lituosepta Cati, 1959*. Revue de Micropaléontologie, 8 (4), p. 233-239, 2 fig., 2 pls., Paris.
6. Heinz R.A. & Isenschmid C. (1988) – *Mikroifazielle und stratigraphische Untersuchungen im Massivkalk (Malm) der Préalpes médianes*. Eclogae geol. Helv., 81 (1), p. 1-62, 19 fig., 7 pls., Basel.
7. Luperto Sinni E. & Masse J. P. (1994) – *Precisazioni micropaleontologiche sulle formazioni di Piattaforma carbonatica del Giurassico Superiore e del Cretaceo basale del massiccio del Gargano (Italia Meridionale) e implicazioni stratigrafiche*. Palaeopelagos, 4, p.243-266, 2 fig., 6 pls., Roma.
8. Pelissié B., Peybernès B. & Rey J. (1984) – *Les grands foraminifères benthiques du Jurassique moyen/supérieur du sud-ouest de la France (Aquitaine, Causses, Pyrénées). Intérêt biostratigraphique, paléoécologique et paléobiogéographique*. In: Benthos '83, 2nd int. Symp. Benthic Foraminifera (Pau, April, 1983), p.479-489, 5 fig., 2 pls., Pau.
9. Peters, K. F. (1861) – *Geologische und mineralogische Studien aus dem südöstlichen Ungarn, insbesondere aus der Umgegend vom Réybánya*. Sitzung-Ber. Akad. Wiss. Wien, Math.-Naturwiss. Classe, 43(1), p. 385-463, Wien.
10. Racoviță, E. G. (1927) – *Observations sur la glacière naturelle dite "Ghețarul de la Scărișoara"*. Bull. Soc. Sci. Cluj, III, p. 75-108, Cluj.
11. Racoviță, G. & Onac, B. P. (2000) – *Scărișoara Glacier Cave. Monographic Study*. Ed. Carpatice, 139 p., Cluj-Napoca.
12. Rusu, T., Racoviță, G. & Coman, D. (1970) – *Contributions a l'étude du complexe karstique de Scărișoara*. Ann. Spéléol., 25, p. 383-408, București.
13. Schmidl, A. A. (1863) – *Das Bihar-Gebirge an der Grenze von Ungarn und Siebenbürgen*. Verl. Förster u. Bartelmus, 442 p., Wien.

NEW DATA CONCERNING THE AGE OF LIMESTONE FROM SCĂRIȘOARA

14. Senowbari-Daryan B., Bucur I.I. & Abate B. (1994) – *Upper Jurassic calcareous algae from the Madonie Mountains, Sicily*. Beitrage zur Palaeontologie, 19, p.227-259, 2 fig., 1 tab., 11 pls., Wien.
15. Septfontaine M. (1981) – *Les foraminifères imperforés des milieux de palte-forme au Mésozoïque: détermination pratique, interprétation phylogénétique et utilisation biostratigraphique*. Revue de Micropaléontologie, 23 (3-4), p. 169-206, 10 fig., 3 pls., Paris.
16. Septfontaine M. (1988) – *Vers une classification évolutive des Lituolidés (Foraminifères) jurassiques en milieu de plate-forme carbonatée*. Revue de Paléobiologie, Vol. Spec. N°2 (Benthos '86), p.229-256, 5 fig., 2 pls., Genève.
17. Septfontaine M., Arnaud-Vanneau A., Bassoullet J.-P., Gusic I., Ramalho M. & Velic I. (1991) – *Les foraminifères imperforés des plates-formes carbonatées jurassiques: état des connaissances et perspectives d'avenir*. Bull. Soc. Vaud. Sc. Nat. (Bull. 312 Lab., Géol., Géophys. et Musée géol. Univ. Lausanne), 80 (3), p.255-277, 2 fig., Lausanne.
18. Silvestru E. & Ghergari L. (1994) - *On the paleokarst in the cave Ghețarul de la Scărișoara (Bihor Mountains, Romania)*. Theor. Appl. Karst., 7, p. 155-161.

PLATE EXPLANATION

Plate I.

Fig. 1 : Coral boundstone. Sample 1334 (3), x 20.

Fig. 2 – 5 : Bioclastic-peloidal grainstones.

2 – sample 1339 (3), x 17;

3 – sample 1335 (1), x 17;

4 – sample 1339 (1), x 17;

5 – sample 1339 (2), x 17.

Fig. 6 : Bioclastic-peloidal grainstone-packstone. Sample 1335 (2), x 20.

Plate II.

Fig. 1 – 3 : *Labyrinthina mirabilis* WEYNSCHENK.

1 – sample 1339 (2), x 38;

2 – sample 1339 (4), x 38;

3 – sample 1337, x 38.

Fig. 4 : *Troglotella incrustans* WERNLI & FOOKES. Sample 1339 (2), x 70.

Fig. 5 : *Andersenolina alpina* (LEUPOLD). Sample 1339 (2), x 70.

Fig. 6 : *Salpingoporella annulata* CAROZZI. Sample 1339 (3), x 70.

Fig. 7, 8 : *Salpingoporella pygmaea* (GUEMBEL)

7 – sample 1335 (1), x 38;

8 – sample 1339 (2), x 38.

Fig. 9 : "*Tubiphytes*" *morronensis* CRESCENTI. Sample 1339 (1), x 38.

Plate III.

Fig. 1 : *Salpingoporella pygmaea* (GUEMBEL). Sample 1334 (1), x 38.

Fig. 2 : *Solenopora jurassica* NICHOLSON. Sample 1339 (1), x 20.


Fig. 3, 4 : *Nipponophycus ramosus* YABE & TOYAMA

3 – sample 1337, x 20;

4 – sample 1339 (3), x 38.

Fig. 5 : *Linoporella capriotica* (OPPENHEIMER). Sample 1339 (2), x 20.

PLATE I



NEW DATA CONCERNING THE AGE OF LIMESTONE FROM SCĂRIȘOARA

PLATE II




PLATE III

