

University of South Florida Digital Commons @ University of South Florida

USF Tampa Graduate Theses and Dissertations

USF Graduate Theses and Dissertations

4-6-2016

Examination of the Use of Accelerated Resolution Therapy (ART) in the Treatment of Symptoms of PTSD and Sleep Dysfunction in Veterans and Civilians

Marian Jevone Hardwick

Follow this and additional works at: https://digitalcommons.usf.edu/etd

Part of the Nursing Commons, and the Psychiatric and Mental Health Commons

Scholar Commons Citation

Hardwick, Marian Jevone, "Examination of the Use of Accelerated Resolution Therapy (ART) in the Treatment of Symptoms of PTSD and Sleep Dysfunction in Veterans and Civilians" (2016). *USF Tampa Graduate Theses and Dissertations.* https://digitalcommons.usf.edu/etd/6092

This Dissertation is brought to you for free and open access by the USF Graduate Theses and Dissertations at Digital Commons @ University of South Florida. It has been accepted for inclusion in USF Tampa Graduate Theses and Dissertations by an authorized administrator of Digital Commons @ University of South Florida. For more information, please contact digitalcommons@usf.edu.

Examination of the Use of Accelerated Resolution Therapy (ART) in the Treatment of

Symptoms of PTSD and Sleep Dysfunction in Veterans and Civilians

by

Marian J. Hardwick

A dissertation submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy Department of Nursing College of Nursing University of South Florida

> Major Professor: Kevin Kip, Ph.D. Kristen Salomon, Ph.D. Constance Visovsky, Ph.D. Cecile A. Lengacher, Ph.D. Janice Zgibor, Ph.D

> > Date of Approval: March 25, 2016

Keywords: Sleep Disturbance, PTSD, Nightmares, Accelerated Resolution Therapy(ART)

Copyright © 2016, Marian J. Hardwick

Acknowledgments

I would like to acknowledge the assistance of my dissertation committee in completing this dissertation. Specifically, I thank my dissertation chair, Dr. Kevin Kip, Ph.D., for his continuous guidance, support, and mentorship through this entire process throughout the completion of this research on Accelerated Resolution Therapy (ART). This includes continuous review and assistance with revisions needed in the process of this dissertation. I am grateful to my other committee members, Dr. Visovsky, Dr. Salomon, and Dr. Lengacher for their honest recommendations and clinical insights related to this dissertation. In addition, I would like to thank Dr. Zgibor for her clinical insights and recommendations. Lastly, I am honored to thank Dr. Kip for statistical support and recommendations.

Table of Contents

List of Tables	iii
List of Figures	vii
Abstract	viii
Chapter One: Background	1
Introduction	1
Brief Historical Review of Etiology of PTSD	3
Comorbidity of Military PTSD and Sleep Disturbance	
The Hyperarousal Symptom of PTSD	
Current PTSD Treatment Approaches	
Purpose of this Study	
Aims	
Aim # 1	9
Hypothesis # 1	
Aim # 2	
Hypothesis # 2	
Aim # 3	
Hypothesis # 3	
Significance of the Study	
Innovation of the Study	
Chapter Two: Literature Review	12
Introduction	12
Conceptual Framework	
Theory of Unpleasant Symptoms Component #1. Influencing Factor	
Predisposing Risk Factor for Development of PTSD	
Psychiatric History and Risk of Suicide	14
Military Service Risk Factors for Development of PTSD	
Length of Deployment and Multiple Deployments	
Exposure to Combat	
Military Sexual Trauma	
Post-Deployment Risk Factors for Development of PTSD	
Sleep Disturbance	
Theory of Unpleasant Symptoms TOUS Component #2. Symptom Experience	
Insomnia	
Nightmares	
Fragmentation and Awakenings	

Non-restorative sleep	
Comorbidities	
Emergence of Accelerated Resolution Therapy (ART)	24
Sleep Disturbance	
Chapter Three: Methods	27
Introduction	27
Research Design	27
Specific Aims # 1 and # 2	
Population	
Setting and Sample	
Sample Size and Power Analysis	29
Instrumentation	29
PTSD Response (DSM-IV)	
PTSD Checklist (PCL)	
Sleep Response – Self-Report	
Pittsburgh Sleep Quality Index (PSQI)	
The PSQI-Addendum (PSQI-A)	
Intake Forms	
Medical History Form	
Demographic Form	
ART Intake Assessment Form	
Sleep Response – Objective Measurement (qEEG)	
Comorbidity Measurement	
Centers for Epidemiological Studies Depression Scale (CES-D)	
State-Trait Inventory for Cognitive and Somatic Anxiety (STICSA)	
Psychiatric Diagnostic Screening Questionnaire (PDSQ)	
Procedures	
Approval	
Recruitment Procedures	
Data Collection	
Description of ART Intervention	
Data Management/Analysis	
Aims # 1	
Hypothesis # 1	
Normality	
Aims # 2	
Hypothesis # 2	
Specific Aims # 3	
Design	
Population	
Data Management/Analysis	
Chapter Four: Results	
Introduction	40

Research Aim # 1	40
Hypothesis # 1	40
Demographic and Clinical Presentation Characteristics	
Medication Use at Baseline	
Assessment of Normality	44
Mean Scores Before and After Treatment with ART	45
Aim Number Two	47
Research Aim # 2	47
Hypothesis # 2	48
Baseline (Pre-ART) Correlations	50
Post-ART Correlations	51
Pre-ART to Post-ART Change Correlations	52
PCL-M and qEEG	52
Aim Number Three	55
Research Aim # 3	55
Hypothesis # 3	55
Comparison of Baseline Demographic Characteristics by Military	
Status	55
Comparison of Baseline Clinical Characteristics by Military Status	57
Pearson Correlations between PCL and PSQI Stratified by Military Status	
PCL and PSQI at Time 1 (Baseline)	
Pearson Correlation for PCL1 and PSQI1	
Pearson Correlation for PCL2 and PSQI2	
Pearson Correlation for PCL3 and PSQI3	59
Pearson Correlation Between Change in PCL and Change in PSQI Overall	
and by Military Status	60
Multiple Linear Regression Analysis	
Magnitude of Acute and Sustained Changes	64
Chapter Five: Discussion, Conclusions and Recommendations	66
Introduction	66
Summary of the Results	66
Aim # 1	66
Aim # 2	67
Aim # 3	68
Scientific Implications of Findings	68
Aim # 1	68
Aim # 2	
Aim # 3	
Implications for Nursing	69
Limitations	
Recommendations for Future Study	70
References	72
Appendices	95

Appendix A: ART Intake Questionnaire	
Appendix B: Medical History Form	
Appendix C: State-Trait Inventory for Cognitive and Somatic Anxiety	
Appendix D: PTSD Checklist – Military Version (PCL-M)	101
Appendix E: Demographics	
Appendix F: PSQI	104
Appendix G: Center for Epidemiologic Studies Depression Scale (CES-D)	
Appendix H: Institutional Review Board Approval	109
About the Author	End Page

List of Tables

Table 1	Descriptive Characteristics for Continuous Demographic and Clinical Presentation Variables	41
Table 2	Frequencies and Percentages of Demographic and Clinical Presentation Variables	42
Table 3	Medications Taken at Baseline	44
Table 4	Normality Assessment for Continuous Sleep Measures at Baseline	45
Table 5	Normality Assessment for Normalized qEEG Sleep Measures at Baseline	46
Table 6	Mean PCL-M and PSQI Scores Before and After Treatment with ART	47
Table 7	Mean qEEG Scores Before and After Treatment with ART (Non-Normalized)	48
Table 8	Paired Sample t-Test for Pre-ART to Post-ART Measures of Sleep Function and PTSD	49
Table 9	Descriptive Statistics of Change from Baseline for qEEG and PSQI	50
Table 10	Pearson Correlations between qEEG and PSQI at Baseline	51
Table 11	Pearson Correlations between qEEG and PSQI Post ART	52
Table 12	Pearson Correlations between Changes from Baseline for qEEG and PSQI	53
Table 13	Pearson Correlations between qEEG and PCL-M at Baseline	54
Table 14	Pearson Correlations between qEEG and PCL-M Post ART	54
Table 15	Pearson Correlations between Changes from Baseline for qEEG and PCL-M	55
Table 16	Independent Sample t-test for Demographic Characteristics by Military Status	56

Table 17	Chi-Square Analyses between Categorical Demographical Variables and Military Status	
Table 18	Independent Sample t-Test for Clinical Characteristics by Military Status	57
Table 19	Results of Multiple Linear Regression with PCL and Military Status Predicting PSQI Score at Baseline	61
Table 20	Results of Multiple Linear Regression with PCL and Military Status Predicting PSQI Score After Treatment Completion with ART	62
Table 21	Results of Multiple Linear Regression with PCL and Military Status Predicting PSQI Score Sat 3-Month Follow-up	63
Table 22	Results of Multiple Linear Regression with Change in PCL and Military Status Predicting Change in PSQI Score	63

List of Figures

Figure 1	Full Conceptual Framework	13
Figure 2	Pared-Down Conceptual Models	14
Figure 3	EEG	31
Figure 4	Scatterplot between PCL and PSQI at baseline	58
Figure 5	Scatterplot between PCL and PSQI after Treatment Completion	59
Figure 6	Scatterplot between PCL and PSQI at 3-Month Follow-up	59
Figure 7	Scatterplot between change in PCL and change in PSQI (total sample)	60
Figure 8	Scatterplot between change in PCL and change in PSQI scores	61

Abstract

Posttraumatic Stress Disorder (PTSD) is a prevalent anxiety disorder that is debilitating to both veterans and civilians following one or more traumatic events. Sleep disturbances are hallmark features of PTSD. Sleep disturbances and PTSD remain two significant PTSD-related issues that continue to plague veterans returning from active duty, thereby preventing full reintegration into society. The same problem exists for civilians. This research was conducted as a previously collected pilot study data and a secondary data analysis. The purpose of the study consisted of: 1) examining the impact of treatment with Accelerated Resolution Therapy (ART) on symptoms of PTSD and sleep disturbances; 2) examining the relationships and treatment response among both subjective and objective measures of sleep function; and 3) comparing the relationship between PTSD and sleep disturbances among military versus civilians, including the effects of treatment with ART.

The study represents one of only a few studies consisting of subjective measures of PTSD (PCL checklist) and sleep quality (Pittsburgh Quality Sleep Index (PSQI)), and objective measurement of sleep function by use of electroencephalography (EEG) testing and based on a 30-minute nap protocol. The aims of this study were to: 1) investigate the effects of ART on comorbid PTSD and sleep disturbances in U.S. veterans measured both subjectively (self-report) and objectively (sleep EEG data) from previously collected pilot study data; 2) assess the relationships between objective and subjective measures of sleep disturbances before and after treatment with ART for symptoms of PTSD in U.S. veterans from previously collected pilot

study data; and 3) compare self-report PTSD and sleep disturbances symptoms between civilians and veterans before and after treatment with ART using a secondary analysis from two previously conducted studies.

For Specific Aims 1 and 2, the methods consisted of previously collected pilot study data of 8 veterans who were treated with ART at the University of South Florida, College of Nursing. For Specific Aim 3, data were pooled from two completed studies of ART directed by Dr. Kevin Kip that included civilians (*n*=75) and veterans (*n*=50) who were treated for PTSD. Data analysis for Aim 1 included the use of paired *t* tests to compare PSQI score and each stage of sleep measured from qEEG (Delta, Theta, Alpha, Beta, Gamma) before and after treatment with ART. For Aim 2, Pearson correlation was used to assess the relationship between objective measurement of sleep disturbances and subjective sleep quality before and after ART. For Aim 3, multiple linear regression models were fit with PSQI (sleep) score as the dependent variable, PCL (PTSD) score as the primary independent variable, along with a main effect term for military status (civilian versus military) and an interaction term (military status * PCL score).

Results for aims 1 and 2 showed the mean age of the sample to be 37.6 years, 87.5% male, 87.5% White (non-Hispanic), 87.5% had experienced prior combat, 50% had experienced 5 or more traumatic memories that impacted their lives, and 87.5% had previous treatment for PTSD. Sample mean scores were above established screening criteria for PTSD (PCL-M = 63.7), sleep disturbance (PSQI = 14.5), and Center for Epidemiologic Studies Depression Scale (CES-D = 28.9). For Aim 1, after treatment with ART, the mean score on the PSQI dropped 4.88 points, mean score on the PCL-M dropped -30.13 points, thereby indicating significant reductions in sleep dysfunction and symptoms of PTSD. Mean Delta 1.5-3.5 Hz waves increased pre/post by 299.89 (*p*=.032), and Theta 4-6.5 Hz waves increased pre/post mean by 83.07

(p<0.001), both indicative of improved sleep quality. Results for Aim 2 showed statistically significant strong inverse correlations between PSQI and Theta 1.5-3.5 Hz waves (r=-0.79) and PSQI and Alpha 8-11 Hz waves (r=-0.89) at baseline. Post-ART, non-significant trends were observed for higher PSQI scores and higher Beta (conscious, alert) waves. For Aim 3, mean age of military participants (n=50) was 41.9 years versus 40.4 years among civilians (n=75, p=.439). For the military cohort, 18% were female compared to 80% among civilians (p<0.001), with lower Hispanic ethnicity among military compared to civilian participants (12% vs. 27%, p=0.04). In multiple regression analysis, change in PCL score was a strong predictor of change in PSQI score, regardless of military status PCL.

In summary, within the setting of PTSD, military participants tend to present with different traumatic exposures and worse sleep quality compared to civilian counterparts. In spite of these differences, the treatment protocol with ART demonstrated similar level of benefit (reduction in symptoms of PTSD and sleep disturbance) for both military and civilian personnel. Thus, nurses caring for individuals with PTSD, whether military or civilian, need to routinely assess sleep disturbances and initiate an open dialogue regarding these conditions. In return, nurses will be able to provide patients with resources to help them better understand and address these concerns, including after experiencing restless nights of sleep. Lastly, nurses should recognize the bi-directional temporal relationship between PTSD and sleep disturbances places. This places a premium on assessing these conditions collectively, rather than as discrete, independent clinical conditions.

х

Chapter One:

Background

Introduction

Posttraumatic stress disorder (PTSD) is a prevalent anxiety disorder that is debilitating to its victims following exposure to one or more traumatic events. According to the Diagnostic and Statistical Manual of Mental Disorders DSM-V (2013), sleep disturbances and nightmares are hallmark features of PTSD. There are four core characteristic symptoms following exposure to traumatic incidents that include intrusion and avoidance symptoms, negative modifications in cognitions and mood, and changes in arousal and reactivity (American-Psychiatric-Association, 2013). Comorbidity rates with respect to PTSD are often >80% and include sleep disturbances, depression, panic disorder, substance abuse, high somatic symptom severity, decreased role functioning, and an increased risk of suicide (Brady, Killeen, Brewerton, & Lucerini, 2000; Kessler, 2000; Oquendo, 2003; Kessler, Berglund, Delmer, Jin, Merikangas, Walters, 2005; Hoge, Terhakopian, Castro, & Messer, Engel, 2007; Krysinska & Lester, 2010). Of these comorbidities, sleep disturbances is very difficult to treat, and may be refractory to treatments that successfully resolve other PTSD symptoms (Spoormaker and Montgomery, 2008, Galovski, et al., 2009). In terms of severity, risk of suicide is elevated in both civilians and veterans who have experienced traumatic events such as natural disaster, violent crime, sexual assault, and critical illness (Krakow et al., 2000; Krakow et al., 2000; Krakow et al., 2001; Krakow et al., 2004; and Lamarche & Koninck, 2007).

Among the civilian population in the U.S., the lifetime and past year prevalence of PTSD have been estimated at 6.8% and 3.5%, respectively (Kessler et al., 2008). U.S. civilians experience trauma somewhat differently than military personnel with the most common sources of trauma being physically or sexually assaulted, being in an accident or fire, witnessing violent death or harm to a close family member or friend including physical or sexual assault, and witnessing a natural disaster (Kilpatrick et al., 2013). By way of comparison, the most prevalent forms of trauma experienced among Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) veterans include: 1) having a friend killed or wounded, seeing dead or seriously injured noncombatants; 2) witnessing an accident resulting in death or serious injury, smelling decomposing bodies; 3) having an experience with an improvised explosive device (IED); 4) receiving an injury that does not require hospitalization, and among women, experiencing military sexual trauma (Tanielian & Jaycox, 2008; Maguen et al., 2012).

Sleep disturbances is a broad term that can include one or more different aspects of sleep problems including but not limited to insomnia, sleep-disordered breathing, trauma- related nightmares, and circadian rhythm sleep disorders. Sleep disturbances encompass disorders of initiating and maintaining sleep (insomnias), disorders of excessive somnolence, disorders of sleep–wake schedule, and dysfunctions associated with sleep, sleep stages, or partial arousals (Cormier, 1990). Evidence suggests that sleep disturbances *appear before the onset of PTSD*, and therefore sleep disturbances could be a risk factor in development of PTSD (Koren, Arnon, Lavie & Klein, 2002; Mellman et al., 2002; and Mellman et al., 2004). This evidence suggests that sleep disturbances to be a <u>core feature</u> (Spoormaker & Montgomery, 2008; and Lydiard & Hammer, 2009). Indeed, evidence shows that sleep disturbances are a frequent residual symptom after successful PTSD treatment, whereas

both sleep disturbances and severity of PTSD symptoms may be alleviated following treatments that focus only on sleep disorders (Jacobs-Rebhun , 2000; and Zayfert & DeViva, 2004). With respect to military PTSD, the most common types of sleep disturbances include trauma-related nightmares in addition to insomnia and inability to maintain sleep including waking up too early, and having non-restorative sleep.

Regarding the high prevalence of nightmares, there remains a lack of successful interventions that target trauma-related nightmares (Cook, *et al.*, 2010) despite evidence- based psychotherapy for PTSD at large. This assumption comes from the notion that there is no direct distinction between nightmares and other reoccurring phenomena (Cook, *et al.*, 2010). Thus, sleep disturbances and trauma-related nightmares remain two significant PTSD-related issues that continue to plague veterans returning from active duty, thereby preventing full re-integration into society.

Brief Historical Review of Etiology of PTSD.

In 1871, symptoms of PTSD started to be described for a certain group soldiers as tachycardia, anxiety, breathlessness, and hyper-arousal (Javidi, & Yadollahie, 2012). After World War I, these symptoms were given a name known as *shell shock*. Presently, according to the Diagnostic and Statistical Manual V, by definition, an individual must be exposed to trauma or a very stressful event for their diagnosis to be considered PTSD (2013). According to Wallace, *et al.* (2011) recurring nightmares related to service, avoiding sleep and frequent nocturnal checking of windows and doors within the home are all behaviors consistent with PTSD. Hyperarousal in the presence of PTSD is when one is in a state of psychological and physiological tension, and manifests as exaggeration of startled responses and sleep disturbances (Krakow, *et al.*, 2000).

Some important traumatic events which predispose to the development of PTSD include violent personal assault, being taken hostage or kidnapped, becoming a prisoner of war, torture, terrorist attacks, a range of war-related and combat exposures, or severe car accidents (Russel, 2011). Of personal assaults, sexual abuse is a leading cause of PTSD in both civilians and military personnel (Perrin et al., 2013). In addition, witnessing a crime, accident or war, as opposed to personal involvement, is also strongly associated with development of PTSD (Perrin et al., 2013). After a traumatic event, many individuals may develop acute symptoms (i.e. Acute Stress Disorder) including but not limited to severe anxiety, dissociative symptoms, dissociative amnesia, poor concentration, sleep disturbances, and de-realization (American Psychological Association, 2013). These symptoms may resolve but, on the other hand, they can also worsen in some individuals and progress to frank PTSD.

Comorbidity of Military PTSD and Sleep Disturbance.

Comorbidity is when two disorders occur in the same individual, simultaneously and the interactions between the illnesses affect the course of the disease as well its prognosis (National Institute on Drug Abuse, 2010). Recent research has reported that approximately 74% of military personnel with sleep disturbances and PTSD symptoms rated their sleep significantly worse in the deployed and post deployed environments, with 40% having a sleep efficiency of <85% (Peterson, Goodie, Satterfield & Brim, 2008). Collectively, this phenomenon represents a major health problem in terms of physical, mental and emotional functioning. Compared to the civilian population, sleep disturbances are significantly higher among combat veterans with mood disorders and symptoms of PTSD (Capaldi et al., 2011). Importantly, sleep disturbances have been linked to individuals with panic disorder (PD) and alcohol dependence (AD) (Lauterbach et al., 2011). Sleep disturbances and nightmares can be so distressing and severe that they can cause

somatic symptoms, health dysfunction, and poor work performance (Lauterbach et al., 2011). At times, individuals with PTSD give up, refuse, or are simply unable to sleep due the nature of their anxiety.

Sleep disturbances impact long-term chronicity of adverse health status. Having sleep complaints at one month can initiate post-trauma sleep conditions and significantly predict PTSD at twelve months (Koren et al., 2002). Untreated sleep symptoms can persist for years and intensify daytime PTSD symptoms and cause comorbid psychiatric problems and may be the reason for the poor clinical outcomes that occur with PTSD. These conditions may contribute to the poor clinical outcomes often observed in PTSD (Germaine et al., 2008).

Whereas sleep disturbance is commonly associated with PTSD, it is poorly understood in terms of etiology and which treatments work best (Nakamura, Lipschitz, Landward, Kuhn & West, 2011). First, there is considerable uncertainty as to the temporal relationship between PTSD and sleep disturbances (i.e. the "chicken or egg" phenomena). Based on quantitative EEG analysis, there have been discrepancies as to whether or not sleep disturbances serve as a platform for future development of PTSD, as opposed to the more conventional hypothesis that PTSD is a primary cause of future sleep disturbances. In reality, both directions of association are plausible and perhaps a major reason for which there exists consistent evidence of a strong association between PTSD and sleep quality. At the broadest level, new, evidence-based treatments are needed that address the major clinical dilemma of comorbid military PTSD and sleep disturbances. This challenge is the motivation for examination of Accelerated Resolution Therapy (ART) in this setting, and the basis for this dissertation.

The Hyperarousal Symptom of PTSD.

Hyperarousal is a core symptom of PTSD (American-Psychiatric-Association, 2013) and

often manifests as being easily startled, feeling tense or "on edge", and having difficulty sleeping, and/or having angry outbursts. The information from traumatic events in the setting of PTSD is stored and processed in the brain to be adaptive (i.e. to be "on guard" for future similar circumstances), yet is also maladaptive in everyday life. This includes, but is not limited to an imbalance produced by insufficient control from the prefrontal cortex and parasympathetic nervous system (physiology) over excessive amygdala and sympathetic activity (Hammer 1999, Morris 2013, Semple 2000, Shin 2006, Zoldaz 2013).

Given this state of hyperarousal, there have been psychophysiological investigations into triggers of symptoms of PTSD. This includes the study of increased arousal from noise and light and physiological reactivity in persons suffering from PTSD. Increased reactivity to both traumarelated cues and unconditioned stimuli, such as loud noises, has been the topic of research for decades. Elevated levels of physiological arousal elicited by audiovisual and imaginal reminders of the original trauma have been reported in studies of combat veterans and traumatized civilians of both sexes (Blanchard et al., 1986; Pitman et al., 1987; Pitman et al., 1990; Shaley et al., 1992; Shaley et al., 1993). In addition, studies with individuals who have PTSD demonstrate greater sympathetic responses, such as increased heart rate to strong stimuli such as loud noises (Paige et al, 1990; Shaley et al., 1992; Shaley 1997; Orr et al., 1995; Orr et al., 1997). Recent psychophysiological studies of PTSD have started to focus on the physiological hyperactivity of patients to unconditioned stimuli, such as the exaggerated response toward loud noises. Exaggerated startle response, such as to loud noises or light, is a criterion for the diagnosis of PTSD as defined in the DSM V (American Psychiatric Association, 2013). In this realm, many individuals with PTSD who have been exposed during combat or military training to very loud battle noises or different assortment of lights are more likely to be startled post- deployment

(Shalev et al., 1998).

Conceptually, the traumatized person stays in a state of hyperarousal in an effort to protect themselves from real life events as well as nightmares that may continuously replay in the minds (Krakow, *et al.*, 2000). The resultant morbidity and mortality related to lack of sleep has been well documented and the manifestations of sleep disturbances concomitant to PTSD include obstructive sleep apnea (OSA), periodic leg movement disorder, sleep terrors, nocturnal anxiety attacks, and sleep avoidance (Germain, Hall, Krakow, Katherine Shear, & Buysse, 2005). Thus, with the ongoing burden and high prevalence of returning military veterans afflicted with PTSD, addressing their sleep disturbances is critical in improving quality of life, restoration of health, and psychological well-being (Ulmer, *et al.*, 2011).

Current PTSD Treatment Approaches.

First-line recommended treatment approaches for adults with trauma-related issues with PTSD vary substantially by setting (American Psychiatric Association, 2004; National Institute for Health and Clinical Excellence, 2005; Foa, Keane & Friedman, 2008; Forbes et al., 2010; Management of PTSD and ASR, 2010). Cognitive behavioral therapies (CBTs) are formally endorsed and frequently used in the treatment of PTSD among veterans. This includes the use of prolonged exposure (PE) therapy, cognitive processing therapy (CPT), and eye movement desensitization and reprocessing (EMDR) (Resick, Schnicke, 1992; Shapiro, 2001; Friedman, 2003; Ballenger et al., 2004; Foa, Hembree & Rothbaum, 2007; Resick et al., 2012). Whereas use of PE and CPT is promulgated as a preferred treatment modality in military and veteran treatment settings, in civilian settings, only 17% of licensed psychologists trained in exposure therapy reported using this modality to treat PTSD, in large part due to a reported 59% treatment dropout rate (Becker, Zayfert & Anderson, 2004). In a similar sobering manner, a recent review

of randomized controlled trials conducted primarily among civilians reported that approximately two-thirds of patients who receive PE therapy or CPT (i.e. standard-of- care), retain their diagnosis post-treatment (Steenkamp, 2015).

Accelerated Resolution Therapy (ART) is a new innovative, brief exposure-based therapy for PTSD that has been recently studied in both civilian and military populations (Kip et al., 2012; Kip et al., 2013). ART is delivered in just 1-5 treatment sessions over an approximate 2week timeframe. The ART protocol does not require homework or skills practice, thereby reducing patient commitment time by more than 50% when compared to other treatments such as PE, CPT, and EMDR. Results from empirical data from both civilians and military personnel treated with ART were based on near-identical treatment and outcome assessment protocols resulting in appropriate comparison of clinical presentation and treatment response (Kip et al., 2012; Kip et al., 2013). These data provide the opportunity to examine relationships between symptoms of PTSD and sleep disturbances, before and after treatment with ART, and among both civilian and military adults.

Purpose of the Study.

The purpose of this study was to investigate, from a previously conducted pilot data study, and through a separate secondary data analysis, the impact of treatment with ART on symptoms of comorbid PTSD and sleep disturbances. This includes both subjective and objective measures of sleep disturbances. In addition, a secondary purpose was to examine the consistency versus discordance in sleep symptoms between civilians and veterans with PTSD, and whether ART appears to have a differential effect on these symptoms by military versus civilian status.

Aims.

The Specific Aims of this dissertation were as follows:

Aim # 1: Investigate the effects of ART on comorbid PTSD and sleep disturbances in U.S veterans measured both subjectively (self- report) and objectively (sleep EEG data), from previously collected pilot study data.

Hypothesis # 1: ART will be effective in improving both subjective and objective measures of sleep disturbances in U.S veterans.

Aim # 2: Assess the relationship between objective and subjective measures of sleep disturbances before and after treatment with ART for symptoms of PTSD in U.S veterans, from previously collected pilot study data.

Hypothesis # 2: There will be a strong association between objective and subjective measures of sleep disturbances in U.S. veterans before treatment with ART, after treatment with PTSD, and based on treatment-related change in PTSD symptoms.

Aim # 3: Compare self-report PTSD and sleep disturbance symptoms between civilians and veterans before and after treatment with ART using a secondary data analysis from two previously conducted studies.

Hypothesis # 3: Civilians and veterans will report similar self-report PTSD and sleep disturbances symptoms before and after treatment with ART.

Significance of the Study

Since the September 11, 2001 terrorist attacks on the World Trade Center, more than 2.6 million American military personnel had been deployed to Iraq, Afghanistan, or both (U.S. Department of Defense). Approximately 21% of the soldiers in Operation Iraqi Freedom/Operation Endurance Freedom (OIF/OEF), after service, will receive a diagnosis of

PTSD (Seal et al., 2010). Approximately 70 to 90% of veterans have difficulty initiating or maintaining sleep, and 70% of civilians who have PTSD have sleep disturbances (Mahe, Rego, & Asnis, 2006; DaViva, Zayfert & Mellman, 2004). Evidence suggests that about 50% achieve some level of PTSD remission yet continue to experience sleep disturbances symptoms (Zayfert, & DeViva, 2004; Belleville, Guay, & Marchand, 2009). Among many contributing factors, military duties and roles in both combat and noncombat environments have changed individual sleep cycles during deployment.

In the backdrop of comorbid PTSD and sleep disturbances, current evidence-based treatments for PTSD have highly variable results and substantial rates of drop out, and similarly, successful interventions for major types of sleep disturbances, including target trauma-related nightmares, remain a major challenge. Importantly, untreated sleep disturbances can persist for decades and exaggerate daytime PTSD symptoms and associated comorbid psychiatric problems (Germaine, *et al.*, 2008). Thus, examination of new therapies that may effectively treat military PTSD and sleep disturbances, in combination, is highly warranted.

Innovation of the Study.

The analyses carried out in this dissertation are innovative in several ways. First, this study represents one of only a few studies with both subjective and objective measurement of sleep function in veterans, and is the only study in which the influence of ART on symptoms of sleep dysfunction by use of electroencephalography (EEG) during a 30-minute nap is examined. Similarly, this is the only study to date in which symptoms of military PTSD and sleep disturbances are compared between civilian and veterans, including before and after treatment with ART. Finally, consistent with the ART protocol, this study examined whether comorbid symptoms of PTSD and sleep disturbances in veterans can be substantially reduced with ART in

just a few treatment sessions. This evidence would provide the rationale for larger-scale followup research, particularly given the lengthy PTSD treatment regimens currently endorsed by the VA and DoD.

Chapter Two:

Literature Review

Introduction

This chapter describes the conceptual framework and review of the relevant PTSD/sleep disturbances literature. This includes review of the epidemiology and etiology of military PTSD and sleep disturbances, risk factors for comorbid military PTSD and sleep disturbances, efficacy of current treatment approaches, and characteristics of the ART intervention proposed in the current study.

Conceptual Framework.

The Theory of Unpleasant Symptoms (TOUS) guides the basis for this dissertation. The TOUS has three major components: (i) influencing factors that affect the symptom experience; (ii) symptoms that the individual is experiencing; and (iii) consequences of the symptom experienced (Lenz et al., 1997). Within this realm, it is not well established as to why some individuals with traumatic experiences go on to develop PTSD, while others with similar experiences do not. However, as reviewed briefly above, many potential factors may predispose to the development of PTSD. Specifically, duration, intensity, quality, and timing of traumatic exposures are risk factors for future development of PTSD (Javidi, & Yadollahie, 2012; American Psychological Association, 2013).

Within the framework of the TOUS, a wide range of exposures and behavioral and environmental factors can influence the development and maintenance of PTSD (Figure 1). Once present, PTSD is often associated with comorbidities including but not limited to symptoms of anxiety, depression, and substance abuse and dependence. With respect to military PTSD, the strength and consistency of potential risk factors varies considerably. Figure 1 depicts the "full" conceptual framework for the TOUS with respect to development of comorbid PTSD and sleep dysfunction. This "full" model is based on a review of the literature and includes the three major etiological components of: (1) Influencing Factors \rightarrow (2) Symptom Experience \rightarrow (3) Symptom Consequences. As depicted in the figure, there exist a large number of predisposing and military service factors that may contribute to the development of PTSD.




Figure 1. Full Conceptual Framework

For the purpose of this dissertation, a "pared down" model of the TOUS was used. Specifically, as seen in Figure 2, this dissertation is limited to the two major etiological components of: (1) Influencing Factors and (2) Symptom Experience, and also a reduced set of risk factors for PTSD. This approach is consistent with the three specific aims of this research, and also was selected based on the specific data variables available in the pilot study (specific aims 1 and 2) and secondary data analysis (specific aim 3).

With this approach, the risk factors for PTSD reviewed below include: (i) <u>predisposing</u>: prior psychiatric history; (ii) <u>military service</u>: length of deployment and multiple deployments, exposure to combat, military sexual trauma, sleep disturbance; and (iii) <u>post-deployment</u>: sleep disturbances.




Figure 2. Pared Down Conceptual Model

Theory of Unpleasant Symptoms Component #1. Influencing Factors:

Predisposing Risk Factor for Development of PTSD.

Psychiatric History and Risk of Suicide. Prior psychiatric history can lead to increased

risk of PTSD. The U.S. Army suicide rate doubled between the years of 2004-2005 and 2008-

2009 and reached an all-time high of 27.9/100,000 in 2012 (Schoenbaum M, Kessler RC, Gilman SE, et al, 2014). Given the demand and strong need, the Army created suicide prevention programs. Many of the responses from participants with mental disorders made a connection of suicide, and the prolonged military operations in Iraq and Afghanistan were the leading causes of high rates of mental health cases among these veterans. Evidence also showed that many soldiers are not seeking treatment for the mental issues due to stigmatization (Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL, 2004; Held P, Owens GP, 2013; & Kim PY, Britt TW, Klocko RP, Riviere LA, Adler AB, 2011).

To address the major challenge of mental illness among veterans, beginning in 2006, the Department of Defense (DoD) authorized post-deployment screening in order to identify soldiers returning from deployment with behavioral health problems (Zinzow HM, Britt TW, Pury CL, Raymond MA, McFadden AC, Warner CH, Breitbach JE, Appenzeller GN, Yates V, Grieger T, Webster, 2007). However, with the more recent movement of identifying these veterans, the stigmatization and underreporting has decreased validation of studies in post-deployment screening. Moreover, the limited focus of these surveys has made it very difficult to estimate the actual extent and correlation of untreated mental disorders and causes (Hourani L, Bender R, Weimer B, Larson G, 2012). Jeon et al. (2014) conducted a study where lifetime suicide attempts (LSA) were significantly more frequent in persons with major depressive disorder (MDD) and who had experienced six types of traumas including military combat, witnessing a violent crime, rape or sexual assault, being badly beaten, being threatened by others, and learning of trauma experienced by others.

Capone (2013) conducted a study regarding other Axis I disorders. A total of 25 participants met criteria for current Major Depressive Disorder (MDD). Rates of other current

Axis I disorders in the Diagnostic and Statistical Manual of Mental Disorders-V (DSM-V) in this sample were relatively low. Four participants had diagnoses of Generalized Anxiety Disorder (GAD) and two had diagnoses of panic disorder. Nine participants endorsed symptoms consistent with current alcohol abuse, and an additional nine participants met criteria for current alcohol dependence. The authors stated being interested in history of Alcohol Use Disorder (AUD), since prior drinking and related impairment may be relevant to current relations between PTSD and alcohol use. As far as a lifetime AUD, 52 participants met criteria for alcohol abuse and 50 met criteria for alcohol dependence at some point during their lifetime.

Military Service Risk Factors for Development of PTSD. Military personnel often have risk factors for the development of PTSD. Elbogen et al. (2014) conducted a risk factor study and found significant relationships (p<0.05) between financial instability, combat experience, alcohol misuse, violence or arrests anger, violence and developing PTSD. Phillips, Leardmann, Gumbs & Smith (2010) conducted a study on predisposing risk factors and discovered adverse childhood experiences, prior violence exposures, pay grade, number of deployments, number of close friends or relatives at follow-up, and race/ethnicity were significantly associated with PTSD. O'Toole, Marshall, Schureck & Dobson (1998) conducted a study about risk factors for military personnel where there was a significant finding that men with PTSD reported a greater incidence of assault or mugging. They were also more likely to have admitted to a pre-enlistment civilian criminal record; these were usually juvenile offences, as major crimes would have prohibited enlistment.

Length of Deployment and Multiple Deployments. The length of time and multiple deployments can play a role in the health of a soldier. Fear et al. (2010) conducted a study about the effects of multiple deployments in which overall, 4.0% of participants reported probable

PTSD, 19.7% reported a common mental disorder, and 13.0% reported alcohol misuse. MacGregor, Han, Dougherty & Galarneau (2012) conducted a study that found that overall, 80% of those with two deployments were home for at least as long as the length of their first deployment. The rate of PTSD overall and other mental health disorders was 1.5% and 6.1%, respectively, with higher rates of PTSD among military personnel with two deployments compared to those with a single deployment (2.1% versus 1.2%) and, conversely, higher rates of other mental health disorders among those with a single deployment compared with two (6.3% versus 5.7%). A breakdown of the other mental health disorder category indicates substance abuse disorder as the most common subcategory, with higher rates among those with one deployment than among those with two (2.4% versus 1.7%). Similarly, mood disorders were slightly higher among those with a single deployment.

Exposure to Combat. There are several reactions soldiers can have from combat. Tracie Shea, Reddy, Tyrka & Sevin (2013) conducted a study that showed those with PTSD significantly differed from those without PTSD on multiple hypothesized risk variables with the exception of perceived preparation and training and unit support. Data on combat experiences were available for 219 participants. Of these, 59% reported receiving small arms fire, 57% having seen dead bodies, 56% with exposure to Intermittent Explosive Device explosions, 55% attacked or ambushed, 54% knowing someone injured or killed, 37% seeing dead or injured Americans, 33% clearing homes or buildings, 30% one or more casualties within their unit, 20% handling dead bodies, 15% being injured, 11% being responsible for the death of the enemy, and 1% being responsible for the death of a noncombatant (Tracie Shea, Reddy, Tyrka & Sevin, 2013).

Military Sexual Trauma. Military sexual trauma is most common among women in the military. It is also beginning to be more common in men in today's society. Decker, Rosenheck, Tsai, Hoff & Harpaz-Rotem (2013) conducted a study on participants who reported military sexual trauma (MST) and were more likely to be currently married and to endorse greater severity of PTSD and psychiatric symptoms. Homeless female veterans who reported or experienced MST significantly endorsed lower self-esteem, quality of life, and did not feel safe in their neighborhoods, and as victims in the past year. Kimerling (2007) conducted a study that reported the association of PTSD with MST to be almost three times stronger among women than among men. The link between MST and adjustment disorders was significantly stronger among men than among women. Alcohol disorders and anxiety disorders were more common among both men and women who reported MST, but the relation to MST was significantly stronger among women than among men. The relation of MST to bipolar disorders and schizophrenia or psychoses was strong among men and women, but significantly stronger among men. Women were more likely to show signs of dissociative, eating, and depressive disorders than men.

Post-Deployment Risk Factors for Development of PTSD.

Sleep Disturbance. Sleep disturbances have played a central role in predicting development of PTSD after exposure to trauma (Schoenfeld, 2012, Harvey & Bryant, 1998; Mellman, David, Bustamante, Torres, & Fins, 2001; Koren, Arnon, Lavie, & Klein, 2002), and have been a hallmark symptom of PTSD. Studies have found that sleep disturbance, such as nightmares, are more prevalent in victims who experienced trauma and developed PTSD before the trauma occurred (Mellman et al., 1995; Van Liempt, 2012). Sleep disturbances can also be residual symptoms after PTSD treatment (Zayfert & DeViva, 2004). Furthermore, having

persistence of sleep disturbances after PTSD treatment can be a negative sign for long-term consequence in various psychopathological and somatic domains of the body (Belleville, Guay, & Marchand, 2011). Sleep disturbances and difficulties due to intrusive thoughts or bad dreams about a certain event seem to have a strong relationship with reactivating symptoms of PTSD (Boe et al., 2010).

Recurring military-related nightmares, such as avoiding sleep because of nocturnal checking 'safety perimeter' behaviors or frequent checking of windows and doors in the home environment, are all-incongruent with sleep initiation. Abnormalities in rapid eye movement (REM) support findings of centrally measured hyperarousal during sleep in PTSD (Mellman, 1995). Some researchers have found co-morbid mild sleep-disordered breathing (SDB) to be common in patients with PTSD while others have not (Breslau et al., 2004; Krawkow et al., 2004). Actigraphic data in PTSD is not common, but a few studies have used this method to estimate total sleep time (TST) and awakenings in veterans with PTSD (Westmeyer et al., 2007). Most studies of returning Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) veterans and civilians' sleep quality have relied on subjective responses to a single-item sleep embedded instrument that have not focused on validated sleep questionnaires from the objective aspect in home environments (Lundin, de Boussard,, Edman, & Borg,, 2006; Seelig, 2006; Lew et al, 2007; Lew et al., 2010; McLay, Klam & Volkert, 2010).

Insomnia in particular appears to be associated with both severity and chronicity of PTSD. Pigeon, Campbell, Possemato, and Ouimette (2013) reported insomnia being associated with significantly higher severity of PTSD and depression at the six-month assessment. Contingency table analyses comparing baseline insomnia status to rates of diagnostic level PTSD six months later revealed that 20 of 53 participants with insomnia (37.7%) had PTSD at six

months compared to only 1 of 19 (5.3%) of those without insomnia (df = 1; X^2 = 7.14; p < .01) (Pigeon, Campbell, Possemato, and Ouimette, 2013). Again, these data reinforce the need for better understanding of the relationships between PTSD and sleep disturbances.

Theory of Unpleasant Symptoms TOUS Component #2. Symptom Experience:

Insomnia. Insomnia is a major symptom experience of PTSD with inadequate treatment options today. Although veterans with PTSD seem to experience poor sleep overall, there is not unexpectedly night-to-night variability in the severity, as has been documented in individuals with primary insomnia (Perlis et al., 2010). One potential source of this variability is the participants' daytime experiences. Self-reported stress levels are known to correlate with sleep disturbances across individuals (Healey et al., 1981). It is reasonable to suspect that, within an individual, days with higher levels of stress may be followed by worse sleep and more nightmares the following night. According to Gehrman et al. (2014), greater pre-treatment PTSD severity, as assessed by the PCL-M, is associated with more awakenings, shorter sleep time, longer sleep latency, and more nightmares over the six weeks of diary completion. In terms of potential treatment options, there are some encouraging indications.

Abramowitz, *et al.* (2008) conducted a randomized controlled study to compare the efficacy of symptom-oriented hypnotherapy (n=17) versus pharmacotherapy with 10mg zolpidem (n=16) among chronic combat-related PTSD patients with insomnia. All participants were assessed for PTSD and sleep disturbances symptoms. There was a significant main effect in the treatment group, F (1, 30)=4.96; p=0.034, with PTSD symptoms as measured by the Posttraumatic Disorder Scale (PDS) being lower in the hypnotherapy group (HT) compared to the zolpidem group (ZT). Cognitive behavioral therapy for insomnia (CBT-I) has been shown to improve sleep in individuals with PTSD, with durable gains at 6 month (Roszell, McFall, Malas,

1991). In addition, overall psychosocial functioning improved following CBT-I. The initial evidence regarding CBT-I and nightmares is promising, but further research (i.e. the present proposed use of ART) is needed (Talbot, et al. 2014).

Nightmares. Nightmares are also highly prevalent symptoms in the setting of PTSD. Up to 90% of individuals with PTSD report nightmares and insomnia (Roszell, McFall, Malas 1991; Ohayon & Shapiro, 2000). Participants with nightmares, compared to those without significant nightmares, also had higher baseline levels of PTSD severity as measured by the PCL checklist (Pigeon, W.R, Campbell, C.E., Possemato, K., and Ouimette, P, 2013). Regarding potential treatment options for nightmares, Rhudy, et al. (2010) conducted a randomized control study (n=30) to assess the impact of Exposure, Relaxation, & Rescripting Therapy (ERRT) on physiological and subjective measures of negative affect and fear in response to personal, nightmare-related imagery. ERRT is a brief, 3-session treatment that directly targets nightmares in trauma-exposed persons. It was predicted that ERRT would decrease physiological and subjective reactivity to nightmare imagery. Physiological reactions to nightmare imagery were lower at post treatment than baseline for the treatment group (p < 0.01), whereas physiological reactions of the control group did not change over time (p > 0.40). Limitations include the relatively small sample size, the fact that procedures did not control for physiological-emotional reactivity due to the personal relevance of the nightmare script, and no objective measurements of sleep disturbances.

Germaine, *et al.* (2007) piloted a single, behavioral intervention session that combined image rehearsal therapy (IRT), stimulus control, and sleep restriction in a sample (n=7) of adult victims of violent crimes with PTSD. Participants received a single, 90-minute intervention session and were instructed to change a nightmare they experienced in any way they wanted, and

to rehearse one or two new dream scenarios during the day, at least three times per day for a minimum duration of 5 minutes. Overall, participants showed small-to-moderate improvements in self-report sleep quality and sleep disturbances. Self-reported dream frequency was markedly decreased post-intervention. Improvements in sleep were accompanied by marked improvements in overall daytime PTSD symptom severity and individual PTSD symptom clusters. This included improvement on the Pittsburgh Sleep Quality Index (PSQI) and nighttime PTSD symptoms (p < 0.1) and daytime PTSD, intrusion, and hyperarousal symptoms (p < .05). Findings from this study are limited due to small sample size and reliance on self-report measurement of sleep disturbances.

Fragmentation and Awakenings. Individuals with PTSD report specific types of sleep disturbances including fragmented sleep and multiple awakenings. A recurrent nightmare among individuals with PTSD is a principal source of awakenings and among the most common and distressing symptoms. Historically, most sleep studies among PTSD subjects have been based upon subjective reports. However, some studies have objectively tested sleep quality and REM fragmentation by use of polysomnography (e.g. Mellman, *et al.*, 2007). Sleep has a restorative function and affects emotional regulation whereas poor sleep adversely affects emotional processing of traumatic experiences (Maher, *et al.*, 2006). Future research using objective measurements of sleep quality, including fragmentation and awakenings, is needed for proposed treatment options.

Non-restorative Sleep.

With sleep's restorative nature, it affects emotion regulation and poor sleep may affect the emotional processing of traumatic experiences for individuals (Horne, 1993; Walker, 2009, Maher et al., 2006). When compared to control subjects, studies using objective sleep measures

have suggested that patients with PTSD have: (1) greater REM density (frequency of rapid eye movements that is characteristic of a dream state); (2) hyperarousal and nonrestorative sleep with frequent brief awakenings of less than one minute across all sleep stages; (3) a greater number of shifts from REM to lighter sleep per hour of sleep; and (4) decreased stage 4 (slow wave) sleep, the most restorative sleep stage (Neylan, *et al.*, 2003, Breslau, *et al.*, 2004). Importantly, the REM state is the only time during sleep when the chemistry of the brain is changing. Somewhat paradoxically, Khawaja et al. (2013) reported that shorter sleep time was associated with more awakenings among veterans with a lifetime history of PTSD, and not surprisingly, such impaired sleep was linked to not awakening refreshed from sleep in the morning and daytime sleepiness.

In terms of treatment, quality of sleep (i.e. restorative sleep) has been shown to be higher in a hypnotherapy (HT) group compared to a group treated with zolpidem (ZT)(p=0.003), and was improved from the first to the second assessment (p < 0.0005) (Krakow et al, 2000). In this trial, findings are limited because the hypnotherapeutic intervention was added to ongoing treatment and thus its sole contribution is difficult to tease out. In addition, only self-report sleep disturbances data were collected.

Comorbidities. There is much comorbidity that may result as a symptom experience to PTSD. Kimbrel et al. (2014) examined the relationship between clinician-based PTSD diagnoses and screening positive on non-PTSD Psychiatric Diagnostic Screening Questionnaire (PDSQ) subscales. As a result, veterans who met criteria for PTSD were significantly more likely to screen positive for depression, χ^2 =28.010(1), p<0.001, panic disorder, χ^2 =38.065(1), p<0.001, social phobia, χ^2 =19.636(1), p<0.001, and generalized anxiety disorder, χ^2 =23.303(1), p<0.001(Kimbrel et al., 2014). There was an absence of a relationship between substance use and trauma. Irwin et al., 2014 conducted a study on relationship and impact of depression and anxiety

on PTSD. The results of this study support the theory that anxiety and depression symptoms may mediate the relationship between PTSD symptoms.

Emergence of Accelerated Resolution Therapy (ART). The emergence of Accelerated Resolution Therapy (ART) is based in large part on the limitations of current evidence-based therapies endorsed for treatment of military PTSD. In brief, evidence-based cognitive-behavioral therapy (CBT) has been recognized as the first-line treatment for PTSD by both the Veterans Administration and the Department of Defense. Presently, the most commonly used and practiced therapies for the treatment of PTSD for veterans are Prolonged Exposure (PE) therapy, Cognitive Processing Therapy (CPT) and Eye Movement Desensitization and Reprocessing (EMDR) (Resick, Schinicke, 1992; Ballenger et al., 2004; Nemorff, 2006; Foa, Hembree & Rothbaum, 2007; Resick et al., 2012).

Prolonged exposure therapy involves 10-12 sessions approximately 90 minutes each with corresponding homework assignments (Foa, Hembree & Rothbaum, 2007). The homework requirement is quite demanding. There are two assignments every day that require 1.5 to 2 hours to complete (Resick et al., 2007). This totals approximately 30 to 35 hours of dedicated treatment time over two weeks. In addition, treatment is far from absolute with dropout rates of 50% in some clinical trials of PE and non-response rates between 20-67% (Minners & Hagenaars, 2002; Hembree et al. 2003; Schurr et al., 2007; Schottenbaurer et al., 2008). Cognitive processing therapy (CPT) for treatment of PTSD consists of 12 sessions lasting 60-90 minutes with practice of learned skills outside of the therapy skills (Resick and Schnicke, 1996). The dropout rates for this therapy are approximately 4-29% with non-response rates between 4-48% (Hembree et al., 2003; Schottenbauer et al., 2008. Eye Movement Desensitization Reprocessing (EMDR) consists of 8-12 weekly 90-minute sessions with dropout rates up to 36% and non- response rates

between 7 and 92% (Friedman, 2003; Schottenbauer et al., 2008). Thus, despite these therapies being universally endorsed as first-line treatment for PTSD, they all portend significant limitations including variable efficacy, frequent dropout rates, and lengthy treatment regimens (Kip et al., 2013).

In response to limitations of the current endorsed treatments for military PTSD, ART is an emerging brief exposure-based psychotherapy that was developed in 2008. The ART protocol uses the evidence-based components of imaginal exposure, imagery rescripting, and smooth pursuit eye movements. ART addresses the sentinel PTSD features of distressing emotions, thoughts, sensations, and images in particular, by viewing the development and maintenance of PTSD symptoms as a consequence of a failure of a traumatized individual to exhibit extinction of the affective component of the memory. To date, two empirical studies of ART have been completed, including a prospective cohort study in civilians and a randomized controlled trial conducted among 57 service members and veterans. Kip et al (2013) reported in their randomized trial the mean pre-/post-change on the PCL-M was -17.2 ± 13.4 in the ART group versus -2.5 ± 6.0 in an Attention Control (AC) group (effect size = 1.39; p < 0.0001). These completed studies have resulted in multiple peer-reviewed publications.

For both completed studies of ART, strong, beneficial clinical effects were reported for treatment of symptoms of PTSD in both civilians in veterans. With respect to comorbidities of PTSD, strong effects were reported for concomitant reductions in symptoms of depression and anxiety, and to a lesser extent, reductions in substance abuse and aggression and higher levels of self-compassion. Of note, improvements in self- reported sleep disturbances were also reported with ART, and serve as the basis for more detailed examination of these effects in Aim #3 of this proposal.

Sleep Disturbance. As a risk factor for the development of PTSD during military service and sustainment of PTSD post-deployment, the relationship between PTSD and sleep disturbances was previously reviewed above.

Chapter Three:

Methods

Introduction

This chapter outlines the research methods and procedures used for this study. The first section discusses the research design; this is followed by the setting and sample, instrumentation procedures, and data analysis procedures. The chapter is divided in 2 sections. The first section pertains to Specific Aims 1 and 2 and the examination of the pilot study data of ART and subjective and objective measures of sleep disturbances. The second section pertains to Specific Aim 3 and the secondary data analysis of two previously completed studies of ART.

Research Design.

The previously conducted pilot study data, which corresponds to specific aims 1 and 2, is a prospective cohort study design that made use of self-report and objective (sleep EEG) pre- and post-measurements of symptoms of PTSD and sleep disturbances before and after treatment with ART. The secondary data analysis, which corresponds to specific aim #3, merged "civilian" and "military data from two completed federally funded studies of ART for treatment of psychological trauma. The original "civilian" study made use of an uncontrolled prospective cohort design. The "military study was a randomized controlled trial of ART versus an attention control regimen. All subjects in the attention control regimen were offered ART after 2 sessions of the control condition, and >90% ultimately received ART. Therefore, for this purpose of this analysis, the data from the military

randomized controlled trial are treated as prospective cohort data with treatment response assessed within subjects both before and after treatment with ART.

Specific Aims # 1 and # 2.

Population.

Setting and Sample. This pilot study was originally designed to enroll a total of 10 subjects and ultimately enrolled 8 subjects (veterans). For purposes of analysis, eligibility required participants to have evidence of PTSD and at least a modest level of sleep disturbances. The inclusion criteria were: (1) at least 18 years of age; (2) a U.S. service member or veteran; (3) ability to read and speak English to complete survey questions; (4) denial of suicidal and homicidal ideation or intent, and no evidence of psychotic behavior or being in psychological crisis (determined by the medical form and discretion of the clinician and Psychiatric Diagnostic Screening Questionnaire (PDSQ); and (5) symptoms indicative of comorbid psychological trauma and sleep dysfunction (determined from the 17-item PCL-M (PTSD checklist) and the Pittsburgh Sleep Quality Index (PSQI)).

For PTSD symptoms, a score of >44 on the 17-item PCL-M were used to define significant symptoms of PTSD (Blanchard, 1996). For sleep dysfunction, a score of \geq 5 on the Pittsburgh Sleep Quality Index (PSQI) (sleep quality screening tool) were used to define significant symptoms of sleep disturbances (Buysse, Reynolds, Monk, Berman, & Kupfer, 1989). Exclusion criteria were: (1) brain injury prohibiting speech, writing, and purposeful actions; (2) major psychiatric disorder (e.g. bipolar disorder) concomitant to symptoms of psychological trauma (determined by the medical form and Psychiatric Diagnostic Screening Questionnaire (PDSQ; (3) currently undergoing substance abuse treatment (e.g. alcohol, opioid, cocaine) (determined by the PDSQ); and (4) any medical condition that, in the judgment of the Principal Investigator and/or ART therapist, may

place the individual at high risk due to a potential heightened emotional reaction (e.g. previous heart attack, seizure disorder).

Sample Size and Power Analysis. The original proposed sample size of 10 subjects was designed to provide 80% power to detect a "large" within-subject effect size (assuming type I error rate of 0.05) of 0.84. To evaluate the relationship between subjective and objective measures of sleep disturbances (assessed at 3 intervals), and using the same assumptions (i.e. desired sample size of 10 subjects) and within-subject correlation of 0.50, a "medium" non-zero order correlation of 0.48 would be detectable at 80% power. Given these power calculations, it should be recognized that the design was clearly intended to be exploratory and of a pilot nature. Similarly, the pilot study was deemed to be of insufficient size to control for covariates in the analysis and as part of power calculations.

Instrumentation. The instruments described below were used to measure response before and after treatment with ART. Of note, with the exception of the objective measurement of sleep response (described below), the same self-report instruments were used for the secondary data analysis (specific aim #3).

PTSD Response (DSM-IV)

PTSD Checklist (PCL) – This is a 17-item self-report measure of the 17 DSM-IV symptoms of PTSD. Respondents rate how much they were "bothered by that problem in the past month." Items are rated on a 5-point scale ranging from 1 ("not at all") to 5 ("extremely"). The PCL can be scored in multiple ways. A total score (range 17-85) can be obtained by summing the scores from each of the 17 items. Cutoff scores for a probable PTSD diagnosis have been validated for some populations, but may not generalize to other populations. Another way to score the PCL is to follow the DSM-IV criteria. It has been suggested that a combination of these two approaches (i.e., the requisite number of

symptoms are endorsed within each cluster AND the total score is above the specified cut point for a specific population) may be best. Cronbach's α reliability has been reported as 0.97 (Blanchard, 1996; King et al., 2013).

Sleep Response – Self Report

Pittsburgh Sleep Quality Index (PSQI): This 15-item self-report developed in 1989 (Buysse et al., 1989) measures the quality and patterns of sleep in adults. It takes approximately 3 to 6 minutes to complete. A global PSQI score greater than five has yielded a diagnostic sensitivity of 89.6% and specificity of 86.5% (kappa = 0.75, p < 0.001) in distinguishing good and poor sleepers. Thus, a score of >5 indicates a positive screening for sleep disturbance. The internal consistency reliability coefficient (Cronbach's alpha) for the PSQI has been reported as 0.88 (Farrahi, Nakhaee, Sheibani, Garrusi, & Amirkafi, 2009).

The PSQI-Addendum (PSQI-A): is used to identify disruptive nocturnal behaviors that are characteristic of PTSD among military personnel (Germain et al., 2005). The PSQI-A is comprised of seven different sleep disturbance items such as memories or nightmares of the traumatic experience, and episodes of terror during sleep. Items are rated on a 0 to 3 point scale, where 0 means not in the past month and 3 means three or more times a week. They are summed to create a total score. Thus, the total score can range from 0 (normal) to 21 (severe). Internal consistency is reported as high, with a Cronbach's alpha of 0.89 (Germaine et al., 2005).

Intake Forms:

Medical History Form: This form provides current and past medical history. This includes whether the participant has been previously been treated for PTSD or any other mental condition, as well as current use of medications.

Demographics Form: This form asks the participant about background information including age, gender, ethnicity, marital status, employment status, years of education, military status, branch of the military served in, number of overseas tours served in the military, and deployment and combat experiences.

ART Intake Assessment Form: This form is used by the clinician form to determine an appropriate treatment approach to be used with ART. It specifically asks the number of traumatic memories that are currently impacting one's life, and the specific traumas that the veteran seeks to resolve through treatment with ART. It also inquires about having a current disability for PTSD or another mental health condition, as well as the length of time that the veteran has lived with one or more traumas.

Sleep Response – Objective Measurement (qEEG)

Use of quantitative electroencephalography (qEEG) has a long history in medical research. In brief, the method is used to obtain basic brain patterns of individuals while in a relaxed state (Teplan, 2002). For the pilot study, qEEG measures were collected on each veteran by use of acquisition of multi-channel surface EEG during a daytime nap, with duration of 30 minutes (generic example in Figure 3).



Figure 3. EEG

This occurred before treatment with ART and after completion of treatment with ART. Surface EEG electrodes were attached to the frontal sites: **F3 F4**, central sites: **C3**,

C4, and parietal sites: P3, P4 of the two hemispheres according to the International10-20 EEG placement sites (Jasper, 1958). Measurement of the following EEG frequency bands was performed: Delta (δ): 1-3.5 Hz; Theta (θ): 4-7.5 Hz; Alpha (α): 8-12 Hz; Sigma (σ): 13-16 Hz; Beta (β): 16.5-25 Hz; and Gamma (γ): > 30 Hz. For analytical purposes, values were expressed as non-normalized (raw) and normalized by use of a square root transformation. The analyses were based on 5-minutes of data recording following the start of transition to a sleep state.

Comorbidity Measurement

Centers for Epidemiological Studies Depression Scale (CES-D): The CES-D is a widely used 20-item self-report scale which measures the current level of depressive symptomatology in the general population, with an emphasis on depressed mood during the past week. The scale measures frequency of depressive feelings and behaviors on a four-point scale ('rarely or none of the time,' to 'most or all of the time'). The possible range is 0-60 with a score of 16 as being indicative of a positive screening of depression. The CES-D incorporates the main symptoms of depression and was derived from five validated depression scales, including the Beck Depression Inventory (BDI). Internal consistency has been reported as relatively high, with a Cronbach's alpha of 0.79 (Radloff, 1977).

State-Trait Inventory for Cognitive and Somatic Anxiety (STICSA)- The 21item self- administered questionnaire in the general population, the State-Trait Inventory for Cognitive and Somatic Anxiety (STICSA) was designed to assess cognitive and somatic symptoms of anxiety as they pertain to one's mood in the moment (state) and in general (trait). There are a total of 20 questions with a four-point summated scale ranging from 1 with no anxiety to 4 with extreme anxiety. The cutoff point of being positive for screening of this questionnaire is a score of 39. The instrument items take an estimated 6 minutes to complete. The STICSA is a reliable and valid measure of state and trait cognitive and somatic anxiety, with coefficients ranging from 0.91 (Kim et al., 2008).

Psychiatric Diagnostic Screening Questionnaire (PDSQ): The PDSQ is a 125item (yes/no) instrument used to screen for Axis I disorders in the general population and serve as a global assessment of psychopathology. It can be quickly hand scored to obtain a total score plus scale scores for 13 disorders: Major Depressive Disorder, Generalized Anxiety Disorder, Panic Disorder, PTSD, Alcohol Abuse/Dependence, Drug Abuse/Dependence, Psychosis, Bulimia/Binge-Eating Disorder, Obsessive-Compulsive Disorder, Somatization Disorder, Social Phobia, Agoraphobia, and Hypochondriasis. Six items on the depression scale provide a measure of suicidal ideation. Internal consistency (Cronbach's alpha coefficient) for the PDSQ has been reported as 0.82 (Zimmerman & Mattia, 1999).

Procedures

Approval. The Institutional Review Board at the University of South Florida previously granted approval for this study (Pro 00011530).

Recruitment Procedures.

Veterans were recruited from multiple sources including community-based organizations and veteran membership organizations within the greater Tampa Bay area, as well as academic programs at USF. Recruitment through the USF Office of Veterans Services was based on the established relationships between Dr. Kip and his research team and retired Lieutenant Colonel Larry Braue, Ed.D., Director of Veteran Services at USF. A recruitment flyer was developed for the study and distributed at community- based events, including local presentations (e.g. Rotary Club) on ART made by Dr. Kip, PI of the parent study.

Upon initial contact, the research coordinator presented the consent form to potential participants based on their inquiry about the study. Potential participants were encouraged to take their time in reviewing the consent form and to discuss study participation with a friend or family member. Veterans who consented to participate signed the informed consent form and were provided a copy for their records. The Informed Consent form stated that partaking in the study would not affect the participant's military benefits including payments, disability status, or diagnosis. Patient confidentiality and security and of all study data was maintained by having password protected computers and locking up all case report forms in Dr. Kip's office.

Data Collection.

Clinical evaluations used for screening and trial eligibility consisted of multiple pre-ART screening instruments including the PCL-M Checklist, Pittsburg Sleep Quality Index (PSQI), Psychiatric Diagnostic Screening Questionnaire (PDSQ), and Accelerated Resolution Therapy (ART) Standard Protocol questionnaire. In addition, participants completed a demographic data form and clinical history on medical and mental health history. The ART Clinical Director, Dr. Hernandez, who conducted the initial intake evaluation with study participants, confirmed their eligibility for the study by reviewing the data.

Description of ART Intervention.

Description of the ART protocol has been published. (Kip et al 2012, Kip et al 2013, Kip et al 2014). In brief, the ART protocol was delivered using the four core elements found in most A-level trauma-focused psychotherapies (U.S. Department of Veterans Affairs 2015). In the first major component of ART, imaginal exposure was used

whereby veterans were asked to recall (verbally or non-verbally) details of the traumatic event while focusing their attention on physiological sensations, thoughts, and emotions. With coaching from the ART clinician, the veteran was composed into a relaxed and alert state of mind, and is then exposed to re-activation of the targeted memory for a very short period of time (30-45 seconds). This short period of exposure to the memory was immediately followed by identification and diminishment (or eradication) of the emergence of any uncomfortable emotional or somatic symptoms. Two complete phases of intermittent exposure to the targeted memory were performed.

In the second major component of ART, imagery rescripting was used whereby the veteran imagined a new way to think about the trauma in order to change (replace) the negative traumatic narrative, including sensory material and images, to positive material. This approach is consistent with memory reconsolidation and the work of Smucker who noted that much of the cognitive-affective disturbance associated with intrusive trauma-related memories is embedded in the traumatic images themselves, and that modifying the traumatic imagery becomes a powerful, if not preferred, means of processing the traumatic material (Smucker 1997).

During both imaginal exposure and imagery rescripting, the veteran followed the therapist's hand back and forth moving their eyes from right to left, with 40 bilateral eye movements performed per set. During both components, the veteran was not speaking, but rather "watching" his or her original or newly imagined scene. Treatment of the scene was considered complete (successful) when an acceptable alternative was imagined to satisfaction, and the physiological and emotional intensity of the memory was altered and minimized. By protocol, the number of planned treatment sessions ranged from 1 to 5 based on the extent of prior trauma history. The individual treatment sessions were

scheduled for approximately one hour with additional time allocated as needed.

Data Management/Analysis.

For aims one and two, qEEG data were gathered from a sample of eight participants and matched with their corresponding PCL-M and PSQI scores from before and after ART. These data were used to assess the differences in self-reported sleep quality (PSQI), Post-Traumatic Stress Disorder symptoms (PCL-M) and qEEG measures taken from Delta, Theta, Alpha, Beta, and Gamma stages of sleep from before to after ART. The same data were used to inform research question two, and quantify the correlational relationships between self-reported (PSQI) and objective (qEEG) measures of sleep quality.

Descriptive statistics were calculated for the primary variables included in Aims 1 and 2. Continuous variables are expressed as mean \pm standard deviation (SD); categorical variables are presented as frequencies and percentages.

Aim # 1: Investigate the effects of ART on comorbid PTSD and sleep disturbances in U.S veterans measured both subjectively (self- report) and objectively (sleep EEG data), from previously collected pilot study data.

Hypothesis # 1: ART will be effective in improving both subjective and objective measures of sleep disturbances in U.S veterans.

Normality. Prior to analysis, normality was assessed for each of the continuous sleep measures using measures of skew and kurtosis for each continuous score. In addition, a student's *t*-test for location was conducted with the null hypothesis that the data are centered on zero. Data are typically considered normal if they meet the following criteria for normality: (a) a skew between -2.0 and 2.0, and (b) kurtosis between -7.0 and

7.1 (Kline, 2011). Next, data regarding the spread and central tendency were assessed for both subjective (i.e., PCL-M and PSQI) and objective sleep measures (i.e., qEEG) before

and after ART.

Paired *t* tests were used to examine the difference in subjective sleep quality (PSQI), subjective PTSD symptoms (PCL-M), and objective sleep disturbances (qEEG) measures from before to after ART. One dependent sample *t*-test was conducted on the self-reported sleep quality (PSQI), one was conducted on self-reported PTSD symptoms (PCL-M), and one was conducted on each stage of sleep measured from qEEG. This resulted in a series of five dependent sample *t*-tests, with one for Delta, one for Theta, one for Alpha, one for Beta, and one for Gamma.

Aim # 2: Assess the relationship between objective and subjective measures of sleep disturbances before and after treatment with ART for symptoms of PTSD in U.S veterans, from the previously collected pilot study data.

Hypothesis # 2: There will be a strong association between objective and subjective measures of sleep disturbances in U.S. veterans before treatment with ART, after treatment with PTSD, and based on treatment-related change in PTSD symptoms.

To assess the correlational relationship between measures of subjective sleep disturbances (as measured by the qEEG), and subjective sleep quality (as measured by the PSQI) before and after ART, a Pearson correlation matrix was created. Both objective and subjective measurements were taken before and after ART was delivered, and a change score was also calculated for each. Each time point was assessed for correlations with other time points to determine if there were significant relationships. Change scores were also created for this analysis, which measured the difference in each score from baseline to post-ART.

Any correlation with a coefficient of over .50 may provide insight into a correlation that may be detected with a larger sample size (Tabachnick & Fidell, 2012). As such,

correlations with an r coefficient of .50 or greater were further examined. An r coefficient of .50 or greater indicates that 25% or more of the variance between two variables are related (Pallant, 2010) and was chosen based on Cohen's (1988) suggestion that this r indicates a medium effect size.

Specific Aim # 3.

Design. Data were pooled from 2 completed studies of ART directed by Dr. Kip in a secondary data analysis. The overall description for the 2 studies, one conducted in civilians (n=75) and the other conducted in service members and veterans (n=50) were treated for PTSD, are as follows:

Population. For the civilian study, a total of 97 persons were screened, of whom, 17 (17.5%) were determined to be ineligible for the study. Among the 80 enrolled participants, 66 (82.5%) completed the full course of treatment including initial post-treatment assessment. For the military study, a total of 63 service members/veterans were assessed for trial eligibility, of whom, 57 (90.5) were clinically eligible and enrolled.

Measures. For assessment of PTSD among veterans, the 17-item PCL-M (Military) Checklist was used. This is a self-report instrument of DSM-IV symptoms of PTSD in response to stressful military experiences, and is used with service members and veterans. The 17-item PCL-C Checklist is the civilian version of the PTSD assessment, and was used for data collected in the civilian study. Cronbach's α reliability has been reported as 0.97 (Blanchard, 1996; King et al., 2013).

For assessment of sleep function, the 15-item self-report Pittsburgh Sleep Quality Index (PSQI) developed in 1989 (Buysse et al., 1989) was used. The PSQI measures the quality and patterns of sleep in adults. It takes approximately 3 to 6 minutes to complete. A global PSQI score greater than five yielded a diagnostic sensitivity of 89.6% and specificity

of 86.5% (kappa = 0.75, p less than 0.001) in distinguishing good and poor sleepers. A score of >5 indicates a positive screening for sleep disturbance. The internal consistency reliability coefficient (Cronbach's alpha) has been reported as 0.88 (Farrahi, Nakhaee, Sheibani, Garrusi, & Amirkafi, 2009).

Data Management/Analysis. To inform aim # 3, secondary data were gathered from the two previous studies described above, and included a total of 125 observations. These data were used to examine the relationship between self-reported PTSD symptoms (PCL-M) and self-reported sleep quality (PSQI) among study participants who received ART. Since the study hypothesis involved comparison of treatment response between civilians and veterans, an initial step was to compare demographic and presenting clinical characteristics between the 2 groups. This included use of student *t*-tests to compare compare compare compared variables.

Civilian versus military status was used in stratified analyses. Pearson correlation coefficients were calculated to examine the strength of association between self-report scores on the PCL (PTSD) checklist and PSQI. These calculations were performed before treatment with ART, after treatment with ART, at 3-month follow-up, and based on the pre-to-post treatment change. This was followed by use of multiple linear regression analysis. With this approach, the PSQI score served as the dependent variable with PCL score as the primary independent variable, along with a main effect term for military status (civilian vs. military) and an interaction term (military status x PCL score). The interaction term (military status x PCL score) was included to examine whether the relationship between PTSD score and PSQI varies (is modified) by military status.

Chapter Four:

Results

Introduction

This chapter describes the results obtained for examination of the 3 specific aims and 3 corresponding hypotheses. As with earlier chapters, the results are presented separately for aims 1 and 2 (pilot study) and then for aim 3 (secondary data analysis).

Research Aim # 1: From previously collected pilot study data investigate the effects of ART on comorbid PTSD and sleep quality in U.S veterans measured both subjectively (self-report) and objectively (sleep EEG data).

Hypothesis # 1: ART will be effective in improving both subjective and objective measures of sleep disturbances in U.S veterans.

Demographic and Clinical Presentation Characteristics

Table 1 provides descriptive characteristics for demographic and clinical presentation variables that were measured on a continuous scale. As seen, the mean age was 37.6 ± 6.5 years. The mean PSQI score was 14.5 ± 4.4 which is much higher than the screening score of ≥ 5 used to indicate impaired sleep function (Buysse, 1989). Similarly, the mean PCL-M score of 63.7 ± 10.4 was high given the potential score range on this instrument from 17 to 85. These high mean values for sleep disturbance and symptoms of PTSD reflect the study inclusion criteria. In addition, the mean depression score of 28.9 ± 9.2 was much higher than the cutpoint of ≥ 16 used to indicate a possible diagnosis of depression (Radloff, 1997).

Variable	Ν	М	SD	Minimum	Maximum
Age in years	8.00	37.63	6.50	28.00	47.00
Pittsburgh Sleep Quality Index					
at baseline	8.00	14.50	4.38	7.00	20.00
PTSD Checklist (PCL-M)					
Score at baseline	8.00	63.75	10.44	46.00	78.00
T-score of Psychiatric					
Diagnostic Screening					
Questionnaire at baseline	8.00	55.63	6.95	46.00	66.00
Raw Score Brief Symptom					
Inventory (BSI) at baseline	8.00	31.00	11.59	14.00	46.00
Adjusted score Center for					
Epidemiologic Studies					
Depression Scale (CES-D) at					
baseline	8.00	28.88	9.23	14.00	41.00
Pain Scale (SF-36) at baseline	7.00	41.79	8.50	22.50	45.00

Table 1. Descriptive Characteristics for Continuous Demographic and Clinical

 Presentation Variables.

Table 2 provides descriptive characteristics for demographic and clinical presentation variables that were measured categorically. As seen, the majority of study participants were male, of white race, a discharged veteran, and had experienced combat-related activity. Seven of the eight participants screened positive for depression at baseline, along with a high prevalence of other comorbidities including generalized anxiety disorder, and agoraphobia. The range of PSQI scores was 7 to 20, confirming that all participants indicated impaired sleep function (i.e. PSQI score >5).

Variable	n	%
Gender		
Male	7	87.50
Female	1	12.50
Race	1	12.00
Black	1	12.50
White	7	87.50
Employment status	,	07.00
Full-time employment	5	62.50
Part-time employment	1	12.50
Unemployed or disabled	1	12.50
Other	1	12.50
Military status		
Reservist	2	25.00
Discharged/veteran	6	75.00
Experience combat-related activity	-	
No	1	12.50
Yes	7	87.50
Head trauma		
No	7	87.50
Yes	1	12.50
Number of traumatic memories impacting life		
1 to 2	1	12.50
3 to 4	3	37.50
5 or more	4	50.00
Currently on disability for PTSD/MH problem		
No	4	50.00
Yes	4	50.00
Previous Treatment for PTSD		
No	1	12.50
Yes	7	87.50
Prior Treatment – Psychotherapy		
Yes	8	100.00
Screen Positive for PTSD (PDSQ) at baseline		
No	1	12.50
Yes	7	87.50
Diagnosis of Depression by CES-D at baseline		
No	1	12.50
Yes	7	87.50
Screen positive for Major Depressive Disorder Suicidality (PDSQ) at baseline		
Yes	8	100.00
Screen Positive for Generalized Anxiety Disorder (PDSQ) at baseline		
No	2	25.00
Yes	6	75.00

Table 2. Frequencies and Percentages of Demographic and Clinical Presentation Variables.

Variable	n	%
Screen Positive for Social Phobia (PDSQ) at baseline		
No	3	35.50
Yes	5	62.50
Screen Positive for Somatization Disorder (PDSQ) at baseline		
No	4	50.00
Yes	4	50.00
Screen Positive for Obsessive Compulsive Disorder (PDSQ) at baseline		
No	8	100
Screen Positive for Panic Disorder (PDSQ) at baseline		
No	5	62.50
Yes	3	35.70
Screen Positive for Agoraphobia (PDSQ) at baseline		
No	1	12.50
Yes	7	87.50
Screen Positive for Hypochondriasis (PDSQ) at baseline		
No	6	75.00
Yes	2	25.00
Screen Positive for Alcohol Abuse (PDSQ) at baseline		
No	7	87.50
Yes	1	12.50
Screen Positive for Drug Abuse/Dependence (PDSQ) at baseline		
No	8	100.00
Pittsburgh Sleep Quality Index at Baseline		
7	1	12.50
11	2	25.00
16	2	25.00
17	1	12.50
18	1	12.50
20	1	12.50

 Table 2 (Continued). Frequencies and Percentages of Demographic and Clinical Presentation Variables.

Medication Use at Baseline. As seen in table 3, the 8 participants reported taking a range of medications. In terms of major classification of medications, respective percentages were: anti-anxiety (100%, anti-depressant (100%%), anti-seizure (20%), sleep (100%), and pain (20%).

ID	Med 1	Med 2	Med 3	Med 4	Med 5	Med 6	Med 7	Med 8
1	Fluoxetine	Vitamin B12	Lisinopril	Prazosin	Inhaler	Clotrimazol e		
2	Depoprevera							
3	Hyroxidine							
4	Prozozin	Citalopram						
5	Cymbalta	Seraquel	Prazosin	Meloxicam				
6	Seroquel	Celebrex	Hydrozine	Gabapentin	Tramadol	Mexazoin		
7	Lipitor	Celexa	Buspar	Lamictal	Vitamin D3	Nuvigil	Micardis	Abilify
8	Paxil	Celexa	Bubroxin	Lisinopril	Amlodine	Topomax	Tranadal	Tynelol
		Traziadone	Modafinil	Tricor	Viagra			
			Dexanphatamine	Oxycodone	Meloxicam			

Table 3. Medications Taken at Baseline.

Assessment of Normality. As stated above, normality was assessed for each of the continuous sleep measures. Only the baseline Beta 25-30 frequency measure had a skew outside of conventional boundaries for normality, although, this variable had a kurtosis well within the -7 to 7 boundaries. Each variable had a significantly different central tendency from that of 0, as indicated by the series of *t* tests. After normalizing the sleep measures, each qEEG score's variance dropped to zero. Results of the assessment of normality are presented in Tables 4 and 5. These data confirm appropriate use of parametric analyses.

Sleep measure	Skew	Kurtosis	Variance	t test for location	<i>p</i> -value
PSQI at baseline	-0.63	-0.62	19.14	9.37	.001
Delta 1.5 – 3.5 baseline	1.00	0.17	49030.08	8.50	.001
Theta 4-6.5 baseline	0.37	-0.74	9827.23	9.46	.001
Alpha 8-11 baseline	0.62	-1.35	146101.87	3.68	.001
Beta 12-15 baseline	0.65	-1.10	6888.37	5.10	.001
Beta 16-20 baseline	0.47	-1.95	1201.68	6.21	. 001
Beta 20-25 baseline	1.32	0.67	5602.21	3.36	.012
Beta 25-30 baseline	2.05	5.04	659.19	4.57	.003
Gamma > 30 baseline	1.51	3.37	95.55	5.84	.001

Table 4. Normality Assessment for Continuous Sleep Measures at Baseline.

**Note*. Student's *t* test for location against the null hypothesis that $\mu_0=0$.

Mean Scores Before and After Treatment with ART. Descriptive data regarding mean scores for both subjective (i.e., PCL-M and PSQI) and objective sleep measures (i.e., qEEG) before and after ART are presented in Tables 6 and 7. As seen in Table 6, the mean PTSD (PCL-M) score decreased substantially from baseline (63.75) to post-treatment (33.6), as did self-report PSQI scores (14.5 vs. 9.6). As seen in table 7, results were more variable with mean values of different EEG wave frequencies apparently increasing or decreasing after treatment with ART. Results for the normalized values of qEEG scores were similar (data not shown).

Normalized qEEG	Skew	Kurtosis	Variance	t test for	<i>p</i> -value
measure				location	
Delta 1.5 – 3.5 baseline	-1.05	2.13	0.00	12.91	.001
Theta 4-6.5 baseline	0.80	-0.03	0.00	8.14	.001
Alpha 8-11 baseline	0.46	-1.12	0.00	5.37	.001
Beta 12-15 baseline	0.00	-1.27	0.00	8.28	.001
Beta 16-20 baseline	-0.55	-1.67	0.00	9.51	.001
Beta 20-25 baseline	0.98	0.11	0.00	4.66	.002
Beta 25-30 baseline	0.94	-0.85	0.00	5.95	.001
Gamma > 30 baseline	0.55	-1.50	0.00	7.59	.001

Table 5. Normality Assessment for Normalized qEEG Sleep Measures at Baseline

*Note. Student's *t* test for location against the null hypothesis that $\mu_0=0$.

For formal testing of aim #1, a series of paired sample *t*-tests were conducted to examine the difference in subjective sleep quality (PSQI), subjective PTSD symptoms (PCL-M), and objective sleep quality (qEEG) measures from before to after ART. Specifically, one dependent sample *t*-test was conducted on the self-reported sleep quality (PSQI), one was conducted on self-reported PTSD symptoms (PCL-M), and one was conducted on each stage of sleep measured from qEEG.

Variable	М	SD	Min	Max
PTSD Checklist (PCL-M) Score at baseline	63.75	10.44	46.00	78.00
PTSD Checklist (PCL-M) Score post-treatment	33.63	13.34	17.00	56.00
Pittsburgh Sleep Quality Index at baseline	14.50	4.38	7.00	20.00
Pittsburgh Sleep Quality Index post-treatment	9.63	5.40	1.00	16.00

 Table 6. Mean PCL-M and PSQI Scores Before and After Treatment with ART.

As seen in table 8, results indicated a significant improvement in self-reported sleep quality (t(7) = 2.47, p = .043), as measured from the PSQI, and a significant reduction in self-reported PTSD symptoms (t(7) = 9.70, p = .001). Of the objective sleep disturbances data (qEEG), two sleep stages had significantly different readings from before to after ART; these included increases in the frequency of Delta (t(7) = -2.67, p = .032) and Theta (t(7) = -6.48, p < .001) waves. Alpha, Beta, and Gamma readings did not differ significantly in their measures from before to after ART. In aggregate, the results suggest that ART reduces symptoms of PTSD, improves subjective reported sleep quality, and appears to increase the frequency of "slow" brain waves during a 30-minute nap protocol, including Delta and Theta waves.

Aim Number Two

Research Aim # 2: From the previously collected pilot study data, assess the relationship between objective and subjective measures of sleep quality before and after treatment with ART for symptoms of PTSD in U.S veterans.

Variable	М	SD	Min	Max
Delta_15_35_Pre	665.26	221.43	444.17	1082.04
Delta_15_35_Post	965.15	340.58	612.60	1394.78
Theta_4_65_Pre	331.52	99.13	198.74	494.48
Theta_4_65_Post	414.59	106.46	296.45	593.82
Alpha_8_11_Pre	497.22	382.11	126.99	1114.44
Alpha_8_11_Post	566.65	586.08	164.65	1837.42
Beta_12_15_Pre	149.52	83.00	51.98	275.91
Beta_12_15_Post	210.46	221.50	69.33	748.08
Beta_16_20_Pre	76.11	34.67	39.40	121.12
Beta_16_20_Post	108.82	76.92	39.47	275.14
Beta_20_25_Pre	88.78	74.85	23.29	232.29
Beta_20_25_Post	132.77	154.14	30.17	486.97
Beta_25_30_Pre	41.48	25.67	18.84	100.46
Beta_25_30_Post	41.51	26.43	21.71	99.89
Beta_gt_30_Pre	35.69	17.28	17.32	73.05
Beta_gt_30 Post	30.57	14.24	17.65	61.23

 Table 7. Mean qEEG Scores Before and After Treatment with ART (Non-Normalized).

Hypothesis # 2: There will be a strong association between objective and subjective measures of sleep disturbances in U.S. veterans before treatment with ART, after treatment with PTSD, and based on treatment-related change in PTSD symptoms.

Table 8. Paired Sample t-Test for Pre-ART to Post-ART Measures of Sleep Function andPTSD.

	ΔM		-	Confid	lence		
Measure	(pre	SD	SE	inter	val	<i>t</i> (7)	р
	to		-	Lower	Upper		
PSQI	-4.88*	5.59	1.98	0.20	9.55	2.47	.043
PCL-M	-30.13***	8.79	3.11	22.78	37.47	9.69	.001
Delta: 1.5 – 3.5 Hz	299.89*	317.48	112.25	-565.31	-34.47	-2.67	.032
Theta: 4 – 6.5 Hz	83.07***	36.24	12.81	-113.36	-52.77	-6.48	.000
Alpha: 8 – 11 Hz	69.43	621.43	219.71	-588.96	450.09	-0.32	.761
Beta: 12 – 15 Hz	60.94	170.9	60.41	-203.80	113.00	-1.01	.347
Beta: 16 – 20 Hz	32.71	65.28	23.08	-87.28	43.16	-1.42	.199
Beta: 20 – 25 Hz	43.99	101.90	36.01	-129.10	67.34	-1.22	.261
Beta: 25-30 Hz	43.99	101.86	36.01	-129.15	41.17	-1.22	.261
Gamma: > 30 Hz	-5.12	7.87	2.78	-1.46	11.70	1.84	.109

*Note. *p<0.05; **p<0.01; ***p<0.001

To assess the correlational relationship between measures of subjective sleep disturbances (as measured by the qEEG), and subjective sleep quality (as measured by the PSQI) before and after ART, a correlation matrix was created. Both objective and subjective measurements were taken before and after ART were gathered, and a change score was also calculated for each. Descriptive statistics for these change scores are presented in Table 9.

Variable	N	М	SD	Sum	Minimum	Maximum
Change in Delta 1.5-3.5 Hz	8	-299.89	317.48	-2399.00	-829.04	125.35
Change in Theta 4-6.5 Hz	8	-83.07	36.24	-664.53	-129.02	-31.49
Change in Alpha 8-11 Hz	8	-69.43	621.43	-555.47	-1057.00	872.83
Change in Beta 12-15 Hz	8	-60.94	170.86	-487.55	-472.17	57.95
Change in Beta 16-20 Hz	8	-32.71	65.28	-261.66	-154.02	51.30
Change in Beta 20-25 Hz	8	-43.99	101.86	-351.93	-254.68	75.79
Change in Beta 25-30 Hz	8	-0.02	8.19	-0.20	-12.52	11.58
Change in Beta > 30 Hz	8	5.12	7.87	40.95	-6.35	14.82
Change in PSQI	8	4.87	5.59	39.00	-1.00	15.00

Table 9. Descriptive Statistics of Change from Baseline for qEEG and PSQI.

Baseline (Pre-ART) Correlations. Table 10 presents the Pearson correlation coefficients between qEEG scores measured at baseline (pre-ART) and self-report PSQI score at baseline. As seen, there were significant, strong inverse correlations between self-report PSQI scores and Delta and Alpha brain wave activity prior to treatment with ART.

Given the sample size of 8 subjects, it should be noted that a typical Pearson Correlation with a power of .80 to reject a false null requires approximately 81 observations (Faul, Erdfelder, Buchner, & Lang, 2012). Thus, any correlation with a coefficient of over .50 may provide insight into a correlation that may be detected with a larger sample size (Tabachnick & Fidell, 2012) and may be meaningful and represent a "medium" effect (Pallant, 2010, Cohen, 1988). In this realm, the correlation coefficient between self-report PSQI score at baseline and Beta 16-20 Hz at baseline was 0.50.

qEEG at baseline	PSQI at baseline			
Delta 1.5-3.5 Hz at baseline	-0.79*			
Theta 4-6.5 Hz at baseline	-0.45			
Alpha 8-11 Hz at baseline	-0.89**			
Beta 12-15 Hz at baseline	-0.40			
Beta 16-20 Hz at baseline	-0.50			
Beta 20-25 Hz at baseline	-0.31			
Beta 25-30 Hz at baseline	0.01			
Beta > 30 Hz at baseline	-0.04			

Table 10. Pearson Correlations between qEEG and PSQI at Baseline.

*p < 0.05; **p < 0.01

Post-ART Correlations. Table 11 presents the Pearson correlation coefficients between qEEG scores measured after completion of treatment with ART and self-report PSQI score after ART. Unlike the results from the baseline assessment, self-report PSQI scores after treatment with ART were inversely associated with Theta wave activity after treatment with ART. There were non-significant trends for higher PSQI scores after treatment with ART to be associated with higher Beta wave activity. Collectively, the baseline and post-ART correlation results seem to suggest that higher PSQI scores are associated with lower slow wave (Delta and Theta) activity, and that higher PSQI scores seem to be associated with higher Beta wave activity.

qEEG after ART	PSQI post ART		
Delta 1.5-3.5 Hz post ART	0.03		
Theta 4-6.5 Hz post ART	-0.75*		
Alpha 8-11 Hz post ART	0.25		
Beta 12-15 Hz post ART	0.47		
Beta 16-20 Hz post ART	0.66		
Beta 20-25 Hz post ART	0.58		
Beta 25-30 Hz post ART	0.60		
Beta > 30 Hz post ART	0.57		

 Table 11. Pearson Correlations between qEEG and PSQI Post ART.

*p < 0.05

Pre-ART to Post-ART Change Correlations. Table 12 presents the Pearson correlation coefficients between qEEG change (pre-ART to post-ART) and PSQI change scores. There were no significant associations and all correlation coefficients were below a value of 0.50.

PCL-M and qEEG. Given the suggestion that higher PSQI scores (poorer sleep quality) may be associated with lower slow wave brain activity, parallel analyses were conducted to examine whether higher PTSD scores may be associated with lower slow wave brain activity. Table 13 (see below) presents the Pearson correlation coefficients between qEEG scores measured at baseline (pre-ART) and self-report PTSD score (PCL-M) at baseline. There were no significant associations, although there was a suggestion that higher PCL- M scores were associated with higher Beta 25-30 Hz activity (r=0.55, p=0.16).

Change in qEEG	Change in PSQI				
Change in Delta 1.5-3.5 Hz	0.21				
Change in Theta 4-6.5 Hz	-0.30				
Change in Alpha 8-11 Hz	0.17				
Change in Beta 12-15 Hz	0.11				
Change in Beta 16-20 Hz	0.30				
Change in Beta 20-25 Hz	0.26				
Change in Beta 25-30 Hz	-0.02				
Change in Beta > 30 Hz	-0.41				

Table 12. Pearson Correlations between Changes from Baseline for qEEG and PSQI.

Table 13 results are consistent with the baseline (pre-ART results). There were no significant associations between post-ART qEEG scores and post-ART PCL-M scores in table 14 (see below). However, there were suggestions of higher PCL-M scores after treatment with ART being associated with higher Beta wave activity.

Finally, there were no significant associations between qEEG change scores and PCL-M change scores in table 15 (see below). Thus, in aggregate, there was minimal evidence of a relationship between PTSD (PCL-M) scores and qEEG scores.

Baseline qEEG	Baseline PCL-M				
Delta 1.5-3.5 Hz at baseline	-0.23				
Theta 4-6.5 Hz at baseline	-0.20				
Alpha 8-11 Hz at baseline	-0.12				
Beta 12-15 Hz at baseline	0.21				
Beta 16-20 Hz at baseline	0.34				
Beta 20-25 Hz at baseline	0.40				
Beta 25-30 Hz at baseline	0.55				
Beta > 30 Hz at baseline	0.38				

 Table 13. Pearson Correlations between qEEG and PCL-M at Baseline.

 Table 14. Pearson Correlations between qEEG and PCL-M Post ART.

qEEG post ART	PCL-M post ART				
Delta 1.5-3.5 Hz post ART	0.48				
Theta 4-6.5 Hz post ART	-0.44				
Alpha 8-11 Hz post ART	0.04				
Beta 12-15 Hz post ART	0.36				
Beta 16-20 Hz post ART	0.58				
Beta 20-25 Hz post ART	0.51				
Beta 25-30 Hz post ART	0.50				
Beta > 30 Hz post ART	0.49				

Aim Number Three

Research Aim # 3: In a secondary data analysis from two previously conducted studies, compare self-report PTSD and sleep disturbance symptoms between civilians and veterans before and after treatment with ART.

Hypothesis # 3: Civilians and veterans will report similar self-report PTSD and sleep disturbance symptoms before and after treatment with ART.

Change in qEEG	Change in PCL-M				
Change in Delta 1.5-3.5 Hz	0.38				
Change in Theta 4-6.5 Hz	-0.39				
Change in Alpha 8-11 Hz	-0.13				
Change in Beta 12-15 Hz	-0.18				
Change in Beta 16-20 Hz	0.00				
Change in Beta 20-25 Hz	0.01				
Change in Beta 25-30 Hz	-0.17				
Change in Beta > 30 Hz	-0.35				

Table 15. Pearson Correlations between Changes from Baseline for qEEG and PCL-M.

Comparison of Baseline Demographic Characteristics by Military Status. As seen in table 16, the mean age of the 125 participants included in the secondary analysis was 41.0 ± 11.1 years. Of note, the mean age was similar between military and civilian participants enrolled in the two studies.

 Table 16. Independent Sample t-test for Demographic Characteristics by Military Status.

Variable	Total Sample (n = 125)		Military $(n = 50)$		Civilian $(n = 75)$			
	М	SD	M(SD)	SD	М	SD	<i>t</i> (123)	р
	10.00	11.05	41.00	10.55	10.25	0.04	-	120
Age	40.98	11.07	41.92	12.75	40.35	9.84	78	.439

As seen in table 17, the majority of participants in the civilian study (80.0%) were female compared to only 18% in the military study (p<0.001). In addition, the civilian study had higher representation of those with Hispanic ethnicity compared to the military study.

		Military	Civilian		
	N (%)	N (%)	N (%)		
Variable	125	50	75	$\chi^{2}(1)$	р
Female gender	69 (55.2)	9 (18.0)	60 (80.0)	46.63	< 0.001
Hispanic ethnicity	26 (21.1)	6 (12.0)	20 (27.4)	4.22	0.04
Married	72 (57.6)	29 (58.0)	43 (57.3)	0.01	0.94
Sexual trauma	26 (20.8)	7 (14.0)	19 (25.3)	2.34	0.13
Employed	72 (59.0)	29 (58.0)	43 (59.7)	0.04	0.85

Table 17. Chi-Square Analyses between Categorical Demographical Variables and Military Status.

Comparison of Baseline Clinical Characteristics by Military Status. Table 18 shows mean PCL and PSQI scores at the 3 time points consisting of baseline (time 1), post-ART (time 2), and at 3-month follow-up (time 3). At baseline, the mean PCL score was similar between military and civilian participants (56.1 vs. 54.2, p=0.43). However, when examining the 4 PTSD subscales at baseline, military participants presented with higher levels of arousal compared to civilian participants (18.0 vs. 15.0, p=0.002). After treatment completion with ART, the mean PCL score was lower in civilian participants compared to military participants (30.7 vs. 40.7, p=0.004). However, at 3-month follow-up, scores were similar (30.0 vs. 33.1, p=026, respectively).

At baseline, military participants presented with higher mean PSQI scores (12.6 vs. 9.4, p=0.002). This relative difference persisted after treatment with ART (9.9 vs. 6.7, p=0.001) and with a trend at 3-month follow-up (8.8 vs. 6.7, p=0.08).

	Total Sample (n=125)			Military (n=50)		Civilian (n=75)			
Variable	М	SD	М	SD	М	SD	t	р	
Scores at three times (1=Baseline, 2=post ART, 3=3 month follow-up)									
PCL1 (n = 125)	54.9	13.8	56.1	15.3	54.2	12.7	-0.79	0.431	
PCL2 (n = 113)	34.9	15.1	40.7	17.8	30.7	11.2	-3.67	0.004	
PCL3 (n = 91)	31.3	12.6	33.1	12.7	30.0	12.4	-1.14	0.26	
PSQI1 (n = 119)	10.7	4.7	12.6	4.3	9.4	4.4	-3.90	0.002	
PSQI2 (n = 101)	8.1	4.9	9.9	4.8	6.7	4.6	-3.37	0.001	
PSQI3 (n = 79)	7.5	4.9	8.8	5.5	6.7	4.5	-1.80	0.08	
PTSD (PCL) subscales – at baseline									
Intrusion	16.2	4.6	16.0	5.2	16.4	4.2	0.46	0.64	
Avoidance	6.6	2.4	6.8	2.5	6.5	2.2	-0.85	0.40	
Arousal	16.2	5.4	18.0	4.6	15.0	5.6	-3.13	0.002	
Numbing	15.9	5.1	15.3	5.7	16.2	4.7	0.98	0.33	

Table 18. Independent Sample t-Test for Clinical Characteristics by Military Status.

Pearson Correlations between PCL and PSQI Stratified by Military Status. PCL and PSQI at Time 1 (Baseline).

Pearson Correlation for PCL1 and PSQI1. Among civilian participants, there was a strong association between PCL and PSQI scores at baseline (r = 0.62, p < 0.0001). These data are depicted in the scatterplot presented in Figure 4 (left side). Among military participants, there was a modest association between PCL and PSQI scores at baseline (r = 0.48, p = 0.0004). These data are depicted in the scatterplot presented in Figure 4 (right side).




Figure 4. Scatterplot between PCL and PSQI at baseline (civilians left, military right)

Pearson Correlation for PCL2 and PSQI2. Among civilian participants, there was a strong association between PCL and PSQI scores after treatment completion with ART (r= 0.66, p < 0.0001). These data are depicted in the scatterplot presented in Figure 5 (left side). Among military participants, there was a similar strong association between PCL and PSQI scores after treatment completion with ART (r = 0.59, p <0.0001). These data are depicted in Figure 5 (right side).




Figure 5. Scatterplot between PCL and PSQI after Treatment Completion (civilians left, military right)

Pearson Correlation for PCL3 and PSQI3. Among civilian participants, there was a strong association between PCL and PSQI scores at 3-month follow-up (r = 0.66, p < 0.0001). These data are depicted in the scatterplot presented in Figure 6 (left side). Among military participants, there was a similar strong association between PCL and PSQI scores at 3-month follow-up (r = 0.71, p < 0.0001). These data are depicted in the scatterplot presented in Figure 6 (left side).




Figure 6. Scatterplot between PCL and PSQI at 3-Month Follow-up (civilians left, military

right)

Thus, results of the association between PCL and PSQI scores were relatively consistent between veterans and civilians. This included a strong association being present at all three time points.

Pearson Correlation Between Change in PCL and Change in PSQI Overall and by Military Status. The change in PCL and PSQI scores from before to after treatment with ART was calculated overall and by military status. Among all participants, there was a modest association between change in PCL and change in PSQI scores (r =0.38, p < 0.0001). The scatterplot for these data are depicted in Figure 7.




Figure 7. Scatterplot between change in PCL and change in PSQI (total sample)

Among civilian participants, there was also a modest association between change in PCL and change is PSQI scores (r = 0.43, p = 0.001). These data are depicted in the scatterplot presented in Figure 8 (left side). Among military participants, there was a similar modest association between change in PCL and change in PSQI scores (r = 0.40, p =0.009). These data are depicted in the scatterplot presented in Figure 8 (right side).




Figure 8. Scatterplot between change in PCL and change in PSQI scores (civilians left, military right)

Multiple Linear Regression Analysis.

As seen in Table 19, at baseline, PCL score was a strong predictor of PSQI score (t = 6.3, p < 0.001). Military status was also associated with PSQI score at baseline (t = 2.7, p=0.009). However, the interaction between PCL score and military status was not significant (p=0.10). The model R^2 value of 0.40 suggests that PCL score, military status, and the interaction between the two accounted for 40% of the variance in PSQI score at baseline. For interpretation, each unit increase in PCL score at baseline was associated with an estimated increase in PSQI score of 0.22 points. Compared to civilian status, being a military member indicated a 7.4-point higher score on the PSQI at baseline.

Table 19. Results of Multiple Linear Regression with PCL and Military Status Predicting	
PSQI Score at Baseline.	

Source	В	SE	β	t	р
PCL score	0.22	0.03	.64	6.31	< 0.001
Militarv member	7.38	2.77	.78	2.67	.009
PCL*Military	-0.08	0.05	50	-1.64	.103

*Note. $F(3, 115) = 25.96, p < .001, R^2 = .40$

The above regression analysis was replicated based on scores obtained at the completion of treatment with ART (time 2). As seen in Table 20, PCL score again was a strong predictor of PSQI score (t = 6.3, p < 0.001). Military status was also associated with PSQI score after completion with ART (t = 2.3, p = 0.02). Again, the interaction between PCL score and military status was borderline significant (p = 0.07). The model R^2 value of 0.46 suggests that PCL score, military status, and the interaction between the two accounted for 46% of the variance in PSQI score after treatment completion with ART. For interpretation, each unit increase in PCL score after treatment with ART was associated with an estimated increase in PSQI score of 0.27 points. Compared to civilian status, being a military member indicated a 4.7-point higher score on the PSQI after treatment with ART.

Table 20. Results of Multiple Linear Regression with PCL and Military Status Predicting

 PSQI Score After Treatment Completion with ART.

Source	В	SE	β	t	р
PCL2	0.27	0.04	.80	6.29	<.001
Military Status	4.71	2.06	.47	2.29	.024
PCL2*Military Status	-0.10	0.06	46	-1.83	.070

*Note. $F(3, 97) = 27.75, p < .001, R^2 = .46$

Table 21 provides results for the regression analysis for scores obtained at 3-month follow-up. As seen in Table 21, PCL score again was a strong predictor of PSQI score (t= 5.7, p<0.001). However, military status was not associated with PSQI score at 3-month follow-up (t = -0.4, p=0.68), and the interaction between PCL score and military status was

not significant (p=0.32). The model R^2 value of 0.49 suggests that PCL score, military status, and the interaction between the two accounted for 49% of the variance in PSQI score at 3-month follow-up.

Table 21. Results of Multiple Linear Regression with PCL and Military Status PredictingPSQI Score Sat 3-Month Follow-up.

Source	В	SE	β	Т	р
PCL3	0.24	0.04	.61	5.72	<.001
Military Status	-0.95	2.32	09	-0.41	.684
PCL3*Military Status	0.07	0.07	.25	1.00	.321

*Note. $F(3, 75) = 24.27, p < .001, R^2 = .49$

Finally, the analysis was replicated using change in PCL and PSQI scores from baseline to treatment completion with ART. As seen in Table 22, change in PCL score after treatment completion with ART was a strong predictor of change in PSQI score (t= 2.9, p=0.003). Military status was not associated with change in PSQI score (t= -0.30, p=0.77), and the interaction between change in PCL score and military status was not significant (p=0.52)

Table 22. Results of Multiple Linear Regression with Change in PCL and Military Status

 Predicting Change in PSQI Score.

Source	В	SE	β	t	р
Change in PCL (T2-T1)	0.10	0.03	.37	2.92	.004
Military Status	-0.35	1.19	05	-0.30	.767
Change in PCL*Military Status	0.03	0.05	.10	0.64	.522

*Note. $F(3, 75) = 24.27, p < .001, R^2 = .49$

In aggregate, the results of the regression analyses indicated that PCL score was strongly associated with PSQI score at all time points, and that military status (compared to civilian status) was generally associated with higher PSQI scores. The interaction between PTSD score and being a military member was non-significant in all models, suggesting a relatively similar relationship between symptoms of PTSD and sleep quality between both civilians and veterans.

Magnitude of Acute and Sustained Changes

The magnitude of change in PCL and PSQI scores was examined as acute changes (i.e., from pre-ART treatment to immediately post-treatment) and as sustained changes (i.e., from pre-ART treatment, to three months following the treatment). The average acute change on the PCL checklist was a reduction of 19.6 points, whereas the average acute change on the PSQI was a reduction of 2.3 points. Sustained changes for the PCL were an average reduction of 22.9 points, whereas average sustained changes for the PSQI were a reduction of 2.6 points. Thus, the magnitude of change was relatively similar immediately following treatment completion and at 3-month follow-up.

For veterans, the mean acute PCL change was 14.3 ± 12.7 points. Among civilians, the corresponding mean acute PCL change was 23.3 ± 13.1 points. By use of a student *t*-test, the magnitude of acute PCL change was significantly greater in civilians compared to veterans (*t*=3.64, p=0.0004). For sustained change, the difference by military status was less pronounced. The mean sustained PCL change among veterans was 20.6 ± 13.8 points compared to 24.5 ± 12.4 points among civilians. By use of a student *t*-test, the magnitude of sustained PCL change was similar between civilians and veterans (*t*=1.40, p=0.16).

Finally, for veterans, the mean acute PSQI change was 2.5 ± 4.1 points. Among civilians, the corresponding mean acute PSQI change was 2.2 ± 3.0 points. By use of a

student *t*-test, the magnitude of acute PCL change was similar between veterans and civilians (*t*=-0.41, p=0.68). For sustained change, the mean PSQI change among veterans was 3.2 ± 4.3 points compared to 2.0 ± 3.4 points among civilians. By use of a student *t*-test, the magnitude of sustained PCL change was similar between civilians and veterans (*t*=-1.28, p=0.21).

Chapter Five

Discussion, Conclusions, and Recommendations

Introduction

The final chapter of this dissertation includes a synthesis of the results from the study with discussion of the findings, conclusions, and implications for nursing and future research recommendations. The focus of this dissertation was to examine the effects of Accelerated Resolution Therapy (ART) on comorbid PTSD and sleep disturbances. This included examination of both subjective and objective measurement of sleep data (U.S. veterans), in addition to examination of results by military versus civilian status. The study was carried out using 2 data sources and approaches. For specific aims 1 and 2, data from a previously completed pilot study among eight U.S. veterans that included objective sleep EEG data were analyzed. For aim 3, a secondary data analysis was conducted between 2 prior treatment studies of ART, which included extensive self- report data on symptoms of PTSD and sleep disturbances. All three specific aims were analyzed considering pre-treatment data, post-treatment data, and change in symptom and EEG measurements as a result of participants being treated with ART.

Summary of the Results

Aim #1. The primary hypothesis for aim #1 was that treatment with ART would be effective in improving both subjective and objective measures of sleep disturbances in U.S veterans. Objective measurement of sleep quality was based on EEG analysis of brain wave activity based on a 30-minute nap protocol conducted prior to treatment with ART and after treatment completion with ART. From this pilot study of 8 U.S. veterans treated with

ART, there was confirmatory evidence that ART resulted in improved subjective reported sleep quality, in addition to evidence of increased Delta and Theta waves which are in the frequency of "slow" brain waves. Despite the small sample of just 8 veterans, improvements in subjective and objective measures of sleep achieved statistical significance. Thus, these data provide supporting evidence that when ART is used as a primary treatment for PTSD, corresponding improvements in sleep quality measured using different approaches may be expected to occur.

Aim #2. The primary hypothesis for aim #2 was that subjectively reported measurement of sleep function would be associated with EEG-objective measures of sleep function before treatment with ART, after treatment with ART, and based on treatmentrelated change in PTSD symptoms. Relatively consistent with the results of aim #1, there was evidence of significant, strong inverse associations between self-report (PSQI) scores and Delta brain wave activity prior to treatment with ART. After treatment with ART, there were non-significant trends for higher PSOI scores after treatment with ART to be associated with higher Beta wave activity. Taking these results together, it appears that higher (worse) self-report sleep scores (PSQI) are associated with less frequent slow wave (Delta) activity, and that higher PSQI scores seem to be associated with higher Beta wave activity. These results are in the expected direction (i.e. poorer self-report sleep quality associated with less frequent "slow wave" and more frequent "fast wave" frequency). However, there were no significant associations between qEEG change (pre-ART to post-ART) and PSQI change scores or qEEG change scores and PCL-M change scores. These results indicate that objective (EEG) measurement of sleep disturbances following treatment for PTSD does not correlate directly with self-report symptoms of sleep disturbances or PTSD.

Aim #3. The primary hypothesis for aim #3 was that civilians and veterans would report similar self-report PTSD and sleep disturbances symptoms before and after treatment with ART. As hypothesized, there were modest to large, statistically significant correlations between self-reported PCL (PTSD) and PSQI (sleep) scores observed among both civilians and veterans. These associations with present at baseline, immediately after treatment completion with ART, at 3-month follow-up, and based on change in scores from pre-treatment to post-treatment. In multiple regression analysis, change in PCL (PTSD) score was a strong predictor of change in PSQI (sleep) score, irrespective of military versus civilian status. Thus, even though military members tended to present with higher (worse) PSQI scores at baseline, treatment with ART and reductions in PTSD were associated with comparable reductions (improvement) in PSQI scores for both civilians and veterans.

Scientific Implications of Findings

Aim #1. When ART is used as a primary treatment for PTSD, consistent improvements in sleep disturbances were observed using subjective and objective methods and among both civilians and veterans. Other methods that may be used to examine objective and subjective sleep quality include use of actigraphy and sleeping diaries (King, Spence, Hickey, Sargent, Elesh, and Connelly, 2015). Both subjective and objective measurements play important roles in estimating sleep disturbances and should be incorporated in clinical studies to help address sleep disorders (Lin Zhang, 2007).

Aim # 2: Results indicate that objective or EEG measurements of sleep disturbances after treatment for PTSD may not correlate directly with self-report symptoms of sleep disturbances or PTSD. Recent findings suggest that objective measurements of sleep function (EEG) may not as effective in capturing manifestations of changes in subcortical circuits during REM and NREM sleep that may contribute to PTSD reliability

(Germain, 2013). However, sleep diaries can be a relatively effective sleep measurement approach (Germain, 2013). Sleep disturbances are considered a hallmark of PTSD, and the present study examined EEG measurements in relation to PTSD and subjective measurement of sleep quality. Human and animal studies have provided evidence for years supporting the notion that sleep plays an important role in PTSD-relevant emotion and memory processing for brain. Sleep diaries and other vigorous objective measurements suggest REM sleep disturbances are implicated in PTSD, and that multiple measurement methods should be used (Germain, 2014).

Aim # 3: Military members presented with higher or worse PSQI scores than civilians at baseline. However, treatment with ART resulted in relatively comparable reductions in symptoms of PTSD as well as reductions (improvement) in PSQI scores for both civilians and veterans. These findings are consistent with earlier work showing that veterans and civilians with PTSD have more severe sleep disturbances than those without trauma (Lewis, Creamer & Failla, 2009).

Implications for Nursing

Whereas additional empirical research is needed in this area, the nurse caring for the individual with PTSD, whether civilian or military, needs to routinely assess sleep disturbances and initiate an open dialogue regarding these conditions. This rapport can make it easier for patients to understand their current symptom experience and address ways in which to heal and maximize wellness. In return, the nurse will be able to provide the patient with the resources needed to help them better understand and address these concerns, including after experiencing restless nights of sleep. While multiple authors (Schoenfeld (2012), Harvey & Bryant (1998); Mellman, David, Bustamante, Torres, & Fins (2001); Koren, Arnon, Lavie, & Klein (2002)) have shown that sleep disturbances

may play a central role in future predicting development of PTSD after exposure to trauma, this also corroborates the need for nurse to be attentive to sleep dysfunction, and the possibility that it may be largely a consequence of recent trauma exposure and early manifestation of PTSD. The presumed bi-directional temporal relationship between PTSD and sleep disturbances places a premium on assessing these conditions collectively, rather than as discrete, independent clinical conditions.

Limitations

Most importantly, there was no control group to compare the results achieved with ART. This uncontrolled pilot study design is standard for an early stage evolving therapy such as ART, but can provide suggestion of effectiveness. Second, for aims 1 and 2, there was a very small sample size of eight participants in the study limiting precision of results. In addition, having seven men and one woman in the aim 1 and 2 analyses limits generalizability, as does the imbalance in aim 3 of having 80% women in the civilian sample compared to 18% in the veterans sample.

Recommendations for Future Study

Based on the existing scientific literature in this area and the findings of the current study, the following are future research recommendations.

1. To continue to examine objective measures of sleep function (in the presence and absence of PTSD) that are non-invasive, provide routine monitoring and export of data, and track very closely with clinical diagnosis and self-report of sleep quality. The EEG approach used in the present study, while useful from a scientific perspective, is not practical for everyday objective monitoring of sleep function. Future studies with an actigraphy will likely be more useful.

2. Establish a clinical recommendation for nurses and advanced practice nurses to

identify those who may be more likely to have sleep disturbances in conjunction with symptoms of PTSD, and therefore, would derive the most benefit from screening and intervention for these symptoms.

References

- Abramowitz E.G., Barak Y., Ben-Avi I. and Knobler H.Y. (2008) Hypnotherapy in the treatment of chronic combat-related PTSD patients suffering from insomnia: a randomized, zolpidem-controlled clinical trial. *Int. Journal of Clinical and Experimental Hypnosis*, 56(3), 270-80.
- Adams R.E. & Boscarino J.A. (2006). Predictors of PTSD and delayed PTSD after disaster: the impact of exposure and psychosocial resources. J Nerv Ment Dis 194,p 485–493
- American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders (5th Ed.) Author, Washington DC.
- American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders. (5th ed.). Washington DC; 2013. Affairs, department of veteran. (2007).
 Military Sexual Trauma. Retrieved from

http://www.ptsd.va.gov/public/pages/military-sexual-trauma-general.asp.

American Psychiatric Association. Practice Guideline for the Treatment of Patients with Acute Stress Disorder and Posttraumatic Stress Disorder. Arlington, VA 2004. Retrieved on

http://psychiatryonline.org/content.aspx?bookid=28§ionid=1670530.

Ballenger JC, Davidson JR, Lecrubier Y, et al. (2004). Consensus statement update on posttraumatic stress disorder from the international consensus group on depression and anxiety. Journal of Clinical Psychiatry;65 Suppl 1:55-62.

- Becker CB, Zayfert C, Anderson E. (2004). A survey of psychologists' attitudes towards and utilization of exposure therapy for PTSD. Behaviour Research and Therapy, 42:277-92.
- Belleville, G., Guay, S., & Marchand, A. (2009). Impact of sleep disturbances on PTSD symptoms and perceived health. *Journal of Nervous and Mental Disease, 197*(2), 126-132. doi: 10.1097/NMD.0b013e3181961d8e Bisson J. Post-traumatic stress disorder. (2007). BMJ 334:789-93.
- Belleville G, Guay S, Marchand A. (2011). Persistence of sleep disturbances following cognitive-behavior therapy for posttraumatic stress disorder. Journal of Psychosomatic Research; 70:318–327.
- Ben-Ezra, M., Palgi, Y., Hamama-Raz, Y., & Shrira, A. (2014). Factors Predicting Risk for
 PTSD Among Highly Exposed Civilians: Two Longitudinal Studies. *Psychiatr Q.*doi: 10.1007/s11126-014-9323-5.
- Blanchard, E.B., Kolb, L.C., Gerardi, R.J., Ryan, P. & Pallmeyer T.P. (1986). Cardiac response to relevant stimuli as an adjunctive tool for diagnosing post-traumatic stress disorder in Vietnam veterans. *Behavior Therapy*, 17, pp. 592–606 Blanchard, E.B.; Jones-Alexander, J.; Buckley, T.C.; Forneris, C.A. (1996).Psychometric properties of the PTSD checklist (PCL). *Behav. Res. Ther*, *34*, 669–673.
- Boe H. J, Holgersen K. H, Holen A. (2010). Reactivation of posttraumatic stress in male disaster survivors: The role of residual symptoms. Journal of Anxiety Disorders;24:397–402.
- Brady KT, Killeen TK, Brewerton T, Lucerini S. (2000) Comorbidity of psychiatric disorders and posttraumatic stress disorder. *Journal of Clincal Psychiatry* 61:22-32.

- Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P (1998) Trauma and posttraumatic stress disorder in the community: the 1996 Detroit Area Survey of Trauma. *Arch Gen Psychiatry*. 55(7):626–632.
- Breslau N., Roth T., Burduvali E., et al. (2004) Sleep in lifetime posttraumatic stress
 disorder: a community-based polysomnographic study. *Arch Gen Psych*, 61(5), 50816.
- Bryant, R.A. (2003). Early predictors of posttraumatic stress disorder. Biol Psychiatry, 53, p 789–795.
- Buysse, D.J., Reynolds, C.F., Monk, T.H., et al. (1989). The Pittsburgh Sleep Quality Index: A new instrument for psychiatric research and practice. *Psychiatry Research*, 28(2), 193-213.
- Capaldi V.F., 2nd, Guerrero M.L. and Killgore W.D. (2011) Sleep disruptions among returning combat veterans from Iraq and Afghanistan. *Mil Med* 176(8), 879-88.
- Capone, C., McGrath, A. C., Reddy, M. K., & Shea, M. T. (2013). Trauma-related correlates of alcohol use in recently deployed OEF/OIF veterans. *J Trauma Stress*, 26(3), 354-360. doi: 10.1002/jts.21817.
- Center, Armed Forces Health Surveillance. (2011). Associations between repeated deployments to Iraq (OIF/OND) and Afghanistan (OEF) and post-deployment illnesses and injuries, active component, U.S. Armed Forces, 2003-2010. Part II. Mental disorders, by gender, age group, military occupation, and "dwell times" prior to repeat (second through fifth) deployments. *MSMR*, *18*(9), 2-11.
- Charles Sturt University. (2013). Trait and state anxiety. *NSW Education and Communities*. Retrieved on

http://www.hsc.csu.edu.au/pdhpe/core2/focus2/focus2/4021/fac2_2_2_1.htm.

- Cook J.M., Harb G.C., Gehrman P.R., Cary M.S., Gamble G.M., Forbes D. and Ross R.J.
 (2010) Imagery rehearsal for posttraumatic nightmares: a randomized controlled trial. *J Trauma Stress* 23(5), 553-63.
- Cormier, R.E., (1990). Sleep disturbances. *Clinical Methods: The History, Physical, and Laboratory Examinations. 3rd edition.* Retrieved on http://www.ncbi.nlm.nih.gov/books/NBK401/.
- DaViva, Zayfert & Mellman. (2004). Factors associated with insomnia among civilians seeking treatment for PTSD: an exploratory study. *Behavioral Sleep Medicine*. 2(3), 162–176.
- Decker, S. E., Rosenheck, R. A., Tsai, J., Hoff, R., & Harpaz-Rotem, I. (2013). Military sexual assault and homeless women veterans: clinical correlates and treatment preferences. *Womens Health Issues*, 23(6), e373-380. doi: 10.1016/j.whi.2013.09.002.
- Elbogen, E. B., Cueva, M., Wagner, H. R., Sreenivasan, S., Brancu, M., Beckham, J. C., & Van Male, L. (2014). Screening for violence risk in military veterans: predictive validity of a brief clinical tool. *Am J Psychiatry*, *171*(7), 749-757. doi: 10.1176/appi.ajp.2014.13101316.
- Farrahi, J., Nakhaee, N., Sheibani, V., Garrusi, B., & Amirkafi, A. (2009). Psychometric properties of the Persian version of the Pittsburgh Sleep Quality Index addendum for PTSD (PSQI-A). *Sleep Breath*, *13*(3), 259-262. doi: 10.1007/s11325-008-0233-3.

- Fear, N. T., Jones, M., Murphy, D., Hull, L., Iversen, A. C., Coker, B., ... Wessely, S. (2010). What are the consequences of deployment to Iraq and Afghanistan on the mental health of the UK armed forces? A cohort study. *Lancet*, *375*(9728), 1783-1797. doi: 10.1016/s0140-6736(10)60672-1.
- Foa EB, Hembree EA, Rothbaum BO. Prolonged Exposure Therapy for PTSD. Emotional Processing of Traumatic Experiences. Therapist Guide. New York: Oxford University Press; 2007.
- Foa EB, Keane TM, Friedman MJ. (2008). Effective treatments for PTSD: Practice guidelines from the International Society for Traumatic Stress Studies. New York: Guilford.
- Forbes D, Creamer M, Bisson JI, et al. (2010). A guide to guidelines for the treatment of PTSD and related conditions. J Trauma Stress, 23:537-52.
- Frans O, Rimmo PA, Aberg L, Fredrikson M (2005) Trauma exposure and post-traumatic stress disorder in the general population. Acta Psychiatr Scand 111(4):291–299
- Friedman MJ. (2003). Post traumatic stress disorder: The latest assessment and treatment strategies. Third ed. Kansas City: Compact Clinicals.
- Foa EB, Hembree EA, Rothbaum BO. (2007).Prolonged Exposure Therapy for PTSD. Emotional Processing of Traumatic Experiences. Therapist Guide. New York: Oxford University Press.
- Galovski T.E., Monson C., Bruce S.E. and Resick P.A. (2009) Does cognitive-behavioral therapy for PTSD improve perceived health and sleep impairment? *J Trauma Stress* 22(3), 197-204.

- Gehrman, P. R., Harb, G. C., Cook, J. M., Barilla, H., & Ross, R. J. (2014). Sleep diaries of vietnam war veteram war veterans with chronic PTSD: the relationships among insominia symptoms, psychological stress, and nightmares. *Behaviotur and Sleep Medicine*. doi: 10.1080/15402002.2014.880344.
- Germain, A. 2014. Sleep disturbances as the hallmark of PTSD: where are we now? *Jorunal of Psychiatry*. April 1; 170(4): 372-382.
- Germaine, A., Hall, M., Krakow, B., Katherine Shear, M., & Buysse, D. J. (2005). A brief sleep scale for Posttraumatic Stress Disorder: Pittsburgh Sleep Quality Index Addendum for PTSD. *Journal of Anxiety Disorders, 19*(2), 233-244. doi: 10.1016/j.janxdis.2004.02.001.
- Germaine A., Shear M.K., Hall M., Buysse D.J. (2007) Effects of a brief behavioral treatment for PTSD-related sleep disturbances: a pilot study. *Behaviour Research & Therapy*, 45, 627-32.
- Germaine A., Buysse D.J. and Nofzinger E. (2008) Sleep-specific mechanisms underlying posttraumatic stress disorder: integrative review and neurobiological hypotheses. *Sleep Medicine Review*, 12(3), 185-95.
- Gilbertson MW, McFarlane AC, Weathers FW, Keane TM, Yehuda R, Shalev AY, Lasko NB, Goetz JM, Pitman RK (2010). Is trauma a causal agent of psychopathologic symptoms in posttraumatic stress disorder? Findings from identical twins discordant for combat exposure. Journal of Clinical Psychiatry 71, 1324–1330.
- Goldberg J, True WR, Eisen SA, Henderson WG (1990). A twin study of the effects of the Vietnam War on posttraumatic stress disorder. Journal of the American Medical Association 263, 1227–1232.

- Hamner MB, Lorberbaum JP, George MS. Potential role of the anterior cingulate cortex in PTSD: review and hypothesis. Depression and Anxiety. 1999;9:1-14.
- Hapke U, Schumann A, Rumpf HJ, John U, Meyer C (2006) Posttraumatic stress disorder: the role of trauma, pre-existing psychiatric disorders, and gender. Eur Arch Psychiatry Clin Neurosci 256(5):299–306.
- Healey, E. S., Kales, A., Monroe, L. J., Bixler, E. O., Chamberlin, K., & Soldatos, C. R.(1981). Onset of insomnia: Role of life-stress events. Psychosomatic Medicine, 43, 439–451.
- Held P, Owens GP. (2013). Stigmas and attitudes toward seeking mental health treatment in a sample of veterans and active duty service members. Traumatology; 19(2): 136–43.
- Hembree EA, Foa EB, Dorfan NM, Street GP, Kowalski J, Tu X. (2003). Do patients drop out prematurely from exposure therapy for PTSD? Journal of Traumatic Stress;16:555-62.
- Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL.(2004). Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. N Engl J Med; 351(1): 13–22.
- Hoge CW, Terhakopian A, Castro CA, Messer SC, Engel CC. (2007). Association of posttraumatic stress disorder with somatic symptoms, health care visits, and absenteeism among Iraq war veterans. *American Journal of Psychiatry* 164:150-3.
- Horne JA. (1993). Human sleep, sleep loss and behaviour. Implications for the prefrontal cortex and psychiatric disorder. Br J Psychiatry;162:413–19.

- Hourani L, Bender R, Weimer B, Larson G. (2012). Comparative analysis of mandated versus voluntary administrations of post-deployment health assessments among Marines. Mil Med; 177(6): 643–8.
- Javidi, H., & Yadollahie, M. (2012). Post-traumatic Stress Disorder. Int J Occup Environ Med, 3(1), 2-9.Jasper, H.H. (1958). The ten twenty electrode system of the international federation. Electroenceph. and Clinical Neurophysiology, 10, 371-375.
- Jeon, H. J., Park, J. I., Fava, M., Mischoulon, D., Sohn, J. H., Seong, S., . . . Cho, M. J. (2014). Feelings of worthlessness, traumatic experience, and their comorbidity in relation to lifetime suicide attempt in community adults with major depressive disorder. *J Affect Disord*, *166*, 206-212. doi: 10.1016/j.jad.2014.05.010.
- Kessler RC. (2000). Posttraumatic stress disorder: the burden to the individual and to society. *Journal of Clinical Psychiatry* 61 Suppl 5:4-12.
- Kessler RC, Berglund PA, Chiu W-T, et al. (2008). The National Comorbidity Survey Replication (NCS-R): Cornerstone in improving mental health and mental health care in the United States. In: The WHO World Mental Health Surveys: Global perspectives on the epidemiology of mental disorders. New York: Cambridge University Press;165-209.
- Kessler RC, Berglund P, Delmer O, Jin R, Merikangas KR, Walters EE. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62:593-602.
- Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995) Posttraumatic stress disorder in the National Comorbidity Survey. Arch Gen Psychiatry 52(12):1048– 1060.

- Kessler, R. C., Rose, S., Koenen, K. C., Karam, E. G., Stang, P. E., Stein, D. J., . . . Carmen Viana, M. (2014). How well can post-traumatic stress disorder be predicted from pre-trauma risk factors? An exploratory study in the WHO World Mental Health Surveys. *World Psychiatry*, 13(3), 265-274. doi: 10.1002/wps.20150.
- Khawaja, I. S., A, M. H., Westermeyer, J., Thuras, P., & Hurwitz, T. (2013). Nocturnal awakening & sleep duration in veterans with PTSD: An actigraphic study. *Pak J Med Sci, 29*(4), 991-996.
- Kilpatrick D, Resnick HS, Milanak ME, Miller MW, Keys KM, Friedman MJ. (2013).
 National estimates of exposure to potentially traumatic events and PTSD prevalence using DSMIV and DSM-5 criteria. J Trauma Stress, 26:537-47.
- Kim PY, Britt TW, Klocko RP, Riviere LA, Adler AB.(2011). Stigma, negative attitudes about treatment, and utilization of mental health careamong soldiers. Mil Psychol; 23(1): 65–81.
- Kim, T. S., Chung, M. Y., Kim, W., Koo, Y. J., Ryu, S. G., Kim, E. J., . . . Chae, J. H.
 (2008). Psychometric properties of the Korean version of the Short Post- Traumatic Stress Disorder Rating Interview (K-SPRINT). *Psychiatry Clin Neurosci, 62*(1), 34-39. doi: 10.1111/j.1440-1819.2007.01774.x.
- Kimbrel, N. A., DeBeer, B. B., Meyer, E. C., Silvia, P. J., Beckham, J. C., Young, K. A., & Morissette, S. B. (2014). An examination of the broader effects of warzone experiences on returning Iraq/Afghanistan veterans psychiatric health. *Psychiatry Res.* doi: 10.1016/j.psychres.2014.12.007.
- Kimerling, R., Gima, K., Smith, M. W., Street, A., & Frayne, S. (2007). The Veterans Health Administration and military sexual trauma. *Am J Public Health*, 97(12), 2160-2166. doi: 10.2105/ajph.2006.092999.

- King, M. W., Street, A. E., Gradus, J. L., Vogt, D. S., & Resick, P. A. (2013). Gender differences in posttraumatic stress symptoms among OEF/OIF veterans: an item response theory analysis. *J Trauma Stress*, 26(2), 175-183. doi: 10.1002/jts.21802.
- King, H.C., Spence, D.L., Hickey, A.H., Sargent, P., Elesh, R., & Connelly, C.D., Auricular acupuncture for sleep disturbance in veterans in with post-traumatic stress disorder: A feasibility study. *Military Medicine 2015*, 180, (5), 582.
- Kip K.E., Elk CA, Sullivan KL, Kadel R, Lengacher CA, Long CJ, Rosenzweig L, Shuman A, Hernandez DF, Street JD, Girling SA, Diamond DM. Brief treatment of symptoms of post-traumatic stress disorder (PTSD) by use of Accelerated Resolution Therapy (ART). Behavioral Sciences *2012; 2(2):* 115-134.
 PMID:23482431.
- Kip, K.E., Hernandez, D.F., Shuman, A., Witt, A., Diamond, D.M., Davis, S., Kip, R.,
 Abhayakumar, A., Wittenberg, T., Girling, S.A., Witt, S. & Rosenzweig, L.,
 Comparison of accelerated resolution therapy (ART) for treatment for symptoms of
 PTSD and sexual trauma between civilian and military adults. *Military Medicine*2015; 180, 9:964.
- Kip K.E., Rosenzweig L, Hernandez DF, Shuman A, Diamond DM, Girling SA, Sullivan KL, Wittenberg T, Witt AM, Lengacher CA, Anderson B, McMillan SC.
 Accelerated Resolution Therapy for treatment of pain secondary to symptoms of combat-related posttraumatic stress disorder. *European Journal of Psychotraumatology* 2014, 5: 24066 http://dx.doi.org/10.3402/ejpt.v5.24066
 PMID: 24959325.

- Kip, K.E., Rosenzweig L, Hernandez DF, Shuman A, Sullivan KL., Long CJ, Taylor J, McGhee S, Girling SA, Wittenberg, T, Sahebzamani, FM, Lengacher CA, Kadel R, Diamond DM. Randomized controlled trial of accelerated resolution therapy (ART) for symptoms of combat-related post-traumatic stress disorder (PTSD). *Military Medicine* 2013; 178: 1298-1309. PMID: 24306011.
- Kip K.E., Shuman A, Hernandez DF, Diamond DM, Rosenzweig L. Case report and theoretical description of Accelerated Resolution Therapy (ART) for militaryrelated post-traumatic stress disorder. *Military Medicine* 2014; 179:31-37. PMID: 24402982.
- Kip, K. E., Sullivan, K. L., Lengacher, C. A., Rosenzweig, L., Hernandez, D. F., Kadel, R.,
 ... Diamond, D. M. (2013). Brief treatment of co-occurring post-traumatic stress and depressive symptoms by use of accelerated resolution therapy((R)). *Front Psychiatry*, *4*, 11. doi: 10.3389/fpsyt.2013.00011.
- Koffel, E., Polusny, M. A., Arbisi, P. A., & Erbes, C. R. (2013). Pre-deployment daytime and nighttime sleep complaints as predictors of post-deployment PTSD and depression in National Guard troops. *J Anxiety Disord*, 27(5), 512-519. doi: 10.1016/j.janxdis.2013.07.003.
- Koren D, Arnon I, Lavie P, Klein E. (2002) Sleep complaints as early predictors of posttraumatic stress disorder: a 1-year prospective study of injured survivors of motor vehicle accidents. Am J Psychiatry;159(5):855–57.
- Krakow B, Artar A, Warner TD, Melendrez D, Johnston L, Hollifield M, Germain A, Koss M (2000) Sleep disorder, depression, and suicidality in female sexual assault survivors. Crisis 21(4):163–170.

- Krakow B, Germain A, Tandberg D, KossM, Schrader R, Hollifield M, Cheng D, Edmond T (2000) Sleep breathing and sleep movement disorders masquerading as insomnia in sexual-assault survivors. Compr Psychiatry 41(1):49–56.
- Krakow B., Hollifield M., Schrader R., Koss M., Tandberg D., Lauriello J., McBride L.,
 Warner T.D., Cheng D., Edmond T. and Kellner R. (2000) A controlled study of
 imagery rehearsal for chronic nightmares in sexual assault survivors with PTSD: A
 preliminary report. *Journal of Traumatic Stress* 13(4), 589-609.
- Krakow B, Johnston L, Melendrez D, Hollifield M, Warner TD, Chavez-Kennedy D, Herlan MJ (2001) An open-label trial of evidence-based cognitive behavior therapy for nightmares and insomnia in crime victims with PTSD. Am J Psychiatry 158(12):2043–2047.
- Krakow B, Haynes PL, Warner TD, Santana E, Melendrez D, Johnston L, Hollifield M, Sisley BN, Koss M, Shafer L (2004) Nightmares, insomnia, and sleep-disordered breathing in fire evacuees seeking treatment for posttraumatic sleep disturbance. J Trauma Stress 17(3):257–268. doi:10.1023/B:JOTS.0000029269.29098.67.
- Krawkow, B., Haynes, P.L., Warner, T.D. Warner *et al.* (2004). Nightmares, insomnia, and sleep-disordered breathing in fire evacuees seeking treatment for posttraumatic sleep disturbance. *J Trauma Stress*, 17 (3), pp. 257–268.
- Krysinska K, Lester D. (2010). Post-traumatic stress disorder and suicide risk: a systematic review. *Archives of Suicide Research* 14:1-23.
- Lamarche LJ, De Koninck J (2007) Sleep disturbance in adults with posttraumatic stress disorder: a review. J Clin Psychiatry 68(8):1257–1270.

- Lauterbach, D., Behnke, C. & McSweeney, L.B. (2011). Sleep problems among persons with a lifetime history of posttraumatic stress disorder alone and in combination with a lifetime history of other psychiatric disorders: a replication and extension. *Comprehensive Psychiatry*. 52(6), 580-586.
- Lenz, E., Pugh, L., Milligan, R., Gift, A., & Suppe, F. (1997). The Middle-Range Theory of Unpleasant Symptoms: An Update. *Advances in Nursing Science*, 19(3), 14-27.
- Lew, H.L., Poole, J.H., Vanderploeg, R.D et al. (2007). Program development and defining characteristics of returning military in a VA Polytrauma Network Site. J Rehabil Res Dev, 44 (7), pp. 1027–1034.
- Lew, H.L., Pogoda, T.K., Hsu, P.T. *et al.* (2010). Impact of the "polytrauma clinical triad" on sleep disturbance in a department of veterans affairs outpatient rehabilitation setting. *Am J Phys Med Rehabil*, 89 (6), pp. 437–445.
- Lewis, V.L., Creamer, M. & Failla, S. (2009). Is poor sleep in veterans a function of posttraumatic stress disorder? *Military Medicine*. 174,9; 948.
- Lin ZHANG, Z.Z., (2007). Objective and subjective measures for sleep disorders. *Neuroscience Bulletin.* 23(4): 236-240.
- Lundin, A., de Boussard, C., Edman, G., Borg, J. (2006). Symptoms and disability until 3 months after mild TBI. *Brain Inj*, 20 (8), pp. 799–806.
- MacGregor, A. J., Han, P. P., Dougherty, A. L., & Galarneau, M. R. (2012). Effect of dwell time on the mental health of US military personnel with multiple combat tours. *Am J Public Health, 102 Suppl 1*, S55-59. doi: 10.2105/ajph.2011.300341.
- Maher MJ, Rego SA, Asnis GM. (2006). Sleep disturbances in patients with post- traumatic stress disorder: epidemiology, impact and approaches to management. CNS Drugs; 20(7):567–90.

- Maguen S, Cohen B, Ren L, Bosch J, Kimerling R, Seal K. (2012). Gender differences in military sexual trauma and mental health diagnoses among Iraq and Afghanistan veterans with posttraumatic stress disorder. Women's Health Issues, 22:e16-e66.
- Management of Post-Traumatic Stress Disorder and Acute Stress Reaction (2010). Department of Veterans Affairs, 2010. Retrieved at http://www.healthquality.va.gov/guidelines/MH/ptsd/.
- McLay, R.N., Klam, W.P, Volkert, S.L. (2010). Insomnia is the most commonly reported symptom and predicts other symptoms of post-traumatic stress disorder in U.S. service members returning from military deployments. *Mil Med*, 175 (10), pp.759– 762.
- Mellman TA, Bustamante V, Fins AI, Pigeon WR, Nolan B (2002): REM sleep and the early development of posttraumatic stress disorder. *Am J Psychiatry*; 159:1696– 1701.
- Mellman TA, Knorr BR, Pigeon WR, Leiter JC, Akay M (2004): Heart rate variability during sleep and the early development of posttraumatic stress disorder. *Biol Psychiatry*; 55:953–956.Z.
- Mellman T. A, Kulick-Bell R, Ashlock L. E, Nolan B. (1995). Sleep events among veterans with combat-related posttraumatic stress disorder. The American Journal of Psychiatry;152:100–105.
- Mellman T.A., Pigeon W.R., Nowell P.D. and Nolan B. (2007) Relationships between REM sleep findings and PTSD symptoms during the early aftermath of trauma. *Journal of Traumatic Stress*, 20(5), 893-901.
- Merriam-Webster Dictionary. (2013). An encyclopedia and Britannia company. Retrieved on http://www.merriam-webster.com/dictionary.

- Milliken CS, Auchterlonie JL, Hoge CW. (2007). Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq war. JAMA; 298(18): 2141–8.
- Minnen AV, Hagenaars M. (2002). Fear activation and habituation patterns as early process predictors for response to prolonged exposure treatment in PTSD. Journal of Traumatic Stress;15:359-67.
- Morris MC, Rao U. Psychobiology of PTSD in the acute aftermath of trauma: Integrating research on coping, HPA function and sympathetic nervous system activity. Asian Journal of Psychiatry. 2013;6:3-21.
- O'Toole, B. I., Marshall, R. P., Schureck, R. J., & Dobson, M. (1998). Risk factors for posttraumatic stress disorder in Australian Vietnam veterans. *Aust N Z J Psychiatry*, 32(1), 21-31.
- Nakamura Y., Lipschitz D.L., Landward R., Kuhn R. and West G. (2011) Two sessions of sleep-focused mind-body bridging improve self-reported symptoms of sleep and PTSD in veterans: A pilot randomized controlled trial. *J Psychosom Res* 70(4), 335-45.
- National Institute of Drug Abuse. (2010). *What is comorbidity?* Retrieved on http://www.drugabuse.gov/publications/research-reports/comorbidity-addictionother-mental-illnesses/what-comorbidity.
- National Institute of Health. (2013). Depression. *Medline Plus*. Retrieved on http://www.nlm.nih.gov/medlineplus/ency/article/003213.htm.
- National Institute for Health and Clinical Excellence. (2005). Post-traumatic stress disorder: the treatment of PTSD in adults and children. NICE Clinical Guidelines, No. 26. London.

- Nemeroff CB, Bremner JD, Foa EB, Mayberg HS, North CS, Stein MB. (2006).
 Posttraumatic stress disorder: A state-of-the-science review. Journal of Psychiatric Research; 40:1-21.
- Norris FH, Murphy AD, Baker CK, Perilla JL, Rodriguez FG, Rodriguez JJ (2003) Epidemiology of trauma and posttraumatic stress disorder in Mexico. J Abnorm Psychol 112(4):646–656.
- O'Brien, B.S & Sher, L. (2013). Military sexual trauma as a determinant in the development of mental and physical illness in male and female veterans. *International Journal of Adolescent Medicine and Health*, 25, 3, 269-274.
- Oquendo MA, Friend JM, Halberstam B, Brodsky BS, Burke AK, Grunebaum MF, Malone KM, Mann JJ. (2003). Association of comorbid posttraumatic stress disorder and major depression with greater risk for suicidal behavior. *American Journal of Psychiatry* 160:580-2.
- Orr SP, Pitman RK, Lasko NB, Herz LR (1993). Psychophysiologic assessment of posttraumatic stress disorder imagery in World War II and Korean Combat Veterans. J Abnorm Psychol, 102:152–159.
- Orr, S.P, Lasko, N, Shalev, A.Y, Pitman, R.K. (1995). Physiologic responses to loud tones in Vietnam veterans with PTSD. *Journal of Abnormal Psychology*, 104, pp. 75–82.
- Orr, S.P, Solomon, Z., Peri, T., Pitman, R.K. & Shalev, A.Y. (1997). Physiologic responses to loud tones in Israeli veterans of the 1973 Yom Kippur war. *Biological Psychiatry*, 41, pp. 319–326.
- Paige, S.R., Reid, G.M, Allen, M.G & Newton, J.E.O. (1990). Psychophysiological correlates of post-traumatic stress disorder in Vietnam veterans. Biological Psychiatry, 27, pp. 419–430.

- Perkonigg A, Kessler RC, Storz S, Wittchen HU (2000) Traumatic events and posttraumatic stress disorder in the community: prevalence, risk factors and comorbidity. Acta Psychiatr Scand 101(1):46–59.
- Perlis, M. L., Swinkels, C. M., Gehrman, P. R., Pigeon, W. R., Matteson-Rusby, S. E., & Jungquist, C. R. (2010). The incidence and temporal patterning of insomnia: A pilot study. Journal of Sleep Research, 19, 31–35.
- Perrin, M., Vandeleur, C. L., Castelao, E., Rothen, S., Glaus, J., Vollenweider, P., & Preisig, M. (2013). Determinants of the development of post-traumatic stress disorder, in the general population. *Soc Psychiatry Psychiatr Epidemiol*. doi: 10.1007/s00127-013-0762-3.
- Peterson A. L., Goodie J.L., Satterfield W. A., & Brim W. L. (2008). Sleep disturbance during military deployment. *Military Medicine*, 5, 173-230.
- Phillips, C. J., Leardmann, C. A., Gumbs, G. R., & Smith, B. (2010). Risk factors for posttraumatic stress disorder among deployed US male marines. *BMC Psychiatry*, 10, 52. doi: 10.1186/1471-244x-10-52.
- Phillips, A. (2013). Perceived stress. *Springer Reference*. Retrieved on http://www.springerreference.com/docs/html/chapterdbid/345488.html.
- Pigeon, W.R, Campbell, C.E., Possemato, K., and Ouimette, P. (2013). Longitudinal relationships of insomnia, nightmares, and PTSD severity in recent combat veterans. *Journal of Psychosomatic Research*, 75(6), 546-550.
- Pitman RK, Orr SP, Forgue DF, de Jong JB, Claiborn JM(1987). Psychophysiology of PTSD imagery in Vietnam combat veterans. Arch Gen Psychiatry, 44:970–975.

- Pittman, R.K., Orr, S.P., Forgue, D.F, Altman, B., de Jong, J.B., Herz, L.R. (1990)
 Psychophysiologic responses to combat imagery of Vietnam combat veterans with post-traumatic stress disorder vs. other anxiety disorders. *Journal of Abnormal Psychology*, 99, pp. 49–54.
- Radloff, L.S. (1977). The CED-D scale: A self-report depression scale for research in the general population. Applied Psychological Measurement, 1, 385-401.
- Resick PA, Schnicke MK. (1992). Cognitive processing therapy for sexual assault victims. Journal of Consulting and Clinical Psychology;60:748-56.
- Resick PA, Schnicke MK. (1993). Cognitive processing therapy for rape victims: A treatment manual. Newbury park: Sage.
- Resick PA, Williams LF, Suvak MK, Monson CM, Gradus JL. (2012). Long-term outcomes of cognitive-behavioral treatments for posttraumatic stress disorder among female rape survivors. Journal of Clinical and Consulting Psychology;80:201-10.
- Richards, A., Metzler, T.A., Ruoff., L.A., Inslicht, S.S., Rao, M., Talbot, L.S., & Neylan,
 T.C. (2013). Sex differences in objective measures of sleep in PTSD and healthy
 control subjects. *Journal of Sleep Research 2013*, December; 22(6): 679-687.
- Rhudy J.L., Davis J.L., Williams A.E., et al. (2010). Cognitive-behavioral treatment for chronic nightmares in trauma-exposed persons: assessing physiological reactions to nightmare-related fear. *Journal of Clinical Psychology*, 66(4), 365-82.
- Roszell, D.K., McFall, M.E. & Malas, K.L. (1991). Frequency of symptoms and concurrent psychiatric disorder in Vietnam veterans with chronic PTSD. Hosp Community Psychiatry; 42:293–6.

- Roy-Byrne P, Arguelles L, Vitek ME, Goldberg J, Keane TM, True WR, Pitman RK (2004). Persistence and change of PTSD symptomatology a longitudinal co- twin control analysis of the Vietnam Era Twin Registry. Social Psychiatry and Psychiatric Epidemiology 39, 681–685.
- Russel, L. (2011). Post-Traumatic Stress Disorder (PTSD). Available from http://www.mental-health-today.com/ptsd/dsm.htm Retrieved on October 18, 2013).
- Schoenbaum M, Kessler RC, Gilman SE, et al. (2014). Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). JAMA Psychiatry; 71(5): 493–503.
- Schnurr, P.P., Spiro, A. & Paris, A. (2000). Physician-diagnosed medical disorders in relation to PTSD symptoms in older male military veterans. Health Psychol 19: pp. 91–97.
- Schnurr PP, Friedman MJ, Engel CC, et al. (2007). Cognitive behavioral therapy for posttraumatic stress disorder in women: a randomized controlled trial. JAMA;297:820-30.
- Schottenbauer MA, Glass CR, Arnkoff DB, Tendick V, Gray SH (2008). Nonresponse and dropout rates in outcome studies on PTSD: review and methodological considerations. Psychiatry;71:134-69.
- Seal, K.H., et al. (2010). Bringing the war back home: mental health disorders among 103,788 US veterans returning from Iraq and Afghanistan seen at Department of Veterans Affairs facilities. *Archives of Internal Medicine*, 167:476-482.
- Seelig, A.D., Jacobson, I.G., Smith, B.*et al.* (2010). Sleep patterns before, during, and after deployment to Iraq and Afghanistan. *Sleep*, 33 (12), pp. 1615–1622.

- Semple WE, Goyer PF, McCormick R, Donovan B, Muzic RF, Jr.,, Rugle L, et al. Higher brain blood flow at amygdala and lower frontal cortex blood flow in PTSD patients with comorbid cocaine and alcohol abuse compared with normals. Psychiatry. 2000;63:65-74.
- Shalev, A., Attias, J., Bleich, A., Shulman, H., Kotler, M., & Shahar, A. (1988). Audiological evaluation of nonalcoholic, drug-free posttraumatic stress disorder patients. *Biol Psychiatry*, 24(5), 522-530.
- Shalev, A.Y, Orr, S.P, Peri, T., Scheiber, S., Pitman, R.K. (1992). Physiologic responses to loud tones in Israeli patients with post-traumatic stress disorder. *Archives of General Psychiatry*, 49, pp. 870–875.
- Shalev AY, Orr SP, Pitman RK (1993). Psychophysiologic assessment of traumatic imagery in Israeli civilian post-traumatic stress disorder patients. Am J Psychiatry, 150:620–624.
- Shalev, A. Y., Peri, T., Canetti, L., & Schreiber, S. (1996). Predictors of PTSD in injured trauma survivors: a prospective study. *Am J Psychiatry*, 153(2), 219-225.
- Shaley, A.Y, Peri, T., Orr, S.P., Bonne, O, Pitman, R.K. (1997). Auditory startle responses in help-seeking trauma survivors. Psychiatry Research, 69, pp. 1–7.
- Shapiro F. (2001). Eye movement desensitization and reprocessing: Basic principles, protocols and procedures (2nd ed.). New York: Guilford.
- Shin LM, Rauch SL, Pitman RK. Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. Annals of the New York Academy of Sciences. 2006;1071:67-79.

Smucker MR. Post-traumatic stress disorder. Northvale, NJ: Jason Aronson; 1997.

- Spoormaker V.I. and Montgomery P. (2008) Disturbed sleep in post-traumatic stress disorder: secondary symptom or core feature? *Sleep Med Rev* 12(3), 169-84.
- Steenkamp MM, Litz BT, Hoge CW, Marmar CR. Psychotherapy for military-related PTSD, A review of randomized clinical trials. *JAMA*. 2015;314:489-500.
- Stein MB, Walker JR, Hazen AL, Forde DR (1997). Full and partial posttraumatic stress disorder: findings from a community survey. Am J Psychiatry 154(8):1114–1119.
- Swinkles, C.M, Ulmer, C.S, Beckham, J.C., Buse, N. & Calhoun, P.S. (2013). The association of sleep duration, mental health, and health risk behaviors among U.S. Afghanistan/Iraq era veterans. *Sleep*, 36, (7):1019-1025.
- Talbot, L. S., Maguen, S., Metzler, T. J., Schmitz, M., McCaslin, S. E., Richards, A., ...
 Neylan, T. C. (2014). Cognitive behavioral therapy for insomnia in posttraumatic stress disorder: a randomized controlled trial. *Sleep*, *37*(2), 327-341. doi: 10.5665/sleep.3408.
- Tanielian T, Jaycox L. (2008). Invisible Wounds of War: Psychological and Cognitive Injuries, Their Consequences, and Services to Assist Recovery. Santa Monica, CA: RAND Corporation.
- Teplan, M. Fundamentals of EEG measurement. *Measurement Science Review* 2002; Volume 2, section 2; 1-11.
- U.S. Department of Veterans Affairs. 2010 VA/DoD Clinical Practice Guideline for Management of Post-Traumatic Stress.

http://www.healthquality.va.gov/guidelines/MH/ptsd/. Accessed May 11, 2015.

Van Liempt S. (2012). Sleep disturbances and PTSD: A perpetual circle? European Journal of Psychotraumatology;3 doi: 10.3402/ejpt.v3i0.19142.

- Wallace, D. M.,Wolhgemuth, W.K. (2011). Insomnia characteristics and clinical correlates in Operation Enduring Freedom/Operation Iraqi Freedom veterans with post-traumatic stress disorder and mild traumatic brain injury: An exploratory study. *Sleep Medicine*. 12(9), 850-859.
- Warner CH, Appenzeller GN, Grieger T, et al. (2011). Importance of anonymity to encourage honest reporting in mental health screening after combat deployment. Arch Gen Psychiatry; 68(10): 1065–71.
- Warner CH, Breitbach JE, Appenzeller GN, Yates V, Grieger T, Webster WG. (2007).Division mental health in the new brigade combat team structure: part II.Redeployment and postdeployment. Mil Med; 172(9): 912–7.
- Westmeyer, J., Sutherland, R.J., Freerks, M. *et al. (2007)*. Reliability of sleep log data versus actigraphy in veterans with sleep disturbance and PTSD. *J Anxiety Disord*, 21 (7), pp. 966–975.
- Wood, A.C, Kuntsi, J. & Asherson, P. (2008). Actigraph data are reliable, with functional reliability increasing with aggregation. *Behavior Research Methods*, 40, (3),873-878.
- Ulmer C.S., Edinger J.D. and Calhoun P.S. (2011) A multi-component cognitivebehavioral intervention for sleep disturbance in veterans with PTSD: a pilot study. *J Clin Sleep Med* 7(1), 57-68.
- U.S Department of Defense. (2001). Remembering September 11, 2001. Retrieved on http://www.defense.gov/home/features/2011/0911_911/.
- U.S. Department of Veterans Affairs. (2010). 2010 VA/DoD Clinical Practice Guideline for Management of Post-Traumatic Stress [Online]. Washington, DC: Available at: http://www.ptsd.va.gov/public/pages/prolonged-exposure-therapy.as.

- Zayfert C, DeViva J. C. (2004). Residual insomnia following cognitive behavioral therapy for PTSD. Journal of Traumatic Stress; 17:69–73.
- Zimmerman, M. & Mattia, J. (1999). The reliability and validity of a screening
 Questionnaire for 13 DSM-IV Axis I disorders (the Psychiatric Diagnostic
 Screening Questionnaire) in psychiatric outpatients. *Journal of Clinical Psychiatry*, 60,(10), 677-683.
- Zinzow HM, Britt TW, Pury CL, Raymond MA, McFadden AC, Burnette CM. (2013). Barriers and facilitators of mental health treatment seeking among active-duty army personnel. Mil Psychol; 25(5): 514–35.
- Zoladz PR, Diamond DM. Current status on behavioral and biological markers of PTSD: A search for clarity in a conflicting literature. Neurosci Biobehav Rev.2013;37:860-95.

Appendices

Appendix A: ART Intake Questionnaire

1) About how many traumatic memories are still impacting your life?□ None□ 1 to 2□ 3 to 4□ 5 or more

Which traumatic memory/memories would you like to resolve with A.R.T.?

2)	Are you currently on disability for post-traumatic stress disorder (PTSD) or any mental health problem?
3)	Do you feel that you are ready to let go of your traumatic memories so that you no longer could see them in the same way (i.e. if there was a way to do this)? □ YES □ NO
	If No, what would prevent you from wanting to let go?
4)	Is there any guilt associated with your traumatic memory or memories? □ YES □ NO If Yes, can you briefly explain what the guilt is about?
5)	How long have you lived with the traumatic memory or memories? □ 0-3 months □ 4 to 6 months □ 7 to 12 months □ 1 to 3 years □ 4 to 6 years □ 7 to 10 years□ 11+ years
6)	I am motivated to move forward in my life:
	Least motivated = 0 Most motivated = 10 (Please circle your answer)
	0 1 2 3 4 5 6 7 8 9 10
7)	Have you been diagnosed with any mental health conditions in the past?

 \Box YES \Box NO

If Yes, what is the condition or conditions?

8) What other problems are impacting your life now beside the traumatic memories and other symptoms related to PTSD?

9)If you were able to rid yourself of the symptoms that occur from your traumatic memories, what percentage of your mental health problems would be resolved (circle one number)?

<u>0% 10% 20% 30% 40% 50% 60% 70% 80% 90% 100%</u>

Name of therapist reviewing form: _______ Signature of therapist: ______ Today's Date: ______

Page 2 of 2

Appendix B: Medical History Form

Π) #		Visit:
D	ate:		
	Medical	History Form	
		Page 1 of 2	
1)	Current and Past Medical Histo	ry (check all that apply)	
	Diabetes	Cancer	Vision
	Problems		
	Lung disease	Thyroid Problem	
	Intestinal Disorder	y	
	Kidney Disease	Liver Disease	Muscle/bone
	Heart Disease		
	Neurological problems	Seizures	
	High Blood Pressure		GYN
	problems, if female	Hearing Problems	
	Spinal cord Injury	-	Severe
	burns	Head trauma	
	Paralysis (if yes, check all t	hat apply)	
	Upper extremity(ies)		
	Lower extremity(ies)		
	Amputation (if yes, check b	below all that apply)	
		Upper limb(s)	
		Upper extremity(ies)	
2)	Have you previously received a	any treatment for Post-Traumatic	Stress Disorder or
,	any other type of mental health	-	_No

If yes, please specify:

If applicable, what type of treatment(s) did you receive (check all that applies):

Individual therapy with a mental health professional Type of therapy (check all the apply) ____Psychotherapy: ____Cognitive Behavior (CBT)

		Months of Treatment:
	Group Psychotherapy	Months of Treatment:
	Individual Psychotherapy	Months of Treatment:
	Prolonged Exposure	Months of Treatment:
	Eye Movement desensitiza	ation and reprocessing (EMDR) Months of Treatment:
Prescri	ption Medication	Months of Treatment:
Other (specify)	Please	
3.) Please list the na	mes of any prescription medic	cations you are currently taking:

```
Medication name
```

Dose (if known)

Frequency (e.g. daily)

Appendix C: State – Trait Inventory for Cognitive and Somatic Anxiety

13. State-Trait Inventory for Cognitive and Somatic Anxiety (STICSA)

Your mood at this moment

Below is a list of statements which can be used to describe how people feel. Beside each statement are four numbers which indicate the degree with which each statement is self-descriptive of your mood at this moment (eg, 1 = not at all, 4 = very much so). Please read each statement carefully and circle the number which best indicates how you feel right now, at this very moment, even if this is not how you usually feel.	Not at all	A little	Moderately	Very much so
In general				
1. My heart beats fast	1	2	3	4
2. My muscles are tense	1	2	3	4
3. I feel agonized over my problems	1	2	3	4
4. I think that others won't approve of me	1	2	3	4
I feel like I'm missing out on things because I can't make up my mind soon enough	1	2	3	4
6. I feel dizzy	1	2	3	4
7. My muscles feel weak	1	2	3	4
8. I feel trembly and shaky	1	2	3	4
9. I picture some future misfortune	1	2	3	4
10. I can't get some thought out of my mind	1	2	3	4
11. I have trouble remembering things	1	2	3	4
12. My face feels hot	1	2	3	4
13. I think that the worst will happen	1	2	3	4
14. My arms and legs feel stiff	1	2	3	4
15. My throat feels dry	1	2	3	4
16. I keep busy to avoid uncomfortable thoughts	1	2	3	4
17. I cannot concentrate without irrelevant thoughts intruding	1	2	3	4
18. My breathing is fast and shallow	1	2	3	4
19. I worry that I cannot control my thoughts as well as I would like to	1	2	3	4
20. I have butterflies in the stomach	1	2	3	4
21. My palms feel clammy	1	2	3	4

Appendix D: PTSD Checklist – Military Version (PCL – M)

PTSD Checklist – Military Version (PCL-M)

Name:	Unit:
Best contact number and/or email:	
Deployed location:	

Instructions: Below is a list of problems and complaints that veterans sometimes have in response to a stressful military experience. Please read each one carefully, put an "X" in the box.

	[⁻ [1				
		Not at all	A little bit	Moderately	Quite a bit	Extremely
1.	Repeated, disturbing memories, thoughts, or images of a stressful military experience?					
2.	Repeated, disturbing dreams of a stressful military experience?					
3.	Suddenly acting or fealing as it a stressful military experience ware happening again (as if you were reliving it)?					
4.	Feeling very upset when something reminded you of a stressful military experience?					
5.	Having physical reactions (e.g., heart pounding, trouble breathing, or sweating) when something reminded you of a stressful military experience?					
6.	Avoid thinking about or talking about a stressful military experience or avoid having feelings related to it?					
7.	Avoid activities or talking about a stressful military experience or avoid having feelings related to it?					
8.	Trouble remembering important parts of a stressful military experience?					
9.	Loss of interest in things that you used to enjoy?					
10.	Feeling distant or cut of from other people?					
11.	Feeling amotionally numb or being unable to have loving feelings for those close to you?					
12.	Feeling as if your future will somehow be aut short?					
13.	Trouble failing or staying asleep?					
14.	Feeling initable or having angry outbursts?					
15.	Having difficulty concentrating?					
16.	Being "super alerf" or watchful on guard?					
17.	Feeling jumpyor easily startled?					

Has anyone indicated that you've changed since the stressful military experience? Yes __ No__

Appendix E: Demographics

ID #	
Date:	_

Visit:

D) Native Hawaiian/Pacific Islander

E) American Indian/Alaskan

DEMOGRAPHICS (Page 1 of 2)

Please complete the following information. This information is only for research staff reference and will remain strictly confidential.

1.) What is your age?_____

2.) Please indicate your gender: A) Male____ B)Female____

- 3.) Please indicate your race:
 - A) Black or African America
 - B) White Native
 - C) Asian

4.) Please indicate your ethnicity:A) Non-HispanicB) Hispanic

- 5.) What is your <u>current</u> marital status? (Circleone)
 - A) Never married D) Divorced or separated
 - B) Married E) Widowed
 - C) Living with someone but not married

6.) What is your current employment status? (CircleOne)

A) Full-time employment	E) Unemployed, disabled
B) Part-Time employment	F) Retired due to pain
C) Unemployed, not interested	G) Retired not due to pain
D) Unemployed, looking for work	H) Other (specify):

- 7.) How many years of education have you completed starting with the first grade? Years
- 8.) Are you currently enrolled in an educational program?
 - A) Yes B) no C) I am planning to be

ID#	
Date:	

Visit:

DEMOGRAPHICS (Page 2 of 2)

9.) If you answered Yes to #8, what type of educational program are you enrolled in?				
A) High School (GO	CSE, etc) B) Vocational Sector	chool		
C) College/Associat	te Level degree D) College/bachelo	ors Level degree		
E) College/Graduate	e Level Degree F) Other			
10.) What is your curre	ent military status? (Circle one)			
A) Active Duty	B) Reservist C) Discharged/Vet	eran		
11.) What branch of the	military have you primarily served in	n? (Circle one)		
A) Army	E) Marine Corps	I) Coast Guard		
B) Army Reserves	F) Marine Corps Reserves	J) Coast Guard		
Reserves				
(1) N	G h E	\mathbf{V} \mathbf{N} \mathbf{C} 1 \mathbf{C} 1		
C) Navy	G) Air Force	K) National Guard		
, .	G) Air Force H) Air Force Reserves	L) National Guard		
, .		/		

A) 1 B) 2 C) 3	D) 4 or more
13.) Where were you deploy	ved? (Circle all that apply.)
A) Persian Gulf	D) Vietnam
B) Iraq	E) Kosovo
C) Afghanistan	F) Other

13a) Length of deployment(s) in months: Shortest___Longest____

14.) During your tour(s) of duty, did you experience any combat-related activities? (For example, patrols, under enemy fire, being surrounded by the enemy, kidnapped or imprisoned, firing upon the enemy, hand to hand combat, pinned down, shot down)

A) Yes B) No

- 15.) What are the principals type(s) of distressing experiences for which you are seeking treatment? (Circle all the apply)
 - A) Witness death or execution
 - C) Witness major injuries (non-lethal)
 - E) Sexual assault

- B) IED blast or combat explosion
- D) Physical assault
- F) Other Specify:

Thank you for your participation.

Appendix F: Pittsburgh Sleep Quality Index (PSQI)

Subjec	t's Initials	ID#	Da	te	Time	AM _PM		
	PITTSBURGH SLEEP QUALITY INDEX							
The f shoul		relate to your usual si accurate reply for the ons.				wers		
1.	During the past m	onth, what time have	you usually gone t	o bed at night?				
		BED TIM	E					
2.	During the past me	onth, how long (in min	utes) has it usually	taken you to fall a:	sleep each n	ight?		
		NUMBER OF M	INUTES					
з.	During the past m	onth, what time have	you usually gotten	up in the morning?	?			
		GETTING UP	TIME					
4.	During the past m different than the r	onth, how many hou number of hours you	rs of <u>actual sleep</u> spent in bed.)	did you get at nigh	nt? (This ma	ay be		
		HOURS OF SLEEP	PER NIGHT					
Forea	ch of the remainin	g questions, check t	he one best respo	nse. Please answ	ver <u>all</u> questi	ions.		
5.	During the past m	onth, how often have	you had trouble sl	eeping because yo	ω			
a)	Cannot get to slee	p within 30 minutes						
	Not during the past month	Less than once a week	Once or twice a week					
b)	Wake up in the m	iddle of the night or e	arly morning					
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week				
c)	Have to get up to	use the bathroom						
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week				

d) Cannot breathe comfortably

	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week	
e)	Cough or shore los	udly			
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week	
f)	Feel too cold				
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week	
g)	Feel too hot				
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week	
h)	Had bad dreams				
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week	
i)	Have pain				
	Not during the past month		Once or twice a week	Three or more times a week	
j)	Other reason(s), p	lease describe			
	How often during the past month have you had trouble sleeping because of this?				
	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week	

6. During the past month, how would you rate your sleep quality overall?

Very good	
Fairly good	
Fairly bad	
Very bad	

During the past month, how often have you taken medicine to help you sleep (prescribed or "over the counter")?

Not during the	Less than	Once or twice	Three or more
past month	once a week	a week	times a week

 During the past month, how often have you had trouble staying awake while driving, eating meals, or engaging in social activity?

Not during the	Less than	Once or twice	Three or more
past month	once a week	a week	times a week

During the past month, how much of a problem has it been for you to keep up enough enthusiasm to get things done?

No problem at all	
Only a very slight problem	
Somewhat of a problem	
A very big problem	

10. Do you have a bed partner or room mate?

No bed partner or room mate	
Partner/room mate in other room	
Partner in same room, but not same bed	
Partner in same bed	

If you have a room mate or bed partner, ask him/her how often in the past month you have had . . .

a) Loud snoring

Not during the	Less than	Once or twice	Three or more
past month	once a week	a week	times a week

b) Long pauses between breaths while asleep

Not during the	Less than	Once or twice	Three or more
past month	once a week	a week	times a week

c) Legs twitching or jerking while you sleep

Not during the	Less than	Once or twice	Three or more
past month	once a week	a week	times a week

Page 4 of 4

d) Episodes of disorientation or confusion during sleep

	Not during the past month	Less than once a week	Once or twice a week	Three or more times a week
e)	Other restlessness	while you sleep; plea	se describe	
	Not during the	Less than	Once or twice	Three or more
	past month	once a week	a week	times a week

Appendix G: Center for Epidemiologic Studies Depression Scale (CES-D)

Center for Epidemiologic Studies Depression Scale (CES-D), NIMH

Below is a list of the ways you might have felt or behaved. Please tell me how often you have felt this way during the past week.

	During the Past Week			
	Rarely or none of the time (less than 1 day)	Some or a little of the time (1-2 days)	Occasionally or a moderate amount of time (3-4 days)	Most or all of the time (5-7 days)
1. I was bothered by things that usually don't bother me.				
 I did not feel like eating; my appetite was poor. 				
 I felt that I could not shake off the blues even with help from my family or friends. 				
 I felt I was just as good as other people. 				
 I had trouble keeping my mind on what I was doing. 				
 I felt depressed. I felt that everything I did was an effort. 				
 I felt hopeful about the future. I thought my life had been a failure. I thought my life had been a failure. I felt fearful. My sleep was restless. I was happy. I talked less than usual. I felt lonely. People were unfriendly. I enjoyed life. I had crying spells. I felt sad. I felt that people dislike me. I could not get "going." 				

SCORING: zero for answers in the first column, 1 for answers in the second column, 2 for answers in the third column, 3 for answers in the fourth column. The scoring of positive items is reversed. Possible range of scores is zero to 60, with the higher scores indicating the presence of more symptomatology.

Appendix H: Institutional Review Board Approval



RESEARCH INTEGRITY AND COMPLIANCE Institutional Review Boards, FWA No. 00001669 12901 Bruce B. Downs Blvd., MDC035 • Tampa, FL 336124799 (813) 974-5638 • FAX(813) 974-7091

4/23/2013

Marian Hardwick, College of Nursing 6147 5th Avenue North Saint Petersburg, FL 33710

RE: Full Board Approval for Initial Review

IRB#: Pro00011530

Title: The Relationship Between Post-Traumatic Stress Disorder and Sleep Disturbance Before and After Treatment with Accelerated Resolution Therapy (ART)

Study Approval Period: 4/8/2013 to 4/8/2014

Dear Dr. Hardwick:

On 4/8/2013, the Institutional Review Board (IRB) reviewed and APPROVED the above application and all documents outlined below.

Approved Item(s):

Protocol Document(s): <u>PTSD</u> and Sleep Dissertation Protocol.docx

Accepted Documents:

Centers for Epidemiological Studies Depression Scale <u>Pittsburg Sleep Quality Index</u> <u>Psychiatric Diagnostic Screening Questionnaire</u> <u>PTSD - C</u> <u>PTSD Checklist - M</u> State- Trait Inventory for Cognitive and Somatic Anxiety

Consent/Assent Document(s)*:

PTSD and Sleep Consent Form.pdf

*Please use only the official IRB stamped informed consent/assent document(s) found under the "Attachments" tab. Please note, these consent/assent document(s) are only valid during the approval period indicated at the top of the form(s). As the principal investigator of this study, it is your responsibility to conduct this study in accordance with IRB policies and procedures and as approved by the IRB. Any changes to the approved research must be submitted to the IRB for review and approval by an amendment.

We appreciate your dedication to the ethical conduct of human subject research at the University of South Florida and your continued commitment to human research protections. If you have any questions regarding this matter, please call 813-974-5638.

Sincerely,

Vjørgensen Mo

E. Verena Jorgensen, M.D., Chairperson USF Institutional Review Board



RESEARCH INTEGRITY AND COMPLIANCE Institutional Review Boards, FWA No. 00001669 12901 Bruce B. Downs Blvd., MDC035 • Tampa, FL 33612-4799 (813) 974-5638 • FAX(813)974-7091

3/11/2016

Marian Hardwick, MS, RN College of Nursing 12901 Bruce B. Downs Blvd., MDC22 Tampa, FL 33612

RE: Expedited Approval for Continuing Review

- IRB#: CR3_Pro00011530
- Title: The Relationship Between Post-Traumatic Stress Disorder and Sleep Disturbance Before and After Treatment with Accelerated Resolution Therapy (ART)

Study Approval Period: 4/8/2016 to 4/8/2017

Dear Mrs. Hardwick:

On 3/10/2016, the Institutional Review Board (IRB) reviewed and APPROVED the above application and all documents contained within including those outlined below.

Approved Item(s): Protocol Document(s): Study Protocol Clean 12-16-2014

Consent/Assent Document(s)*: Consent form (clean) version 3.0 2-24-2015.pdf

*Please use only the official IRB stamped informed consent/assent document(s) found under the "Attachments" tab on the main study's workspace. Please note, these consent/assent document(s) are only valid during the approval period indicated at the top of the form(s) and replace the previously approved versions.

The IRB determined that your study qualified for expedited review based on federal expedited category number(s):

(8) Continuing review of research previously approved by the convened IRB as follows: (a) where (i) the research is permanently closed to the enrollment of new subjects; (ii) all subjects have completed all research-related interventions; and (iii) the research remains active only for

long-term follow-up of subjects; or (b) where no subjects have been enrolled and no additional risks have been identified; or (c) where the remaining research activities are limited to data analysis.

As the principal investigator of this study, it is your responsibility to conduct this study in accordance with USF HRPP policies and procedures and as approved by the USF IRB. Any changes to the approved research must be submitted to the IRB for review and approval by an amendment. Additionally, all unanticipated problems must be reported to the USF IRB within five (5) calendar days.

We appreciate your dedication to the ethical conduct of human subject research at the University of South Florida and your continued commitment to human research protections. If you have any questions regarding this matter, please call 813-974-5638.

Sincerely,

Vjørgensen MD

E. Verena Jorgensen, M.D., Chairperson USF Institutional Review Board



RESEARCH INTEGRITY AND COMPLIANCE Institutional Review Boards, FWA No. 00001669 12901 Bruce B. Downs Bbrd., MDC035 • Tampa, FL 33612-4799 (813) 974-5638 • FAX(813)974-7091

3/14/2013

Kevin Kip, Ph.D. College of Nursing 12901 Bruce B. Downs., Blvd MDC 22 Tampa, FL 33612-4766

RE: Full Board Approval for Initial Review

IRB#: Pro00011641

Title: Accelerated Resolution Therapy (ART) for Rapid Resolution of Symptoms of Psychological Trauma

Study Approval Period: 3/14/2013 to 3/14/2014

Dear Dr. Kip:

On 3/14/2013, the Institutional Review Board (IRB) reviewed and APPROVED the above application and all documents outlined below.

Approved Item(s):

Protocol Document(s):

ART SAMHSA Protocol with Appendices 1-25-2011

As the principal investigator of this study, it is your responsibility to conduct this study in accordance with IRB policies and procedures and as approved by the IRB. Any changes to the approved research must be submitted to the IRB for review and approval by an amendment.

We appreciate your dedication to the ethical conduct of human subject research at the University of South Florida and your continued commitment to human research protections. If you have any questions regarding this matter, please call 813-974-5638.

Sincerely,

inka Ph.D.

John Schinka, Ph.D., Chairperson USF Institutional Review Board

View: Study - Pro00011	1641	
------------------------	------	--



1.15	уре			
		Reviewer	Date Created	Date Modified
1 fc IF U tc re P	RB Staff Change Request .1.2 Please revise to include the blowing additional statement: "Note to RB: This study was previously approved inder IRB# Pro00000894 and expired due to a lapse in filing a continuing review. The esearch project is being resubmitted for urpose See More	Anna Davis		
_	Change Request Completed - Trudy			

Study Identification Information

You must complete all of the required questions on this page to create your Human Research Application. As you continue through the application, you will automatically be guided to the appropriate pages needed to complete your submission.

1.1.1 * Study Title (this title must be the same as the title on your protocol, Investigators Brochure and most cases, informed consent document): Accelerated Resolution Therapy (ART) for Rapid Resolution of Symptoms of Psychological Trauma

> * Short Title (this title is used throughout the site to identify the study): ART SAMHSA

1.1.2 * Brief Study Description:

Note to IRB: This study was previously approved under IRB# Pro00000894 and expired due to a lapse in filing a continuing review. The research project is being resubmitted for purposes of data analysis only

This is a 1-year, prospective exploratory pilot study whereby consenting individuals with symptoms of psychological trauma (N=100) will be provided with Accelerated Resolution Therapy (ART) by trained ART clinicians. Study participants will complete questionnaires of symptoms of psychological trauma, psychological wellness, trauma-related guilt, depressive symptoms, anxiety

1.1

symptoms, anger, sleep quality, positive challenge/growth, self-compassion, and alcohol use. These will be obtained before treatment with ART, immediately following treatment with ART, and at 2- and 4-months follow-up. Within-subject changes in these measures will be compared by use of graphical techniques, paired t tests, and mixed models. Because ART is designed for rapid resolution of psychological distress, participating subjects are anticipated to undergo a minimum of two and a maximum of five ART sessions. The primary study endpoint will be the magnitude of change in symptoms of psychological trauma, as measured by repeated assessments using the PTSD Checklist (PCL-Civilian). Secondary endpoints will include self-report measures of psychological wellness, trauma-related guilt, depressive symptoms, anxiety symptoms, anger, sleep quality, positive change/growth, self-compassion, and alcohol use.

1.1.3 * Is this research being conducted to fulfill an educational requirement (such as a dissertation or thesis)?

Yes <i>No

1.1.4 * Does the University of South Florida and/or any of its senior officials have a potential conflict of interest related to this research (e.g. an ownership interest in an entity related to the research; a patent, trademark, copyright or licensing agreement in the test article or method being studied)?

🗆 Yes 💌 No

If you answered Yes to this question, before proceeding with the completion of this application please contact the USF Conflict of Interest Program at 813-974-5638 or coi-research@usf.edu for assistance in determining whether there is an institutional conflict of interest. Please note that the USF COI Committee may require human subjects research with a related institutional conflict of Interest to be reviewed by an external IRB.

1.1.5 * Principal Investigator / Student Investigator: Kevin Kip

> You are listed automatically as a Study Coordinator. If there is someone else on the study that will assist with the IRB regulatory processes, they should be listed as a Study Coordinator and/or Secondary Study Coordinator.

- 1.1.6 Study Coordinator / Primary Regulatory Specialist: Sue Girling
- 1.1.6a Secondary Study Coordinator / Regulatory Specialist: Independent of the second study wittenberg

The PI does not need to be listed as a Co-Investigator or Key Personnel.

If yes, please add Co-Investigators:

First Name	Last Name	Organization	Profile
Kevin	Kip	College of Nursing	00001020
Helene (Laney)	Rosenzweig	College of Nursing	00003016
Amy	Shuman	College of Nursing	00003771

1.1.8 * Are there any Key Personnel on this study?

• Yes 🔍 No

If yes, please add Key Personnel/Study Staff and assign roles for their participation on the study:

Name	Organization	Roles on Study	Other Role On Study
Sue Girling	College of Nursing	Obtains Informed Consent	
Marian Hardwick	College of Nursing	Manuscript Preparation	

1.1.9 Is this study a resubmission of a study previously reviewed and/or approved by the USF IRB or another IRB?

🖲 Yes 🔍 No

If yes, please provide the Title and USF IRB/Pro Number if previously submitted to the USF IRB. If submitted to an external IRB, include the name of the IRB, date of submission and outcome of the review. Accelerated Resolution Therapy (ART) for Rapid Resolution of Symptoms of Psychological Trauma (Pro00000894)



RESEARCH INTEGRITY AND COMPLIANCE

Institutional Review Boards, FWA No. 00001669 12901 Bruce B. Downs Blvd., MDC035 • Tampa, FL 33612-4799 (813) 974-5638 • FAX(813)974-7091

2/16/2016

Kevin Kip, Ph.D. College of Nursing 12901 Bruce B. Downs Blvd., MDC 22 Tampa, FL 33612-4766

RE: Expedited Approval for Continuing Review

IRB#: CR3 Pro00011641

Title: Accelerated Resolution Therapy (ART) for Rapid Resolution of Symptoms of Psychological Trauma

Study Approval Period: 3/14/2016 to 3/14/2017

Dear Dr. Kip:

On 2/16/2016, the Institutional Review Board (IRB) reviewed and APPROVED the above application and all documents contained within including those outlined below.

Approved Item(s): Protocol Document(s): ART SAMHSA Protocol with Appendices 1-25-2011

The IRB determined that your study qualified for expedited review based on federal expedited category number(s):

(8) Continuing review of research previously approved by the convened IRB as follows: (a) where (i) the research is permanently closed to the enrollment of new subjects; (ii) all subjects have completed all research-related interventions; and (iii) the research remains active only for long-term follow-up of subjects; or (b) where no subjects have been enrolled and no additional risks have been identified; or (c) where the remaining research activities are limited to data analysis.

As the principal investigator of this study, it is your responsibility to conduct this study in accordance with USF HRPP policies and procedures and as approved by the USF IRB. Any changes to the approved research must be submitted to the IRB for review and approval by an amendment. Additionally, all unanticipated problems must be reported to the USF IRB within five (5) calendar days.

We appreciate your dedication to the ethical conduct of human subject research at the University of South Florida and your continued commitment to human research protections. If you have any questions regarding this matter, please call 813-974-5638.

Sincerely,

PAn

Kristen Salomon, Ph.D., Vice Chairperson USF Institutional Review Board



DIVISION OF RESEARCH INTEGRITY AND COMPLIANCE Institutional Review Boards, FWA No. 00001669 12901 Bruze B. Downs Blvd. MDC035 • Tamps, FL 336124799 (\$13) 974-5638 • FAX (\$13) 974-5618

October 29, 2010

Kevin Kip, PhD College of Nursing MDC 22

RE: Full Board Approval for Initial Review

IRB#: Pro00000210 Title: Accelerated Resolution Therapy (ART) for Psychological Trauma Study Approval Period: 10/15/2010 to 10/15/2011

Dear Dr. Kip,

On 10/15/2010 the Institutional Review Board (IRB) reviewed and APPROVED the above application and all documents outlined below. Please note that your approval for this study will expire on 10/15/2011.

Approved Items: Protocol Document(s):

ART Protocol 10-27-2010	10/28/201	0 4:07 PM	0.	01
Consent/Assent Document(s)				
Appendix B2. ART Informed Con	nsent with	10/29/2010	10:34	0.01
HIPAA 10-27-2010.pdf.pdf		AM		0.01

Please note, if applicable, the **informed consent/assent documents are valid during the period indicated by the official, IRB-Approval stamp located on the form.** Valid consent must be documented on a copy of the most recently IRB-approved consent form. As the principal investigator of this study, it is your responsibility to conduct this study in accordance with IRB policies and procedures and as approved by the IRB. Any changes to the approved research must be submitted to the IRB for review and approval by an amendment. We appreciate your dedication to the ethical conduct of human subject research at the University of South Florida and your continued commitment to human research protections. If you have any questions regarding this matter, please call 813-974-9343.

Sincerely,

Anite Kalal

Krista Kutash, PhD, Chairperson USF Institutional Review Board

Cc: Anna Davis, USF IRB Professional Staff

If you answered Yes to this question, before proceeding with the completion of this application please contact the USF Conflict of Interest Program at 813-974-5638 or coi-research@usf.edu for assistance in determining whether there is an institutional conflict of interest. Please note that the USF COI Committee may require human subjects research with a related institutional conflict of interest to be reviewed by an external IRB.

1.1.5 * Principal Investigator / Student Investigator: Kevin Kip

> You are listed automatically as a Study Coordinator. If there is someone else on the study that will assist with the IRB regulatory processes, they should be listed as a Study Coordinator and/or Secondary Study Coordinator.

- 1.1.6 Study Coordinator / Primary Regulatory Specialist: Sue Girling
- 1.1.6a Secondary Study Coordinator / Regulatory Specialist: @

The PI does not need to be listed as a Co-Investigator or Key Personnel.

1.1.7 * Are there any Co-Investigators/Faculty Advisors involved in this study? If you are a student, you <u>must</u> list your Faculty Advisor as a Co-Investigator.

 ● Yes ○ No

If yes, please add Co-Investigators:

First Name	Last Name	Organization	Profile
Diego	Hernandez	College of Nursing	00007318
Helene (Laney)	Rosenzweig	College of Nursing	00003016

> If yes, please add Key Personnel/Study Staff and assign roles for their participation on the study:

Name	Organization	Roles on Study	Other Role On Study
Marian Hardwick	College of Nursing	Manuscript Preparation	
Hongdao Meng	School of Aging Studies	Data Analysis	
Mariangeli Miranda	College of Nursing		Data Entry Specialist
Trudy Wittenberg	College of Nursing	Addresses Regulatory Issues	

1.1.9 Is this study a resubmission of a study previously reviewed and/or approved by the USF IRB or another IRB?

Yes No

If yes, please provide the Title and USF IRB/Pro Number if previously submitted to the USF IRB. If submitted to an external IRB, include the name of the IRB, date of submission and outcome of the review.



RESEARCH INTEGRITY AND COMPLIANCE Institutional Review Boards, FWA No. 00001669 12901 Bruse B. Downs Blod., MDC035 • Tampa, FL 33612-4799 (\$13) 974-5638 • FAX(\$13)974-7091

9/21/2015

Kevin Kip, Ph.D. College of Nursing 12901 Bruce B. Downs Blvd., MDC 22 Tampa, FL 33612-4766

RE: Expedited Approval for Continuing Review

IRB#: CR5_Pro00000210

Title: Accelerated Resolution Therapy (ART) for Psychological Trauma

Study Approval Period: 10/15/2015 to 10/15/2016

Dear Dr. Kip:

On 9/17/2015, the Institutional Review Board (IRB) reviewed and APPROVED the above application and all documents contained within including those outlined below.

Approved Item(s): Protocol Document(s): <u>ART</u>

The IRB determined that your study qualified for expedited review based on federal expedited category number(s):

(8) Continuing review of research previously approved by the convened IRB as follows: (a) where (i) the research is permanently closed to the enrollment of new subjects; (ii) all subjects have completed all research-related interventions; and (iii) the research remains active only for long-term follow-up of subjects; or (b) where no subjects have been enrolled and no additional risks have been identified; or (c) where the remaining research activities are limited to data analysis.

As the principal investigator of this study, it is your responsibility to conduct this study in accordance with USF HRPP policies and procedures and as approved by the USF IRB. Any changes to the approved research must be submitted to the IRB for review and approval by an amendment. Additionally, all unanticipated problems must be reported to the USF IRB within five (5) calendar days.

We appreciate your dedication to the ethical conduct of human subject research at the University of South Florida and your continued commitment to human research protections. If you have any questions regarding this matter, please call 813-974-5638.

Sincerely,

Le l'Ara-4

Kristen Salomon, Ph.D., Vice Chairperson USF Institutional Review Board

About the Author

Marian J. Hardwick grew up in Saint Petersburg, Florida. She obtained her first B.S at Florida State University in Health Education and taught Health and Anatomy I Honors at Seminole High for one year. She went back to school where she received her second B.S and attended an accelerated nursing program known as the VA Nursing Academy at the University of South Florida in Tampa, Florida in 2008 to 2009. She worked as a charge nurse in the psychiatric unit at James A. Haley VA Hospital for two years before returning to receive her MSN in Nursing Education and becoming a part- time educator in 2010. She entered the BS-Ph.D. program in 2010.

Marian became a McKnight Scholar that paid her full tuition for four years as well as a Jonas scholar her last two years of school. She is also a member of the Doctoral Nursing Student Organization at the University of South Florida. She is particularly interested in the impact that accelerated resolution therapy (ART) has on veterans with PTSD and civilians who have sleep disturbance.